

**Biosystematic investigations in the family
of duckweeds (*Lemnaceae*) (vol. 4)**

The family of *Lemnaceae* – a monographic study

Volume 2

(Phytochemistry; physiology; application; bibliography)

by Elias LANDOLT and Rikael KANDELER

1987

Best Available Copy

CONTENTS

1. Phytochemical characteristics	9
1.1. General chemical composition and inorganic components	9
1.1.1. Water content	9
1.1.2. Variation of mineral content	12
1.1.3. Status of some elements within the Lemnaceae	12
1.1.4. The mineral content of Lemnaceae in comparison with other water plants	12
1.1.5. Variation of mineral content due to different measuring techniques and different environmental conditions	13
1.1.6. Variation of the mineral content during different seasons and during lifetime	16
1.1.7. Variation of mineral content in different species and clones	16
1.2. Organic components	20
1.2.1. General remarks	20
1.2.2. Proteins and amino acids	21
1.2.2.1. Proteins	21
1.2.2.2. Amino acids	22
1.2.3. Carbohydrates	25
1.2.4. Lipids and fatty acids	28
1.2.5. Organic phosphorus compounds and nucleic acids	30
1.2.6. Chlorophylls, carotenoids, phytochrome and cytochrome	31
1.2.7. Flavonoids	35
1.2.8. Lignins	41
1.2.9. Oxalates and other organic anions	42
1.2.10. Vitamins and plant growth substances	44
1.2.11. Enzymes	47
1.2.12. Other organic substances	56
2. Physiological characteristics	58
2.1. General	58
2.2. Cultivation	59
2.2.1. Aseptic culture technique	59
2.2.2. Nutrient solutions	61
2.2.3. Cultivation and preservation techniques	65
2.2.4. Tissue cultures	66
2.3. Vegetative growth	67
2.3.1. Measurements	67
2.3.1.1. Growth rate	67
2.3.1.2. Dry weight of fronds	69
2.3.1.3. Frond size and other frond characteristics	69
2.3.2. Effects of gaseous substances	71
2.3.2.1. General remarks	71
2.3.2.2. Carbon dioxide	71
2.3.2.2.1. Influence of the CO ₂ concentration	71
2.3.2.2.2. Intensity of CO ₂ uptake	73
2.3.2.2.3. Mode of CO ₂ uptake	74
2.3.2.2.4. CO ₂ uptake from the water	74
2.3.2.3. Sulfur dioxide (H ₂ S)	76
2.3.2.4. Hydrogen sulfide (O ₂)	77
2.3.2.5. Ozone and ozonated hexene	77
2.3.2.6. Further gases	78
2.3.3. Effects of dissolved chemical compounds	79

Publications on "Biosystematic investigations in the family of duckweeds (Lemnaceae)" published in the "Veröffentlichungen des Geobotanischen Institutes ETH, Stiftung Rüböl, Zürich"

- vol. 1 (No. 70). LANDOLT E. (ed.), 1980: Key to determination; cytological variation; amino acid composition and sugar content; bibliography; list of studied material.
- vol. 2 (No. 71). LANDOLT E., 1986: The family of Lemnaceae - a monographic study. Vol. 1 of the monograph: Morphology; karyology; ecology; geographic distribution; systematic position; nomenclature; descriptions.
- vol. 3 (No. 80). LUEPOLD A., 1983: Das Wachstum von Wasserlinsen (Lemnaceae) in Abhängigkeit des Nährstoffangebotes, insbesondere Phosphor und Stickstoff. (Growth of duckweeds (Lemnaceae) depending on nutrient supply, especially phosphorus and nitrogen).
- vol. 4 (No. 95). LANDOLT E. and KANDELER R., 1987: The family of Lemnaceae, a monographic study. Vol. 2 of the monograph: Phytochemistry; physiology; application; bibliography.

2.3.3.1. Absorption of dissolved compounds from the water	79
2.3.3.2. Concentration of the nutrient solution	79
2.3.3.3. pH of the nutrient solution	81
2.3.3.3.1. Optimal pH for growth	81
2.3.3.3.2. Lower and higher pH limit for growth	82
2.3.3.3.3. Special pH effects	83
2.3.3.4. Inorganic substances	86
2.3.3.4.1. General remarks	86
2.3.3.4.2. Monovalent alkali metals (K, Na, Rb, Cs, Li)	87
2.3.3.4.2.1. Potassium (K)	90
2.3.3.4.2.2. Rubidium (Rb), Lithium (Li), and Caesium (Cs)	90
2.3.3.4.2.3. Sodium (Na)	92
2.3.3.4.3. Bivalent alkali metals (Ca, Mg, Sr, Ba)	99
2.3.3.4.4. Nitrogen (N), Phosphor (P), and sulfur (S)	99
2.3.3.4.4.1. Nitrogen (N)	99
2.3.3.4.4.1.1. General remarks	99
2.3.3.4.4.1.2. Sources of nitrogen utilizable for Lemnaceae	99
2.3.3.4.4.1.3. The effect of different nitrogen supply	99
2.3.3.4.4.1.4. The effect of nitrate and ammonium; uptake mechanisms	104
2.3.3.4.4.1.5. Utilization of organic nitrogen	112
2.3.3.4.4.2. Phosphorus (P)	113
2.3.3.4.4.3. Sulfur (S)	116
2.3.3.4.5. Carbon (C)	118
2.3.3.4.6. Chlorine (Cl) and other halogens (F, Br, J)	121
2.3.3.4.7. Metallic trace elements	122
2.3.3.4.7.1. General remarks	122
2.3.3.4.7.2. Boron (B) and aluminum (Al)	123
2.3.3.4.7.3. Molybdenum (Mo) and tungsten (W)	125
2.3.3.4.7.4. Iron (Fe)	125
2.3.3.4.7.5. Manganese (Mn)	126
2.3.3.4.7.6. Zinc (Zn)	127
2.3.3.4.7.7. Copper (Cu)	128
2.3.3.4.7.8. Cobalt (Co) and nickel (Ni)	130
2.3.3.4.7.9. Chromium (Cr)	130
2.3.3.4.7.10. Lead (Pb) and cadmium (Cd)	131
2.3.3.4.7.11. Arsenic (As), selenium (Se) and vanadium (V)	133
2.3.3.4.7.12. Silver (Ag) and mercury (Hg)	134
2.3.3.4.7.13. Germanium (Ge) and silicone (Si)	134
2.3.3.4.7.14. Other minerals	135
2.3.3.5. Organic substances; toxins	135
2.3.3.5.1. General remarks	135
2.3.3.5.2. Energy suppliers and building material	135
2.3.3.5.2.1. Carbohydrates	135
2.3.3.5.2.2. Amino acids	139
2.3.3.5.3. Chelating agents	142
2.3.3.5.3.1. General remarks	142
2.3.3.5.3.2. Ethylenediaminetetraacetic acid (EDTA)	143
2.3.3.5.3.3. EDDHA, SA, ASA, BEA, 8-HQ	144
2.3.3.5.4. Vitamins	146
2.3.3.5.5. Hormones and hormone-like substances	148
2.3.3.5.5.1. Auxins, auxin antagonists and effectors of auxin metabolism (BZ, 2,4-D, IAA, IBA, NAA, PAA, PCIB, FPA, TIBA)	148
2.3.3.5.5.2. Cytokinins (adenine, BA, iP, kinetin, purine, thiokinetin, zeatin)	150

2.3.3.5.5.3. Gibberellins (GA)	155
2.3.3.5.5.4. Absciscic acid (ABA)	156
2.3.3.5.5.5. Ethylene and ethylene releasing substances (ethrel, ACC)	160
2.3.3.5.6. Toxic substances and phytochemical substances not classified	161
2.3.3.5.6.1. General remarks	161
2.3.3.5.6.2. Herbicides	161
2.3.3.5.6.3. Algicides and fungicides	170
2.3.3.5.6.4. Insecticides and other pesticides	170
2.3.3.5.6.5. Detergents	171
2.3.3.5.6.6. Further toxins and other plant active substances	172
2.3.4. Influence of temperature	176
2.3.4.1. Temperature dependence of metabolic processes	176
2.3.4.2. Influence of constant and varying temperatures	181
2.3.4.3. Optimum temperature	183
2.3.4.4. Minimum temperature	184
2.3.4.4.1. Minimum temperature for survival	184
2.3.4.4.2. Minimum temperature for growth	185
2.3.4.5. Maximum temperature	186
2.3.4.5.1. Short-term maximum temperature	186
2.3.4.5.2. Long-term maximum temperature	188
2.3.5. Influence of light and other radiations	188
2.3.5.1. Light (including UV radiation)	188
2.3.5.1.1. General remarks	188
2.3.5.1.2. Light as source of energy in photosynthesis	188
2.3.5.1.2.1. Light saturation, minimal and maximal light intensity	189
2.3.5.1.2.2. Light effect in relation to temperature	194
2.3.5.1.2.3. Light effect in relation to nutrient supply	195
2.3.5.1.2.4. Differences in the light effect between different species and clones	196
2.3.5.1.2.5. Relation between light effect and the age of the fronds	197
2.3.5.1.2.6. Indirect effects of light intensity	197
2.3.5.1.2.7. Duration of light	198
2.3.5.1.3. Light effect via phytochrome system	200
2.3.5.1.4. Light effects not completely explainable by chlorophyll or phytochrome action	201
2.3.5.1.5. Heterotrophic growth	204
2.3.5.2. Radiations outside the visible range	205
2.3.5.2.1. X-Rays (γ -radiation)	208
2.3.5.2.2. Radionuclides and other radioactive substances	209
2.3.5.2.3. Laser radiation	210
2.4. Development	211
2.4.1. Vegetative development	211
2.4.1.1. Morphogenesis of fronds	217
2.4.1.2. Morphogenesis of roots	219
2.4.1.3. Abscission	219
2.4.1.4. Senescence and rejuvenation	220
2.4.1.5. Chloroplast development	228
2.4.1.6. Movement of chloroplasts	231
2.4.2. Dormancy	235
2.4.2.1. Induction of turions and turion-like fronds	235
2.4.2.2. Dormancy and germination of turions	239

2.4.2.3. Dormancy and germination of seeds	241
2.4.3. Flowering	243
2.4.3.1. Flower induction	243
2.4.3.1.1. The hormonal state	243
2.4.3.1.2. Some heavy metals, chelating agents, and phenolic substances	251
2.4.3.1.3. Significance of nitrogen metabolism	254
2.4.3.1.4. Significance of carbohydrates and energy-generating processes	258
2.4.3.1.5. Photoperiodism and effects of coloured light	260
2.4.3.1.6. Effects of temperature	264
2.4.3.1.7. Effectors of membrane transport	264
2.4.3.1.8. Some further factors	265
2.4.3.2. Flower and fruit development	267
2.5. Metabolism	270
2.5.1. Generation of energy	270
2.5.1.1. Photosynthesis	270
2.5.1.1.1. Factors limiting or restricting photosynthesis	274
2.5.1.1.2. Steps of the photosynthetic process	281
2.5.1.2. Respiration	284
2.5.2. Membrane transport	284
2.5.2.1. Membrane potential, proton extrusion and proton co-transport processes	284
2.5.2.2. Uptake and release of electrolytes	287
2.5.2.2.1. Alkaline cations	287
2.5.2.2.2. Ammonium	289
2.5.2.2.3. Alkaline-earth cations	290
2.5.2.2.4. Anions	290
2.5.2.2.5. Leakage	292
2.5.2.3. Uptake, transport and release of organic substances	293
2.5.2.3.1. Sugars and amino acids	293
2.5.2.3.2. Phytohormones of other growth substances	294
2.5.2.3.3. Further organic substances	295
2.5.2.3.4. Leakage	296
2.5.3. Location and accumulation of minerals	298
2.5.3.1. Location in organs and tissues	298
2.5.3.2. Location within the cell	298
2.5.3.3. Accumulation	300
2.5.4. Water relations	302
2.5.5. Carbon metabolism	304
2.5.5.1. Starch	304
2.5.5.2. Organic acids	305
2.5.5.3. Myo-Inositol	307
2.5.5.4. Apiose and apogalacturonans	308
2.5.6. Assimilation of phosphorus, nitrogen, and sulfur	311
2.5.6.1. Assimilation of phosphate	311
2.5.6.2. Storage of phosphate	311
2.5.6.3. Acquisition of phosphate at P deficiency	312
2.5.6.4. Nitrate reduction	314
2.5.6.5. Assimilation of ammonia, urea, and other exogenous organic N compounds	316
2.5.6.6. Assimilation of sulfur	320
2.5.7. Amino acids, proteins, and nucleic acids	324
2.5.7.1. Biosynthesis of amino acids	324
2.5.7.1.1. Lysine	324
2.5.7.1.2. Homocysteine, methionine	324

2.5.7.1.3. Threonine, isoleucine	327
2.5.7.1.4. Valine, leucine	328
2.5.7.1.5. Other amino acids	328
2.5.7.2. Methionine metabolism	328
2.5.7.3. Biosynthesis and turnover of proteins	332
2.5.7.3.1. Dependence on light	332
2.5.7.3.2. Proteins of the chloroplast	334
2.5.7.3.3. Other proteins	336
2.5.7.3.4. Protein metabolism in general	337
2.5.7.4. RNA metabolism	339
2.5.7.4.1. Ribosomal RNA	339
2.5.7.4.2. Poly(A) RNA	342
2.5.7.4.3. Inhibitor studies	343
2.5.7.4.4. Ribonucleases	343
2.5.7.5. DNA structure and function	344
2.5.7.5.1. Chloroplast DNA	344
2.5.7.5.2. Mitochondrial DNA	346
2.5.7.5.3. Nuclear genes	347
2.5.7.5.4. In vitro transcription, DNA repair, plasmid uptake	347
2.5.8. Further compounds	349
2.5.8.1. Phospholipids and galactolipids	349
2.5.8.2. Isoprenoids	350
2.5.8.3. Chlorophyll	351
2.5.8.4. Cinnamic acids	352
2.5.8.5. Flavonoids	354
2.5.9. Circadian rhythms in metabolism and photoperiodism	361
3. Application and economic importance	369
3.1. Introduction	369
3.2. Biomass, productivity, energy content, and nutritive value	371
3.2.1. Productivity	371
3.2.2. Biomass	373
3.2.3. Energy content	374
3.2.4. Content of proteins and other nutritive substances	375
3.3. Cultivation and harvest	379
3.3.1. Cultivation	379
3.3.2. Harvest	380
3.4. Utilization as food plant	382
3.4.1. General remarks	382
3.4.2. Human nutrition	384
3.4.3. Mammalian feed	384
3.4.4. Bird feed	386
3.4.5. Feed-stuff for fish and other cool-blooded animals	387
3.5. Utilization in waste water	390
3.5.1. General remarks	391
3.5.2. Removal of nutrients	390
3.5.3. Removal of heavy metals and other toxicants	394
3.6. Utilization as test and indicator plant	400
3.7. Utilization for production of energy and chemical compounds	402
3.7.1. Energy production	402
3.7.2. Production of pharmaceutical compounds	402
3.7.3. Enzyme production	403
3.8. Special utilizations	406
3.8.1. Cosmonautic plant	406
3.8.2. Manure	406

3.8.3. Reduction of water losses in arid regions	407
3.8.4. Reduction of mosquito breeding	408

script - Acknowledgements	410
---------------------------	-----

ography	413
---------	-----

ey of subfamilies, genera, sections and species in the <u>Lemnaceae</u>	581
eviations	582
of the figures	585
of the tables	588
x of subjects	590
ents of vol. 1 (E. LANDOLT, 1986.	635
eröff.Geobot.Inst.ETH.Stiftung Rübel, Zürich, 71)	

1. PHYTOCHEMICAL CHARACTERISTICS

1.1. GENERAL CHEMICAL COMPOSITION AND INORGANIC COMPONENTS

1.1.1. Water content

Lemnaceae contain between 86% and 97% water depending on growth conditions. The main variation lies between 91% and 96% (e.g. VERNADSKY and VINOGRADOFF 1931, KRZECZOWSKA et al. 1975, MUZTAR et al. 1978, AMADO et al. 1980, BONOMI et al. 1981, MONETTI and BASTELLI 1983). Favourable growth conditions (fast growth rate) result in a greater amount of water. On the other hand, resting fronds, especially turions, store a lot of starch. Therefore they have a much higher dry weight percentage than normally growing fronds. The water content in Lemnaceae fronds is lower at higher temperatures than at lower ones (MESTAYER et al. 1984). However, no distinct differences between different species have been observed. 8 species (2 species of each genus) cultivated under identical conditions varied only between 3.9% and 4.1% dry weight (AMADO et al. 1980). The differences between different species found by VERNADSKY and VINOGRADOFF (1931) must be attributed to different environmental conditions.

1.1.2. Variation of mineral content

In table 1.1 the variations in mineral content for the different elements are listed. The results are not split up according to the species since the culture conditions in the different investigations varied considerably. In general, the variations of the mineral content between plants cultivated under different culture conditions are much greater than between different species. DYKJJOVA (1979) summarized the known results of measurements of mineral content of Lemnaceae and other water plants. Much more data have been published since. Some of the papers deal with Lemnaceae growing in waters charged with heavy metals from industrial wastes. For the abilities of Lemnaceae to accumulate heavy

metals see chapter 2.5.3.3 and 3.5.3. A detailed enumeration of all literature indications as well as of the growth conditions and the species of Lennaceae used was not possible for table 1.1 due to difficulties in making the many and complex data comparable. The results of the following publications have been incorporated within the data of table 1.1.:

AMADO et al. (1980), ALLENDY (1967, 1968, 1981), BAUMEISTER and ERNST (1978), BONOMI et al. (1981), BOYD (1968), CLARK et al. (1981), COMGILL (1970), CULLEY and EPPS (1973), CULLEY et al. (1978), DENTON (1966), DY-KYJOWA (1979), ERNST and MARQUENIE-VAN DER WERFF (1978), FANKHAUSER et al. (1976), GLANDON and McNABB (1978), GUTHRIE and CHERRY (1979a,b), HAKONSON and WHICKER (1975), HARVEY and FOX (1973), HUNTER (1976), HUTCHINSON and CZYRSKA (1975), KARPATI and POMOGYI (1979), KARPATI et al. (1985), KHAKIMOVA et al. (1971), KHAKIMOVA and GALKINA (1973), KOVACS et al. (1984), KRZECZOWSKA et al. (1975), LEDL et al. (1981), LIEBERT (1980b), MOHETI and BASTELLI (1983), MUDROCH and CAPOCIANCO (1979), MUZZAR et al. (1978), ORNES (1979), PEDKOVA and LUBIANOV (1969), PEVERLY (1985), PIISPANEN and LAHDESMAKI (1983), POLAR and KUECKEZEZAR (1986), REAY (1972), RIEMER and TOTH (1968), RODGERS et al. (1978), RUSOFF et al. (1980), SANKARAN (1972), SETO et al. (1979), SILVEY (1967), STRAUSS (1973, 1976), SUCKCHAROEN (1980), SUTTON and ORNES (1975), TAN (1970), THELLIER and LE GUEL (1967a,b), TRIDECH et al. (1981), VAN DER WERFF (1981), VARENKO and CHUICO (1971), VARENKO and LUBIANOV (1973), VAVRUSKA (1966), VERNAUSKI and VINOGRADOFF (1931), WELMER and ARMSTRONG (1979), WENTSEL and BERRY (1975), WOLVERTON and McDONALD (1981).

The ratio of some elements in Lennaceae is shown in table 1.2, it is also dependent on the environmental conditions (see chapter 1.1.5).

Table 1.2. Mineral ratios in Lennaceae, according to many authors (see list in the text)

K/Na	0.67 - 37
K/Ca	1 - 2
Ca/Mg	0.05 - 20
C/N	6.7 - 8.6

Table 1.1. Variation of content of elements in % of the dry weight (according to many authors, see list in the text)

Ag	0.3 - 50 x 10 ⁻⁶	Mg	0.04-2.8
Al	0.000-11.4	Mn	0.003-6.4
As	0.2-23.5 x 10 ⁻³	Mo	0.2-0.4 x 10 ⁻³
B	0.02-3.25	N	0.8-7.8
Ba	0.03-0.11	Na	0.03-1.3
Br	0.25-0.65 x 10 ⁻²	Nb	0.2 x 10 ⁻³
C	30.5-43.7	Ni	0.7 x 10 ⁻⁴ - 0.2
Ca	0.18-4.5	P	0.03-2.8
Cd	<0.1 x 10 ⁻⁴ - 6.7	Pb	0.2 x 10 ⁻⁴ - 0.2
Ce	0.2 x 10 ⁻³	Pr	0.4 x 10 ⁻⁴
Cl	0.08-4.29	Ra	traces
Co	0.9 x 10 ⁻⁴ - 1.1	Rb	0.0054
Cr	0.3-17.8 x 10 ⁻³	S	0.33-7.0
Cs	0.4-50 x 10 ⁻³	Sb	0.0015-0.012
Cu	0.2 x 10 ⁻³ - 3.2	Se	0.0018-0.12
F	0.2 x 10 ⁻³	Si	0.41-5.35
Fe	0.007-3.2	Sn	0.2-3.6 x 10 ⁻²
Ga	0.9 x 10 ⁻⁴	Sr	0.008-0.11
H	5.4	Ti	0.0018-0.32
Hg	0.04-18 x 10 ⁻⁴	V	0.3-10 x 10 ⁻³
J	0.4-25 x 10 ⁻⁴	Y	0.4 x 10 ⁻⁴
K	0.03-7.0	Zn	0.004-0.14
La	0.9 x 10 ⁻⁴	Zr	0.9 x 10 ⁻⁴
Li	0.8-6 x 10 ⁻³		

to be characteristic. The ratio is 7 in L. minor compared with 17 in Eichhornia (WOLVERTON and McDONALD 1981). This is explained by the fact that Lemnaceae have only thin cell walls and a very reduced vascular system (scarcely any sclerenchymatous tissue). STRAUSS (1973) pointed out the unusually high content of magnesium compared with Ca accumulated by L. minor. The Mg/Ca ratio was always above 1, under certain conditions it reached values up to 20. Measurements of other authors do not fully confirm these results. Mostly the Ca content was much higher (up to 20 times) than the Mg content (cf. COMGILL 1970, DYKJOWA and KVERT 1978, DYKJOWA 1979). The content of inorganic anions is especially high in L. trisulca. It amounts to 7% S (in form of SO_4^{2-}), 1.4% Cl^- (JANAUER 1982).

The great amount of phosphorus in Lemnaceae is explained by the ability to form inositolphosphates (see chapter 1.2.5) and condensed inorganic phosphates (BIELESKI 1968a). In several Lemnaceae species NIEMEYER (1975) and INHUELSEN and NIEMEYER (1975) demonstrated linear oligophosphates with two to seven phosphate residues, cyclic condensed metaphosphates (tri-, tetra-, penta- and hexa-metaphosphates) and high molecular condensed phosphates. The saturation of these poly-anions by cations is undoubtedly the basis of the unusual ability to accumulate heavy metals. The high accumulation of boron by Lemnaceae is explained by the ability to bind the absorbed boron in the cell walls in form of easily hydrolysable boromonoesters and of very stable borate diesters (DUVAL et al. 1980, THELLIER et al. 1979).

1.1.5. Variation of mineral content due to different measuring techniques and different environmental conditions

Washing of the Lemnaceae before analyzing yields different results compared with unwashed samples (MUZTAR et al. 1978). If washed under acid conditions, an additional amount of SO_4^{2-} , PO_4^{3-} , Ca^{2+} , K^+ , and Na^+ is washed away (up to more than 50%). Evidently, these elements are in- crusted on the surface of the frond (ALLENBY 1981). For this reason the upper limits of mineral content in table 1.1 may well be too high.

The concentration of the minerals in the nutrient solution has a primary influence on the mineral content within the frond. The N and P content rose from 2.8 to 5.6% N and 0.4 to 1.4% P, respectively between low and

1.3. Status of some elements within the Lemnaceae

The total amount of nitrogen within the frond of L. minor c. 12% are of the water soluble amino acids and proteins. A very small amount was identified as ammonium (BRAUER et al. 1971). WEINER and ARM- (1979) measured 3.8 mg P per g dry weight in Lemna sp. from Wis- in. 13% of the phosphorus was organic. This organic P was bound to (total to 37%). More than 70% of the total sulfur content of L. minor is incorporated into organic thio compounds, and more than 25% is present as free sulfate in the vacuoles (THOIRON et al. 1981). The iron is present in L. gibba in the ferric form (GOODMAN and DEKOCK 1982, GOODMAN et al. 1982). Most of the iron is antiferromagnetically-coupled at low temperatures. The most common compound found was ferritin, an iron-storage protein. Boron is found as borate in the cytoplasm and in the vacu- of L. minor; also borate diester is present (THELLIER et al. 1979). UHANN et al. (1986) developed a method of analytical ion microscopy demonstrate the distribution of nitrogen in L. gibba.

1.4. The mineral content of Lemnaceae in comparison with other water plants

According to DYKJOWA (1979), a medium to great amount of most minerals typical for Lemnaceae compared with other water and marsh plants (150 species tested). The content of the following elements was not, or rare- reached in other plant families: N, P, Mn, Zn, Ni, Si, Ba, Rb, Pb, and B. The K/Ca ratio varying between 1 and 2 in Lemnaceae is very for monocotyledons. In L. minor, the content of Ca, Na, and Zn is or, the content of Fe lower than in Eichhornia (BONOMI et al. 1981). accumulation of boron in L. minor is ten times higher than in Cera- yllum demersum (GLANDON and McNABB 1978). VARENKO and CHUICO (1971) found an especially great amount of Zn and Co in S. polyrrhiza com- pared with other water plants. Mn showed the second highest and Cu only lithium value. A very high accumulation of Cd, Zn, Cu, and Pb was ob- served by VAN DER WERFF and PRUIT (1982) in fronds of S. polyrrhiza and ibba. Only Elodea and Callitriche out of many investigated water plants showed similar or slightly higher content of these elements. A relatively low ratio of C/N for normal growing fronds of Lemnaceae seems

high N and P content in the culture medium. The maximum content in the frond was reached at a concentration of about 4 mg of N and P, respectively (REJMANKOVA 1981, see fig. 1.1). AEBLI (1986) got similar results with *L. minor*. The P content varied between 0.05% of the dry weight (at a concentration of 0.025 mg P per liter culture solution), 0.1% (0.1 mg/l), 0.37% (0.4 mg/l), and 1% (1.55 mg/l). At higher P concentrations in the nutrient solution, the P content of the fronds did not enhance considerably.

The content of heavy metals is correlated with the concentration of the metal in the nutrient solution (fig. 1.2).

The concentration of a mineral is also dependent on the concentration of other minerals. For instance, the potassium content of *L. gibba* and *L. minor* can rise from 2.5% of the dry weight under normal conditions up to 5% under low nitrogen supply (ERICSSON et al. 1982). The copper content is raised to about 130% if Cd is added to the culture solution together




Fig. 1.1.1. The N and P content of *Lemna gibba* in relation to the N and P content of the nutrient solution (from REJMANKOVA 1981)

with copper. On the other hand, the Cd content is lowered by the simultaneous addition of Cu, As, or Zn; the Mn content is lowered by Cu, Cd, and Zn in the solution, and the Zn content by Cu and As (NASU et al. 1985). The C/N ratio is significantly higher in N-deficient cultures and in turions. It can reach up to 20. The calcium content of *L. gibba* is enhanced by the addition of ABA to the medium. It is lowered by BA (DEKOCK and HALL 1981). On the other hand, ABA at a concentration of 10 ppm diminishes the potassium content of *L. gibba* from 60 mg to 15 mg per g dry weight (LIEBERT 1977). The lowest P content under which growth of *L. minor* is still possible lies between 0.036 and 0.053% of the dry weight (FEKETE et al. 1976).




Fig. 1.1.2. The Cd, Cr, Cu, Pb, and Zn content of *Spirodela polyrrhiza* in relation to the concentration of the metals in the nutrient solution (data for Cd, Cu, Pb, and Zn from VAN DER WERFF 1981, for Cr from STAVES and KNAUS 1985).

The upper limit of mineral content which still allows an optimal growth rate in L. minor proved to be 0.08% of the dry weight for boron (FRICK 1985a), 0.05% for iron, 0.18% for manganese, 0.03% for copper, 0.11% for inc. and 0.56% for molybdenum (FRICK 1985b).

1.1.6. Variation of the mineral content during different seasons and during lifetime

AKHIMOV et al. (1981) measured the nitrogen content of L. minor during different seasons. In winter the nitrogen content as well as the content of sucrose and mannose are distinctly higher than during the summer; for the cellulose content it is vice versa. In table 1.3, another example is given. Main changes in this example are within the organic compounds. The content of certain elements during lifetime is not stable. FISCHER (1949) showed that the Ca content of the fronds of L. minor rises and the K content diminishes with age. LIEBERT (1980b) measured 80 mg Ca per dry weight in old fronds and old cultures whereas the Ca content of young fronds and of young cultures was only 28 mg per g dry weight.

1.1.7. Variation of mineral content in different species and clones

In table 1.4, differences in mineral content between S. polyrrhiza, L. minor, and L. trisulca are shown (KARPATI and POMOGYI 1979). S. polyrrhiza and L. minor were collected in the same pond, L. trisulca originated from a neighbouring one. Table 1.5a gives an example of three Lemna species cultivated under the same conditions (ERICSSON et al. 1982). KOLES (1986) put some results from literature together of minimum and maximum nitrogen and phosphorus content in different species (table 1.5b). Since the study conditions of the different authors were not identical, the results are only partly comparable. DOCAUER (1983) measured the mineral content which is necessary to achieve half of the maximum growth rate in five different Lemnaceae species (table 1.6). From tables 1.4 to 1.6 it can be drawn that there are distinct differences between different species. Since only one clone of each species was investigated, it is not known if the differences are species specific or only characteristics of different clones. The relatively big differences

Table 1.3. Variation of chemical composition of Wolffia globosa from Japan (named as W. arrhiza) between summer and winter (after SEKINE 1979) in % of the dry weight

	protein	fat	carbo- hydrate	cellu- lose	minerals	H ₂ O	chloro- phyll
summer	40-45	10-14	25-30	1-2	6-8	3	1-2
winter	8-10	18-20	60-65	1-2	6-8	3	1-2

Table 1.4. Differences in mineral content between Spirodela polyrrhiza, Lemna minor, and L. trisulca (according to KARPATI and POMOGYI 1979). Content in % of the dry weight (converted from the values of the fresh weight by the factor 20)

Species	K	Na	Ca	Mg	Mn	Zn	Fe	P	N
<u>S. polyrrhiza</u>	0.27	0.12	0.31	2.17	0.54	0.05	0.74	0.97	3.99
<u>L. minor</u>	1.53	0.02	0.18	1.92	0.03	0.05	0.06	0.83	8.74
<u>L. trisulca</u>	0.26	0.05	0.34	2.90	0.55	0.04	0.44	1.81	4.79

Table 1.5a. Mineral content of three Lemna species cultivated under identical conditions (according to ERICSSON et al. 1982) in % of the dry weight

Species	Elements				
	N	K	P	Ca	PO ₄
<u>L. gibba</u>	4.6	5.3	1.1	0.57	0.35
<u>L. minor</u>	4.7	5.7	0.86	0.47	0.40
<u>L. aequinoctialis</u>	4.1	4.9	0.74	0.33	0.30

in table 1.3 may be attributable to local environmental differences rather than purely to differences in species. From the results of ERICSON et al. (1982) it seems that L. aequinoctialis generally has a lower level of minerals than the other two species investigated. The investigations of DOCAUER (1983) indicate that the Wolffias need to accumulate

Table 1.5b. Differences of minimum and maximum content of N and P between different species (modified from KOLES 1986)

Species	content in % of the dry weight			
	N min	N max	P min	P max
<u>Spirodela polyrrhiza</u>	1.7 (a)	7.9 (b) -	0.3 (c)	2.1 (d)
<u>S. punctata</u>	1.7 (d)	7.2 (e)	0.6 (f)	2.4 (g)
<u>Lemna gibba</u>	1.5 (d)	7.7 (b)	0.7 (d)	2.6 (d)
<u>L. obscura</u>	0.8 (h)	4.6 (i)		
<u>L. minor</u>			0.4 (k)	2.4 (g)

References
 SUTTON and ORNES 1977 f CULLEY et al. 1981
 ORON et al. 1985 g ALLENBY 1981
 SUTTON and ORNES 1975 h TRIDECH et al. 1981
 CULLEY et al. 1981 i HARVEY and FOX 1973
 HILLMAN and CULLEY 1978 k MUZTAR et al. 1976

Table 1.6. Mineral content of Lemnaceae species grown at concentrations of minerals to accomplish half of the maximum growth rate (in moles per mg wet weight) (from DOCAUER 1983)

Species	NO ₃ ⁻	PO ₄ ³⁻
<u>S. polyrrhiza</u>		
<u>L. minor</u>	0.14	0.011
<u>L. turionifera</u>	0.12	0.008
<u>W. borealis</u>	0.14	0.006
<u>W. columbiana</u>	0.17	0.018
	0.08	0.016

more phosphates to reach a good growth rate than S. polyrrhiza and the two Lemnas. W. borealis has an extremely high and W. columbiana an extremely low level of nitrate when half the growth rate is reached.

Of the three investigated species S. polyrrhiza, S. punctata and L. gibba, S. punctata accumulates most Cr and L. gibba least at low Cr concentrations. At higher concentrations (1 ppm and more) there is no longer any difference between the species (STAVES and KNAUS 1985). In a comparison between S. polyrrhiza and L. minor, VAN DER WERFF (1981) observed more Cd, Cu, and Pb and less Zn in L. minor than in S. polyrrhiza at concentrations between 0.1 to 10 mM in the solution.

1.2.2. Proteins and amino acids

1.2.2.1. Proteins

The protein content of *L. minor* is composed of 49% albumine, 1.7% globuline, 32.1% gluteline, and 11.2% prolamine, according to MACIEJEWSKA-POTAPCZYK et al. (1970). The corresponding values for *S. polyrrhiza* amount to 25-55% albumine, 0.7-5% globuline, 15-30% gluteline, and 18-49% prolamine (included other remaining proteins), the variation depending on the preceding growth conditions (BYTNIIEWSKA and POTAPCZYK 1981).

The total protein content of the *Lemnaceae* varies between 6.8 and 45% of the dry weight (e.g. WHITE 1939, KESER 1955, MACIEJEWSKA-POTAPCZYK et al. 1970, BHANTUNNAVIN and MCGARRY 1971, TULGANOW 1972, CHANG et al. 1977, 1978, PORATH et al. 1979, AMADO et al. 1980, ROBINETTE et al. 1980, RUSOFF et al. 1980, BYTNIIEWSKA and POTAPCZYK 1981, APPENROTH et al. 1982).

The total content of protein in *Lemnaceae* is, under favourable conditions, one of the highest within the plant kingdom. ANTANINENE and TRAI-NAUSKAITE (1985) found *L. trisulca*, in Lithuania, one of the four species with the highest protein content out of 74 macrophyte species tested.

The protein content is much dependent on the growth conditions. It rises with higher light intensity (WHITE 1939); also it is distinctly higher at 24°C than at 18°C (LEHMANN et al. 1981). Young cultures contain more protein than old ones (CHANG et al. 1978) and fronds more than roots (LEHMANN et al. 1981). Fronds of *L. aequinoctialis* with a size of 5-6% of the final one only reach about 10% of that of the full-grown fronds (DAYKO et al. 1980b). The mineral content of the nutrient solution (especially the nitrogen content) greatly influences the protein content of three nutrient solutions tested (Hoagland, Hutner 1/5, and Pirson-Seidel). Hoagland gave rise to the highest protein content (AMADO et al. 1980). This is probably due to the fact that the nitrogen content is highest in the Hoagland solution (see also KRZECOWSKA et al. 1975). Indeed, the nitrogen content of the nutrient solution (at least for lower concentrations, see fig. 1.1) is in the first place positively correlated with the protein content of the fronds (CULLEY and EPPS 1973, SUTTON and ORNES 1975, SAID et al. 1979, REJMANKOVA 1979). ORON et al. (1987) measured the protein content of *L. gibba* at different concentrations of

2. ORGANIC COMPONENTS

2.1. General remarks

A variation of the different main groups of organic components in *Lemnaceae* is summarized in table 1.7. The data were taken from the same authors as in table 1.1. CHANG et al. (1978) measured much higher content of lipids (up to 26.2%) than other authors, probably due to different analyzing methods. JOHRI and SHARMA (1980) report 47% ash in *L. minor*. This great amount can scarcely be explained. The percentage of ash remains relatively constant under different temperature conditions of growth (NESTAYER et al. 1984). The carbohydrates are not always defined and measured in the same way. In starving cultures and in turions, the content of carbohydrates (especially starch) can reach much higher values. REID and BYLESKI (1970a) observed 75% starch in *S. punctata* grown in P-deficient cultures.

The density of the culture did not exert any influence on the content of ash, carbohydrate, and protein, nor on the amount of the cell wall fraction in *L. minor* (TUCKER 1981). Differences between species may occur. However, no investigations have been made with more than one clone (except the amino acid studies of AMADO et al. 1980, see chapter 1.2.2). CHANG et al. (1978) measured the crude fibre content of *S. polyrrhiza*, *S. punctata*, *L. aequinoctialis*, and *W. globosa* (named as *W. arrhiza*) with 7.4%, 6.1%, 7.0%, and 7.4%, respectively.

1 to 2.6% of the total carbon fixed by *L. minor* is released to the water (BAKER and FARR 1987).

Table 1.7. Variations of organic components in *Lemnaceae* (data from many authors) in % of the dry weight

proteins	6.8 - 45.0
lipids	1.8 - 9.2
crude fibres	5.7 - 16.2
carbohydrates	14.1 - 43.6
ash	12.0 - 27.6

ammonium. It amounted to c. 17% at an ammonium concentration of 5 mg/l and to c. 28% at 10 mg/l. BORNKAMM (1970a) measured the protein content of L. minor and L. trisulca in solutions with 0.5 and 1.0 mM N. In the first-named concentration the protein content was 7.1 and 5.0%, respectively, in the second concentration 18.2 and 9.9%, respectively. According to MALEK and COSSINS (1983b), the protein content of nitrogen- and sulfur-deficient cultures of S. polyrrhiza was only 8 to 31% of that of the controls. Higher levels of Ca in the medium consequently lowered the protein content of S. polyrrhiza (LECHEVALLIER 1977a). Exposure to ozone resulted in a 25% reduction of the protein content of L. minor within two hours (CRAKER 1972). The density of the culture, however, has no influence on the protein content of L. minor.

The differences in protein content between different species are generally smaller than between different clones of the same species. In 5 clones of L. trisulca cultivated under identical conditions, AMADO et al. (1980) measured the following protein content in % of the dry weight: 6.8, 15.4, 17.7, 18.0, 23.0. The total variation of the protein content of all 94 clones of 28 species was between 6.8 and 37.0%. All stress situations cause a loss of soluble protein per frond due to a decrease in the rate constant of protein synthesis and an increase in the rate constant of protein degradation (DAVIES 1978, COOKE et al. 1979b) (see also chapter 2.4.1.4). Extensin, a component part of stress-produced proteins was analyzed in L. minor (BIGGS and KOSSUTH 1985). Stresses which increased levels of extensin include temperature extremes, intense photosynthetically active radiation, UV-B-radiation, and strong osmoticum.

GTB-binding proteins were detected in the extract of L. aequinoctialis containing membrane components (HASUNUMA and FUNADERA 1987). NECHUSHTAI et al. (1987a,b) isolated two chlorophyll-protein complexes of the photosystem I complex of L. gibba.

1.2.2.2. Amino acids

KESER (1955) detected 28 amino acids in S. polyrrhiza. L. minor also contained 28 amino acids which were however partly different from those of S. polyrrhiza. Table 1.8 gives a survey of the content of principal amino acids in Lemnaceae, according to different authors. MACIEJEWSKA-POTAPCZYK et al. (1970) and TULCANOW (1972) investigated L. minor; CHANG

et al. (1977, 1978) S. polyrrhiza, L. aequinoctialis, and W. globosa (named as W. arhiza); BYTNIEMSKA and MACIEJEWSKA-POTAPCZYK (1980) S. polyrrhiza; PORATH et al. (1979) S. polyrrhiza, L. gibba, L. minor, and W. arhiza; RUSOFF et al. (1980) S. polyrrhiza, S. punctata, L. gibba, and W. columbiana; APPENROTH et al. (1982) W. arhiza, and AMADO et al. (1980) 94 clones of 28 species of all 4 genera of Lemnaceae. In addition to the amino acids mentioned in table 1.8, KESER (1955) noted within the free amino acids of S. polyrrhiza and L. minor glutathione, taurine, β -alanine, γ - and α -aminobutyric acid but no oxyproline. Oxyproline was only observed in the protein.

The percentage of the different amino acids is very different depending on growth conditions, age of cultures, and methods of analysis. Lack of nitrogen alone lowers the content of free amino acids (especially asparagine) in S. polyrrhiza and L. minor and raises the content of glutamine. K- and P-deficiency in the solution had no influence, however (KESER 1955). The percentage of the content of different amino acids of S. polyrrhiza in N- and S-deficient cultures varies between 7% for methionine (controls 100%) and 1793% for asparagine (MALEK and COSSINS 1983b). Urea as nitrogen source leads to distinctly lower portions of asparagine, aspartic acid, arginine, glutamine, glutamic acid, and alanine (COOK 1968). Raising CO₂ supply enhances the content of alanine, arginine, glutamic acid, aspartic acid, glutamine, and asparagine and lowers that of glycine, serine, and histidine (MOELLER et al. 1977). HUBALD et al. (1979) investigated the influence of asparagine, glutamic acid and glycine on amino acid and protein content of L. gibba. The light quality (white, red, blue) has no influence on the composition of amino acids of the soluble proteins in W. arhiza (APPENROTH et al. 1982). CHANG et al. (1978) demonstrated the effect of growth substances and age of culture on the composition of amino acids in Lemnaceae. PERRY (1963) observed different protein constituents in turions and normal fronds of S. polyrrhiza. Differences in the amino acid content between different species do not show up, however, differences between clones are frequent (AMADO et al. 1980).

In general, Lemnaceae, if compared with other plant families, contain a relatively great amount of leucine, threonine, valine, isoleucine, and phenylalanine whereas the content of cysteine, methionine, tyrosine, and taurine is relatively low.

Table 1.9. Carbohydrate content in Lemnaceae, in % of the dry weight

Elements	Species						
	L. minor						
	L. tri- sulca	L. aequi- noctialis	L. minus- cula	"duck- weed"			
Authors	1	2	3	4	5	6	7
soluble sugars		4.8	1.8-13.6	1.3		1	
starch		4.8	5.8- 8.7	0.25	4.6-13.8	3.0	
hemicelluloses			0.9- 9.5				21.7
celluloses			4.1-16.6			14.0	10.0
apiose	4.8						
xylose	2.5						
glucose	14.0						
mannose			0.2- 4.0				
sucrose			0.1- 1.5				

References

- 1 DUFF and KNIGHT 1963
- 2 MUZTAR et al. 1979
- 3 RAKHIMOV et al. 1981
- 4 JANAUER 1982
- 5 CHANG et al. 1978
- 6 STEUBING et al. 1980
- 7 WOLVERTON and McDONALD 1981

tween the different species could be observed (table 1.10). In addition to the neutral sugars mentioned, small amounts of glucosamine and galactosamine were detected as well as a sugar which probably was identical with apiose. Apiose was observed in Lemnaceae at a content of 4-8% of the dry weight (BELL et al. 1954, DUFF and KNIGHT 1963, DUFF 1965, VAN BEUSEKOM 1967, HART and KINDEL 1970a,b). The apiose was isolated mostly from the pectin fraction. According to HART (1969), 83% of the apiose is incorporated in the cell wall where it is present up to 20%, as a component of the polysaccharide apiogalacturonan (BECK 1964, BECK and KANDLER 1965, HART 1969, MASCARO 1975). Among others, PICKEN and MENDICINO (1967), GUSTINE (1969), and GRISEBACH et al. (1972a,b) report on the biosynthesis of apiose and xylose in *L. minor* (see chapter 2.5.5.4). Apiose occurs, within the monocotyledons, in some water and shore plants, especially in species of Hydrocharitaceae, Potamogetonaceae, and Zannichelliaceae which live submerged in brackish water (VAN BEUSEKOM 1967). No apiose was found in the families Araceae and Najadaceae. MUZTAR et al. (1978) analyzed the cell wall of *L. minor*. They observed 30.3% of the dry weight hemicelluloses and 19.2% celluloses (beside 2.4% lignin). According to these authors, *L. minor* has by far the highest hemicellulose content of 11 water plants studied.

The sugar content of the three carbohydrate components pectins, hemicelluloses and celluloses is given in table 1.11 (data from DEKOCK et al. 1979).

Table 1.11. Content of different sugars within the 3 carbohydrate groups of pectins, hemicelluloses and celluloses in *L. gibba*, in % of the total amount (after DEKOCK et al. 1979). The pectin fraction and to a lesser degree the hemicellulose fraction contain components of starch

carbohydrate groups	pectins	hemicelluloses	celluloses
sugars			
rhamnose and fucose	11	0	0
galactose	22	11	0
arabinose	18	8	0
glucose	7	41	75
xylose and apiose	40	38	25
mannose	2	2	0

Table 1.10. Percentage of sugars in Lemnaceae (total amount of neutral sugars is 100%) (after AMADO et al. 1980)

glucose	55.3-57.3	arabinose	8.2-11.8
galactose	13.3-16.4	fucose	2.4- 5.6
mannose	1.9- 3.9	rhamnose	1.9- 2.6
xylose	7.1-14.2		

AMBROSE (1978) investigated the carbohydrates released into water by different Lemnaceae (L. gibba, L. minor, L. trisulca). She always found glycerol and arabinose. L. gibba and L. trisulca emitted in addition fructose, glucose, and some unidentified sugars. Chemical analysis of L. aquinoctialis (named as L. minor) revealed the presence of the reducing sugars cardenolide and digitoxose (YONG and THO 1976).

1.2.4. Lipids and fatty acids

The lipid content of Lemnaceae varies between 2 and 9% (LECHEVALLIER 1966, 197a7, BHANTUNAVIN and MCGARRY 1971, SU et al. 1973a, HILLMAN and CULLEY 1978b, STEUBING et al. 1980). CHANG et al. (1978) analysed between 6.3 and 26.2% lipids. The highest lipid content was measured in M. globosa cultures supplied with gibberellins. Perhaps the striking deviation of the result of CHANG et al. from other results can be explained by different methods of analysis. Like other organic compounds of Lemnaceae, the lipid content is conditioned by the age of the cultures (CHANG et al. 1978) and by the composition of the nutrient solution (LECHEVALLIER 1977a, GRENIER et al. 1979). 63 percent of the lipids of Lemnaceae belong to the phospholipids (with palmitic acid, linoleic acid and linolenic acid), 31% to the galactolipids (mostly linolenic acid), and 6% are neutral lipids (GRENIER et al. 1979). The indications of CHECHENKIN (1955) that no high saturated acids occur in L. minor, are not confirmed. The acid lipids of S. polyrrhiza contain 65% linolenic acid, 15% palmitic acid and 8% linoleic acid; quantitative determinations of fatty acids have also been made for single phospholipids, lipids, and sulfolipids (LECHEVALLIER 1966, 1967). The addition of polyethylene glycol to the nutrient solution results - among other things - in a reduced lipid content of S. polyrrhiza. The relative content of linoleic acid was enhanced and that of linolenic acid lowered (LECHEVALLIER 1977b). A similar shifting of the ratio of linoleic acid/linolenic acid showed up in S. punctata after the addition of glucose to the nutrient solution (GROB and EICHENBERGER 1969). In both cases, the changes of lipid composition occur in connection with enhanced frond senescence (yellowing of chloroplasts). There is another effect of sucrose in younger cultures of S. polyrrhiza, the lipid as well as the chlorophyll and protein content per chloroplast is highest with 2% sucrose supply

(LECHEVALLIER et al. 1971). Also the addition of benzyladenine to the culture medium changes the lipid composition of Lemnaceae. The percentages of total palmitic and α -linolenic acids increase and the percentage of linoleic acid decreases in L. minor after supply of benzyladenine. At 2 and 5 ppm BA the proportion of phospholipids increases greatly as compared with galactolipids (especially the diacylgalactosylglycerol) and total neutral lipids. Also the α -linolenic acid content of total neutral lipids is considerably enhanced (BERUBE et al. 1982). Benzyladenine (10^{-3} mM) increases the total phospholipid content by 20% in S. polyrrhiza. The degree of unsaturation of polar lipid fatty acids rises slightly due to greater amounts of linoleic and linolenic acids (LEPABIC 1980). Atrazine in sublethal doses stimulates the lipid metabolism of chloroplasts in L. minor (higher desaturation of fatty acids) (GRENIER and BEAUMONT 1983). Lipids of chloroplasts are described by BAHL et al. (1971) in S. polyrrhiza. EICHENBERGER (1975) studied the lipids of microsomes in L. minor. The hydrocarbons of S. polyrrhiza and L. trisulca were investigated comparatively by LECHEVALLIER (1969). He observed distinct differences between the two species: the hydrocarbons of S. polyrrhiza mostly contain long saturated and unsaturated chains (C_{17} to C_{29}) which could be identified partly as squalenes and partly as alkanes and alkenes; the hydrocarbons of L. trisulca show shorter chains (C_{12} to C_{23}) and only slight amounts of squalenes and few unsaturated chains (alkenes). Sterols of both species are frequent in form of campesterol, stigmasterol, and β -sitosterol. The proportion of stigmasterol is high in S. polyrrhiza (c. 70%) and low in L. trisulca (c. 25%), the percentage of β -sitosterol is high in L. trisulca (c. 55%) and low in S. polyrrhiza (c. 10%) (LECHEVALLIER 1970). D'HARLINGUE et al. (1976) measured the ratio of campesterol, stigmasterol and β -sitosterol from S. polyrrhiza held in solutions without sugar or with 2% sucrose. The ratios amounted to 26:43:30 and 20:48:30, respectively. The fatty acid composition of L. minor was studied by PREVITERA and MONACO (1983). The triacylglycerol fraction contained mainly 16:1 and 16:3 fatty acids and a hydroxy C_{16} acid. The same authors (1984) analyzed sitosterol, phytadiene, lycopersene, phytol, and (4R)-4-hydroxy-isophytol in L. minor. Different oxygenated fatty acids have been isolated from L. trisulca by MONACO and PREVITERA (1987), e.g. (12S)-hydroxyhexadeca-8Z,10E,14Z-trienoic acid and a prostaglandin-like C_{16} fatty acid.

1.2.5. Organic phosphorus compounds and nucleic acids

After 9 days supply of radioactive phosphate to S. punctata, BIELESKI (1968a) was able to extract phosphorus compounds and to identify them by means of chromatographic and autoradiographic methods. Most of the organic-bound phosphate (5.1 $\mu\text{mol P}$ per g fresh weight) occurs in the RNA. The dominant phospholipid is phosphatidyl choline (1.6 $\mu\text{mol P}$ per g fresh weight). In addition, there are phosphatidyl ethanolamine, phosphatidyl glycerol, phosphatidyl inositol, phosphatidyl serine, and phosphatidic acid. Glucose-6-phosphate (0.5 $\mu\text{mol P}$ per g fresh weight) and phosphoglyceric acid (0.25 $\mu\text{mol P}$ per g fresh weight) are the quantitatively prominent phosphate-esters. Furthermore, the author was able to demonstrate and determine quantitatively most of the nucleotides, sugar phosphates, and some more products of the intermediary metabolic processes as well as several inositol phosphates. The inositol phosphates and especially inositol hexaphosphate (phytic acid) have also been noted in several other Lemnaceae species: S. polyrrhiza, L. gibba, L. minor, W. gladiata (INHUELSEN and NEMEYER 1978, SCHEINER et al. 1978, ROBERTS and LOEWUS 1968). WEIMER and ARMSTRONG (1979) measured 0.37% myo-inositol hexaphosphate in Lemna spp. Phytic acid is produced mainly in turions and resting fronds as a phosphate deposit. Factors which induce the formation of turions and resting fronds (nitrogen deficiency, sugar supply, short-day conditions at temperatures of 8-10°C) consequently enhance the accumulation of phytic acid (SCHEINER et al. 1978). Phosphatidylcholine and phosphoethanolamine bases as intermediates in the synthesis of the first substance have been demonstrated in L. aequinoctialis by MUDD and DATKO (1986b).

D'HARLINGUE et al. (1976) measured the content of NADP and NAD (about 3 to 4 μmol per mg dry weight) in S. polyrrhiza. About 75% of the pyridine nucleotides were oxidized. Pyridine nucleotides of the same species have been investigated by LECHEVALIER et al. (1977). HASANUMA (1986) reports on cyclic nucleotides in L. aequinoctialis. VAN EE and PLANTA (1982) isolated and characterized polyribosomes and non-ribosomal ribonucleoprotein particles in S. punctata. TOBIN and KLEIN (1975) report on the isolation and translation of mRNA in L. gibba. A similar RNA in Spirodela was detected by ROSNER et al. (1977a). Further studies on RNA were made by DYER and BOWMAN (1976), BOWMAN and DYER (1979), GRESSEL (1978), EICHORN (1984, 1986), EICHORN and AUGSTEN

(1984), KEUS et al. (1984a,b), VANDENBERGHE et al. (1984), DE HEIJ et al. (1985), KARLIN-NEUMANN et al. (1985). Investigations on RNA metabolism and on DNA structure and function are summarized in chapters 2.5.7.4 and 2.5.7.5.

The banding pattern of chloroplast DNAs have been used to distinguish different clones of L. aequinoctialis (BEPPU et al. 1985) and to characterize different species of Lemna (BEPPU in lit. 1986).

GEGER (in lit. 1986) measured the DNA content of the chromosomes of some Lemnaceae which varied according to the species (see table 3.2 in LAN-DOLT 1986).

1.2.6. Chlorophylls, carotenoids, phytochrome, and cytochrome

Results on chlorophyll and carotenoid content from the literature are put together in table 1.12. The content of both substances is dependent on culture conditions and species. Some of the variations in table 1.12 are probably also caused by different analyzing techniques. Very low chlorophyll contents must be attributed to a partial destruction during analysis. Low chlorophyll content was observed at low nitrogen and phosphorus concentration of the nutrient solution (REJMANKOVA 1979). The content was 1.2% of the dry weight in solutions with 375 mg N per liter and 154 P per liter and 0.4% if the N and P content was only 1/25 (for L. minor). Low copper concentration (FILBIN and HOUGH 1979) and low boron concentration (EICHORN and AUGSTEN 1974) resulted also in a low chlorophyll content. According to ROMBACH (1976), the chlorophyll and carotenoid content in L. minor rises with increasing light intensity, whereas REJMANKOVA (1979) observed lower chlorophyll content in S. polyrrhiza and L. minor at high photosynthetically active radiation. The chlorophyll content is lower in solutions with sugar than without (in S. polyrrhiza) (D'HARLINGUE et al. 1976). In S. polyrrhiza, it rises with age up to the 15th day of frond growth (GAPONENKO and STAZHETSKII 1969). Also it is higher in normal fronds of W. arhiza (0.9%) than in resting fronds (0.6%) or turions (0.16%) (GODZIEMBA-CZYŻ 1970). According to ANTONIELLI and CAGIOTTI (1976), the chlorophyll content is higher in L. trisulca than in S. polyrrhiza and L. gibba. REJMANKOVA (1979) reports a higher chlorophyll content in S. polyrrhiza (0.5-1.2%) than in L. minor (0.2-0.9%). SINHA and SAHAI (1975) compared the rela-

Table 1.12. Chlorophyll and carotenoid content in Lemnaceae, in % of the dry weight.

* the original data were measured in relation to fresh weight
 + consisting of 0.034 β -carotene and 0.096 lycopene and xanthophyll
 " about 1/4 of it β -carotene and 3/4 mixed xanthophylls
 L.: Lemnaceae, L.m.: L. minor, S.p.: S. polyrrhiza, W.a.: W. arrhiza

authors	species	chlorophyll total	chlorophyll a	chlorophyll b	carotenoid
1	L.			0.006-0.01	
2	W.a.	4-7	0.5		
3	L.m.				
4	L.m.	0.3	1.6	0.25	
5	S.p.			0.27	
6*	L.m.	0.6	1.7	0.4	
7	S.p.	0.2	0.4	0.2	
8	L.m.	0.02			0.13+
9	L.m.				0.08
10	L.	0.2-1.2		0.35-0.63 0.11-0.24	0.15-0.21"
11	W.a.				

- References
- 1 CULLEY and EPPS 1973
 - 2 EICHHORN and AUGSTEN 1974
 - 3 FILBIN and HUGH 1979
 - 4 KAVANAGH 1941
 - 5 LECHEVALLIER 1977b
 - 6 McLAREN and SMITH 1976
 - 7 MONEGER 1968e
 - 8 MUELLER and LAUTNER 1954
 - 9 MUZTAR et al. 1979
 - 10 REJMANKOVA 1979
 - 11 TOT 1962

tive chlorophyll content of 4 Lemnaceae species under given culture conditions. It amounted to 1.3, 1.28, 1.06, and 1, respectively, for S. polyrrhiza, L. aequinoctialis (named as L. minor), W. globosa (named as W. arrhiza), and W. microscopica.

The ratio of chlorophyll a to b varies between 1 and 3.7. It is higher for L. minor (1.7-3.7) than for S. polyrrhiza (1.0-2.2) (REJMANKOVA 1979). Other values are 2.2 for S. polyrrhiza (LECHEVALLIER et al. 1976), 3 for L. minor (ARO 1982), and 4 for W. arrhiza (TOT 1962). The ratio is low in solutions with a low nitrogen level and at high temperatures (37° compared with 22°C) (REJMANKOVA 1979). It increases with the age of the fronds (up to the 15th day) (GAPONENKO and STAZHETSKII 1969). However it does not change very much with changing light intensities (ARO 1982).

The ratio of chlorophylls to carotenoids was determined in L. minor by McLAREN and SMITH (1976) as 6.4. MUZTAR et al. (1979) analysed the carotenoid content of different water plants. For L. minor they noted the following components (in % of the dry weight): Carotene (0.015), xanthophylls (0.056 : 0.003 as monohydroxy pigment, 0.039 as dihydroxy pigment, and 0.014 as polyhydroxy pigment). In comparison to terrestrial plants (Medicago sativa, Zea mays) the total content of carotenoids is 10 times higher in Lemnaceae. BONOMI et al. (1981) noted, in L. minor, β -carotene (0.03), and xanthophyll (0.05). MONEGER (1968a,d) observed, in S. polyrrhiza, the following carotenoids (in % of the dry weight): β -carotene (0.04), lutein (0.04), lutein-epoxide (0.01), violaxanthin (0.01) and neoxanthin (0.003). In L. gibba, the following carotenoids were demonstrated by SIEFERMANN (1971, 1972): β -carotene, lutein, neoxanthin, zeaxanthin, antheraxanthin, and violaxanthin. The carotenoid content is dependent on the light. It increases with light duration after dark growth. With short illumination, the maximum content of carotenoids is formed under green, red, and blue light (MONEGER 1968a,c, MONEGER and JACQUES 1968). SIEFERMANN (1971, 1972) observed more zeaxanthin and, to a lesser extent, more antheraxanthin and less violaxanthin with increasing light intensities. Lutein isolated from L. aequinoctialis revealed an inhibiting effect on BEA induced flowering of the same species (FUJIOKA et al. 1986c).

To investigate possible differences in carotenoid composition of Lemnaceae, Prof. Dr. C.H. Eugster and Dr. R. Buchecker of the Institute of Organic Chemistry, University of Zurich, in 1970 kindly analysed the ca-

rotinoids of 5 Lemnaceae species (*S. polyrrhiza*, *L. gibba*, *L. valdiviana*, *W. hyalina*, *W. brasiliensis*) (unpublished results). By thin-layer chromatography it could be demonstrated, that all species contain the same carotinoids of which the following have been identified: β -carotene, lutein, zeaxanthin (only few), and neochrome. Of the green and grey pigments (chlorophylls and possibly phaeophytins) 9 different substances could be recognized. In the composition of chlorophylls, *L. valdiviana* was distinctly different from *L. gibba*, *W. hyalina*, and *W. brasiliensis*, which all had 4 identical chlorophylls; in contrast, *L. valdiviana* showed 5 different chlorophylls of higher polarity (EUGSTER and BUCHECKER, unpublished). *L. valdiviana* is the only investigated species which is able to live submerged. A comparison with other submerged species would be desirable.

LAM and WALKIN (1985) studied a photosynthetic mutant of *L. aequinoctialis* lacking the cytochrome b_6-f complex but with a similar amount of the light harvesting chlorophyll-protein complex of photosystem II as in the wild-type *Lemna*. Further studies on the light harvesting chlorophyll a-b protein were made by KOHORN et al. (1986) and MAENPAEAE and ARO (1986). THORNEER et al. (1986) detected six different chlorophyll-proteins in *L. gibba* by electrophoretic separation.

Investigations of BIGGS and KOSSUTH (1980) with UV irradiation show that the carotinoids as well as the flavonoid pigments play an important role in photoprotection mechanisms of *S. punctata* and *L. minor* (and other higher plants).

In several heterotrophically cultivated *Spirodela* and *Lemna* species (with etiolated fronds), phytochrome has been demonstrated spectrophotometrically (ROMBACH 1965, ROMBACH and SPRUIT 1968, PORATH and BEN-SHAUL 1973, ROMBACH 1978). After illumination of plants, dark reversion of Pfr (half-life of 2-6 hours) as well as decay of phytochrome have been observed. The phytochrome content of plant parts with and without meristem tissues was determined on a dry weight basis, but did not differ significantly in *L. minor* (ROMBACH and SPRUIT 1968). The phytochromes extracted from etiolated tissues of duckweed do not show any cross-reactions with 6 investigated monoclonal anti-rye phytochrome antibodies and 6 monoclonal anti-pea phytochrome antibodies (SAJI et al. 1984).

1.2.7. Flavonoids

Whereas anthocyanins are often easily visible (*Spirodela*, some species of *Lemna*), other flavonoids can only be recognized by chemical tests or analysis. The anthocyanins which are dissolved in the cell sap are not identical with the brownish red pigment which can be observed as small dots in special pigment cells of dead fronds of *Spirodela*, *Wolffiella*, and *Wolffia*. According to HEGELMAIER (1968) this pigment can be dissolved in boiling potassium carbonate in *Wolffiella* and *Wolffia* but not in *Spirodela*. There must be a chemical difference between the pigments of the *Wolffiaceae* and *Spirodela*, which has never been verified chemically. These pigments also belong to the flavonoid group and develop in *S. punctata* and *W. borealis* after UV irradiation or drying out from phlobaphene-like compounds by oxidation and polymerisation (WITZTUM 1974a). They are probably identical with the flavonols observed by McCURE and ALSTON (1966).

THIMANN and collaborators investigated the anthocyanin of *S. punctata* and the conditions which lead to its production (see chapter 2.5.8.5). They determined the anthocyanin to be petunidin-3-glucoside (NG and THIMANN 1962), but later on they corrected this statement into the 3,5-diglucoside of petunidin (NG et al. 1964). In *S. polyrrhiza* KRAUSE and STRACK (1979a) identified cyanidin 3-monoglucoside (as was already found by McCURE 1964 as the major anthocyanin of *S. polyrrhiza*), and in addition malonylcyanidin 3-monoglucoside. The anthocyanin cyanidin-bioside was found by REZNIK and NEUHAEUSEL (1959) in *L. minor* and *L. trisulca*. In *L. trisulca*, cyanidin-glycoside could be detected in addition. These chemical compounds exist as colourless pseudobases in the living plants and become red if treated with mineral acids. The anthocyanin in *S. polyrrhiza* is zwitterionic by acylation with aliphatic dicarboxylic acids (HARBORNE 1986). In *S. punctata*, JURD et al. (1957) identified the glycoflavones saponarin, isosaponarin, vitexin, and the flavone apigenin. McCURE studied the flavonoids of *Lemnaceae* extensively and published his results in a series of papers (McCURE 1964, 1967a,b, 1968, 1970, 1975; McCURE and ALSTON 1963, 1964a,b, 1966). As an example of a relatively well marked biochemical differentiation within a whole family which is morphologically not very distinctly structured and therefore difficult to handle, the results of McCURE received broad attention and recognition. That is why they will be discussed here in more detail. A

survey of the results is given in table 1.13. In this table, only groups of flavonoids but not single identified flavonoids are stated. However, McCURE and ALSTON (1966) looked upon these single compounds as especially typical for a Lemnacean species. They distinguished 40 identified flavonoids, 7 unidentified flavonoids, and 27 possible flavonoids. The 5 groups of flavonoids distinguished by McCURE and coworkers are: 1) glycoflavones: orientin, vitexin, isosaponarin, homo-orientin, luto-narin, lucenin, vicenin, saponaretin, and 4 acylated glycoflavones; 2) anthocyanins: petunidin-3,5-diglucoside, cyanidin-3-glucoside, and two additional not identified compounds (another anthocyanin, malonylcyanidin-3-glucoside, was identified by KRAUSE and STRACK 1979a); 3) flavonols: 11 quercetin compounds and 4 kaempferol compounds; 4) flavones: 5 apigenin compounds and 4 luteolin compounds.

According to McCURE and ALSTON (1966), the only variation of the flavonoid composition within a species occurred in L. aequinoctialis (named as L. perpusilla) where a single flavonoid was not found in all samples. Later (1967a) McCURE showed that clones from Africa and Asia differed in at least two flavonoids from American clones. Even mutants of the clone no. 6746 show some variability in the presence of a certain flavonoid. A morphological examination of the living clones of McCURE (see table 1.13) showed that the delimitation of species by McCURE was not the same as that of the present author. Especially surprising are his results with the two species L. valdiviana and L. minuscula which are extremely difficult to identify morphologically. However, they look very different in their flavonoid composition. McCURE investigated, according to the present author, 5 clones of L. valdiviana under the name of L. minima and 11 clones of L. minuscula and in addition 2 clones of L. valdiviana under the name of L. valdiviana. Eleven flavonoids were detected in "L. minima" and 4 in "L. valdiviana" (in addition 3 possible flavonoids); the 2 groups had only 1 flavonoid in common. This big difference in flavonoid composition without any intermediate pattern can only be explained if it is assumed that this difference is caused by a

Table 1.13 (p. 37). Flavonoids of Lemnaceae (according to McCURE and ALSTON 1966, McCURE 1970).
The species name of the key clone of McCURE is underlined.

species name according to McCURE	identification by LANDOLT (unpublished)	number of clones	glyco-flavones	antho-cyanins	flavonols	flavones	compounds possible	flavonoids	number of unidentified
<u>S. intermedia</u>	<u>S. intermedia</u>	1	4	2	5	3	1	1	1
<u>S. biperforata</u>	<u>S. biperforata</u>	1	4	2	2	3	1	1	1
<u>S. polyrrhiza</u>	<u>S. polyrrhiza</u>	21	8	4	10	34	12	12	12
<u>S. oligorrhiza</u>	<u>S. oligorrhiza</u>	9	4	4	4	11	5	5	5
<u>L. gibba</u>	<u>L. gibba</u>	12	4	4	4	10	12	12	12
<u>L. minor</u>	<u>L. minor</u>	34	8	5	5	6	34	34	34
<u>L. obscura</u>	<u>L. obscura</u>	6	5	5	5	3	6	6	6
<u>L. trisulca</u>	<u>L. trisulca</u>	5	5	5	5	3	5	5	5
<u>L. perpusilla</u>	<u>L. perpusilla</u>	8	3	3	3	5	8	8	8
<u>L. trinervis</u>	<u>L. trinervis</u>	1	3	3	3	2	1	1	1
<u>L. valdiviana</u>	<u>L. valdiviana</u>	2	4	4	4	11	5	5	5
<u>L. minima</u>	<u>L. minima</u>	11	4	4	4	11	5	5	5
<u>L. oblonga</u>	<u>L. oblonga</u>	5	3	3	3	5	5	5	5
<u>L. lingulata</u>	<u>L. lingulata</u>	5	3	3	3	5	5	5	5
<u>W. oblonga</u>	<u>W. oblonga</u>	3	4	4	4	11	5	5	5
<u>W. oblonga</u>	<u>W. oblonga</u>	2	4	4	4	11	5	5	5
<u>W. gladiata</u>	<u>W. gladiata</u>	1	4	4	4	11	5	5	5
<u>W. microscopica</u>	<u>W. microscopica</u>	1	4	4	4	11	5	5	5
<u>W. papulifera</u>	<u>W. papulifera</u>	6	4	4	4	11	5	5	5
<u>W. borealis</u>	<u>W. borealis</u>	1	4	4	4	11	5	5	5
<u>W. arthiza</u>	<u>W. arthiza</u>	2	4	4	4	11	5	5	5
<u>W. globosa</u>	<u>W. globosa</u>	1	4	4	4	11	5	5	5
<u>W. columbiana</u>	<u>W. columbiana</u>	5	4	4	4	11	5	5	5
<u>Spirodela Lemna Wolffia</u>	<u>Spirodela Lemna Wolffia</u>	32	9	4	12	6	5	5	5
<u>number of clones</u>	<u>number of clones</u>	17	9	4	12	6	5	5	5
<u>number of identified flavonoids</u>	<u>number of identified flavonoids</u>	12	9	4	12	6	5	5	5
<u>number of unidentified compounds</u>	<u>number of unidentified compounds</u>	20	9	4	12	6	5	5	5
<u>number of unidentified compounds</u>	<u>number of unidentified compounds</u>	20	9	4	12	6	5	5	5

single gene. McCURE had identical conditions for the growth of his clones which is of course a precondition for reliable results. To test the influence of different conditions, he studied the formation of flavonoids in 7 clones of S. punctata under 26 different conditions. Out of 8 compounds (including 13 flavonoids) recognized in the chromatograms, only 9 (8 of which flavonoids) could be observed under all conditions (McCURE and ALSTON 1964a). However all compounds except one could be detected if there was sufficient material for investigation (McCURE 1975). BALL et al. (1967) investigated the chromatographical spot patterns of S. polyrrhiza and S. punctata in relation to growth conditions and to different clones. They reported differences between two clones of the same species under identical conditions, as well as within the same clone under different environmental conditions. Especially N and P deficiency and day-length caused deviations of the pattern. The investigations were repeated by H. CLARK (in PARKS et al. 1972). He found many more constituents in his extracts than the former authors reported, thus showing that certain substances might be overlooked depending on the methods used and the amount of plant material investigated. One clone of S. polyrrhiza was studied by McCURE (1968) to identify the formation of flavonoids under different culture conditions. Cyanidin glucoside is only formed under white or blue light but not under red or in the dark. Contrary to anthocyanins, glycoflavones are also produced in the dark, though not in the same quality as in the light. Under optimal light conditions, S. intermedia contains 58 µmol vitexin per 100 g fresh weight, 12 µmol orientin, and 36 µmol cyanidin glucoside; in the dark, the values are 17, 4, and 0, respectively. The cyanidin glucoside production seems to be phytochrome-regulated. In S. polyrrhiza, light-dependent anthocyanin synthesis is most effective under blue and red light. Far red is as only 10% or less of the effectiveness of blue or red light (MANCINI 1977). The author also sees some evidence for an involvement of photosynthesis in anthocyanin synthesis of S. polyrrhiza (compare chapter 2.5.8.5).

In S. polyrrhiza, SAUNDERS and McCURE (1976) observed a quantitatively different flavonoid pattern in chloroplasts compared with whole fronds (much more luteolin-7-glucoside) whereas in S. intermedia no differences showed up. Differences in the flavonoid pattern also occur between normal fronds and turions of S. polyrrhiza (REZNIK and MENSCHICK 1969, ENSCHICK 1970). The flavonoid content was influenced by the nitrogen

source. Turions showed 9 different flavonoids (including one anthocyanin) in addition to the ones mentioned by McCURE and ALSTON (1966). The additional substances are characterized by a higher degree of glycosylation. HORSEL et al. (1972) demonstrated, in L. minor, the existence of a flavonol transforming enzyme. They think it possible that the lacking accumulation of flavonols in L. minor is due to the action of this enzyme. McCURE (1975) mentions some biosynthetic steps leading to the construction of flavonoids in S. intermedia under the influence of external factors. The synthesis of the C₁₅ ring, the closure of the heterocyclic ring, and the 3-hydroxylation are stimulated by a low intensity of red light but proceed also in darkness. Introduction of the 3'-hydroxyl group needs low-energy blue or white light whereas the oxydation of the heterocyclic ring is only performed in high-energy blue or white light. The anthocyanin content of S. intermedia is highest in full-grown fronds whereas the orientin content is highest in still growing fronds. The vitexin content does not change between the two growth stages. Certain factors in the nutrient medium also influence the content of 3',4'-hydroxylated flavonoids (cyanidin-3-glucoside and orientin) but not of the 4'-hydroxylated flavonoid (vitexin) (McCURE 1973, 1975). Kinetin (at 10⁻³ mM) promotes and gibberellin (at 10⁻² mM) inhibits flavonoid accumulation in the light (McCURE 1973).

In contrast to McCURE and ALSTON (1966), VEEN (1975) found variations of the flavonoid pattern in L. gibba and L. minor not only between the species but also between different clones of the same species. The flavonoid pattern was similar but not identical to that of McCURE and ALSTON (1966) for the same species. Also contrary to McCURE, SU et al. (1973a) detected in L. minor flavones as well as flavonols. According to McCURE L. minor contains neither flavones nor flavonols. However, there is a slight possibility that SU et al. confused L. minor with S. punctata which looks similar and has flavones and flavonols according to McCURE. Since SU et al. had their material from Minnesota where S. punctata does not occur and since the material has been checked by taxonomists, the identification can scarcely be doubted. The following conclusions do not consider the results of SU et al. and might have to be revised. The conclusions of the extensive studies on flavonoids of Lemnaceae in respect to evolution and taxonomy of the family are as follows:

1) The genus Spirodela shows the highest differentiation in flavonoid formation. The investigated species contain not only anthocyanins but also 3 additional flavonoid groups.

2) Glycoflavones are evidently typical for the Lemnoideae. In the Wolffioidae they occur very rarely (only in W. arhiza, W. columbiana, and W. globosa).

3) Flavonols are characteristic for species with pigment cells (which are only visible in dead fronds) (cf. WITZTUM 1974a). The only species which has flavonols but no pigment cells in the vegetative fronds is W. microscopica. Wolffia and Wolffiella species have pigment cells in the anthers and on the stigma. Since W. microscopica flowers under the culture conditions applied the presence of flavonols is understandable. It is to be expected that all Wolffia and Wolffiella species contain flavonols when flowering.

4) It does not seem very likely that Wolffia has a biphyletic origin (one group deriving from Lemna and the other from Wolffiella) as was proposed by TURNER (1967) established only by the presence or absence of flavonols.

5) Anthocyanins are restricted to the genus Spirodela and to the most differentiated sections of Lemna (sections Lemna and Hydrophylla). Contrary to the indications of McCURE, anthocyanins do exist in L. minor (REZNIK and NEUHAEUSL 1959, VEEN 1975). S. punctata differs in the anthocyanins from all other species.

6) The genus Wolffiella as far as investigated has no flavonoids other than flavonols, however only a few species have been investigated.

7) It is not yet clear how far the flavonoid pattern of a clone can help in identifying the species within morphologically poorly differentiated groups. More clones of each known species must be investigated. In general, most flavonoids are spread throughout the whole plant kingdom. They can therefore not be used in the taxonomy of higher categories.

Possible physiological significance of flavonoids for Lemnaceae is, according to McCURE (1975): "light absorbance, chelating, and antioxidant properties, ability to hydrogen-bind with proteins, possible participation in electron transport systems, steric mimicry of certain hormones, and frequent antimicrobial effectiveness".

1.2.8. Lignins

According to WOLVERTON and McDONALD (1981), a mixture of Lemna and Spirodela contained 2.7% lignin related to the dry weight. DYKE and SUTTON (1977) measured 1.7% lignin from L. obscura (named as mixture of L. minor and L. gibba). From cell walls of L. minor MUZZAR et al. (1978) analysed 2.4% acid-detergent lignin per wall dry weight. STEUBING et al. (1980) report a much higher content of lignin (17% of the dry weight) in L. minuscula (named as L. valdiviana) from Chile. Since L. minuscula has practically no sclerenchymatic tissue and only very few vascular cells (less than L. minor), it must be concluded that some precursor compound was included in the lignin by the analysis method used. STAFFORD (1964) observed lignin reactions in 15 plant species. He did not get any reaction with L. minor (as well as Elodea) in phloroglucinol or with other histochemical tests. Only trace amounts of a blue colour in the phenol test showed up, but no p-hydroxycinnamic acid or ferulic acid were de-

Table 1.14. Content of lignin components in Lemnaceae, in μmol per 300 g dry weight sample (from BLAZEY and McCURE 1968)

species	name according to BLAZEY and McCURE 1968	p-hydroxy-benzaldehyde	vanillin	syringaldehyde
<u>S. intermedia</u>	<u>S. intermedia</u>	9.1	4.1	4.9
<u>S. polyrrhiza</u>	<u>S. polyrrhiza</u>	6.6	1.7	1.1
	<u>S. biperforata</u>	5.3	0.1	0.0
<u>S. punctata</u>	<u>S. oligorrhiza</u>	2.8	1.0	0.6
<u>L. gibba</u>	<u>L. gibba</u>	3.0	0.0	0.0
<u>L. minor</u>	<u>L. minor</u>	3.0	0.2	0.0
<u>L. obscura</u>	<u>L. obscura</u>	4.5	0.0	0.0
<u>L. trisulca</u>	<u>L. trisulca</u>	4.3	0.0	0.0
<u>L. acquinoctialis</u>	<u>L. perpusilla</u>	5.5	1.0	0.0
	<u>L. trinervis</u>	3.3	0.4	0.0
<u>L. valdiviana</u>	<u>L. valdiviana</u>	6.8	0.0	0.0
	<u>L. minuscula</u>	4.9	0.1	0.0
<u>W. oblonga</u>	<u>W. oblonga</u>	4.2	0.0	0.0
<u>W. microscopica</u>	<u>W. microscopica</u>	6.1	2.1	0.0
<u>Hordeum vulgare</u>		10.5	11.7	9.7
<u>Elodea densa</u>		4.3	0.0	0.0

pectable in the acidified ether extracts. The lignification of 11 Lemna-caea species (1 clone of each species) was investigated by BLAZEY and McCURE (1968). Beside p-hydroxybenzaldehyde they found vanillin and syringaldehyde (see table 1.14). Vanillin and syringaldehyde are supposed to originate from lignin, however p-hydroxybenzaldehyde is known from fractions which do not originate from lignin. As is recognizable from table 1.14, there are 3 species (S. polyrhiza, L. aequinoctialis, and L. valdiviana) which are represented with 2 clones. A certain variability shows up between the 2 different clones, but, in general, the differences between different species are more significant. The quantitative and qualitative formation of lignins probably also varies with the growth conditions. The results of BLAZEY and McCURE (1968) show that the lignification is highest in Spirodela which is the only genus containing syringaldehyde. This is not surprising because Spirodela develops more vascular cells than Lemna and the Wolffioideae. Vanillin is also present in some species of Lemna. The relatively high content of vanillin in W. microscopica must be attributed to the flowering status of the investigated material. It is known that W. microscopica flowers regularly under culture conditions and that the flower organs are the only parts of the Wolffioideae which contain vascular cells. Furthermore, BLAZEY and McCURE (1968) found the anther walls of W. microscopica rich in sclereids which can be stained by the phloroglucinol-HCl technique. It is surprising that L. gibba, L. obscura, and L. trisulca with noticeable vascular cell rows in the nerves do not contain vanillin. This suggests that p-hydroxybenzaldehyde originates at least partly from lignin. Anyway, it has to be considered that probably p-hydroxybenzaldehyde is, especially in monocotyledons, beside vanillin and syringaldehyde a decomposition product of lignin since in this plant group p-coumaryl alcohol (beside coniferyl alcohol and sinapyl alcohol) is incorporated into lignin to a relatively high degree.

1.2.9. Oxalates and other organic anions

L. minor contains up to 4% (of the dry weight) oxalic acid which is precipitated to 80% as raphide crystals and stored in special cells (idioblasts) (BORNKAMM 1965). In Spirodela, oxalates are present in form of raphides and druses besides the free oxalate content. Oxalates remains

dissolved in the cell sap within the Wolffioideae, where it never forms crystals. Evidently, Lemnaceae do not produce oxalate oxidases; Spirodela and Lemna are dependent on calcium to neutralize accumulations of oxalic acid (LOETSCH and KINZEL 1971). The amount of oxalate is correlated with the protein content.

Low temperatures favour oxalate formation. In L. minor cultivated at 20°C BORNKAMM (1970c) measured 2.3% oxalate, at 25°C 2.4%, and at 30°C 0.8%. Dark-grown L. minor forms 4 times as many crystal cells as do light grown plants (FRANCESCHI 1985). The oxalate is formed by glycolate oxidase (BORNKAMM 1965, 1969a, FRANCESCHI 1985).

WROBLEWSKI (1976) investigated the chemical composition of raphides of L. minor by means of analytical electron microscopy. He distinguishes between two kinds of crystals (table 1.15): smaller ones with hexagonal or rectangular sectional view, and bigger ones with an hourglass-like sectional view. This is the only report on two different forms of crystals in L. minor. Maybe the small crystals are young stages of a raphide. It is interesting to note that there is not only a quantitative difference between the two different crystal forms but also a difference in chemical composition (table 1.15). AL-RAIS et al. (1971) measured up to 2% Mg in the raphides of Lemnaceae. The oxalate of raphides is present in the monohydrate form, the oxalate of druses (not measured in Lemnaceae) contains calcium oxalate x 2.25 H₂O. According to WATTENDORF (1985) the crystalline Ca oxalate occurs in Lemna as the stable monohydrate (whewellite) as well as the dihydrate. As a rule, the dihydrate does not show up in crystal idioblasts, but is found dissolved in all living cells. L. minor is able to incorporate Sr (but not Ba, Cd, Co, and Mn) into the raphides (FRANCESCHI and SCHUEREN 1986).

Table 1.15. Relative chemical composition of raphide idioblasts in L. minor (from WROBLEWSKI 1976)

Crystals	Chemical elements			
	Mg	S	Cl	Ca
big form	0.41	1.34	-	41000.0
small form	-	1.94	0.84	40.1

The presence of malate in Lemnaceae was demonstrated by KOPP et al. (1974a), KANDELER and HELDWEIN (1979), HELDWEIN and KANDELER (1981) and many other authors. The accumulation of malate only proceeds during photosynthesis and seems to be connected with nitrate metabolism. The malate content of L. aequinoctialis is about 10 times higher in the light (0.003 mmol per g fresh weight) than in darkness (KANDELER and HELDWEIN 1979). JANAUER (1982) identified, from L. trisulca, 0.1 mval organic anions per g dry weight. The following anions could be demonstrated in the sequence of decreasing content: fumarate, malate, glycolate, and citrate.

1.2.10. Vitamins and plant growth substances

CLARK and THOMAS (1934) and CLARK et al. (1938) successfully fed rats nourished with a vitamin-free food with S. polyrrhiza. In this way they demonstrated marked amounts of vitamins A, B₁, and C within S. polyrrhiza. Sixty-six to 110 mg per kg fresh weight carotene, an indicative of vitamin A, was determined by CULLEY and EPPS (1973) in Lemnaceae. FRAHM (1938) proved that S. polyrrhiza contains vitamin B₁. The presence of vitamin C in L. minor was corroborated by RAKHIMOV and RAKHIMOVA (1975). NAKAMURA (1960) noted the vitamins A, B₂, B₆, and C in W. globosa (named as W. arrhiza). MUZAFFAROV et al. (1971) isolated the vitamins B₁, B₂, B₆, E, and PP from L. minor. Especially the content of vitamin E (0.02-0.04 mg per g fresh weight) and of vitamin PP (0.04-0.06 mg per g fresh weight) is remarkable. Nicotinic acid was isolated by FUJIOKA et al. (1986a) from flowering plants of L. gibba and L. aequinoctialis. It is supposed that nicotinic acid influences the flowering process by interacting with other plant hormones (FUJIOKA et al. 1986b).

THIMANN and SKOOG (1940) achieved auxin-like effects on Avena coleoptiles by applying ether extracts of S. punctata (named as L. minor). In a further publication, THIMANN et al. (1942) demonstrated that the yield of auxins from S. punctata is greatly increased by a short incubation with chymotrypsin or trypsin; ficin has a similar though smaller effect. Therefore, a major part of the auxin seems to be bound to protein or a protein-like substance in S. punctata. SARGENT (1957) isolated four growth-stimulating substances and one growth-inhibiting substance from L. minor. The mainly active substance was identified as IAA. According

to SLOVIN and COHEN (1985b), the IAA level in Lemnaceae is unusually high. They measured, in L. gibba, per kg fresh weight 6.4 µg free IAA, 51.5 µg esterified IAA and 9.2 µg amide IAA. WITZTUM et al. (1978) measured up to 2.2 mg IAA per kg fresh weight in S. punctata; the addition of sucrose enhances the IAA content in S. punctata, UV irradiation lowers it (WITZTUM et al. 1978). The endogenous IAA level of L. gibba was also measured by COHEN et al. (1986). It is difficult to explain the great difference in IAA content between the results of SLOVIN and COHEN (1985b) and WITZTUM et al. (1978).

PIETERSE et al. (1971a) and PIETERSE (1972) observed two gibberellin-like substances in several species of Lemnaceae. Preliminary determinations of the gibberellin content in L. aequinoctialis and L. gibba cultivated under identical conditions showed higher values in the short-day plant L. aequinoctialis than in the long-day plant L. gibba (LADENBURGER, BAUER and KANDELER, unpubl. results). In L. aequinoctialis, substances were found with auxin-like and gibberellin-like activities as well as neutral and acidic inhibitors. The endogenous level of gibberellins increases during the night if the plants are treated with a short far-red irradiation at the end of the day. Supposedly, the content of gibberellins is dependent on the active form of phytochrome (PEKIC and NESKOVIC 1982). BEZEMER-SUBRANDY and VELDSTRA (1971a) analyzed cytokinins in alkaline hydrolysates of t-RNA of L. minor. In n-butanol extracts of W. arrhiza some cytokinin-like compounds can be separated by thin-layer chromatography and quantitated by the Amaranthus bioassay (EICHHORN and AUGSTEN 1980, EICHHORN 1986). The fraction which has the same R_f value as zeatin dominated in all investigated extracts. The authors found 98-244 µg zeatin equivalent per kg fresh weight. Cultivation of plants under blue light increased the cytokinin level (in comparison to white light) if the population was growing fast (good supply of phosphate). Red light on the other hand raised the content of the zeatin-like compound if the plants were growing slowly (low supply of phosphate). Another factor which modifies the cytokinin content in Lemnaceae is nitrate (LOEPPERT, GRUNTZEL and KANDELER, unpubl. results). Omission of nitrate from the medium lowers the level of cytokinins in L. aequinoctialis within 24-48 hours (GRUNTZEL 1982).

The levels of ABA which have been determined in Lemnaceae differ by several orders of magnitude. WITZTUM and KEREN (1978a,b) found 2-14x10⁻⁴ g ABA per g fresh weight in S. polyrrhiza, HUBER and SANKHLA (1979) 5x

3-6 g ABA per g fresh weight in L. minor, and MERTEN (1979) 2-9x10⁻⁷ g ABA per g fresh weight in winter material of L. gibba and L. aequinoctialis, and 4-42x10⁻⁹ g ABA per g fresh weight in summer material of the same species. Accumulation of ABA within the fronds or release to the surrounding medium may be one of the causes for the differing results. Release of ABA to the nutrient medium was observed by MERTEN (1979) and UKS et al. (1975, 1980). HUBER (1985) detected an increasing amount of xanthoxin and a decrease of ABA in pentachlorophenol-treated L. minor. The formation of xanthoxin, which is supposed to be a stress indicator, must be coupled with a membrane damage of the bleached fronds permitting the release of unsaturated fatty acids and herewith the peroxidation of xanthophylls (e.g. violaxanthin).

NUJOKA et al. (1983a, 1985) established the presence of benzoic acid (BEA) in L. aequinoctialis and L. gibba (30-44 µg per kg fresh weight). A positive correlation was found between the endogenous level of BEA and flowering of Lemna. BEA has also been demonstrated by TAKIMOTO and KAKAHASHI (cited from CLELAND and BEN-TAL 1983).

WATTOO et al. (1983, 1984d, 1986a) measured less than 0.05 ppm ethylene in S. punctata when cultivated in Hutners solution. If 0.02 mM Cu is added to the culture solution, the production of ethylene is 15- to 30-fold. The authors think that the stimulation of ethylene production in (II)-supplemented solutions is caused by O₂ mediated damage of cellular membranes. FAERBER (1984), SCHARFETTER et al. (1984) report on the ethylene production of L. gibba. FAERBER et al. (1986) and KANDELER and FAERBER (1987) observed ethylene production under uncrowded and crowded conditions. Under uncrowded conditions, the production of ethylene amounted to 0.25-0.5 nl per g fresh weight and hour in L. gibba and L. minor, and 1.5-2.0 nl per g fresh weight and hour in S. polyrrhiza. Under crowded conditions it increased slowly in L. gibba and immediately in S. polyrrhiza and L. aequinoctialis. The addition of ABA did not result in a higher ethylene production. The production and release of ethylene in L. minor from aminocyclopropane-1-carboxylic acid (ACC) was studied by FUHRER (1985).

OVANELLI et al. (1981) demonstrated spermidine in L. aequinoctialis, and FLORES and GALSTON (1982) found a great amount of agmatine and spermidine in vegetative and flowering fronds of L. gibba. ROTH et al. (1985) and COLLIER et al. (1986) isolated 3.0 ng per g dry weight of a somatostatin-like substance from L. gibba. Somatostatin

is a hormone known from hypothalamus of vertebrates. The role of the hormone in Lemna as well as in bacteria and in spinach where it has also been found is not known. The authors conclude that vertebrate-type peptide hormones which are spread through many organisms have early evolutionary origins. Similarly, an insulin-related substance has been found in Lemna gibba G3 (and also in spinach) by COLLIER et al. (1987). The substance stimulates glucose oxidation and lipogenesis in isolated adipocytes from young rats. The role of this insulin-like compound in plants is unknown.

1.2.11. Enzymes

Occurrence, properties, and regulation of more than hundred enzymes have been established within members of the Lemnaceae (table 1.16). The known enzymes are listed in alphabetical order with indications of the investigated species and references (as numbers). The enzyme content is dependent on the culture conditions. APPENROTH et al. (1986) showed that the light quality affects the activity of alanine aminotransferase and aspartate aminotransferase in W. arifolia. Blue light stimulates the activity of both enzymes and red light decreases it. The significance of the enzymes for metabolism and developmental processes will be discussed in chapter 2.5. PERRY (1963) showed that in S. polyrrhiza many enzymes in normal fronds are essentially absent from turions.

GILES (1977-1979, unpublished results) examined 11 Lemnaceae species for 7 enzyme systems and studied the banding pattern on different gels: general protein (GP), tetrazolium oxidase (TO), malate dehydrogenase (MDH), glutamate dehydrogenase (GDH), alcohol dehydrogenase (ADH), peroxidase (PER), and glutamate oxaloacetic transaminase (GOT). In the staining reactions (see table 1.17) and in the banding pattern (fig. 1.3), genera and species as well as different clones of the same species showed differences. There are probably biochemical differences in the structure of the enzymes in different groups resulting in different response to the treatments. The lack of reaction of Wolffia species might be due partly to relatively unfavourable conditions during culture and partly to the submerged life form. Unfortunately, GILES did not follow up the investigations. The following preliminary conclusions might be drawn from her results (see p. 55):

Table 1.16. Enzymes determined in members of Lemnaceae
(for references see at the end of the table.)

name of enzyme	test species	references
o-acetylserine sulphydrylase	L. aequinoctialis	137
adenosine 5'-phosphosulfate sulfotransferase (APSSase)	L. minor	10a,b,11,12a, 114,133b,150, 151
adenosine 5'-triphosphate-ase (ATPase)	S. polyrhiza	12b
ATP-phosphohydrolase	S. punctata	50
ATP-sulphydrylase	L. minor	133b
alanine aminotransferase	W. arrhiza	66
	L. minor	12a,117,133c, 30b,155
	L. gibba	59
alanine dehydrogenase	L. minor	66
alanine glyoxylate aminotransferase	W. arrhiza	63
	L. minor	108
alcohol dehydrogenase	L. gibba	39
	3 Wolffia spp.	1b,39
aldolase	L. minor	20,60,111,112
(di) aminopimelate decarboxylase	S. punctata	149
β -amylase	L. aequinoctialis	122
	S. polyrhiza	51
	L. gibba	142
	L. minor	46a
D-apsosyltransferase	L. minor	89a,100,101,102
ascorbic acid oxidase	L. minor	52,53
aspartate aminotransferase	W. arrhiza	1b,66
aspartokinase	L. minor	157
carbonic anhydrase	L. minor	153
	L. trisulca	153
	W. columbiana	153
catalase	L. minor	38,52,53
cellulase	L. minor	120
β -cyanoalanine synthase	L. trisulca	140
cystathionine δ -synthase	L. aequinoctialis	40,41,136-139
cystein synthase	L. minor	10b
cytochrome oxidase	Lemna sp.	16
flavonol-converting enzyme	L. minor	54
β -D-fructofuranosidase (invertase)	W. arrhiza	66
D-fructose-6-phosphate 1-phosphotransferase	"duckweed"	109
D-galacturonosyl transferase	L. minor	83
glucose-6-phosphate dehydrogenase	L. gibba	91
	L. minor	60,61
	L. aequinoctialis	103
glutamate dehydrogenase (see under NAD)	W. arrhiza	28,29,30a

Table 1.16. (continued)

name of enzyme	test species	references
glutamate glyoxylate aminotransferase	L. minor	108
glutamate oxaloacetic transaminase	S. intermedia	39
	S. punctata	39
	L. gibba	39
	L. minor	39
	L. valdiviana	39
	W. hyalina	39
	L. minor	108
glutamate pyruvate aminotransferase	L. minor	56,105,107
glutamate synthase	L. minor	56,104-107
glutamine synthetase	L. gibba	30b
glutathion peroxidase		
glyceraldehyde 3-phosphate dehydrogenase (see under NADP)		
glycolate oxidase	L. minor	9,32,33,33a,38 52,53
α -glycolyldase, β -glycolyldase	L. trisulca	9
4-hydroxyphenylpyruvate dioxygenase	W. arrhiza	66
	L. gibba	86,92
hydroxypyruvate reductase	L. minor	1a,33a
invertase	L. gibba	144
isocitric dehydrogenase (see under NADP)		
malate dehydrogenase (see under NADP)		
malic enzyme	L. minor	57,60
mercaptopyruvate sulfotransferase	L. minor	115
myo-inositol kinase	L. gibba	7
myo-inositol-1-phosphate kinase	L. gibba	7
myo-inositol-1-phosphate synthase	L. gibba	96,119
myo-inositol-triphosphate kinase	L. aequinoctialis	87
NAD (and NADP) glutamate dehydrogenase	L. gibba	39,121
	L. minor	26,27a,39,48, 59,60-65,105, 113,130,135 23,24,25
	L. aequinoctialis	39
	L. valdiviana	39
	S. intermedia	39
	S. punctata	39
	W. hyalina	39
	3 Wolffia spp.	39
NADH cytochrome-c-reductase	S. punctata	27b
NADP glyceraldehyde 3-phosphate dehydrogenase	L. gibba	44,45,46,91,93, 116,158
	L. minor	57

Table 1.16. (continued)

name of enzyme	test species	references
NADP isocitric dehydrogenase	L. gibba	91
NADP malate dehydrogenase	L. minor	60, 61
	S. polyrrhiza	18, 19
	L. minor	58-61
	11 spp.	39
NAD and NADP nitrate reductase	S. polyrrhiza	131
	S. punctata	34, 35, 36a, 148
	L. gibba	2
	L. minor	56, 60, 62, 97, 98,
		99, 123, 127, 128,
		129, 132
	L. trisulca	89b
NADP 6-phosphogluconate dehydrogenase	L. aequinoctialis	21, 132, 135
nitrite reductase	L. gibba	91
S. nuclease	S. punctata	34
pectinase	L. trisulca	89b
peroxidase	L. gibba	67
	L. minor	120
	L. minor	125, 126
	11 spp.	39
phenylalanine ammonia-lyase	S. polyrrhiza	42, 43
	S. punctata	81b
	L. gibba	42, 43
	L. aequinoctialis	40, 43
phosphatases (acid)	S. punctata	5, 72-81a
	L. gibba	91
	L. minor	133a
phosphatases (alkaline)	S. punctata	5, 75-80
	L. minor	85
	W. arrhiza	65
phosphodiesterase	S. punctata	78
phosphoenolpyruvate carboxylase	L. minor	57, 60, 88
phosphofructokinase	L. minor	60
o-phosphohomoserine sulphydrylase	L. aequinoctialis	137
o-phosphoric acid-monoester phosphohydrolase (acid and alkaline)	W. arrhiza	66
phosphorylase	L. gibba	142
polyphenolase	L. minor	52, 53
protein kinases	L. minor	71, 143
pyrophosphatase (alkaline)	L. aequinoctialis	13, 68-70, 152
pyrophosphate phosphohydrolase	L. minor	133a
pyruvic kinase	W. arrhiza	65
ribonucleases	L. gibba	94
RNA polymerase	S. punctata	73-81
	L. gibba	94

Table 1.16. (continued)

name of enzyme	test species	references
ribulose-1,5-biphosphate carboxylase	S. punctata	50, 133c
	L. gibba	37, 124, 133c,
		141, 156
	L. minor	36b, 57
	L. valdiviana	133c
	11 spp.	14
ribulose-5-phosphate kinase	L. minor	57
serine glyoxylate aminotransferase	L. minor	108
sulphydrylase	L. aequinoctialis	17
superoxide dismutase	L. gibba	145
tetrazolium oxidase	9 spp.	39
threonine synthase	L. minor	118
	L. aequinoctialis	41
thylakoid superoxide dismutase	S. punctata	84
trehalase	L. aequinoctialis	147a
tyrosine ammonia-lyase	S. polyrrhiza	43
UDP apiose / UDP-xylose synthase	L. aequinoctialis	43
UDP D-glucuronic acid cyclase	L. minor	47, 90, 110, 154
UDP D-glucuronic acid	L. minor	47
carboxy-lyase	L. minor	47
urease	S. punctata	6, 15
uridine kinase	L. gibba	95
D-xylosyltransferase	L. minor	102, 110, 154
enzyme system producing C ₆ -aldehydes (from unsaturated fatty acids)	S. polyrrhiza	49

References

- 1a ANDERSON and EMES 1985)
 1b APPENROTH et al. 1986)
 2 AUGSTEN and HUBALD 1980)
 3 BAKSHI et al. 1978)
 4 BAKSHI et al. 1979)
 5 BIELESKI 1974)
 6 BOLLARD et al. 1968
 7 BOLLMANN et al. 1980
 8 BORNKAMM 1965
 9 BORNKAMM 1969a
 10a BRUNOLD and SCHMIDT 1976
 10b BRUNOLD and SCHMIDT 1978
 11 BRUNOLD and SUTER 1983
 12a BRUNOLD and SUTER 1984b
 12b BUCZEC and SULEY 1986
 13 CHAPMAN et al. 1975
 14 CHEN and WILDMAN 1981
 15 COOK 1968
 16 DALY and BROWN 1954
 17 DATKO et al. 1977
 18 DAVIDSON and SIMON 1981a
 19 DAVIDSON and SIMON 1983
 20 DAVIES 1978
 21 DEVI and MAHESHWARI 1979
 22 DROTAR et al. 1985
 23 DUKE 1975
 24 DUKE et al. 1975
 25 DUKE and KOUKARI 1977
 26 EHMKE and HARTMANN 1976
 27a EHMKE and HARTMANN 1978
 27b EICHENBERGER 1975
 28 EICHHORN 1983
 29 EICHHORN and AUGSTEN 1977a
 30a EICHHORN and AUGSTEN 1977b

Table 1.16 (continued) References

- 30b ELLIS 1969
 31 EMES and ERISMANN 1981
 32 EMES and ERISMANN 1982
 33 EMES and ERISMANN 1984
 33a ERISMANN et al. 1986
 34 FERGUSON 1969a
 35 FERGUSON 1970
 36a FERGUSON and KNYPL 1974
 36b FERREIRA and DAVIES 1987a,b
 37 FLORES and TOBIN 1985
 38 FREDERICK et al. 1973
 39 GILES unpubl. results
 40 GIOVANELLI et al. 1984
 41 GIOVANELLI et al. 1986
 42 GORDON 1977
 43 GORDON and KOUKKARI 1978
 44 GOTO 1978
 45 GOTO 1979a
 46 GOTO 1979b
 46a GUNIA 1987
 47 GUSTINE et al. 1975
 48 HARTMANN and EHMKE 1976
 49 HATANAKA et al. 1978
 50 DE HEIJ et al. 1984
 51 HENSEN 1954
 52 HILLMAN 1954
 53 HILLMAN 1955
 54 HOESL et al. 1972
 55 HUBALD and AUGSTEN 1979
 56 HUBER 1982
 57 HUBER and SANKHLA 1979
 58 HUBER and SANKHLA 1981
 59 HUBER et al. 1982
 60 HUMPHREY et al. 1977
 61 JEFFERIES et al. 1969a,b
 62 JOY 1968
 63 JOY 1969b
 64 JOY 1971
 65 JUNGnickEL and VOELKSCH 1979
 66 JUNGnickEL et al. 1982
 67 KARLIN-NEUMANN et al. 1985
 68 KATO et al. 1983
 69 KATO et al. 1984
 70 KATO and FUJII 1985
 71 KEATES and TREWAWAS 1974
 72 KNYPL 1976
 73 KNYPL 1977a
 74 KNYPL 1977b
 75 KNYPL 1978
 76 KNYPL 1979a
 77 KNYPL 1979b
 78 KNYPL 1982
 79 KNYPL and JANAS 1980
 80 KNYPL and KABZINSKA 1977
- 81a KNYPL and SOBOLEWSKA 1978
 81b KNYPL et al. 1986
 82 LASSOCINSKI 1982
 83 LEINBACH 1975
 84 LEWINSOHN and GRESSEL 1984b
 85 LIEDTKE and OHMANN 1969
 86 LOEFFELHARDT and KINDL 1979
 87 LOEWUS and LOEWUS 1971
 88 MARQUES et al. 1985
 89a MASCARO and KINDEL 1977
 89b MELZER and EXLER 1982
 90 MENDICINO and ABOU-ISSA 1974
 91 MIYATA and YAMOMOTO 1969
 92 MUEHL and LOEFFELHARDT 1982
 93 MUELLER and ZIEGLER 1969
 94 NAKASHIMA 1979a
 95 NAKASHIMA and TSUDZUKI 1976
 96 OGUNYEMI et al. 1978
 97 OREBAMJO and STEWART 1974
 98 OREBAMJO and STEWART 1975a
 99 OREBAMJO and STEWART 1975b
 101 PAN 1974
 102 PAN and KINDEL 1977
 103 POSNER 1971
 104 RHODES et al. 1975
 105 RHODES et al. 1976
 106 RHODES et al. 1979
 107 RHODES et al. 1980
 108 RUTTER and ERISMANN 1985
 109 SABULARSE and ANDERSON 1982
 110 SANDERMANN and GRISEBACH 1970
 111 SARAWEK and DAVIES 1977a
 112 SARAWEK and DAVIES 1977b
 113 SCHEID et al. 1980
 114 SCHMIDT 1975
 115 SCHMIDT 1984
 116 SCHMIDT-CLAUSEN et al. 1969
 117 SCHMUTZ and BRUNOLD 1984
 118 SCHNYDER et al. 1975
 119 SCHWARCZ et al. 1974
 120 SCHWEBEL 1973
 121 SHEPHARD and THURMAN 1973
 122 SHIMURA and VOGEL 1966
 123 SIMS et al. 1968
 124 SLOVIN and TOBIN 1981
 125 SPILLER et al. 1973
 126 STAFFORD 1964
 127 STEWART 1968
 128 STEWART 1972a
 129 STEWART 1972b
 130 STEWART and RHODES 1977a,b
 131 STEWART and SMITH 1972
 132 STEWART et al. 1974
 133a STROTHER 1984

Table 1.16 (continued) References

- 133b SUTER et al. 1986
 133c TAKEMOTO and NOBLE 1986
 134 TANAKA et al. 1986
 135 TEIXEIRA and DAVIES 1974
 136 THOMPSON et al. 1981
 137 THOMPSON et al. 1982a
 138 THOMPSON et al. 1982b
 139 THOMPSON et al. 1983
 140 TIMOFEEVA and KRAEVA 1983
 141 TOBIN and SUTTIE 1980
 142 TREICHEL 1974b
 143 TREWAWAS 1973
 144 ULLRICH-EBERIUS et al. 1978
 145 VAUGHAN et al. 1982
- 146 VAUGHAN et al. 1983
 147a VELUTHAMBI et al. 1981
 147b VELUTHAMBI et al. 1983
 148 VIJAYARAGHAWAN et al. 1982
 149 VOGEL and HIROVONEN 1971
 150 VON ARB and BRUNOLD 1980
 151 VON ARB et al. 1985
 152 VAN LOON et al. 1975
 153 WEAVER and WETZEL 1980
 154 WELLMANN and GRISEBACH 1971
 155 WILSON and KANG 1973
 156 WIMPEE et al. 1983
 157 WONG and DENNIS 1973
 158 ZIEGLER et al. 1969

Table 1.17. Enzyme reactions to 7 different stainings of 11 Lemnaceae species (mostly 2 clones per species) (according to GILES, unpubl. results)

ADH alcohol dehydrogenase
 GDH glutamate dehydrogenase
 GOT glutamate oxaloacetic transaminase
 GP general protein
 MDH malate dehydrogenase
 PER peroxidase
 TO tetrazolium oxidase

++ strong reaction
 + medium reaction
 (+) weak reaction
 - no reaction

Species	Enzymes						
	GP	TO	MDH	GDH	ADH	PER	GOT
<i>S. intermedia</i>	++	++	(+)	+	-	+	+
<i>S. punctata</i>	++	++	+	+	-	+	+
<i>L. gibba</i>	++	++	+	+	(+)	+	+
<i>L. minor</i>	++	++	+	+	(+)	+	+
<i>L. valdiviana</i>	++	++	+	+	-	+	+
<i>W. hyalina</i>	++	++	+	+	-	-	+
<i>W. neotropica</i>	++	-	(+)	+	-	-	-
<i>W. gladiata</i>	++	-	(+)	-	-	-	-
<i>W. australiana</i>	++	++	+	+	+	+	-
<i>W. arrhiza</i>	++	++	+	+	+	+	-
<i>W. globosa</i>	++	++	+	+	+	+	-

	GP	MDH	GDH	ADH	PER	GOT
Spirodela	7178 7820 6725 7132 7613 6861	7376 7555 7225 7250	7590 7452 7211 7267	7014 7347 8272 7234		
intermedia						
punctata						
Lemna						
gibba						
minor						
Lemna						
valdiviana						
Wolffella						
hyalina						
neotropica						
Wolffella						
gladiata						
Wolffella						
australiana						
Wolffella						
arrhiza						
globosa						

Fig. 1.3. Banding patterns for six different enzyme systems of eleven Lemnaceae species (GILES, unpubl. results)

- 1) The clones of a species mostly behave rather similarly. The results of the two clones each of *S. punctata* and *W. hyalina* are remarkably similar. On the other hand, two clones of *L. valdiviana* differed for three enzymatic systems. In general, differences between different species are bigger than between the clones of one species.
- 2) The bands of MDH and of PER are best suited to show species specific differences. Differences between the genera are not very pronounced except for the bands of MDH. In this enzyme system, all genera may be characterized by a special band pattern (only the two submerged species of *Wolffiella* do not show any banding).
An extension of the investigations to more clones, more species, and more growth conditions is desirable.

CHEN and WILDMAN (1981) electrofocused fraction-1-protein of 11 species of Lemnaceae (*S. polyrrhiza*, *S. punctata*, *L. gibba*, *L. minor*, *L. aequinoctialis*, *W. neotropica*, *W. brasiliensis*, *W. borealis*, *W. australiana*, *W. arrhiza*, *W. globosa*). P-1-protein is the enzyme (ribulose-1,5-biphosphate carboxylase-oxygenase) in green plants which catalyzes the fixation of CO₂ during photosynthesis. CHEN and WILDMAN found, within the family of Lemnaceae, 4 types of large subunit polypeptid clusters within

Table 1.18. Polypeptid composition of large (LS) and small (SS) subunits of Fraction-1-protein for different species of Lemnaceae (after CHEN and WILDMAN 1981)

species	no.	LS				SS							
		I	II	III	IV	1	2	3	4	5	6	7	8
<i>S. polyrrhiza</i>	7003												
<i>S. punctata</i>	7606		x			x							
<i>L. gibba</i>	-			x						x			
<i>L. minor</i>	60				x								x
<i>L. aequinoctialis</i>	6746												
<i>W. neotropica</i>	7225		x			x				x		x	
<i>W. brasiliensis</i>	7303			x							x		
<i>W. borealis</i>	7566		x								x		
<i>W. australiana</i>	7211												
<i>W. arrhiza</i>	7193		x										
<i>W. globosa</i>	7243		x			x							

he enzyme macromolecule and 8 individual small subunit polypeptides of different isoelectric point (table 1.18). Each species can be characterized by a specific combination. Taxonomically related species do not always have a similar pattern. Since only one clone of a species has been investigated, it is not known if a given combination is typical for a species or only for a clone.

...2.12. Other organic substances

U et al. (1973a) analyzed, in a Lemnaceae species (called L. minor), annins and saponins as well as traces of alkaloids and steroids. In contrast, McCURE (1970) found neither alkaloids nor terpenes in Lemnaea.

Cardiac glycoside-like substances (belonging to the steroids) were tated to occur in L. aequinoctialis (named as L. minor) (YONG and THO 976). Acetylcholine was identified from L. gibba extracts by paper chromatography, pharmacological activity on frog muscle, and sensitivity to acetylcholinesterase (HOSHINO and OOTA 1978).

n root cells of a Lemnaceae species (called L. minor) WROBLEWSKI (1976) observed amorphous inclusions which he interprets as tannin idioblasts. annins are known from the brownish red pigment cells of Spirodela and certain species of Wolffiella and Wolffia. In the living fronds, the colourless content of these cells consists of a phlobaphene-like compound (WITZTUM 1974a; see also chapter 1.2.7). It is not quite clear if the tannin idioblasts are identical with the pigment cells or if they also occur outside these cells and even in species (e.g. genus Lemna)

ble 1.19. Relative chemical composition of tannin idioblasts of L. minor (from WROBLEWSKI 1976)

	Mg	S	Cl	Ca
Big granules within membrane limited vacuoles	0.7	3.6	1.1	10.4
Medium coarse material within vacuoles	0.0	0.8	1.2	3.4
Finely granular material within vacuoles	0.8	1.0	1.0	4.3

which do not contain pigment cells. A confusion between L. minor and S. punctata cannot be excluded. From the vacuoles of turions of S. polyrrhiza VINTEJOUX (1978) isolated tannin-like polyphenolic substances associated with proteic and polysaccharidic compounds. Other polyphenolic substances are located in intercellular spaces, cell walls, zones of the cytoplasm, and along the plasmalemma and the tonoplast. WROBLEWSKI (1976) analyzed the chemical composition of tannin idioblasts. He detected the following elements (table 1.19).

According to REZNIK and NEUHAEUSEL (1959), L. minor and L. trisulca contain caffeic acid, L. trisulca also ferulic acid. In S. polyrrhiza KRAUSE (1978) analyzed the following derivatives of cinnamic acids: 1-feruloylglucose, 1-sinapoylglucose, 5-p-coumaroylquinic acid and 5-caffeoylquinic acid. 4 known isoprenoids (lycopersene, sitosterol, phytol, and trans-1,3-phytadiene) and a novel diterpene ([4R]-4-hydroxyisophytol) have been detected in L. minor by PREVITERA and MONACO (1984) (see chapter 2.5.8.2).

HCN tests (Picrin test and Feigl's test) were made at the Geobotanical Institute in Zürich with clones of S. intermedia (1 clone), S. punctata (2 clones), L. ecuadoriensis (1 clone), L. turionifera (1 clone) and proved to be negative.

No methane emission nor methane in the fronds could be detected in Lemnaceae by SEBACHER et al. (1985).

2. PHYSIOLOGICAL CHARACTERISTICS

2.1. GENERAL

Members of the family of Lemnaceae are suitable organisms to investigate physiological processes and effects of different chemical substances. The utilization of Lemnaceae is therefore wide-spread. Many laboratories throughout the world are involved in studies with Lemnaceae. The amount of papers reporting on physiological properties of Lemnaceae is steadily mounting. The specific literature is so voluminous, diverse, and complex that a complete survey on all work done can scarcely be achieved. A concise appreciation of the older literature has been made by HILLMAN (1961a). A more recent survey of the environmental requirements of Lemnaceae (and other water plants) was made by STEPHENSON et al. (1980). For most details of different methods and experimental conditions, the original literature must be referred to.

Sometimes, it is difficult to compare results of different authors due to different species used or to deviating experimental conditions and methods. This might explain some of the non-conformable results. The identity of the Lemnaceae species in the physiological literature is not always unequivocal. Changes in nomenclature have been made in this work without being mentioned. Apparent misidentifications are corrected by reference to the original name in brackets. Many of the chemical compounds used are named by customary abbreviations. These abbreviations are explained at the end of the volume (p. 582).

2.2. CULTIVATION

2.2.1. Aseptic culture technique

Aseptic culturing of Lemnaceae is generally used today. Methods to obtain sterile fronds have been described by CLARK (1932), SAEGER (1930), LANDOLT (1957), HILLMAN (1961a), BOWKER et al. (1980) and many others. The following method is used at the Geobotanical Institute of the Swiss Federal Institute of Technology (SPIT) in Zurich and is applicable to every species of Lemnaceae:

1) Cleaning of the new clone in water and cultivation under near optimal conditions for one to two days. In this way a good number of healthy fronds will be achieved and spores of microorganisms will eventually germinate, thus becoming more sensitive to the sterilization technique. An addition of sugar may be of advantage in some cases in which spores cannot be brought to germination otherwise. However, the cultivation time must be restricted to 1 or 2 days to prevent too much growth of microorganisms.

2) Dipping of single fronds in a 0.5% solution of hypochlorite (e.g. NaOCl; bleaching agents such as Chlorex, Purex, Eau de Javelle, diluted 10 times) for 1/2 to 5 minutes.

3) Short washing in aseptic H₂O and transferring to an aseptic nutrient solution.

4) After several days, transferring of the fronds which propagate and float in still clear (not cloudy) solution to a new aseptic nutrient solution containing 1% sugar, 0.5% casamino acids and 0.004% yeast extract. This will show up eventual contaminations at once.

If all the fronds die or if the solutions become cloudy or covered by fungi, the treatment has to be begun again. Normally about 1-10% of the fronds succeed in staying alive and become aseptic. Some species (e.g. Wolffiella) need more attempts than others.

Table 2.1. Composition of some nutrient solutions in mg/l used for Lemnaceae growth.

* originally CaCO_3 dissolved in HNO_3

- 1) Hutner 1/5 3) Hillman 5) Bonner-Devirian
2) Hoagland 4) Pirson and Seidel

Substances	1	2	3	4	5
KNO_3		1000	1515	400	85
NH_4NO_3	40				
KH_2PO_4		140	680	200	20
K_2HPO_4	80				
KCl					60
$\text{Ca}(\text{NO}_3)_2 \cdot 4 \text{H}_2\text{O}$	40 *	600	1180		240
$\text{CaCl}_2 \cdot 6 \text{H}_2\text{O}$				1200	
$\text{MgSO}_4 \cdot 7 \text{H}_2\text{O}$	100	500	490	300	40
$\text{FeSO}_4 \cdot 7 \text{H}_2\text{O}$	5		5.4	5	
ferritartrate		5			4
tartaric acid			3		
$\text{MnCl}_2 \cdot 4 \text{H}_2\text{O}$		2	3.6	0.3	
$\text{MnSO}_4 \cdot \text{H}_2\text{O}$	3				0.1
$\text{ZnSO}_4 \cdot 7 \text{H}_2\text{O}$	13	0.2	0.2		1
H_3BO_3	3	3	3	0.5	1
MoO_3		0.07			
$\text{Na}_2\text{MoO}_4 \cdot 2 \text{H}_2\text{O}$	5		0.12		0.025
$\text{CuSO}_4 \cdot 5 \text{H}_2\text{O}$	0.8	0.08	0.08		0.03
$\text{CoSO}_4 \cdot 7 \text{H}_2\text{O}$	0.2				
Na_2 salt of EDTA	100				

2.2.2. Nutrient solutions

Some of the most used nutrient solutions are put together in table 2.1. The mineral molarities of these and further solutions are compared in table 2.2. The different nutrient solutions may be characterized as follows:

- 1) Solution of Hutner. The solution of HUTNER (1953) is too concentrated for most species of Lemnaceae. According to BOIGIANO (1979), it is especially the high boron content which is toxic. Therefore, the solution was diluted five times for most of our experiments. The diluted Hutner solution is characterized by rather low concentration of the main nutrients, a comparatively high concentration of trace elements, and the addition of the chelating agent EDTA. Chlorine is absent. To supply the plants with Cl, it is recommended to exchange $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ for $\text{Mn Cl}_2 \cdot 6 \text{H}_2\text{O}$ (Cl content 0.04 mM).

- 2) Solution of Hoagland. The solution of Hoagland as used by JACOBS (1947) contains a large amount of nitrogen and potassium, but no sodium, and only few trace elements. To cultivate Lemnaceae on this solution for a longer time, it is necessary to add some NaCl (c. 0.1 mM) to the solution. The iron is added to the solution as tartrate which works as a chelating agent.

- 3) Solution of Hillman (M solution). HILLMAN (1961c) modified the solution of Hoagland by increasing the content of potassium, nitrogen, phosphorus, and calcium (and sodium).

- 4) Solution of Pirson and Seidel. The disadvantage of the solution of PIRSON and SEIDEL (1950) is the absence of sodium, zinc, molybdenum, copper, and cobalt. In addition, due to the lack of a chelating agent the absorption of trace elements especially iron is hampered.

- 5) Solution of Bonner-Devirian. The Bonner-Devirian solution as described in HILLMAN (1959b) has rather low concentrations of elements (especially trace elements except chlorine).

- 6) Solution of Eyster. EYSTER (1966) tried to optimize the amount of necessary elements by measuring growth rates and other characteristics of *S. polyrrhiza* successively varying the different elements. The solution he subsequently proposed to be optimal is distinguished by high content of nitrogen and potassium and relatively small amounts of trace elements using EDTA as a chelating agent.

- 7) Solution of Datko 4. DATKO et al. (1980a) worked out a solution for

Table 2.2. Comparison of the molar nutrient content (in mmol/l or mM) of 7 different nutrient solutions
1 Hutner 1/5, 2 Hoagland, 3 Hillman, 4 Pirson and Seidel, 5 Bonner-Devirian, 6 Eyster, 7 Dakko, 8 Docauer

Elements	N	P	S	Ca	Mg	K	Na	Fe	Mn	Zn	B	Mo	Co	Cu	Cl	EDTA
1	1.4	0.4	0.4	0.2	0.4	0.9	0.2	0.02	0.02	0.08	0.03	0.05	0.001	0.0004	0	0.3
2	15	1	2	4	2	11	0	0.02	0.01	0.0015	0.05	0.005	0	0.0004	0.02	0
3	23	5	2	8	2	50	0.05	0.02	0.0015	0.05	0.005	0	0	0.0004	0.04	0
4	4	1.5	1	1	1	6	0	0.02	0.002	0	0.01	0	0	0	10	0
5	3	0.15	0.2	1	0.2	2	0.01	0.02	0.001	0.007	0.001	0.001	0	0.0001	1	0
6	26	0.5	2	0.5	16.5	0.1	0.01	0.01	0.01	0.1	0.001	0.001	0.001	0.001	0.13	0.1
7	5.8	0.4	0.4	1.4	0.8	1.5	0.11	0.06	0.01	0.003	0.07	0.006	0.005	0.003	1.2	0.08
8	0.58	0.04	0.55	0.45	0.55	0.74	1.49	0.036	0.004	0.002	0.009	0.001	0.0002	0.000002	1.58	0.16

optimal growth of L. aequinoctialis. The solution has a rather low content of the main nutrients and a relatively high content of trace elements (including chlorine).

- 8) Solution of Docauer. DOCAUER (1983) developed a medium which is comparable to natural waters (of Michigan). His solution is relatively poor in P and N and rich in Na and Cl. It also contains EDTA.

If supplemented with the missing elements, the nutrient solutions mentioned work well for most species. Optimization of some of the solutions might be achieved by adding the minimum nutrients or by consistently using the chelating agent EDTA. The content of macro-nutrients of 1/5 Hutner and of Bonner-Devirian is at the lower level of the optimum and should be enhanced. KRZECOWSKA et al. (1975) compared the frond size and protein content of L. minor grown in solutions of Hutner, Hoagland, and Pirson and Seidel (at different concentrations). The protein content was highest in fronds grown in Hoagland solution and lowest in Pirson and Seidel solution. The sequence is in accordance with the content of nitrogen in the different solutions. The frond size (not measured in the solution of Pirson and Seidel) was biggest in 0.4 Hutner and in 0.7 Hoagland solution respectively. It did not differ significantly between these two solutions.

A true analysis of optimization of the necessary amount of macro- and micro-nutrients as proposed by HOMES (1963) is still missing. Since the optimum amount of an element for best growth lies well within a wide variation (see EYSTER 1966, LUBOEND 1983, BEYER 1983), the differences between the solutions do not necessarily result in differences of growth rates. But for long-lasting experiments without regular changing of the solutions, it is important to have a well balanced medium with the concentration of the elements near the upper limit for optimal growth.

The pH of the solution is best between 5.5 and 6 (see chapter 2.3.3). In general, the growth rate is not influenced significantly by smaller variations of the pH. However, if ammonium is the only source of nitrogen, the pH might sink rapidly during the experiment and hamper growth or ultimately cause the fronds to die (chapter 2.3.3.3).

Some difficulties may be caused by undisturbed solutions staying at constant temperatures. As a consequence, the solution layer just below the absorbing lower surface of the fronds becomes poorer in nutrients since the absorption is faster than the diffusion from lower layers (EVANS

1972). Occasional movements of the culture vessels help to distribute the nutrients better. Also the carbon dioxide supply will be facilitated.

It is known that different species of the Lemnaceae have different needs of nutrients (LANDOLT 1957 and unpubl., ZIMMERMANN 1981, LUBOEND 1983, BEYER 1983, DOCAUER 1983). In addition, submerged species (e.g. L. tri-sulca, W. lingulata, W. oblonga, W. gladiata, W. denticulata) need a source of carbonate to achieve optimal growth (see chapter 2.3.3.4.5) since the diffusion of CO₂ in the water is too weak for good growth.

2.2.3. Culture and preservation techniques

Cultivation of Lemnaceae is possible in Erlenmeyer flasks or test tubes containing a suitable nutrient solution (chapter 2.2.2). The disadvantage of this simple method is that the medium has to be changed within relatively short periods in order to avoid exhaustion of certain nutrients and to keep growth rates constant. Installation of steady-state culture might be preferable. This technique is described by ERISMANN and FINGER (1968), EICHHORN and AUGSTEN (1969), STRASSER (1971), ERISMANN and BRUNOLD (1973), DAYKO et al. (1978a), TSUDZUKI and KONDO (1979), and ERICSSON et al. (1982). WUESTLING and BOEHM (1980) developed a culture technique of submersed cultivation of W. arrhiza.

The preservation of clones of Lemnaceae is realized by cultivating in test tubes with an agar medium (except for most Wolffiella species which are difficult to grow for any length of time on agar). If 1% sugar and 0.004% yeast extract is added to the medium the cultures are able to grow at low light intensities, and possible infections can be detected quickly. The optimal temperature for storing Lemnaceae is dependent on the species. It should be kept as low as possible to keep growth and evaporation rate low. Species from tropical and subtropical regions need a temperature of about 20°C. The reinoculation to new test tubes with fresh medium has to be done every three months. Within this time the agar dries out and the Lemnaceae fronds begin to die.

It takes about 4 to 6 weeks to get a constant growth rate of Lemnaceae previously kept on agar and transferred to nutrient solution.

Methods of lyophilization or deep freezing are not worked out but urgently needed in order to keep large collections of living Lemnaceae strains for longer periods. In our own experiments, some species of Lemnaceae (e.g. L. minor) were able to survive lyophilization but died after a few days under normal conditions due to the difficulty of cleaning the fronds of glycerol. An adequate method has not been found up to now.

2.2.4. Tissue cultures

Full-grown fronds of *L. aquinoctialis* transferred to a solution with 10 mg/l 2iP and 0.1 mg/l 2,4-D were able to form callus tissue out of the meristematic region around the node (CHANG and HSING 1978). *L. gibba* needed a ratio 10 2,4-D to 1 2iP for callus production (CHANG and CHIU, 1976, 1978). IAA and NAA instead of 2,4-D did not show any effect. After 4 months of growing, the callus began to form green protuberances which later took on frond-like structure (CHANG and HSING 1978). SLOVIN and COHEN (1985c) generated primary callus derived from the meristematic region of *L. gibba* using 2,4-D or IAA-L-alanine as an auxine source. The same authors (1986) investigated chemical mutagenesis and soma-clonal variation in *L. gibba*. Plants regenerated from primary callus exhibit major variations in size and shape of the fronds. Streptomycin treatment produced a number of mutations including a highly gibbous line. NMU mutagenesis resulted in a number of white, yellow, and pale green lines. IV was found to be ineffective.

2.3. VEGETATIVE GROWTH

2.3.1. Measurements

2.3.1.1. Growth rate (multiplication rate)

The growth of *Lemnaceae* is nearly exponential provided that the nutrient and CO₂ supply is sufficient and steady. Contrary to bacteria and other unicellular organisms, fronds of *Lemnaceae* do not propagate by mere doubling. According to species and culture conditions each frond forms up to 20 daughter fronds which become visible within different intervals (several hours to a few days). The determination of the growth rate therefore needs a sufficient number of fronds (at the beginning at least 15-20 fronds) in order to guarantee a more or less statistically based regular growth rate. The simplest method to note the growth rate is to count the visible fronds; daughter fronds are included as soon as they emerge from the pouch of the mother frond. In *Lemnaceae* the preconditions of cultivation have a much longer lasting effect on the growth rate than in unicellular organisms since the formation of the new buds takes place many days before their appearance. As a rule, the experimental conditions should be kept constant for at least 4 weeks before beginning the growth rate measurements. Since the appearance of new daughter fronds is enabled by the elongation of the cells, short-time change in the culture conditions (e.g. short fluctuation of temperature, replacement of the nutrient solution) may show up in a short-term change of growth rate.

The counting of fronds in the closed culture vessel may sometimes be difficult if the number of fronds is very great. Periodical photographs of the cultures might help.

The growth rate is expressed as the difference of the logarithms of the final frond number (F_d) and the initial number (F_o) divided by the number of days of growth. If \log_{10} is used, the result must be multiplied by 1000 to achieve a full value. The growth rate is then represented by the formula:

$$K = \frac{\log_{10}(F_d) - \log_{10}(F_o) \times 1000}{d}$$

A K of 300 means a doubling of frond number within 24 hours, a K of 150 a doubling within 2 days. If the logarithmus naturalis (\ln) is applied the multiplication rate k is expressed as

$$k = \frac{\ln(F_d) - \ln(F_0)}{d} = \frac{K}{1000 \times 0.43} = \frac{K}{430}$$

A doubling of fronds within 24 hours results in a k of 0.7.

The average doubling time T_d (BYSTER 1966) is $\frac{K}{300} = 0.0033 \times K = 1.4 \times k$.

As soon as some factors (e.g. nutrients, light, CO_2) become scarce or if some waste substances concentrate too much, the growth is no longer exponential and the growth rate not steady. Therefore, the experiments with Lemnaceae have to be made in frequently changed solutions or in steady-state culture (see chapter 2.2.3). To be sure that growth is measured within the exponential phase, the counting should be done several times successively. In this way, it should be possible to calculate a regression line according to the following formula:

$$g = \ln(F_0) + bt \quad (\text{cf. LUEOEND 1980})$$

g corresponds to the \ln of the frond number at a time t. b is approximately identical with k.

Different authors (e.g. PIRSON and GOELLNER 1953, HENSSEN 1954, BORNKAMM 1966a, TILLBERG et al. 1979) observed seasonally different growth rates or other varying characteristics of Lemnaceae. To minimize seasonal effects it is advisable to check the chemical composition of the air and the light intensity periodically. For investigations which last for more than one season a control experiment with the same clone and under the same conditions should be performed periodically to detect possible deviations due to different seasonal conditions or annual periodicities.

The maximal growth rates of Lemnaceae are species specific. Highest growth rates measured amount to total K of 300-315 ($k = 0.70-0.74$) in L. aequinoctialis (LANDOLT 1957, DATKO et al. 1980a) and of 280-340 in L. microscopica (VENKATARAMAN et al. 1970). This growth rate corresponds to a doubling time of frond numbers between 20 and 24 hours. Lowest maximal growth rates are observed in Lemnaceae in submerged species (L. trisulcata, W. oblonga, W. gladiata, W. denticulata; LANDOLT 1957

and unpubl.). The limiting factor of submerged species in solutions without sugar is the carbon dioxide supply (cf. chapter 2.3.2.1.). The cellular multiplication rate in meristematic tissue of L. minor was measured by ROMBACH (1976) as 2.7 to 3.0 times higher than the multiplication rate of fronds.

2.3.1.2. Dry weight of fronds

The dry weight of a Lemnaceae frond is species specific, clone specific, and dependent on the environmental conditions. Especially under conditions which favour the storage of starch (e.g. high light intensity and high CO_2 concentration), the dry weight of the frond is enhanced. There is no direct correlation between dry weight of the frond and growth rate. If enough light is present, the accumulation of starch takes place even under poor nutrient conditions or low temperatures which result in slow growth. Under constant culture conditions, the increase of dry weight within a certain period is a measure for the growth rate. However, TILLBERG et al. (1979) report on big fluctuations of mean fresh weight in L. gibba at different times even under the same controlled conditions.

Highest dry weight of a Lemnaceae frond was observed in S. polyrrhiza with 0.6 mg (LANDOLT 1957). Possibly, S. intermedia is able to reach values of frond weight up to 1 mg dry weight. The average dry weight of the smallest Lemnaceae frond (W. angusta or W. globosa) might be as low as 0.01 mg.

2.3.1.3. Size of frond and other frond characteristics

Size of frond, length/width ratio, thickness of frond, number and length of roots, number of cohering fronds, formation of anthocyanin, number and size of papules etc. are further characteristics of Lemnaceae which have been measured to check the influence of certain experimental factors. Especially the size of the frond is a frequently used characteristic in experimental studies. A simple method of measuring the size of the frond is by photography. Cultures of Lemnaceae are pictured from above using a light from below shining through the culture. On a photoelectric cell the light intensity which is proportionate to the total frond area can be measured at once (ASHBY and OXLEY 1935). There is also

the possibility of taking a photograph. In the black-white negative, the fronds are white and the surroundings black. The light which penetrates the negative is proportional to the total size of the frond and can be measured by a photo cell (GORHAM 1941). The size of S. polyrhiza fronds reached 39 mm² in its highest (LANDOLT 1957) under the following conditions: 32°C, 1000 lux, 1/3 Hutner solution with 1% sucrose. In herbarium samples sizes of up to 60 mm² could be observed in S. polyrhiza. The smallest full-grown fronds of W. globosa show a size of less than .1 mm² (own observations).

2.3.2. Effects of gaseous substances

2.3.2.1. General remarks

Gaseous substances are absorbed by Lemnaceae through the open stomata from the air or through the lower epidermis by diffusion from the nutrient medium. The normal composition of the air allows optimal growth for Lemnaceae if circulation of air is guaranteed. Toxic gases in the air influence the growth of Lemnaceae in many ways (see chapters 2.3.2.3 to 2.3.2.6.).

NEY (1960) calculated a gas exchange rate of 2.0 mg per 100 cm² and minute for L. minor. This corresponds to a production of 11 l oxygen per m² and hour. WARD et al. (1963) measured an oxygen production of 4.4 l per m² and hour.

2.3.2.2. Carbon dioxide

2.3.2.2.1. Influence of the CO₂ concentration

Autotrophic growth of L. minor was not possible at a CO₂ concentration of 40 ppm (0.004%) but was at 65 ppm (MUELLER et al. 1977). Under suboptimal conditions of CO₂ (65 and 100 ppm), the fronds are light green and develop 1-3 cm long roots. At 330 ppm CO₂, a concentration which corresponds to the normal air composition, they are dark green, have 4-6 cm long roots and a much faster growth. A supply of 9000 ppm CO₂ (0.9%) does not enhance the growth rate but raises the dry weight, the content of starch, free sugars, free nitrate, and free amino acids; the roots become longer and twisted like a corkscrew. The starch content at 9000 ppm CO₂ is 5 times higher than at 100 ppm. BJOERNDAL and NILSEN (1985) compared growth of L. gibba at a high photon flux rate in normal and in CO₂ enriched air (1500 ppm CO₂). The high CO₂ content results in an increase of the net assimilation rate and in a decrease of the photosynthetic area per unit dry weight. However, it did not affect the growth rate. A significant increase of dry weight and growth rate of L. gibba was observed by ANDERSEN et al. (1985) if CO₂ was enriched up to 6000 ppm. This high CO₂ optimum was explained by the presence of non-functional stomata. The response to high CO₂ was less or absent following four days growth in 2% O₂. The photosynthetic rate was increased by

CO_2 enrichment up to 1500 ppm showing only small increases with further CO_2 enrichment up to 5000 ppm. Figure 2.1 shows the different responses of *Lemna* multiplication at two concentrations of CO_2 and at several light intensities. EYSTER (1966) improved the dry weight increase by aeration with CO_2 enriched air (best results at 50000 ppm = 5%). Aeration with normal air did not bring any better results than no aeration at all. Some authors (e.g. ASHBY and OXLEY 1935, at light intensities of 5000 lux, and LANDOLT 1957 at 2500 and 6000 lux) did not achieve higher growth rates when air was bubbled through the culture flasks. On the other hand, GORHAM (1945) achieved a 10-15% higher growth rate of *S. polyrrhiza* when he aerated the cultures with 4% CO_2 (40000 ppm) at light




Fig. 2.1. Light response curves for photosynthetic rates of *Lemna gibba* measured in air and in CO_2 enriched air at 5000 ppm CO_2 (from ANDERSEN et al. 1985)

intensities of 3000 lux; at 1000 lux no growth acceleration was observed. GORHAM used test tubes for his cultures. Possibly, the diffusion of CO_2 within the tubes was not sufficient to supply the necessary CO_2 . A steady intermixing of the nutrient solution as a consequence of the aeration might also lead to a better supply of nutrients. In a similar way, STEINBERG (1946) was able to enhance the growth rate in crowded cultures by aeration. The growth rate of *L. aquinoctialis* was enhanced by aeration with 0.3% CO_2 ; no further acceleration was achieved with 5% CO_2 (DATKO et al. 1980a). The differences in the results with aeration are probably due to different measuring of growth rates. The growth rate measured as frond increase is not much affected by aeration with normal or CO_2 enriched air except at high light intensities and under crowded conditions. In non-aseptic cultures (as used by ASHBY and OXLEY 1935) on the other hand, the CO_2 might be continuously replaced even at high light intensities due to the respiration of heterotrophic organisms. However, if the frond weight and the photosynthetic rate are considered, the values increase with increasing CO_2 content. At very high light intensities, *L. gibba* is photoinhibited. The percentage of inhibition rises with the lowering of the CO_2 content from less than 30% (1000 ppm CO_2) to 50% (350 ppm corresponding to normal air) and 70% (100 ppm) (NILSEN and DANIELSEN 1985).

2.3.2.2.2. Intensity of CO_2 uptake

GAPONENKO and STAZHETSKII (1969) measured a CO_2 uptake per area in *S. polyrrhiza* at light saturation and 0.5% CO_2 of 3 g per m² and hour. LINDEMAN (1972) recorded the following CO_2 uptake for *L. minor* (table 2.3):

Table 2.3. CO_2 uptake of *Lemna minor* at different light intensities and 5% CO_2 (according to LINDEMAN 1972)

Light intensity in W per m ²	CO_2 uptake in g CO_2 per m ² and hour
10	0.6
70	3.1
100	3.6

2.4. CO₂ uptake of *Spirodela punctata*, *Lemna minor* and *L. valdiviana* at a light intensity of 240 m²·s⁻¹ and at different CO₂ concentrations (from LOATS et al. 1981).

Concentration CO ₂ in ppm	CO ₂ uptake in mg CO ₂ per g fresh weight and hour		
	<i>S. punctata</i>	<i>L. minor</i>	<i>L. valdiviana</i>
300	105	160	140
500	150	205	215
1000	160	250	300

et al. (1981) measured the CO₂ uptake per fresh weight for three different species (table 2.4).

CO₂ content (1%) enhances the netto photosynthesis rate of *L. minor* the first days. Afterwards growth is impeded and the fronds eventually die if nitrate is used as a nitrogen source. In ammonium solution fronds become smaller but reproduce further at the same rate (ANN and MARTI 1986).

2.3. Mode of CO₂ uptake

are generally thought to be C₃ plants. This is true for *S. punctata*, *L. minor*, and *L. trisulca* (ERISMANN 1972, BAUER et al. 1976, and WETZEL 1977), and *L. gibba* (RAGHAVENDRA and DAS 1976) (compare 2.5.1.1). According to RAGHAVENDRA and DAS (1976) and HOUGH and (1977), *Lemnaceae* lack Kranz anatomy. WEDGE and BURRIS (1982) that *Lemnaceae* are not typical for C₃ plants in respect to light ion and optimal temperature of CO₂ assimilation which are both than in C₃ plants, also the low CO₂ compensation point is characteristic rather for C₄ plants. The CO₂ compensation point in axenic *L. minor* is attained at 0.95 ppm CO₂ in the air (FILBIN and HOUGH 1985). These authors found that the activity of the PEP carboxylase of the C₄ acids in early fixation products are very low and within the range normal for C₃ plants.

and SAHA (1986) measured dark carbon dioxide fixation in *S. punctata*. It amounts to 74 pmole per g and hour at 25°C. Submerged aquatic fix 20 to 100 times more CO₂ in the dark. The fixed carbon was

found to about equal parts in the alcohol-soluble and -insoluble fractions.

2.3.2.2.4. CO₂ uptake from the water

In aseptic cultures of some *Lemnaceae* species, the CO₂ content is a minimum factor because the diffusion of CO₂ in water is extremely slow. This results in a growth stop of *L. trisulca* and a very slow growth rate of *Wolffiella* species as *W. gladiata*, *W. denticulata* if sucrose or another carbon source is not added to the solution (Hutner solution does not contain carbonate) (LANDOLT 1957 and unpubl., see also chapter 2.3.3.4.5). Under natural conditions, CO₂ will scarcely be a minimum factor for submerged species because heterotrophic microorganisms produce CO₂ by respiration. Moreover, the submerged species mentioned generally grow in waters containing carbonate.

S. polyrrhiza and *L. minor* which float on the surface of the water are capable of utilizing carbon compounds from the air as well as from the water (WOHLER 1966). WOHLER measured a C uptake from the water in the field between 11 a.m. and 2 p.m. in mg C per g dry weight and hour for different *Lemnaceae*: 1.6 for *S. polyrrhiza*, 1.4 for *L. minor*, 1.9 for *L. trisulca*, and 2.7 for *W. gladiata*. Also SATAKE and SHIMURA (1983) observed in *S. polyrrhiza* CO₂ fixation from air and water. The carbon assimilation from water is comparable to that from the air under normal pH conditions (see table 2.5.). FILBIN and HOUGH (1985) state that in nature (small hard water lake in southern Michigan) 86% of carbon fixed

Table 2.5. Carbon uptake from water in *Spirodela polyrrhiza* in mg C per g dry weight and hour (after SATAKE and SHIMURA 1983). To compare the results of table 2.5 and tables 2.6 the values of table 2.5 have to be multiplied by a factor of ca. 2800.

pH	uptake in water	uptake in air
9.2	0.0044	0.0093
7.0	0.033	0.0079
5.0	0.051	0.0050
4.0	0.059	0.0035

Table 2.6. Carbon uptake from water in *Spirodela polyrrhiza* in nmol C per g fresh weight and minute (after ESHEL and BEER 1986)

nm not measured

pH	uptake in water	uptake in air
8.0	24.8	518
7.2	22.7	600
5.9	5.6	495
4.9	nm	479

by *L. minor* originated from aqueous inorganic carbon whereas in carbonate-free Hutter solution (in the laboratory) under saturating illumination, 63% of the carbon used in photosynthesis came from the atmosphere. The results of SATAKE and SHIMURA (1983) are questioned by ESHEL and BEER (1986) who point to some principal as well as methodological faults in these investigations. In contrast to SATAKE and SHIMURA (1983) they found, in *S. polyrrhiza*, only up to 5% of the carbon used in assimilation as coming from the water (table 2.6). In the water they measured the ratio of free CO_2 to bicarbonate ions at pH 5 as 1 : 0.04 and at pH 7 as 1 : 3.8 (see also fig. 2.9).

ALBERGONI and BASSO (1985) described an apparatus for the simultaneous measurements of photosynthetic ^{14}C fixation by a large number of leaf samples (tested with fronds of *L. minor*).

2.3.2.3. Sulfur dioxide (SO_2)

SO_2 concentrations in the air of up to 0.3 ppm do not have any direct influence on the growth of *L. minor*. However, the starch content of the fronds diminishes from 0.15 ppm SO_2 upwards. A concentration of 0.6 ppm SO_2 resulted initially in a reduction of the growth rate of about 25% and in signs of chlorosis. Interestingly, the tested clone of *L. minor* was able to adapt to the high SO_2 concentration within 50 days, and the normal growth rate was achieved again but the fronds remained smaller (FANKHAUSER et al. 1976). Similarly, LOATS et al. (1981) were not able to detect an influence of 0.5 and 0.75 ppm SO_2 on the growth rate of *L.*

minor and *L. valdiviana*. In contrast, the growth rate of *S. punctata* was slightly reduced. The rate of photosynthesis did not change in *L. minor* but became diminished in the other two species. The dry weight per frond was enhanced in *S. punctata* and *L. valdiviana* but not in *L. minor* under SO_2 influence. SCHAEFER et al. (1975) observed an inhibition of sulfate uptake in *L. minor* at SO_2 concentrations of 0.15, 0.32, and 0.61 ppm. YOON et al. (1981) investigated the effect of 0.5 ppm SO_2 on chromosomes of *L. minor*. SO_2 induced chromosomal nondisjunction and lowered the mitotic index. The same was found by STOECKLI et al. (1975). The authors conclude from their results that SO_2 in the air leads to mutagenous effects.

2.3.2.4. Hydrogen sulfide (H_2S)

Growth rate of *L. minor* was reduced by 15% under the influence of 6 ppm H_2S . The gas impedes the absorption of SO_4^{2-} from the nutrient solution (BRUNOLD and ERISMANN 1970, 1974, 1975). The photosynthesis rate is lowered by 10 to 14% if the H_2S content is enhanced from 6 to 60 ppm. At 120 ppm H_2S the reduction amounts to 88% but is still reversible if the H_2S aeration is stopped. The content of sulfate and cysteine within the frond is enhanced after treatment with H_2S .

2.3.2.5. Ozone (O_3) and ozonated hexene

WANGERMANN and LACEY (1952) reported a damaging effect of UV radiation on *L. minor* which they attribute to the action of developing ozone. Ozone damages *L. minor* and reduces the chlorophyll, protein, and RNA content. The sensibility towards ozone is dependent on the composition of nutrients in the medium. A high amount of nitrogen increases the loss of chlorophyll caused by ozone, a high content of iron diminishes it. The damages of ozone are less if the nutrient solution contains no copper salts (CRAKER 1971, 1972). A concentration of 0.1 ppm ozone lowers growth rate and suppresses flowering in *L. aequinoctialis* (FEDER and SULLIVAN 1969). According to TODD et al. (1956) and ERICKSSON and WEDDING (1956) the ozonated hexene (0.2 ppm) called synthetic smog has a stronger reducing effect on the photosynthesis rate of *L. minor* than ozone (1 ppm). Hexene alone in comparable concentrations had no effect. It is assumed that ozone and ozonated hexene effects the semipermeability

ty of the cell membranes. 100 ppm DPX, an ozone protectant, is able to counterbalance the toxic effect of 0.5 ppm O₃ on L. aequinoctialis (FERDER et al. 1980).

2.3.2.6. Further gases

Carbon monoxide. According to HILLMAN (1954) CO lowers the growth rate of L. minor. However, DALY and BROWN (1954) report that Lemna takes up more oxygen under the influence of CO.

Nitrogen. Photorespiration of L. gibba is blocked and photosynthesis and nitrate reductase activity is reduced by aeration with 100% N₂ (JARCZYK et al. 1984).

Hydrogen fluoride (HF). HF which dissolves in water is taken up passively by turions of S. polyrrhiza by permeation of the undissociated HF through biomembranes and a subsequent dissociation to the impermeable F⁻ ion (KRONBERGER 1983).

Illuminating gas. SAEGER (1933b) reports on the toxicity of illuminating gas on S. polyrrhiza.

Tobacco smoke. 5 blows of 35 ml tobacco smoke into an Erlenmeyer flask brought death to L. gibba plants within 48 hours. The effective substances within the smoke have been detected as dimethylnitrosamine and methylbenzopyrene. Benzanthrane, the concentrate of smoke and the hydrocarbon fraction of the smoke did not show any effect on the growth rate (BHALLA and SABHARWAL 1974).

2.3.3. Effects of dissolved chemical compounds

2.3.3.1. Absorption of dissolved compounds from the water

Water and nutrients are mainly absorbed by the lower epidermis of the fronds of the Lemnaceae which are immersed in water. The roots of Spirodela and Lemna are not very effective in taking up water and dissolved compounds. If the lower epidermis of L. minor is made impermeable with lanolin, the growth rate is greatly reduced. No effect is observed if only the upper surface is covered with lanolin. Cutting off the roots does not influence the growth rate (GORHAM 1941). BLACKMAN and ROBERTSON-CUNNINGHAME (1955) were able to reduce the toxic effect of 2,4-D on L. minor by covering the lower surface of the frond with lanolin regardless of whether the roots were intact or removed. According to SARGENT (from HILLMAN 1961a), there is no transport of water through the root of L. minor if the frond is lifted above the surface of the water with the root tip still dipping in the water. However, FERNANDEZ et al. (1972) report that in S. intermedia the absorption of 2,4-D is achieved mostly through the roots (for explanation see chapter 2.5.2.3.2).

2.3.3.2. Concentration of the nutrient solution

Lemnaceae generally are able to grow within a wide range of nutrient concentrations. Even in deionized water they have been observed growing after 2-3 days (SHARP 1978). But, though Lemnaceae may remain alive in very diluted solutions for days or weeks, using the accumulated nutrients of the older fronds, a long-lasting constant growth rate is only possible in solutions with relatively high nutrient content. LANDOLT (1957) showed that different species behave differently in respect to the minimal, optimal, and maximal concentration of nutrients in the medium. In an experiment with 9 species (LANDOLT unpubl. results) the growth rate was compared in Hutner solution 1/5 and 1/20. S. polyrrhiza, L. gibba, L. aequinoctialis, W. hyalina, W. neotropica, and W. borealis developed as fast in 1/5 Hutner as in 1/20. However W. lingulata, W. oblonga, and W. gladiata had a much higher growth rate in 1/20 Hutner than in 1/5 and looked more healthy. A different utilization of nutrient concentration by 2 species (L. gibba and W. arhiza) is shown in fig. 2.2 (from LANDOLT 1957). At higher concentrations, L. gibba is much faster




Fig. 2.2. Growth rate of *Lemna gibba* and *Wolffia arrhiza* in relation to the concentration of Hutner solution (from LANDOLT 1957)

Table 2.7. Comparison of growth rates of 15 species of Lemnaceae grown in Hutner solution 1/3 and 1/100. The ratio of both rates is given. Each value corresponds to a single clone (LANDOLT unpubl. results)

Species	growth rate
<i>L. gibba</i>	2.3; 1.6
<i>L. turionifera</i>	2.1; 1.8
<i>W. hyalina</i>	1.8
<i>S. punctata</i>	1.6
<i>L. minor</i>	1.6; 1.4; 1.3
<i>S. polyrrhiza</i>	1.4
<i>L. aequinoctialis</i>	1.2
<i>W. arrhiza</i>	1.2
<i>W. borealis</i>	1.1
<i>L. minuscula</i>	1.1; 1.1; 1.1
<i>W. neotropica</i>	1.0
<i>W. linguata</i>	1.0
<i>W. gladiata</i>	1.0
<i>W. columbiana</i>	1.0
<i>W. globosa</i>	1.0

growing than *W. arrhiza*. In 1/30 or 1/100 Hutner, the growth rate of *W. arrhiza* surpassed that of *L. gibba*. In a similar way, CHANG et al. (1977) reported that *W. globosa* (named as *W. arrhiza*) was still able to grow in 1/50 Hutner solution and had its optimum between 1/10 and 1/20 whereas *S. polyrrhiza* and *L. aequinoctialis* achieved good growth at 1/5 and 1/10 but not in lower concentrations. LANDOLT (1957 and unpubl. results) compared growth rates of 15 species in Hutner solution 1/3 and 1/100. The ratios of growth rates K 1/3 to K 1/100 are shown in table 2.7.

The different behaviour of *L. gibba*, *L. turionifera*, and *W. hyalina* on the one hand, and of *L. minuscula* and most *Wolffia* species on the other, is remarkable. It seems that low concentrations can be well utilized by species with great absorbing surface in relation to dry weight (thin fronded and submerged species).

The normal Hutner solution is too concentrated for most species (containing very high amounts of trace elements). HILLMAN (1960a) was able to neutralize the toxic effect of the full Hutner solution on *W. columbiana* by adding kinetin.

Spirodela and *Lemna* form longer roots in diluted solutions; in the same way, the appendage of *W. hyalina* and the conical projection of *W. microscopica* prolongate; also, the fronds get larger and thinner (LANDOLT, unpubl. results). HILLMAN (1954) made a similar observation with *L. minor*. He was able to reverse the effect by adding sucrose to the medium. Lactose or mannitol had no effect. The exact reason of the prolongation effect of low nutrient concentration is not known. Similar effects occur in solutions where only nitrate or phosphate are in low concentrations (see chapter 2.3.3.4.4.).

2.3.3.3. pH of the nutrient solution

2.3.3.3.1. Optimal pH for growth

The influence of the pH on the nutrient absorption of water plants is complex. Especially the solubility of phosphate and of certain trace elements (e.g. Fe, Mo, Zn, Mn) changes with different pH. If chelating agents are used, the limits of pH within which an optimal ion uptake is assured are greatly extended. Good long-lasting growth of 15 species is possible between pH 5 and 8 (higher pH was not investigated) without

significant reduction of the growth rate (LANDOLT, unpubl. results), if EDTA is present in the solution. DATKO et al. (1980a) invariably noted the same growth rate of L. aequinoctialis between pH 4.8 and 7.0. Similar good growth was noted by EYSTER (1966) between pH 4 and 7 for S. polyrrhiza. The weight of the frond is generally higher at high pH (7-8.5) than at lower pH (CHANG et al. 1977). PIETERSE (1972) achieved good growth of W. gladiata in 1/3 Hutner solution between pH 5.5 and 9.0. The growth was stopped at higher and at lower pH and the fronds became yellowish. The relatively high lower pH limit for this submerged species is probably due to the difficulty to get sufficient CO₂ at low pH from the water (cf. chapter 2.3.3.4.5). If sugar is added, W. gladiata is able to grow at pH 4 (see chapter 2.3.3.3.2).

Especially in earlier publications, the pH tolerances for good growth of Lemnaceae often appear much narrower. These results are probably based on cultures with inadequate nutrient solutions (without chelating agent) as is the case in the study of HICKS (1932b) who checked pH tolerances in Knop nutrient solution which yielded good growth of different species only within the following pH limits: S. polyrrhiza (6.4-7.6), L. minor (6.1-6.7), L. trisulca (6.1-6.7), L. minuscula (named as L. cyclostata) (6.1-6.7), W. gladiata (5.0-6.4), W. borealis (6.4-7.6), W. columbiana (6.4-7.0). In experiments with purified waste water, best growth for S. polyrrhiza was observed at pH 6, slightly less at pH 7.8 and 5. No growth occurred at pH 3. McLAY (1976) estimated pH optima for S. punctata with 7.0, for L. minor with 6.2, and for W. australiana with 5.0.

2.3.3.3.2. Lower and higher pH limit for growth

YOSHIMURA (1950) was able to culture S. polyrrhiza at a pH as low as 3.8. When he added 1-10 mM CaCl₂, MgCl₂, or Al₂(SO₄)₃ the plants still developed at a pH of 3.2 to 3.4. It is supposed that these salts have an antagonistic effect on the H⁺ ions. Three species of Lemnaceae from New Zealand (S. punctata, L. minor, and W. australiana) were grown in Hoagland solution by McLAY (1976). The solution was changed daily. S. punctata propagated between a pH near 3 and 10.5, L. minor and W. australiana (named as W. arnhiza) between a pH of 3-4 and 10.5. LANDOLT (unpubl. results) investigated the lower pH limit for growth of 30 Lemnaceae species in 1/10 Hutner solution (table 2.8). To lower the pH successively, NH₄⁺ ions were used as sole nitrogen source. The fronds take up NH₃ by

diffusion and, therefore, the solution rapidly become acidified. The pH at which the cultures die is supposed to represent the lower pH limit. For each clone 3 replications were made. The variations within these replications rarely exceeded 0.1.

The lower limits of pH tolerance vary between the species, but also between clones of the same species. Species with a generally high limit (3.7-4.0) are: L. gibba, L. dispersa, W. hyalina, W. microscopica, L. trisulca, S. polyrrhiza. The following species tolerate rather low pH (2.8-3.2): W. gladiata, S. punctata, W. lingulata, W. oblonga, W. denticulata, W. angusta. The first group contains mostly species which do not colonize waters poor in bases; species of the second group, however, are found occasionally in waters with a low concentration of Ca and Mg.

In nature, Lemnaceae can be observed in waters which have at least temporarily a pH of as low as 3.5 and as high as 10 (LANDOLT and WILDI 1977, LANDOLT unpubl.). It is assumed that organic substances in most of the waters with Lemnaceae have a certain chelating effect.

DOCAUER (1983) measured pH after cultivating species in nutrient solution with NO₃⁻ as sole nitrogen source. His cultures did not die but finally reached a pH of 10.5 (for W. columbiana), 9.5 (for L. turionifera, W. borealis) and 9.0 for S. polyrrhiza.

2.3.3.3.3. Special pH effects

KOPP et al. (1974b) investigated the influence of pH on the absorption of nitrate and ammonium in L. minor. The growth rate in a medium with nitrate as sole nitrogen source is the same at pH 7.1 and 7.7 but 15% lower at pH 6.1. In nutrient solutions with ammonium as sole nitrogen source, the growth rate is similar at pH 6.2 and 7.0 but 15% lower at pH 4.7. The nitrogen content per mg dry weight is correlated with the growth rate but the starch content per mg dry weight is higher at lower pH.

A lower pH facilitates the absorption of growth regulators as the weak acids PAA and TIBA in L. minor (BLACKMAN and SARGENT 1959, BLACKMAN et al. 1959) (compare chapter 2.5.2.3.2). In a similar way the uptake of phenols and 2,4-D is easier at a low pH than at a higher one (SIMON et al. 1952, SIMON and BLACKMAN 1953). The lethal dose of 2,4-D for L. minor is 45 ppm at a pH of 4.6 and 457 ppm at a pH of 6.1. In contrast, weak bases such as BA, are more active at a higher pH (6.2) than at a lower pH (4.2) (HILLMAN 1954, 1955).

Table 2.8. Lower pH limits for growth of Lennaceae (LANDOLT unpubl. results)

Species	no. of clones	lower pH limits clonal values	mean value
<i>S. intermedia</i>	7361, 7342, 8410, 7792, 7291, 7178, 7357, 7702, 7355	3.3, 3.3, 3.3, 3.4, 3.5, 3.5, 3.5, 3.6, 3.7	3.5
<i>S. polyrhiza</i>	7222, 7551, 7110, 7549, 7235	3.5, 3.6, 3.7, 3.7, 3.8	3.7
<i>S. punctata</i>	6725, 7111, 7488, 7248, 7760	3.0, 3.0, 3.0, 3.0, 3.1	3.0
<i>L. gibba</i>	6861, 7218, 7135, 6745, 7245, 7611, 7107, 7705, 6566, 7240, 7021, 7179, 7007, 7533, 7320, 7922, 7262, 7257, 7922a	3.4, 3.5, 3.6, 3.6, 3.7, 3.8, 3.8, 3.9, 3.9, 4.0, 4.0, 4.0, 4.1, 4.1, 4.1, 4.2, 4.2, 4.2, 4.6	3.9
<i>L. disperma</i>	7276, 7818, 7268, 7782, 7259, 7223, 7190	3.6, 3.6, 3.8, 3.9, 3.9, 3.9, 4.0	3.8
<i>L. minor</i>	6591, 6570, 7022, 6580, 7189, 7436, 7538, 7295, 7210, 7008, 6578, 7194, 6728, 7018, 6579	3.1, 3.1, 3.1, 3.2, 3.2, 3.2, 3.3, 3.3, 3.3, 3.3, 3.3, 3.3, 3.4, 3.6, 3.6	3.3
<i>L. japonica</i>	7182, 7951, 7471, 8339, 8653	3.3, 3.4, 3.4, 3.4, 3.4	3.4
<i>L. obscura</i>	7982, 8107, 7134, 7133, 7352, 7599, 7856, 7143,	3.3, 3.3, 3.3, 3.4, 3.4, 3.4, 3.5, 3.5	3.4
<i>L. turionifera</i>	6601, 6619, 6853, 8194, 8117, 8000, 7693, 7137, 6735	3.1, 3.2, 3.4, 3.4, 3.4, 3.4, 3.4, 3.4, 3.5	3.4
<i>L. trisulca</i>	6624, 7794, 7615, 8137, 7315	3.7, 3.8, 3.8, 3.8, 4.0	3.8
<i>L. perpusilla</i>	8539, 8473, 7507, 8017, 8612	3.0, 3.1, 3.3, 3.3, 3.3	3.2
<i>L. aequinoctialis</i>	7122, 8715, 7476, 6746, 7304, 7610, 7643	2.9, 2.9, 3.2, 3.3, 3.4, 3.4, 3.4	3.2
<i>L. valdiviana</i>	7005, 7227, 7703, 8375, 7614	3.2, 3.2, 3.3, 3.4, 3.4	3.3

Table 2.8. (continued)

Species	no. of clone	lower pH limits clonal values	mean value
<i>L. minuscula</i>	8370, 6711, 6717, 7921,	3.1, 3.2, 3.2, 3.3	3.2
<i>W. neotropica</i>	7290, 7279, 7609	3.0, 3.0, 3.3	3.1
<i>W. Welwitschii</i>	7644, 7468	3.2, 3.3	3.3
<i>W. hyalina</i>	7378, 7426, 8640, 7555	3.7, 3.9, 4.0, 4.2	4.0
<i>W. linguata</i>	8041, 7655, 7360, 7725, 8175	2.9, 3.0, 3.1, 3.2, 3.4	3.1
<i>W. oblonga</i>	7166, 7923, 7732, 7997, 8393	2.9, 3.0, 3.2, 3.2, 3.2	3.1
<i>W. gladiata</i>	8261, 7852, 8066, 8350, 9392	2.8, 2.8, 2.9, 2.9, 2.9	2.9
<i>W. denticulata</i>	8221, 7454	3.1, 3.2	3.2
<i>W. microscopica</i>	8359	4.1	4.1
<i>W. brasiliensis</i>	7466, 7925, 8319, 7500, 7897	3.4, 3.4, 3.4, 3.5, 3.5	3.4
<i>W. borealis</i>	7788, 8592, 7566, 7690, 7867	3.4, 3.4, 3.5, 3.5, 3.5	3.5
<i>W. australiana</i>	7605, 7543, 7267, 7773	3.4, 3.4, 3.5, 3.5	3.5
<i>W. angusta</i>	8713, 8714, 7480, 7274	3.1, 3.1, 3.2, 3.2	3.2
<i>W. arrhiza</i>	7323, 7687a, 7014, 7347, 7651	3.4, 3.4, 3.5, 3.5, 3.6	3.5
<i>W. columbiana</i>	7463, 7467, 7574, 7310, 7616	3.4, 3.4, 3.5, 3.6, 3.6	3.5
<i>W. globosa</i>	8245, 6592, 8341, 7233	3.3, 3.4, 3.5, 4.4	3.5

2.3.3.4. Inorganic substances

2.3.3.4.1. General remarks

The requirement of a certain nutrient is partly dependent on the culture conditions. Especially the concentration of other nutrients is important. Certain antagonists might neutralize each other; some elements might be replaced by others; in addition, it is possible that pH and organic substances influence the uptake of trace elements.

Apart from C, O, and H, the following elements proved to be essential for Lemnaceae: K, Ca, Mg, N, P, S, Cl, Fe, Mn, Mo, and B. In addition, a promoting effect is attributed to Na, Li, Cu, Co, Ni, Ga, and Sr. Lemnaceae have a certain capacity to prevent the net uptake of most elements (e.g. K, Ca, Na, P, N, S, Cl). If the concentration of an element in the water reaches a certain optimal level, the content of the element no longer rises within the frond (ALLENBY 1981).

A survey of necessary elements for growth of *S. polyrrhiza* is given in table 2.9, distinguishing between the minimal amount which is needed for constant growth rate, the optimal amount necessary for a maximal growth rate and the maximal amount which still permits a steady growth rate.

Table 2.9. Nutrient requirements of *Spirodela polyrrhiza* in mM (according to EYSTER 1966)

Elements	minimal amount	optimal amount	maximal amount
H as NO ₃ as NH ₄	<0.5	10 - 30	100
P	<0.5	2.5 - 5	75
S	0.005	0.2 - 1	50
K	0.001	0.5 - 20	50
Na	0.005	1 - 20	50
Ca	<0.00001	0 - 60	>60
Mg	0.2	0.5 - 10	50
Cl	0.01	0.2 - 4	50
Fe	0.0001	0.01 - 1	>10
Mn	0.0005	0.01 - 0.1	1
Zn	0.00001	0.001 - 0.06	1
B	0.002	0.02 - 0.5	5
Mo	<0.00001	0.1 - 1	10
Cu	<0.00001	0 - 0.3	3
	<0.00001	0.001	0.1

The effects of the different nutrients on growth of *S. polyrrhiza* are presented in figs. 2.3 and 2.4. It is evident that with other species (or clones) and under other conditions, the values will be different.

The effect of some inorganic toxic substances will be shown in chapter 2.3.3.5.6.6.

The addition of 50% deuterium oxide to water has an inhibiting effect on the growth of *L. minor*. 80% of deuterated water prevents growth (BOUDET et al. 1975). Deuterium oxide impedes protein synthesis and enhances protein degradation (2-3 times at 20% and 9-10 times at 50%) in *L. minor* (COOKE et al. 1979a, 1980b).

The effects of the different nutrients on growth of *S. polyrrhiza* are presented in figs. 2.3 and 2.4. It is evident that with other species (or clones) and under other conditions, the values will be different.

The effect of some inorganic toxic substances will be shown in chapter 2.3.3.5.6.6.

The addition of 50% deuterium oxide to water has an inhibiting effect on the growth of *L. minor*. 80% of deuterated water prevents growth (BOUDET et al. 1975). Deuterium oxide impedes protein synthesis and enhances protein degradation (2-3 times at 20% and 9-10 times at 50%) in *L. minor* (COOKE et al. 1979a, 1980b).

2.3.3.4.2. Monovalent alkali metals (K, Na, Rb, Cs, Li)

2.3.3.4.2.1. Potassium (K)

Good regular growth of *L. minor* was achieved with 0.5-10 mM K (STRAUSS 1976). In *S. polyrrhiza*, the optimal amount of K was found to be between 1 and 20 mM (EYSTER 1966). At 40 mM K the growth rate is much retarded, and at 60 mM it is very low (fig. 2.3). At 0.005 mM K the growth rate of *S. polyrrhiza* was found to be very low too, and the fronds showed chlorotic symptoms. According to WHITE (1936a, 1938, 1939) the requirement for potassium in *L. minor* is dependent on the given light intensity. The lowest optimal amount of K varies between 0.05 mM at 600 lux (60 ft-c) and 5 mM at 3000 lux (300 ft-c).

Lack of potassium expresses itself in a reduced dry weight per frond in *S. polyrrhiza* (EYSTER 1966), in a decrease of the size of the frond, length of the root, photosynthesis rate (WHITE 1938, 1939), cell length, and time of plasmolysis (PIRSON and SEIDEL 1950) in *L. minor*, and in a higher Mg content in *L. gibba* (LIEBERT 1986b) (table 2.10).

Table 2.10. The effect of low potassium content on characteristics of Lemnaceae

- + higher values a) *S. polyrrhiza* 1 EYSTER 1966
 - lower values b) *L. gibba* 2 LIEBERT 1986a
 c) *L. minor* 3 PIRSON and SEIDEL 1950
 4 WHITE 1938, 1939

Characteristics	low K	species	authors
frond area	-	a	1
dry weight	-	a	1
root length	-	c	4
cell length	-	c	3
photosynthesis rate	-	c	4
time of plasmolysis	-	c	4
Mg content	+	b	2




Fig. 2.3. The effect of the concentrations of potassium, calcium, magnesium, nitrate, phosphate, and sulfate on the growth rate of *Spirodela polyrrhiza* (after data in EYSTER 1966)

Potassium is absorbed in exponentially growing *L. minor* at a constant rate. If the K content of the solution is reduced during an experiment, the uptake of K will be faster at first but will finally become slower than in the solution with a higher K content. If the K content is raised the process is inverted (YOUNG and SIMS 1972). Studies with ATP and CCCP suggest that the K uptake is an active process, and is located at the plasmalemma where the ATPase activity can be shown (YOUNG and SIMS 1973).

The influence of Ca and Mg on the uptake of K in *L. minor* was studied by AYADI et al. (1974). The daily periodical uptake of K (as well as of Mg, PO_4 , and NO_3) in continuous light is dealt with in chapter 2.5.9. The absorption of K in concentrations below 1 mM is linear to the logarithm of the content of K ions, above 1 mM it is much faster. Ca reduces the K uptake in concentrations of K above 1 mM (AYADI 1971).

ECHLIN and HAYES (1986) studied the distribution of elements (K, Na, Mg, Ca, P, S, Cl) within the roots. They concluded that potassium is the only element which might be not only taken up locally by the roots but transported from the roots to the aerial parts of the plants.




Fig. 2.4. The effect of the concentrations of iron, manganese, zinc, and chloride on the growth rate of *Spirodela polyrrhiza* (after data in EYSTER 1966)

2.3.3.4.2.2. Rubidium (Rb), Lithium (Li), and Caesium (Cs)

Rubidium (contrary to Caesium) is able to replace potassium for longer times. Growth rate and photosynthesis of L. minor do not change, however the length of the root and the size of the frond are reduced (BIERHUIZEN 1954). The uptake and distribution of rubidium into the root tips of L. minor was studied by ECHLIN et al. (1982c). AYADI et al. (1971), LASSAL-LES et al. (1973) and YOUNG and SIMS (1973) investigated the reciprocal competition between K and Rb during the nutrient uptake in L. minor. The influence of Ca on the Rb uptake of L. minor was demonstrated by AYADI and THELLIER (1970). Substitution of K at a level of 70% or more by Rb extends the period of uptake rhythm in L. gibba G₃ (KONDO 1984b). The need of lithium for growth of Lemnaceae has never been proved. However, an addition of 0.15-0.72 mM Li to the nutrient solution promoted the growth of L. minor (STRAUSS 1976). Similar results are reported for S. polyrrhiza and L. minor by WIEWIORKA and SAROSIEK (1986). If 3.6 mM Li was added, the growth declined. Addition of 0.2 mM Li or more prolonged the rhythm of K uptake in L. gibba and lowered the amplitude of the rhythm. Na (contrary to Rb, Ca, or Mg) at a very low concentration (20 mM) almost completely cancelled the effect of Li (KONDO 1984a). BIELENSKI et al. (1984a,b) report on the transport of Li in L. gibba. LiCl (1 mM) inhibits flower production in the long-day plant L. gibba and stimulates it in the short-day plant L. aequinoctialis under long-day conditions (KANDELER 1970).

Caesium is not essential for Lemnaceae. The uptake of Cs was studied in L. minor by BERGAMINI et al. (1979). It is faster at pH 5 than at higher pH (EL-SHINAWY and ABDEL-MALIK 1980).

2.3.3.4.2.3. Sodium (Na)

It is controversial if Lemnaceae need sodium for constant growth or not. However, a stimulation of the growth rate has often been observed. According to STRAUSS (1976), L. minor needs 5-10 mM Na (0.3-0.6 o/oo NaCl) for good growth. HALLER et al. (1974) achieved best growth of L. obscura (named as L. minor) by adding 14-43 mM Na to the solution. At lower Na concentrations the growth rate was considerably lower. STANLEY and MEADWELL (1976a) working with L. obscura (named as L. minor) also obtained a maximal growth rate with 25-60 mM Na. At 100 mM (6 o/oo NaCl) growth was

greatly reduced and at 150 mM Na (8.8 o/oo NaCl) nearly stopped. HALLER et al. (1974) observed a slight retardation in growth rate at 110 mM Na and attained the lethal concentration at 280 mM Na (16.6 o/oo NaCl). HOWARD-WILLIAMS (1979) investigated the growth rate of S. polyrrhiza, L. aequinoctialis, and W. hyalina in solutions with different contents of Na (added as NaCl together with Na₂CO₃ in the following concentration of Na in mg per l: 5, 20, 100, 250, 500, 750, and 1000). Best growth was achieved for S. polyrrhiza at 250 mg/l, for L. aequinoctialis at <100 mg/l and for W. hyalina at 1000 mg/l. It is not clear if this effect is due to the sodium or to the chloride concentration. However, the high optimal concentration of NaCl for W. hyalina is remarkable.

EYSTER (1966) was not able to promote growth of S. polyrrhiza by adding NaCl to the Na free solution. He measured only dry weight (and not growth rate). At a concentration of 60 mM Na and higher the total dry weight was distinctly lower than in solutions without Na.

In many chemical substances and in the glass of the Erlenmeyer flasks used, Na impurities are present. Therefore, it is difficult to prove the necessity of sodium for Lemnaceae growth. In addition, it is not always certain if a positive effect of NaCl is not due to the Cl (see chapter 2.3.3.4.6). According to HUBER and SANKHIA (1979) an addition of 17-85 mM NaCl to the solution stimulates different photosynthetic enzyme activities in L. minor. However, photosynthetic fixation of CO₂ is impeded at NaCl concentrations of 43-85 mM Na. The Hill reaction of isolated chloroplasts of L. minor shows a promotion at 17 mM NaCl but a slight or strong reduction at 43 and 85 mM, respectively. The internal Na content within the fronds is enhanced sixfold after addition of 85 mM NaCl, the chlorine content only twofold. The ABA level rises to 150% after addition of NaCl.

The addition of NaCl (20, 40, and 80 mM) to the nutrient solution decreased the activities of nitrate reductase, glutamate synthase and glutamine synthetase in L. minor. The activity of glutamate dehydrogenase was slightly stimulated (HUBER 1982). The author points to the fact that the ability of L. minor to assimilate ammonium via the glutamine synthetase / glutamate synthase pathway is inhibited under salt stress conditions. The direct entry of ammonium to glutamate may still be possible (HUBER 1982).

Sodium can partly replace potassium in Lemnaceae (BIERHUIZEN 1954, STRAUSS 1976), however, the sodium ion competes with the potassium ion and slows down the uptake rate of potassium (YOUNG and SIMS 1973).

2.3.3.4.3. Bivalent alkali metals (Ca, Mg, Sr, Ba)

The optimal amount of calcium for Lemnaceae growth depends greatly on the content of magnesium in the solution and vice versa. Best growth with *S. polyrrhiza* was achieved with 0.5-10 mM Ca and 0.2-4 mM Mg (EXSTER 1966). Under optimal supply of Mg the lethal dose of Ca is 50 mM; the same limit was observed for Mg under optimal supply of Ca. The minimum amount is 0.1 mM Ca in solutions with 20 mM Mg, and 0.01 mM Mg in solutions with 10 mM Ca. Fronds in solution lacking calcium, are very small, yellowish to brown, and grow in clusters. The highest dry weight per frond is produced in solutions with 5 mM Ca and 0.4 mM Mg. STRAUSS (1973) observed the highest growth rate of *L. minor* at 0.2-20 mM Ca and at 0.05-10 mM Mg concentrations. The requirements for Ca and Mg by *S. polyrrhiza*, *L. gibba*, *L. minor*, and *L. minuscula* were investigated by ZIMMERMANN (1981). Good growth for all species was achieved with the simultaneous supply of 0.01-15 mM Ca and the same amount of Mg (higher concentrations have not been studied). 0.002 mM Ca and Mg did not suffice for growth. With a constant supply of 0.3 mM Mg, 0.1 mM Ca was not enough for growth. A slow growth rate was obtained with 0.6 mM Ca for *S. polyrrhiza*, *L. minor*, and *L. minuscula* and with 3 mM Ca for *L. gibba*. Specific differences between the different species showed up if the Ca content was kept at 0.3 mM and the Mg content varied. A slow growth rate was achieved at 0.001 mM Mg in *S. polyrrhiza*, *L. minor*, and *L. minuscula*, at 0.002 mM Mg in *L. gibba*. At 5 mM Mg *L. gibba* and *L. minuscula* still grew fairly well whereas *L. minor* and *S. polyrrhiza* showed a much reduced growth rate. A concentration of 7 mM Mg was lethal for all species. The results of ZIMMERMANN suggest that

- 1) Lemnaceae have more need for Ca than for Mg
- 2) Ca and Mg behave as antagonists and are able to detoxicate the effects of each other; especially a high Mg content is toxic if the Ca content is low; if the ratio of Ca/Mg is smaller than 1/20, the fronds of all investigated species die; a reduction of growth rate already takes place at a ratio of 1/10.
- 3) There are differences between species of Lemnaceae in the need of Ca and Mg.

A much more elaborate investigation on Ca and Mg requirements and tolerances of 30 Lemnaceae species is in progress (BEYER 1983 and unpubl. results). Two examples of optimal, maximal, and minimal content of Ca and

Mg in nutrient solutions needed or tolerated for long lasting growth are given in figs. 2.5 and 2.6. There are some differences between species and clones. But they are not as big as could have been expected from the different occurrence of the species in nature. A survey of the maximum and minimum Ca and Mg requirement of each species is given in table 2.11. BOLOGIANO (1979) observed optimal growth rate at Ca concentrations of 0.015 mM for *L. aquinoctialis* and of 0.007 mM for *L. minor*. In solutions with low Ca concentration he noted marked influence of relatively high amounts of EDTA on growth and appearance of Lemna cultures. He assumes that EDTA removes Ca from the solution.

The frond of *L. minor* is larger at Ca concentrations higher than 7 mM whereas fronds of *S. polyrrhiza* and *L. gibba* are smaller and fronds of *L. minuscula* remain unaltered at these concentrations. At concentrations below 0.5 mM Ca, the fronds are distinctly smaller in *S. polyrrhiza* and *L. gibba*, slightly smaller in *L. minuscula* and not changed in *L. minor* compared with a medium concentration. The length/width ratio of the fronds is not varied under different concentrations of Ca and Mg (ZIMMERMANN 1981). The fronds of *L. gibba* are thickest at 1.5 mM Ca; the Mg concentration did not influence the thickness of the frond. At high concentrations of Ca and Mg, clusters of fronds are formed in all species (incomplete disconnections of the daughter fronds). The variation in root length with concentration of Ca and Mg is not uniform. The roots are small in all species at high concentrations of Ca and Mg. At lower concentrations, the length of the root is dependent on the ratio of Ca/Mg and species specific (ZIMMERMANN 1981). *W. arhiza* becomes bigger if the Ca concentration of the Hoagland solution used is doubled. The fronds become smaller if the Mg content is raised. The length/width ratio is highest at the high Ca concentration and lowest at the high Mg concentration (SAROSIEK and WOZAKOWSKA-NATKANIEC 1984). JEFFRIES et al. (1969a) observed a reduction of growth rate of *L. minor* at Ca concentrations from 2 mM upwards. Different clones behaved differently. However, all clones grew more or less better at Ca concentrations of 0.3 mM than of 1 mM. The discrepancy of these results with those of previously mentioned authors in which much higher Ca concentrations still made an optimal growth possible, cannot be explained. Since the culture conditions are not described, it is difficult to discuss likely reasons (e.g. unbalanced nutrient solution with either a low Mg or K content).




Fig. 2.5. Requirement and tolerance of *Spirodela intermedia* (no. 8410) for Ca and Mg (from BEYER, unpubl. results). Temperature 26.6°C, light intensity 14000 lux, continuous illumination. The inner curve circumscribes conditions for optimal growth rate (>90% of the maximal growth rate), the next two curves connect the points with 80% and 60% of the maximal growth rate.




Fig. 2.6. Requirement and tolerance of *Wolffia angusta* (no. 8713) for Ca and Mg (from BEYER, unpubl. results). (Explanations see fig. 2.5)

Table 2.11. Maximum and minimum requirement of Ca and Mg (in mg/l) for long-lasting growth of Lemnaceae species (BEYER, unpubl. results)

Species	no. of clones	Ca		Mg	
		min	max	min	max
<i>S. intermedia</i>	7291	0.8	800	0.2	400
	7342	0.8	800	0.1	400
	7357	0.8	800	0.1	800
	7361	0.4	800	0.1	400
	7820	0.8	800	0.1	400
<i>S. polyrhiza</i>	8410	0.4	800	0.1	400
	7890	0.8	800	0.1	400
	8403	0.8	800	0.1	400
	7181	0.8	800	0.1	800
	7111	0.4	1600	0.05	800
<i>S. punctata</i>	7449	0.4	800	0.05	800
	7487	0.4	800	0.1	800
	7021	0.8	1600	0.1	800
<i>L. gibba</i>	7262	0.4	800	0.1	400
	8128	0.8	800	0.2	400
	7777	0.4	800	0.1	800
<i>L. disperma</i>	7761	0.8	800	0.1	800
	7223	0.8	800	0.1	400
	7189	0.8	1600	0.1	800
<i>L. minor</i>	7210	0.8	1600	0.1	800
	7567	0.4	800	0.1	800
	7427	1.6	1600	0.05	800
<i>L. japonica</i>	8896	0.4	1600	0.05	1600
<i>L. ecuadoriensis</i>	7780	0.8	800	0.1	800
<i>L. obscura</i>	7720	0.4	800	0.05	800
	7134	0.8	800	0.1	800
<i>L. perpusilla</i>	8017	0.8	800	0.05	400
	8473	0.4	800	0.05	400
	8539	0.4	800	0.05	400
<i>L. aequinoctialis</i>	7122	0.8	800	0.05	800
	7300	0.8	800	0.1	400
	7743	0.8	800	0.05	400

Table 2.11 (continued)

Species	no. of clones	Ca		Mg	
		min	max	min	max
<i>L. valdiviana</i>	7005	0.8	800	0.1	400
	7227	0.4	800	0.05	800
	7546	0.8	800	0.05	800
<i>L. minuscula</i>	6726	0.8	800	0.05	800
	7180	0.4	1600	0.05	800
	7724	0.4	800	0.05	800
<i>W. hyalina</i>	7376	3.2	3200	0.1	800
	7555	0.8	1600	0.05	800
<i>W. neotropica</i>	7225	0.8	800	0.05	800
	7290	0.8	800	0.05	400
	7609	0.8	800	0.05	400
<i>W. microscopica</i>	8359	3.2	800	0.05	200
<i>W. australiana</i>	7631	0.8	1600	0.1	800
<i>W. angusta</i>	7274	0.8	800	0.05	800
	7480	0.8	1600	0.05	800
<i>W. arrhiza</i>	7246	0.8	1600	0.05	800
	8272	0.8	1600	0.05	800
<i>W. globosa</i>	7700	1.6	1600	0.05	800

The content of protein nitrogen, of lipochromes, and lipids of *S. polyrhiza* is reduced at high concentrations of Ca (75 or 150 mM) (LECHEVALIER 1977a).

The Ca uptake was studied by STELZ et al. (1975). *L. minor* is able to absorb maximally 0.216 meq Ca per g dry weight. It behaves as a typical calcicole species. The Ca uptake of *L. gibba* is hampered by BA and stimulated by ABA (DECOCK and HALL 1981). The Ca and Mg content of the medium has an influence on the K uptake (AYADI et al. 1974). According to JEFFERIES et al. (1969a), an increase of the Ca concentration raises the K uptake of *W. arrhiza* for a short time. After 15 days the original uptake rate is reached again.

The activity of the malate dehydrogenase is influenced by the Ca concentration of the nutrient solution. Different clones of L. minor showed different growth rates and different reactions of enzyme activities if the Ca concentration was increased. At higher Ca concentrations the enzymes partly agglomerate to aggregates of a higher molecular weight. STEWART and SIMS (from JEFFRIES et al. 1969a) found out that allosteric enzymes of L. minor (such as glucose-6-phosphate dehydrogenase, NADP isocitrate dehydrogenase, and NAD glutamate dehydrogenase) react differently towards varying Ca concentrations. According to NAKASHIMA and TSUBZUKI (1977), Ca is probably necessary in L. gibba to fix the uridine-transporting proteins to the cell membrane. BORNKAMM (1965) and LOETSCH and KINZEL (1971) are of the opinion that Ca plays an important role by precipitating the oxalic acid thereby preventing its accumulation within the cell.

According to NAKASHIMA (1979b), Mg has three different functions in maintaining the RNA synthetic activity of isolated nuclei of L. gibba:

- 1) Mg is a cofactor of RNA polymerase,
 - 2) Mg is a stabilizer for the binding of RNA polymerase I to the nuclei,
 - 3) Mg is a stabilizer of factors stimulating the RNA synthetic activity.
- The exchange of Ca and K ions in L. minor and Ca and Na ions was studied by DEWARTI et al. (1977, 1978). Of the three ions, Ca has by far the highest affinity to the carboxyl groups of the cell wall. Similar results were obtained by MORVAN et al. (1984) who investigated the cell wall selectivity of L. minor for Ca, Mg, K, and Na. Ca was shown to have much higher affinities to the cell wall compared to other ions. SCHREINEMAKERS (1984) suggests a coupled uptake mechanism of Ca and Zn and of Mg and Mn.

The effect of Ca and Mg on the radiosensitivity of W. arrhiza was studied by SAROSIEK and WOZAKOWSKA-NATKANIEC (1984). If the Mg and Ca content of the Hoagland solution is doubled, W. arrhiza is less sensitive to γ -irradiation. The DL 100 rises from 800 to at least 1300 mC kg⁻¹. Strontium (Sr) which is not an essential element for Lemnaceae becomes incorporated into the calcium oxalate raphide crystals in L. minor if added to the nutrient solution (FRANCESCI and SCHUEREN 1986). The growth rate and root length of L. minor and S. polyrrhiza are reduced with rising Sr concentrations up to 100 mg/l (c. 1 mM). S. polyrrhiza is more sensitive to Sr than L. minor (WIEWIORKA and SAROSIEK 1986). Barium (Ba) inhibited growth of L. minor at concentrations of 0.1 mM and

higher. If river water of Illinois river was used in the nutrient solution instead of deionized water, no toxicity of Ba was observed even at 0.5 mM Ba (WANG 1986b).

2.3.3.4.4. Nitrogen (N), phosphorus (P), and sulfur (S)

2.3.3.4.4.1. Nitrogen (N)

2.3.3.4.4.1.1. General remarks

MOLEK (1979) reports that the requirements of nitrogen in Lemnaceae are species specific and dependent on culture conditions. To build up 1 mg dry weight of either S. polyrrhiza or W. arrhiza, 0.05-0.08 mg NO₃-N is needed. EYSTER (1966) measured a minimum amount of NO₃-N for optimal growth of S. polyrrhiza as 0.028 mg / mg dry weight. At high concentrations of N, the nitrogen requirement rose to 0.5 mg / mg dry weight. ALLISON et al. (1948) were not able to demonstrate escaping N₂ from cultures of L. minor supplied with NO₃⁻ or with asparagine.

2.3.3.4.4.1.2. Sources of nitrogen utilizable for Lemnaceae

Lemnaceae are able to take up nitrogen in form of nitrate, nitrite, ammonium, urea, or certain amino acids. However, main sources of nitrogen for most species are nitrate and ammonium. The submerged-growing L. tri-sulca is able to use the following nitrogen sources in order of decreasing effectiveness (decreasing growth rates): urea, aspartic acid, nitrate, glutamic acid, arginine, ammonium, casein hydrolysate. Nitrite supports a relatively high growth rate after an initial lag of 7 days (HOLST and YOPP 1979). Fresh weight and chlorophyll content were highest with casein hydrolysate before nitrite and arginine. According to JOY (1969a) and SIMS et al. (1968), arginine is not a very suitable nitrogen source for L. minor and not at all for S. punctata. Further organic nitrogen sources for Lemnaceae are enumerated in chapter 2.3.3.4.4.1.5.

2.3.3.4.4.1.3. The effect of different nitrogen supply

EYSTER (1966) observed good growth of S. polyrrhiza between 5 and 50 mM NO₃-N. However at concentrations of 5 mM and lower the fronds looked

yellowish and produced much anthocyanin on the lower surface. For the same species, already 2.4-5 mM $\text{NH}_4\text{-N}$ were sufficient for good growth. At 1 mM $\text{NO}_3\text{-N}$ or $\text{NH}_4\text{-N}$, a slow growth rate was noted and some turions were formed; on the other hand, the growth rate slowed down at 100 mM $\text{NO}_3\text{-N}$ or 75 mM $\text{NH}_4\text{-N}$. L. aequinoctialis shows a reduced growth rate at 50 mM N, wilting effects at 400 mM and dead fronds at 1200 mM (FERNANDEZ and BALDOS 1981, FERNANDEZ et al. 1983). According to INGEMARSSON et al. (1983, 1984), L. gibba is able to take up $\text{NO}_3\text{-N}$ from solutions of 0.01 mM upwards.

At low nitrogen concentrations, the growth rate of Lemnaceae is directly proportional to the amount of N added (fig. 2.7). A reduction of the N supply results in a reduced net photosynthesis rate of L. gibba. This effect is due to a reduction in chlorophyll content (observed in L. gibba, L. minor, L. aequinoctialis) (ERICSSON et al. 1982). The respiration rate also decreases concomitantly (INGEMARSSON et al. 1983, 1984). LUEOEND (1980, 1983) investigated the growth rates and other characteristics of 1 clone of 4 Lemnaceae species (S. polyrrhiza, L. gibba, L.




Fig. 2.7. Relation between growth rate and nitrogen supply in Lemna gibba (from ERICSSON et al. 1982)

minor, and L. minuscula) at ten different N concentrations: 0.0003, 0.0016, 0.008, 0.04, 0.2, 1.0, 5.0, 25, 125, and 625 mM. All four species died at the two highest concentrations of N. Best growth (at least 80% of the maximum growth rate) was achieved for: S. polyrrhiza at 0.2, 1.0, 5.0 and 25 mM, L. minor, L. gibba and L. minuscula at 0.04, 0.2, 1.0, 5.0 and 25 mM N (as NH_4NO_3). Fifty percent of the maximum growth rate is still reached at N concentrations of 0.0016 mM (L. minuscula), 0.08 mM (L. minor, L. gibba), and 0.04 mM (S. polyrrhiza). Below 0.2 mM N, S. polyrrhiza sometimes forms turions.

ERICSSON et al. (1982) confirmed the results of LUEOEND which impress that L. gibba is much less able to utilize low nitrogen concentration than L. minor (and L. aequinoctialis). Table 2.12 shows the different

Table 2.12. Nitrogen requirement and nitrogen tolerance of different species of Lemnaceae: minimum amount of nitrogen (to result in half of the maximum growth rate), optimum amount of nitrogen (to achieve at least 80% of the maximum growth rate), and maximum amount of nitrogen (to reach at least half of the maximum growth rate) from DOCAUER 1983 (1), EYSTER 1966 (2), FERNANDEZ and BALDOS 1981 (3), INGEMARSSON et al. 1984 (4), LUEOEND 1983 (5), MUELLER 1983 (6), STANLEY and MEAD-WELL 1976a (7), WHITE 1937a (8).

* at lower N concentrations, turions were formed

Species	requirement of N in mM		
	minimum	optimum	maximum
<u>S. polyrrhiza</u>	0.04 * (5) 0.006 (1)	5-30 (2) 1-25 (5) -10 (6)	120 (2) 100-120 (6)
<u>S. punctata</u> <u>L. gibba</u>	0.01 (4) 0.008 (5)	-20 (6) 0.04-25 (5) -50 (6)	80-120 (6) 150-200 (6)
<u>L. obscura</u> <u>L. turionifera</u> <u>L. minor</u>	0.005 (1) 0.005 (1) 0.008 (5) 0.008 (8)	0.02- 5 (6) 0.2 -25 (5)	60-100 (7) 150-200 (6) 100-150 (6)
<u>L. aequinoctialis</u> <u>L. minuscula</u> <u>W. borealis</u> <u>W. columbiana</u>	0.0016 (5) 0.01 (1) 0.003 (1)	0.04- 1 (5) 0.01(1)-5(6)	450 (3) 30- 50 (6)

reactions of Lemnaceae species in relation to the N concentration. The results are only partly comparable. It is known that different clones of one species may behave differently. The lower limit of the optimum nitrogen supply for L. gibba, for instance, varies between 0.008 and 1 mM depending on the clone investigated (DANN 1982, LANDOLT and DANN 1983). Also, a species does not show exactly the same reactions under different culture conditions. For instance WHITE (1937a) was able to demonstrate that increasing light intensities enhance the optimal N concentration for L. minor. Finally, differences can be observed between different N sources (ammonium versus nitrate) (e.g. LUEOEND 1980, EYSTER 1966). From table 2.12 we can summarize the following:

- 1) the minimum need of N for growth of different species decreases in the following sequence: S. polyrrhiza, L. gibba, W. borealis, L. minor, L. turionifera, W. columbiana, L. minuscula;
- 2) the maximum amount of N tolerated for constant growth decreases in the following order: L. aequinoctialis, L. gibba, L. turionifera, L. minor, S. polyrrhiza, S. punctata, L. obscura, L. minuscula.

The maximum frond size is reached at medium concentrations (0.04-1 mM N for S. polyrrhiza, L. gibba, L. minor and L. minuscula) (LUEOEND 1983). DANN (1982) reported greatest size of L. gibba at concentrations of 0.2-25 mM depending on the clone investigated. The greatest dry weight per frond in S. polyrrhiza was achieved at 25 mM N (EYSTER 1966). The N concentration does not significantly influence the length/width ratio of S. polyrrhiza, L. gibba, L. minor and L. minuscula (LUEOEND 1983) nor the gibbosity of L. gibba (DANN 1982, LANDOLT and DANN 1983). The length of the root is characteristic for the carbon/nitrogen ratio, according to WHITE (1937b). The longest roots are found if the rate of photosynthesis is high and the nitrogen supply low. At very low N concentrations, the roots become smaller again. This root reduction is reported by LUEOEND (1980, 1983) at the following concentrations and lower: 0.0003 mM N (for L. minor) and 0.008 mM N (for S. polyrrhiza, L. minuscula and L. gibba). This reduction is probably due to the fact that photosynthesis is already disturbed at very low N concentrations. DANN (1982) found the longest roots in most clones of L. gibba (6 of 10) at 0.04 mM N. In two clones, the maximum showed up at 0.008 mM and in the remaining two clones at 0.2 mM. Many other authors report on the influence of N concentrations on certain characteristics of Lemnaceae (table 2.13). Low N concentrations result in a reduction of growth rate, size of frond,

chlorophyll content, respiration rate, and starch content (WHITE 1936b, WHITE and TEMPELMANN 1937 with L. minor). Long roots, long root cells, reduced photosynthesis rate, reduced respiration have been observed by PIRSON and GOELLNER (1953) with L. minor. In addition, HUMPHREY et al. (1977) reported for L. minor a reduced glycolysis, a great amount of free amino acids and a high rate of protein degradation in solution with low nitrogen supply. These authors suppose a change of enzyme systems at

Table 2.13. The effect of low nitrogen supply on characteristics of Lemnaceae

+ increase, - decrease, 0 no effect,
* at very low N concentrations the roots are shorter again (6)

a	<u>S. polyrrhiza</u>	1	DANN 1982
b	<u>S. punctata</u>	2	ERIKSSON et al. 1982
c	<u>L. gibba</u>	3	EYSTER 1966
d	<u>L. minor</u>	4	HUMPHREY et al. 1977
g	<u>L. aequinoctialis</u>	5	INGEMARSSON et al. 1984
h	<u>L. minuscula</u>	6	LUEOEND 1983
		7	PIRSON and GOELLNER 1953
		8	THORNSTEINSSON et al. 1985
		9	WHITE 1936b, WHITE and TEMPLEMAN 1937
		10	TANAKA 1986

Characteristics	effect	species	author
area of frond	-	a,d,h	3,6,9
dry weight	-	a	3
length/width ratio	0	a,c,d	6
	+	h	6
gibbosity	0	c	1
length of root	+	a,c,d,h	6,7,9
length of root cells	+	d	7
flowering	+	g	10
turion formation	+	a	3,6
nitrogen content	-	c,g	2
starch content	-	d	9
content of free sugars	0	c	8
anthocyanin content	+	a,c	3,6
chlorophyll content	-	c,d,g	2,9
content of free amino acids	+	d	4
respiration rate	-	c,d	5,7,9
photosynthesis rate	-	c,d,g	2,7
glycolysis rate	-	d	4

low N concentrations. The content of free sugars is not affected (THORSTEINSSON et al. 1985). In L. gibba, the ratio root length / dry weight of the fronds is higher under lower N concentration (INGEMARSSON et al. 1984).

The contents of nitrogen and chlorophyll in the fronds of L. gibba, L. minor, and L. aequinoctialis are proportional to the nitrogen supply. With limited nitrogen, the net CO_2 assimilation rate per frond area is reduced (ERICSSON et al. 1982). The nitrogen content of Lemnaceae, the rate of nitrate uptake as well as the nitrate reductase activities are not dependent on the nitrate concentration in the nutrient solution when medium concentrations of NO_3^- are administered (MUELLER 1983). However, the enzyme activity was species specific. Grown on a medium containing 5 mM NO_3^- , the species showed the following enzyme activities in $\mu\text{mol NO}_3^-$ per g dry weight: 40-53 for S. polyrrhiza and S. punctata, 25-45 for L. turionifera and L. minor, and 10-28 for L. gibba. Maximum nitrate reductase activity in L. minor was achieved at 0.25 mM N, half maximum rate at 0.1 mM (STEWART 1972a).

L. minor does not show aldolase activity in N deficient solutions because an accumulation of low-molecular proteins hampers this process (SARAMEK and DAVIES 1977b). If nitrogen is lacking, proteins are transformed more frequently, and newly formed amino acids are utilized (DAVIES and HUMPHREY 1978). N deficiency reduces the life-span of L. minor fronds (BOESZOEHRNENYI and BOESZOEHRNENYI 1957), it stimulates turion formation of S. polyrrhiza and results in progressive senescence of the fronds (MALEK and COSSINS 1983a).

The optimal nitrogen supply for turion germination of S. polyrrhiza is 1 mM (as KNO_3), according to STBASAKI and ODA (1979). No germination could be observed in solutions without nitrate.

In mixed cultures of natural ponds, the optimal supply of the Lemnaceae with nitrogen is reached between 4 and 8 mg per liter (0.3-0.6 mM N). Higher N concentrations in the water do not raise the N content of the fronds which reaches about 6% of the dry weight (see fig. 1.1, p. 14) (REJMANKOVA 1979, 1981).

2.3.3.4.4.1.4. The effect of nitrate and ammonium; uptake mechanisms

ULLRICH (1987) concludes from a literature survey that Lemnaceae take up NO_3^- ions by nitrate/ H^+ cotransport, and NH_4^+ ions by an ammonium

import which is antagonized by the plasmalemma proton extrusion pump. Therefore the uptake of NO_3^- ions results in an elevation, the uptake of NH_4^+ ions in a reduction of the pH of the solution. However, JUNG-NICKEL and IBRAHIM (1983) report that L. minor is not able to use nitrate in a solution if the pH becomes neutral or weakly alkaline. This is not in accordance with the fact that L. minor and other Lemnaceae species are fast-growing in weakly alkaline solutions if supplied only with nitrate as a nitrogen source (e.g. KOPP et al. 1974a). Several publications deal with the different effect of NO_3^- and NH_4^+ as a nitrogen source on morphological characteristics and metabolism of Lemnaceae. LUEOEND (1980, 1983) investigated the effect of NO_3^- and NH_4^+ as sole nitrogen sources for S. polyrrhiza, L. gibba, L. minor, and L. minuscula. The form of nitrate (KNO_3 , $\text{Ca}(\text{NO}_3)_2$, or NaNO_3) did not produce any difference in the growth rates. A 10-20% higher growth rate was achieved if NH_4Cl , $(\text{NH}_4)_2\text{SO}_4$, or NH_4NO_3 were used. The greatest difference was observed in S. polyrrhiza. The roots of all species were distinctly longer in solution with NO_3^- than with NH_4^+ . Investigations of LANDOLT (unpubl.) resulted in the same direction for roots of S. polyrrhiza, L. gibba, L. disperma, L. minor, L. obscura, L. turionifera, and L. aequinoctialis. The projections of W. microscopica but not the appendages of W. hyalina behaved in the same way. Also for S. punctata it is known that roots become longer in solutions with nitrate than with ammonium (FERGUSON and BOLLARD 1969).

According to MUELLER et al. (1977), the difference in growth rate of L. minor cultivated in nitrate or ammonium solutions depends on the CO_2 concentration of the air. At low concentrations (100 ppm), no difference was observed whereas at high concentrations (9000 ppm) the culture with ammonium showed a slightly higher growth rate. HUBALD and AUGSTEN (1977a) achieved a greater dry weight but only an insignificantly higher growth rate in L. gibba supplied with ammonium than with nitrate. FELLER and ERISMANN (1973) recorded a somewhat higher ash content for L. minor with nitrate. The malate and iron content was enhanced seriously and the nitrogen content lowered (ca. 20% less). If ammonium was the sole nitrogen source, the following compounds were at a higher level in S. punctata (FERGUSON 1966, 1969a, 1970, FERGUSON and BOLLARD 1969): ammonium, arginine, asparagine, and glutamine. Similarly, ERISMANN and KIRK (1969) and MUELLER et al. (1977) observed more glutamine and alanine and less aspartic acid and glutamic acid in L. minor. In cultures with ammonium

as a nitrogen source, *L. minor* produces more free sugar and shows higher specific activity of glycolate oxidase than in cultures with nitrate (EMES and ERISMANN 1982). If supplied with ammonium, one of the primary assimilation products of *L. minor* is glutamic acid (60%) (BAUER et al. 1973). The protein content of *L. minor* is 20-30% higher in ammonium than in nitrate solutions (DICHT et al. 1976). Lowering the temperature from 25°C to 15°C resulted in lower net C fixation and lower oxidation/carboxylation ratio of *L. minor* if supplied with ammonium as the only nitrogen source. In nitrate solutions, only the net C fixation was reduced. Glyoxylate decarboxylation is stimulated by ammonium added to the solution (FUHRER and ERISMANN 1984). Ammonium and glutamine in the medium both activate the glutamine synthetase in *L. minor* (RHODES et al. 1975). The NAD dependent L-glutamate dehydrogenase of *L. gibba* was increased 5 times if the nitrogen source was ammonium or glutamate instead of nitrate (SHEPARD and THURMAN 1973).

The form of the nitrogen has also a different effect on the anthocyanin formation in different species (LANDOLT, unpubl. results). *S. polyrrhiza* and *L. turionifera* show a similar degree of pigmentation in solutions with ammonium and nitrate, *L. gibba* and *L. dispersa* have more anthocyanin in nitrate solutions, and *L. obscura* is most intensely coloured in ammonium solutions. 0.1 mM NH_4^+ added to the solution inhibits flowering of *L. gibba* (KANDELER 1969a) and *L. aequinoctialis* (HILLMAN and POSNER 1971). Many publications deal with the mode of uptake of nitrate, nitrite, and ammonium. Since the uptake of the different forms of nitrogen results in a change of pH, it would be desirable to keep the pH at a constant level in all investigations. This was not done in most studies. This might explain some of the deviating results. According to FERGUSON and BOLLARD (1969) and GROB et al. (1973), *S. punctata* takes up NH_4^+ first, if NH_4^+ and NO_3^- are offered simultaneously. Within the first 10 days 80% of the ammonium but only 20% of the nitrate is used. The preference for ammonium is explained by the fact that the activity of nitrate reductase is inhibited by ammonium. INGEMARSSON et al. (1984) investigated the nitrogen utilization in exponentially growing nitrogen-limited cultures of *L. gibba*. The successive uptake of NH_4^+ and NO_3^- from the solution supplied daily with NH_4NO_3 is shown in fig. 2.8. FELLER and ERISMANN (1971) reported similar results with *L. minor*. Ammonium was taken up first, whether the plants were cultivated in ammonium or nitrate containing solutions before. PORATH and POLLOCK (1982) corroborated

ated these results with *L. gibba*. *S. polyrrhiza* also prefers ammonium if both ammonium and nitrate are supplied (TATKOWSKA and BUCZEK 1983). According to FERGUSON (1966, 1969a, 1970) and FERGUSON and BOLLARD (1969), nitrite can be absorbed together with ammonium. Arginine, asparagine, and glutamine do not influence the activity of nitrate reductase. The nitrate reductase is also present in the frond when nitrogen is lacking in the solution. However, for maximal activity NO_3^- ions are important. Inactivation of nitrate reductase by ammonium was also observed by STEWART et al. (1974) and OREMBAJO and STEWART (1974, 1975a,b). PORATH and AGAMI (1986) noted a correlation between the ammonium (but not the nitrate) content of the water and the occurrence of Lemnaceae in the Negev




Fig. 2.8. Depletion of ammonium and nitrate in the medium by *Lemna gibba* adapted to a relative nitrogen addition rate of 0.25 per day (from INGEMARSSON et al. 1984a)

desert. Since nitrogen and phosphorus were the only chemical parameters studied by the authors the results that the nitrate content is of no importance for the growth of Lemnaceae in nature is not conclusive. JOY (1969a) noted, for L. minor, that the level of nitrate and nitrite reductase remains low in solutions with ammonium or with amino acids. STEWART (1972b) reported similar results. Nitrate reductase in S. punctata is only formed if nitrate or nitrite are present in the solution. High levels of nitrate reductase activity can be induced by the addition of ascorbate to the solution (FERGUSON and KNYPL 1974). Whereas in L. minor nitrate is able to stimulate the formation of nitrate as well as nitrite reductase, the effect of nitrite is restricted to the formation of nitrite reductase (JOY 1969a). No nitrate reductase was observed in W. arhiza either when nitrate as the only nitrogen source was in the solution (SWADER and STOCKING 1971). The nitrate reductase activity of W. microscopica was studied by BAKSHI (1983). GROB et al. (1973) showed that in cultures of S. punctata stimulated by glucose, both NH_4^+ and NO_3^- are absorbed simultaneously (20% NO_3^- and 40% NH_4^+ after 3 days). MELZER and MUELLER (1984) studied nitrate reductase activities of 6 Lemnaceae species using an in vivo test. The activity was strongly enhanced in L. gibba with increasing nitrate concentration in the medium while it was inhibited in L. minuscula. The behaviour of S. polyrrhiza, S. punctata, L. minor and L. turionifera was somewhere between (MUELLER 1983). The nitrate reductase activity of L. trisulca is very low even if it is transferred to water rich in nitrate (MELZER 1980). An enhancement of the nitrate concentration from 0.03 to 0.8 mM did not raise the activity of the nitrate reductase. An addition of 0.3 mM ammonium inhibited the nitrate reductase activity to about 50% after 24 hours. No inhibition was observed in solutions with 0.0006 mM ammonium (MELZER and EXLER 1982, MELZER and MUELLER 1984).

The nitrogen uptake in L. gibba was studied in detail by ULLRICH and co-workers. They observed a different behaviour of nitrogen-starved fronds and fronds supplied with sufficient nitrogen. In fronds cultivated on nitrate solution previously, the NO_3^- uptake was slow and not inhibited by the addition of ammonium. However, nitrogen-starved fronds absorbed nitrate rapidly and ammonium still faster if given separately. An addition of ammonium to the nitrate solution immediately reduced the nitrate uptake to about one third until the ammonium was consumed. Phosphate uptake was inhibited by ammonium similarly to nitrate uptake, but only in

fronds starved of P and N simultaneously not of P alone. The inhibition of the nitrate uptake seems to be due to a decrease of pH rather than of membrane potential depolarisation, and occurs when this decrease cannot be overcome by the H^+ extrusion pump (ULLRICH et al. 1984a,b). LESCH and ULLRICH (1984) observed two phases of ammonium uptake in L. gibba at a pH of 6.5. At the beginning of uptake, the rate is high and scarcely stimulated by light nor impeded by blockers of energy metabolism. The uptake following later is generally lower, in light much higher than in darkness, and sensitive to blockers. The authors conclude that the ammonium uptake is performed by carrier transport as well as by diffusion (at least at pH 7 and higher). The uptake during the first phase of the carrier transport is limited by the bypass through the plasmalemma, during the second phase by the utilization of ammonium within the metabolism. Nitrate reductase activity in S. punctata is 8 times higher in the light than in darkness. Sucrose, glucose, and fructose increase the activity in the light as well as in darkness; DCMU inhibits it completely in the light. Under these conditions ammonium stimulated the nitrate reductase activity whereas amino acids decreased it (VIJAYARAGHAVAN et al. 1982). Contrary to these findings and to results with green algae which absorbed up to 50 times more nitrate in the light than in darkness, the nitrate uptake is not stimulated significantly in L. gibba by light (ULLRICH 1979). The uptake of nitrate in W. arhiza was studied by SWADER and LIN (1975). They detected two different uptake mechanisms: one for low concentrations of nitrate (0.5-0.75 mM N) and another for higher concentrations. The NO_3^- uptake is an active process, 6-methylpurine blocks the nitrate reductase synthesis which results in a reduction of nitrate uptake (TROGTSCH and ULLRICH 1984). According to INGEMARSSON et al. (1985), the uptake of nitrate by Lemna is influenced by the intracellular nitrate concentration. As long as there is a potential for NO_3^- storage in the vacuole, the uptake exceeds the reduction and the nitrate is divided between the readily converted metabolic pool and the storage pool. Once the vacuolar pool is full, the uptake equals assimilation. The nitrate assimilation at low pH is stimulated by Mo (YOSHIMURA 1943).

The complexity of the preferential uptake of ammonium against nitrate is shown in a preliminary experiment with twelve species (LANDOLT unpublished results). Each species was cultivated in solutions with NO_3^- or NH_4^+ as sole nitrogen sources, and with NH_4NO_3 at the same N concentra-

Table 2.14. pH values of culture solutions with nitrate, ammonium-nitrate, and ammonium after cultivation of 14 and 40 days. Initial pH 5.5 (LANDOLT unpubl. results)

Species	no. of clone	pH values	
S. polyrrhiza	7010	7.7	7.2
L. gibba	7135	7.2	7.2
L. dispersa	7223	7.7	7.2
L. minor	7782	7.4	7.6
L. obscura	7856	7.7	7.6
L. turionifera	6853	7.4	7.6
L. aequinoctialis	6746	7.7	7.6
L. minuscula	6600	7.6	7.6
W. hyalina	7376	8.1	8.0
W. microscopica	8359	7.7	8.0
W. australiana	7211	8.0	8.2
W. angusta	7476	8.2	8.2
total		7.2 - 8.2	4.4 - 6.7
NO ₃ ⁻	14	4.6	4.8
	40	4.8	4.6
NH ₄ NO ₃	14	7.0	5.8
	40	5.8	5.8
NH ₄ ⁺	14	4.0	4.2
	40	4.2	4.2
NH ₄ ⁺	14	3.7	3.7
	40	3.7	3.7

tion. The pH which was originally 5.5 was measured after 14 and 40 days. In the solutions with NO₃⁻ alone, the pH rose first to ca. 8 (table 2.14), in solutions with NH₄⁺ it was lowered down to near 4, and in solutions with NH₄NO₃ it behaved somewhere between. If the pH increased in this last solution, it was assumed that preferably NO₃⁻ was used, if it declined, NH₄⁺ was supposed to be taken up first. Only 1 clone of each species was used. The two replicates of each clone did not vary significantly. The correlation of the pH with the form of nitrogen supplied might partly be explained by the uptake of other ions too. But since all species were cultivated under the same conditions the differences might be species specific. According to the pH in the solution after 14 days, the tested species can be divided into three groups (in brackets the pH difference):

- 1) Species with a pH only slightly lower in the NH₄NO₃ solution than in the NO₃⁻ solution (pH difference 0.0-0.5). It is supposed that these species use a great deal of NO₃⁻ beside NH₄⁺ from the beginning under the given conditions: W. angusta (0.0), W. microscopica (0.4), L. obscura (0.4), W. australiana (0.5).
- 2) Species with a slightly lower pH in the NH₄NO₃ solution than in the nitrate solution (pH difference 0.7-0.8). These species probably take up both ammonium and nitrate to a moderate extent, S. polyrrhiza (0.7), L. minor (0.7), L. aequinoctialis (0.8).
- 3) Species with a distinctly lower pH in the NH₄NO₃ solution than in the nitrate solution (pH difference 1.0-1.9). These species seem to absorb ammonium preferentially: L. turionifera (1.0), L. dispersa (1.2), L. minuscula (1.3), L. gibba (1.4) and W. hyalina (1.7). It takes 20 to 30 hours for L. minor to change from ammonium uptake to nitrate uptake (KOPP et al. 1974b). The NO₃⁻ uptake is not linked with the potassium uptake in L. gibba through a common ion permeating system in the cell membrane. NO₃⁻ and K⁺ uptake are separately controlled by oscillators (KONDO 1982c). HILLMAN and LAMM (1981) report a peculiar behaviour of L. aequinoctialis (clone No. 6612) towards nitrogen utilization. In solutions with KNO₃ as sole N source they observed from time to time chlorotic fronds which occurred spontaneously. If the chlorotic fronds were transferred to solutions with nitrogen in a reduced form (e.g. aspartic acid, glutamic acid, glutamine, NH₄Cl, NaNO₂), they became green. The inability of nitrate assimilation can appear in the light or darkness and in different

nutrient solutions. The phenomenon does not seem to be a result of a somatic mutation.

A mutant of *L. aequinoctialis* incapable of ammonium ion uptake and of photosynthesis was detected by POSNER (1962b) and investigated by MONSE-LTSE et al. (1984).

2.3.3.4.1.5. Utilization of organic nitrogen

Lemnaceae are able to utilize many organic nitrogen compounds besides ammonium, nitrate, and nitrite (see also chapter 2.3.3.5.2.2). From more than 160 compounds tested by BOLLARD (1966), the following could be used successfully by *S. punctata* as sole nitrogen source:

- protein amino acids: L-alanine, L-aspartic acid, L-asparagine, L-glutamic acid, L-glutamine, glycine, and 6 other rarer amino acids and substituted amino acids;
- peptides: glycyl-L-alanine and 6 further compounds;
- amides: L-serine amide;
- amines: putrescine;
- guanidine derivatives: agmatine, β -guanido propionic acid;
- urea and urea derivatives: 7 compounds.

No purines and pyrimidines could be used successfully. The absorption of urea in *S. polyrrhiza* is much slower than that of nitrate or ammonium (EYSTER 1966). The optimal amount of urea for a constant (but rather slow) growth lies between 30 and 150 mg (corresponding to 15-60 mM N). If the supply with nitrate nitrogen is sufficient, an addition of 15 mM urea reduces the growth rate of *S. polyrrhiza*. *S. punctata* is only able to use urea if the pH is below 4.3 (BOLLARD et al. 1968). If urea is used as nitrogen source, the concentrations of many amino acids in the fronds of *S. punctata* are lower than if ammonium is supplied. Asparagine and glutamine are taken up by *S. punctata* faster than nitrate, and aspartate and glutamate at about the same rate (FERGUSON 1970). Glutamic acid and asparagine allow only a slow, and glycine a moderate growth of *L. gibba* if applied as sole nitrogen source. L-arginine and L-aspartic acid together as well as L-asparagine alone (in concentrations of 1 mM N) are able to promote the growth rate in solutions supplied with nitrate (HUBALD and AUGSTEN 1977a). The growth rate of *L. minor* was found to be higher in nutrient solutions with a mixture of amino acids than with nitrate and ammonium (JOY 1969a). According to the same author,

subsequent better growth was attained with the following mixtures: glutamic acid-aspartic acid-arginine, casamino acids, casamino acids-nitrate. The positive influence of the amino acid mixture might be due to inadequate nutrient or light conditions. Certain (branched-chain) amino acids as leucin, valine, and isoleucine inhibit the growth rate of *S. polyrrhiza* if used alone (BORSTLAP 1974, VAN MAZIK 1975). Contrary to the results with *Spirodela* and *L. minor*, *L. trisulca* grows better with urea than with nitrate or ammonium. Also aspartic acid and glutamic acid are easily used as nitrogen sources (HOLST and YOPP 1979). The permeability for urea of roots of *L. minor* is exceptionally high compared with other plants (MARKLUND 1936).

2.3.3.4.2. Phosphorus (P)

Phosphorus is taken up mostly as phosphate. According to FITZGERALD (1971), there is also the possibility for *L. minor* to use P as pyrophosphate, triphosphosphate and as P bound in teeth of sharks. A utilization

Table 2.15. Phosphorus requirement and phosphorus tolerance for growth of some Lemnaceae species (minimum amount of phosphorus to result in half of the maximum growth rate, optimum amount of phosphorus to achieve maximum growth rate, maximum amount of phosphorus to achieve still half of the maximum growth rate), from AEBLI 1986 (1), DOCAUER 1983 (2), EYSTER 1966 (3), LUEOEND 1983 (4), FERNANDEZ et al. 1983 (5).

* forming turions at lower concentrations

Species	supply of P in mM		
	minimum	optimum	maximum
<i>S. polyrrhiza</i>	0.00046 (2)	0.1 -1.0 (3)	1.75 (4)
<i>L. gibba</i>	0.0028 * (4)	0.014 -0.35 (4)	10 (2)
<i>L. minor</i>	0.00056 (4)	0.0028-1.75 (4)	8.76 (4)
<i>L. turionifera</i>	0.00045 (2)	0.014 -0.35 (4)	1.75 (4)
<i>L. aequinoctialis</i>	0.00011 (4)	0.01 ->0.2 (1)	
<i>L. minuscula</i>	0.00035 (2)		<80 (5)
<i>W. borealis</i>	0.00011 (4)	0.0028-0.35 (4)	1.75 (4)
<i>W. columbiana</i>	0.00552 (2)		
	0.00148 (2)		

of P from lake mud is possible only to a small amount (less than 1%). P released from pond sediments is either more or less easily available for L. minor than KH_2PO_4 depending on the type of sediment (PEKETE et al. 1976).

Minimum, optimum, and maximum P content of the nutrient solution for different Lemnaceae species are put together in table 2.15. The need for phosphorus rises in the following order: L. minuscula, L. turionifera, L. minor, S. polyrhiza, L. gibba, W. columbiana, W. borealis. Relatively high optimum and maximum values are found in L. gibba and S. polyrhiza. Interestingly, the requirement of phosphorus for the different species is in no correlation to the nitrogen requirement. The ratio of the half saturation constants for nitrate and for phosphate have been calculated by DOCAUER (1983) as follows: L. turionifera (16.9), S. polyrhiza (13.9), L. minor (11.8), W. borealis (3.0) and W. columbiana (1.6). The high P requirements of W. borealis and W. columbiana especially if compared with the N requirements are remarkable.

The influence of the low P concentration on characteristics of Lemnaceae is surveyed in table 2.16. Interestingly, some species behave differently. The length/width ratio becomes higher at low P concentrations in L. minuscula and L. gibba, it does not change in S. polyrhiza and L. minor and it becomes lower in L. turionifera. The root length is greatest at 0.003 mM P in L. gibba, at 0.0005 mM P in L. gibba and L. minuscula, and at 0.00002 mM P in L. minor (LUEOEND 1983). Remarkably, L. minor develops longer roots than L. gibba in solutions with a low phosphorus content whereas L. gibba has longer roots than L. minor in solutions with a low nitrogen content. At high concentration of P (as well as of N), the fronds of Lemnaceae remain together longer and often form clusters (LUEOEND 1983). The stipe between mother and daughter fronds becomes longer and distinctly visible. P and N shortage mostly result in similar reactions of the plants, they have, however, an opposite effect in comparison to K shortage.

The number, size, and dry weight of S. polyrhiza turions is higher in solution with a low phosphorus content (0.5 mM) than in solution with a high P content (1.5 mM). Turions formed under low phosphorus conditions are extremely dormant (JUNGnickel 1986a). DUDLEY (1983, 1987) was able to encourage turion formation of L. turionifera by additional phosphorus.

According to LINDENMAN (1951, 1952, 1972), P shortage lowers the photo-

synthesis rate of L. minor under light saturation. If phosphate is added, the photosynthesis is normalized after a short time. Dark-red light is able to partly compensate the inhibiting effect of P shortage (LINDENMAN 1973). The lack of phosphate results in a 50-fold increase in phosphate activity of all cell extracts of S. punctata (REID and BIELESKI 1970b) (compare chapter 2.5.6.3). In L. minor, STROTHER (1984) showed that acid phosphatase activity increases in phosphorus-deficient solutions; alkaline pyrophosphatase activity decreases during senescence;

Table 2.16. The effect of low phosphorus supply on characteristics of Lemnaceae

* at very low P concentrations the roots are shorter again (5)
+ increase, - decrease, 0 no effect
1 only in solution with sugar

- | | | | |
|---|--------------------------|----|-----------------------------|
| a | <u>S. polyrhiza</u> | 1 | BORNKAMM 1965 |
| b | <u>S. punctata</u> | 2 | EYSTER 1966 |
| c | <u>L. gibba</u> | 3 | JUNGnickel 1978 |
| d | <u>L. minor</u> | | AUGSTEN and JUNGnickel 1983 |
| e | <u>L. turionifera</u> | 4 | LINDENMAN 1951, 1952, 1972 |
| f | <u>L. trisulca</u> | 5 | LUEOEND 1983 |
| g | <u>L. aequinoctialis</u> | 6 | PIRSON and GOELLNER 1953 |
| h | <u>L. minuscula</u> | 7 | REID and BIELESKI 1970a,b |
| i | <u>W. arizhiza</u> | 8 | THORNSTEINSSON et al. 1985 |
| | | 9 | DOCAUER 1983 |
| | | 10 | STROTHER 1984 |

Characteristics	effect	species	author
area of frond	-	a,d,h	2,5
dry weight	-	a,d	1,2
length/width ratio	0	a,d	5,9
		e	9
gibbosity	+	c,h	5
length of root	0	c	5
length of root cells	+	a,b,c,d,h	5,6,7
turion formation	+	d	6
starch content	+	a,i	2,3,5
content of free sugars	+	b,c	7,8
anthocyanin content	-	c	8
protein content	+	a,c,f	2,3,5
content of glutamine and asparagine	0	d	3
oxalate content	+	b	7
respiration rate	-	d	1
photosynthesis rate	-	d	6
phosphatase activity	+	b,d	4,6,7
		b,d	7,10

this decrease is enlarged by phosphorus-deficiency. Apparently, a certain amount of phosphorus is necessary for flower induction of L. gibba under long-day conditions (CLELAND and TANAKA 1986). JUNGNIKE (1978) calculated the P requirement to form a frond of 4 different Lemnaceae species. The following amount of P is needed: 0.00034 mg for S. polyrrhiza, 0.00018 mg for L. gibba, 0.00014 and 0.00004 mg for two different clones of L. minor and 0.00001 mg for W. arhiza. According to EYSTER (1966), S. polyrrhiza requires a minimum content of 3.3 mg P to form 1 g dry weight. FEKETE and RIEMER (1973) reached the maximal P content (1.9%) in fronds of L. minor with a P concentration of 0.1 mM in the medium. Of 10 species of water plants, L. minor accumulated most P of all vascular plants (measured with ^{32}P), but much less than the filamentous alga Oedogonium (VERMAAK et al. 1976).

S. punctata is able to transport PO_4^{3-} from older to younger fronds if kept under P starvation (KNYPL 1979b). A mutual exchange of phosphorus between the plants takes place (KNYPL 1980). BIELESKI (1968a,b) studied the uptake and the incorporation of radioactive P in S. polyrrhiza. He analysed more than 50 different chemical P compounds. In P starvation cultures, the growth stops after 14 days, and the portions of the different P compounds change. Old fronds become chlorotic and young ones accumulate anthocyanin and starch; roots become longer and the photosynthesis rate is reduced, but the nitrogen balance does not change. The P uptake under P shortage in L. gibba was investigated by ULLRICH-EBERIUS et al. (1981), ULLRICH-EBERIUS and NOVACKY (1984). The uptake of P is stimulated by light. It performed best at a pH of 6 and slowed down with rising pH. According to EL-SHINAWY and ABDEL-NALIK (1980) the uptake is much faster still at pH 5 than at pH 6. After P starvation, the uptake is better. The P uptake is supposed to be directed by the intracellular P content. The monovalent phosphate uptake proceeds by H^+ cotransport (see chapters 2.5.2.1 and 2.5.2.2.4).

The optimal supply of P in a mixed culture of Lemnaceae in natural waters is supposed to vary between 0.13 and 0.26 mM. At higher P concentrations (up to 1.3 mM), the P content of the fronds (about 1%) did not rise any longer (see fig. 1.1, p. 14) (REJMANKOVA 1979).

2.3.3.4.4.3 Sulfur (S)

Sulfur is mostly taken up as sulfate. Lemnaceae are also able to absorb

sulfide, thiosulfate, sulfite, cysteine, and cysteic acid. Other substances containing sulfur investigated (e.g. dithionate, tetrathionate, thiourea, methionine, sulfanil amide, sulfanilic acid) either did not have any effect on growth of S. punctata at concentrations of 1 mM S, or the growth rate was reduced (FRASER 1974).

Best growth of S. polyrrhiza was achieved with 0.5 to 20 mM S (EYSTER 1966). At a sulfur concentration of 0.05 mM, the growth rate was distinctly lower, the fronds bigger, heavier, and more intensely pigmented by anthocyanins. At 0.01 mM the fronds looked chlorotic. From 60 mM S upwards the plants stopped growth, appeared small and partly chlorotic, and clustered together. Sulfur deficiency in the medium stimulates tumor formation in S. polyrrhiza (MALEX and COSSINS 1983a). The increase of the sulfate content from 0.014 to 1 mM did not change the growth rate of L. aequinoctialis (DATKO et al. 1980a). This difference in S requirement between S. polyrrhiza and L. aequinoctialis is striking even if we consider the different culture conditions. The sulfur requirement to form one g dry weight of S. polyrrhiza was estimated by EYSTER as 0.1-0.5 mg at minimum. For optimal growth the amount of sulfur needed per g dry weight is 10 mg. DATKO et al. (1978a) calculated the amount of sulfur necessary to build up 1 g protein in L. aequinoctialis as 9-12 mg. The requirement is higher if the molybdenum content of the solution is relatively high (DATKO et al. 1980a). Molybdate and sulfate are taken up by a common mechanism (DATKO and MUDD 1984a).

THOIRON et al. (1969) investigated the sulfate uptake of L. minor under different temperature conditions. The authors postulate the existence of two different systems of sulfate absorption. The uptake of sulfate in L. aequinoctialis was studied by DATKO et al. (1978a,b) within the range of 0.0003 and 1 mM S. No volatile S compounds could be detected. Also, no S was transferred from the fronds back to the solution even at the highest SO_4^{2-} concentrations. The content of S-containing amino acids increased only 1 1/2 to 2 times between the lowest and the highest S concentration of the medium, whereas the anorganic sulfate increased to the 30fold. At least 2 components are involved in the uptake of SO_4^{2-} , one saturating and one linear (non-saturating). 55% of newly taken up SO_4^{2-} enters a compartment with a slow turnover (vacuole?), 45% remains in a compartment (cytoplasm?) in which it is rapidly metabolized to organic components. The saturating system contributes 99% of SO_4^{2-} uptake under standard conditions. In nature, the same system shares at least

65-70% (DATKO and MUDD 1984a). The down-regulation of the sulfate uptake of the saturating system by high concentrations of SO_4^{2-} or cystine is more than compensated by unregulated uptake via the non-saturating uptake system (DATKO and MUDD 1984b). The kinetics of SO_4^{2-} uptake in *L. gibba* was studied by LASS and ULLRICH-EBERIUS (1984a,b). Further details are given in chapters 2.5.2.1 and 2.5.2.2.4.

The sulfide uptake (and also the uptake of gaseous H_2S) was investigated by ERISMANN et al. (1967), BRAENDLE and ERISMANN (1968), BRUNOLD and ERISMANN (1970, 1974, 1975, 1976). Two kinds of sulfide uptake could be detected, one of which is dependent on photosynthesis and leads mostly to fixation of S within the chloroplasts. Sulfide in the medium slightly inhibits photosynthesis and reduces the SO_4^{2-} uptake. This could be demonstrated by gassing the cultures with H_2S .

TAKEMOTO and NOBLE (1984, 1986) studied the growth response of *S. punctata*, *L. gibba*, and *L. valdiviana* to sulfite enrichment under light-limited and light-saturated conditions. Under low light intensity (35 $\mu\text{mol m}^{-2}\text{s}^{-1}$ photosynthetic photon flux density), *L. gibba* and, to a lesser extent, *S. punctata* showed reduced growth rates at 5 and 10 mM K_2SO_3 . Inhibition was less pronounced under high light intensities (200 $\mu\text{mol m}^{-2}\text{s}^{-1}$ photosynthetic photon flux density). No growth reduction was observed for *L. valdiviana*. Since rates of growth and photosynthesis of *L. valdiviana* were higher than of *S. punctata* and *L. gibba*, the authors assume that rapid metabolic and 'mitotic activity are important to resist sulfite action. In a later paper (TAKEMOTO et al. 1986), it was demonstrated that the thiol content in the three investigated Lemnaceae species increased by 30-40% with sulfite treatment under low light intensity. H_2S was emitted by all three species, however, emission rates in *L. valdiviana* which is sulfite-tolerant were up to 4 times higher than in the other species. The thiol content and the H_2S emission of all species was enhanced at high light intensity. The increase of emission rate of H_2S was greater in *L. valdiviana* and *S. punctata* (55-60%) than in *L. gibba* (17%). It was proposed that thiol production and H_2S emission are important sulfite detoxification processes in duckweeds.

2.3.3.4.5. Carbon (C)

Lemnaceae are able to use carbonate and bicarbonate in the solution as carbon source if CO_2 is a minimum factor (WOHLER 1966, WETZEL and MANNY

1972). Under normal conditions, Lemnaceae floating on the surface of the water take up most of the C needed via gaseous CO_2 from the air. Only a small amount of inorganic carbon originates from water soluble carbon, according to BOWKER et al. (1980) who studied *L. minor*. FILBIN and HOUGH (1985) report that *L. minor* takes, from a hard water lake (with relatively high pH), on an average 86% carbon from aqueous inorganic carbon. Under laboratory conditions, the C uptake from water was lower (37%). SATAKE and SHIMURA (1983) measured in *S. polyrrhiza* at low pH (5 and 4) up to 10 times more C uptake from the water than from the air. However, ESHEL and BEER (1986) questioned the results of SATAKE and SHIMURA. These authors did not take into account the different solubility of carbon compounds in water. The dissolved inorganic carbon form varies as a function of pH (fig. 2.9, from ESHEL and BEER 1986). ESHEL and BEER observed under laboratory conditions at a pH of 5.9 about 1% C uptake from the water and at a pH of 7.2 and 8.0 about 4-5% (see also table




Fig. 2.9. The relative distribution of dissolved inorganic carbon among its various chemical forms as a function of pH (from ESHEL and BEER 1986)

2.6). Certainly, some of these differences are due to different growing conditions. For more details see chapter 2.3.2.2.

LANDOLT (1957) reported that L. trisulca grows in aseptic cultures of Hutner 1/5 solution only if sucrose is added. If distilled water is replaced by tap water, a slow growth without sucrose is possible. BATA (1973) succeeded in growing L. trisulca by adding kinetin. At high pH, the growth took place without kinetin. BATA concluded that L. trisulca is only autotrophic at higher pH. Also LANDOLT (1957) was originally of the opinion, that L. trisulca is heterotrophic for some special agents. The explanation of the "heterotrophy" of L. trisulca is the fact that Hutner solution does not contain any carbonate in contrast to the solution used by BATA (1973) and to tap water. The diffusion of CO_2 in the water is 10000 times slower than in the air. Submerged species in aseptic cultures do not get enough CO_2 for good growth. In nature, the CO_2 in the water is constantly replaced by respiring heterotrophic organisms in addition to the carbonate which is present in the water. A preliminary experiment demonstrated the correctness of the assumption (LANDOLT unpubl. results). If NaHCO_3 or Na_2CO_3 is added to Hutners solution containing no sugar, L. trisulca looks healthy and grows steadily at pH 6 and higher. Also W. gladiata and other Wolffiella species which grow extremely slowly in solutions without sucrose show a more intense green colour and a much faster growth if carbonate is added to the solution. Already STEWMANN NIELSEN (1944, 1947) observed that the CO_2 content in water (0.015 mM) is not sufficient to enable photosynthesis of the submerged species Myriophyllum spicatum, Ceratophyllum demersum and Elodea canadensis. Therefore, these species are able to utilize bicarbonate. They reach 30% of the maximal growth rate with 0.5 mM HCO_3^- and 50% with 1 mM. The bicarbonate remains in solution only at higher pH (6.5 to 9); at lower pH CO_2 is formed which escapes. Therefore, a relatively high pH is necessary for a good growth of the submerged species mentioned. For L. trisulca and W. gladiata, a high carbonate content of the water or a high content of organic substances is a precondition for the occurrence in nature (see also vol. 1, chapters 4.2.4.8 and 4.2.4.12, LANDOLT 1986). If no bicarbonate is in the water, the photosynthetic rates of submerged species are likely to be limited by the rate of transport processes outside the photosynthetic cells (RAVEN 1981). KHCO_3 (1 mM) is able to induce flowering in the long-day plant L. gibba under short-day conditions but inhibits flowering under long-day conditions (KANDELER 1964a,b).

2.3.3.4.6. Chlorine (Cl) and other halogens (F, Br, J)

The indispensability of chlorine was demonstrated by MARTIN and LAVOLLEY (1958a,b, 1959) and LAVOLLEY (1963a,b, 1965). Chlorine can partly be substituted by bromine. The minimum requirement of Cl rises with higher light intensity. The Cl^- ions are not only used for photosynthesis but also for other independent processes. This was concluded from the fact that Cl is also necessary for heterotrophic growth in darkness. The minimum requirement of Cl for best growth is species specific (MARTIN 1963a, 1965). It amounts to 0.015 mM Cl for S. polyrrhiza, 0.001 mM for L. minor and 0.003 mM for L. aequinoctialis. EVSTER (1966) achieved best growth for S. polyrrhiza in Cl concentrations between 0.01 and 1 mM. At 10 mM Cl the growth rate is distinctly lower and at 100 mM the fronds die. At low Cl concentrations (0.001 mM), the fronds are relatively narrow and the older ones have red stripes. At still lower concentrations, they become very small, light green, and form clusters. The greatest dry weight per frond is reached at 0.01 mM Cl. At higher concentrations it reaches almost the same amount.

In an experiment of our own, the positive effect of NaCl addition on growth of Lemnaceae has been confirmed (LANDOLT, unpubl. results). Lemnaceae species are normally kept in our laboratory either on Hoagland agar medium (which does not contain Na and only 0.02 mM Cl) or on 1/5 Hutner agar medium (which contains only 0.02 mM Na if the sodium free complexon 2 (EDTA) is used, and 0.04 mM Cl. If 0.3 mM NaCl is added to either of the 2 solutions, the growth rate of all 4 investigated species (S. polyrrhiza, L. minor, W. Welwitschii and W. microscopica) is at first much faster than in the normal solutions. After two weeks, the species were transferred to the normal solutions again. S. polyrrhiza and L. minor kept the same elevated growth rate whereas W. polyrrhiza and W. microscopica slowed down their growth rate significantly. It is not quite sure if this effect is due to the Na^+ or Cl^- content or whether it is a combined effect. It seems more probable, that the Cl content is decisive. This would mean that for long-lasting good growth a Cl content of 0.02 mM is not sufficient (Cl concentration of Hoagland solution). On the other hand, the usual impurities of the chemicals with Cl are sufficient for a slow growth. Also it seems evident that W. Welwitschii and W. microscopica need more NaCl than S. polyrrhiza and L. minor.

Iodine cannot replace chlorine, in contrast to bromine. It is more toxic in solutions without chlorine than in solutions with chlorine (in Cl concentrations from 10^{-4} mM upwards). Iodine is probably not a necessary element for L. minor (MARTIN and LAVOLLEY 1959).

The absorption of chlorine in L. minor is 10 to 20 times faster than that of iodine (THELLIER et al. 1967). The authors are of the opinion that the uptake of chlorine is more complicated than the iodine uptake. In the first case, they assume different uptake systems for different Cl concentrations in the solution and for different environmental conditions; in the second case, they observed only one uptake system for all conditions.

Sodiumfluoride inhibits the growth rate of S. punctata and L. minor at concentrations of 5 mM; it is lethal at 20 mM (BONG et al. 1980).

2.3.3.4.7. Metallic trace elements

2.3.3.4.7.1. General remarks

Of all the natural elements which are not mentioned in the previous chapters of the present volume, a few are essential for the growth of Lemnaceae, others stimulate growth but are not indispensable and still others are indifferent or toxic, according to the concentrations used. The proof of the essentiality of a trace element is rather difficult because the required amounts are small; very often a sufficient quantity of a trace element is available from impurities in the culture vessels, in the chemicals, or in the water. Furthermore, the Lemnaceae are very economical with trace elements, transferring them from mother to daughter fronds. The example of HENSSEN (1954) might illustrate the situation. The author used a nutrient solution without zinc and molybdenum, both elements proved to be essential for Lemnacean growth. She still recorded growth of S. polyrrhiza after a year of cultivation without these elements. However, some deformities could be observed.

The availability of many trace elements is dependent on the pH of the solution. If chelating agents (e.g. EDTA, tartrate, citrate) are added, the uptake of trace elements becomes easier, and the toxicity is not so pronounced (e.g. Fe, Mn, Zn, Cu) (cf. FLY 1935, SIELING 1937, HUTNER et al. 1950, BITCOVER and SIELING 1951). Therefore, the comparison of results within this chapter is difficult. Some of the elements listed in

Table 2.17. Tolerance of Lemna minor towards different micronutrients (concentrations in mg per l) (from FRICK 1985b)

Growth	Fe	Mn	B	Cu	Zn	Mo
no inhibition	27.9	54.9	17.3	3.2	6.5	54.4
complete inhibition	100.5	274.5	86.5	6.3	65.3	271.8

the following chapters are not necessary for the growth of Lemnaceae. However, all elements are mentioned, of which the effects on Lemnaceae were studied. In general, Lemnaceae react relatively sensitively to different heavy metals compared to other plants (e.g. Elodea) (BROWN and RATTIGAN 1979), see also chapter 3.6.

FRICK (1985b) tested the tolerance of L. minor to several micronutrients (table 2.17). A combination of the high concentrations of different micronutrients (except Mo) were not inhibitory, even though separate 2-, 3-, and 4-way combinations were.

Some metal ions (e.g. Cu, Ag, W) promote flowering of Lemnaceae. TANAKA et al. (1986a) suggest that this effect is due to the suppression of nitrate assimilation by these ions.

2.3.3.4.7.2. Boron (B) and aluminum (Al)

Boron was shown to be essential for S. polyrrhiza (EYSTER 1966), S. punctata (THIMANN and EDMONSON 1949), L. minor (STEINBERG 1946) and W. arrhiza (EICHORN and AUGSTEN 1974). The minimal amount for good growth of L. minor is 0.005 mM B (STEINBERG 1946), and for S. polyrrhiza 0.1 mM B (EYSTER 1966). At lower concentrations the growth of S. polyrrhiza was slower, and signs of starvation occurred (yellowish colour with darker stripes). Even at a B concentration of 0.0001 mM, a distinct growth rate could be observed although the cultures were kept in boron-free solution for several weeks before. GLANDON and McNABB (1978) did not note any difference in growth rate of L. minor in solutions with 0.001, 0.01, and 0.1 mM B. The B content of the fronds amounted to 0.1, 0.6, and 0.8 mg per g dry weight, respectively. Similar results were obtained by FRICK (1985a). Best growth of W. arrhiza was achieved with 0.008 mM B. At

0.004 mM B, signs of starvation occurred (EICHORN and AUGSTEN 1974). The toxic limit of B for S. polyrrhiza is 10 mM (EYSTER 1966), for L. minor 5 mM (THELLIER and LE GUIEL 1967a), and for W. arhriza somewhat higher than 0.02 mM (EICHORN and AUGSTEN 1974). It is not clear if the different requirements and tolerances are species specific or due to different growth conditions and (partly) due to different concentration intervals between the investigated solutions.

Shortage of boron results for W. arhriza in a reduction of dry weight, chlorophyll, and protein content as well as in an elevation of the dark respiration rate, of the photosynthesis rate, and of the carbohydrate content. With boron deficiency, the ratio of glycolysis/pentoses shunt pathway is changed in favour of the latter (EICHORN and AUGSTEN 1974). The protein content of L. minor is reduced, the starch content raised (from 3% to 6.5% of the dry weight) and the fronds clustered if boron is lacking (SCHOLZ 1962). The shortage of boron has a greater effect with nitrate than with ammonium as a nitrogen source. EYSTER (1966) therefore raises the question if boron plays some role in the uptake or assimilation of the nitrate. High light intensities aggravate the boron starvation symptoms, low light intensities moderate them (TANAKA 1966).

In clay suspensions, B concentrations of as low as 0.01 mM inhibit growth of L. minor. However, the toxic effect can be neutralized by addition of Ca (FOX and ALBRECHT 1958). The highest boron content in the frond (4600 ppm B on a dry weight basis) was achieved with 1 ppm B (0.1 mM) and relatively high Ca content. Also BOLGIANO (1979) reports on a more pronounced toxicity of high boron concentrations if the Ca concentration is low. Compared with Ceratophyllum demersum, L. minor takes up much more boron (up to 45 times) (GLANDON and McNABB 1978). The boron uptake is active, according to THELLIER and LE GUIEL (1967a). There seem to be at least four disparate functions of boron (according to FRICK 1985a): in nucleic acid metabolism, in carbohydrate biosynthesis and transport, in interactions with plant growth substances, and in membrane integrity.

Borate does not considerably influence the absorption of sulfate in L. minor (THELLIER and TROMEUR 1968).

Aluminum is not essential for Lemnaceae. It has a toxic effect on S. polyrrhiza and L. minor at a concentration of 1 mM (WIEWIORKA and SAROSIEK 1986).

2.3.3.4.7.3. Molybdenum (Mo) and tungsten (W)

The minimum amount of molybdenum required by L. minor is 0.0002 mM (STEINBERG 1946). EYSTER (1966) was not able to demonstrate the necessity of Mo for S. polyrrhiza, probably due to impurities in his chemical substances. However, YOSHIMURA (1943) proved that S. polyrrhiza (as well as different Lemna species) needs at least 5×10^{-7} mM Mo. Ten mM Mo are toxic to S. polyrrhiza (EYSTER 1966). YOSHIMURA (1943) obtained effects of toxicity of Mo already at 1 mM for S. polyrrhiza and Lemna spp.

Shortage of Mo resulted in flowering of L. aequinoctialis (YOSHIMURA 1943). This result was corroborated by TANAKA et al. (1979b). Root growth of L. gibba is inhibited at 0.1 mM Mo and completely stopped at >1 mM (LIEBERT 1986a).

Mo stimulates the nitrate assimilation at low pH, enhancing the reduction of nitrate and preventing the accumulation of nitrate in the fronds. If Mo is lacking in the solution, the Lemnaceae cultures die in a medium with NH_4NO_3 since they only take up ammonium and in this way lower the pH to a toxic value. According to DATKO et al. (1980a) and DATKO and MUDD (1982), Mo competes with sulfur. Therefore, an addition of higher amount of sulfate or cystine is able to detoxicate the Mo. At low sulfate concentration in the solution, Mo is toxic in concentrations as low as 0.05 mM for L. aequinoctialis. DATKO and MUDD (1980) suppose that Mo inhibits formation of reduced sulfur in L. aequinoctialis.

Tungsten (W) is said to have a similar effect on Lemnaceae as Mo (YOSHIMURA 1943). However, tungstate is a competitive inhibitor of molybdate function and was able to induce flowering in L. aequinoctialis similarly to Cu, Hg, Ag, and cyanide ions (TANAKA et al. 1979b, 1986a). It is assumed by these authors that the ions mentioned inhibit nitrate reductase activity and consequently nitrate assimilation, and that this suppression causes day-length-independent flowering.

2.3.3.4.7.4. Iron (Fe)

The fronds of S. polyrrhiza become chlorotic and red underneath below a concentration of 0.001 mM Fe (EYSTER 1966). Also YOSHIMURA (1950) noted chlorotic fronds of S. polyrrhiza under shortage of iron. The fronds accumulated anthocyanins and oxalate and formed turions. Optimal growth of S. polyrrhiza was observed by EYSTER (1966) between 0.01 and 0.1 mM

Fe. Under optimal conditions of Fe, the fronds reach up to 11.5 mm in length. The minimum amount of Fe for good growth of L. minor was determined by STEINBERG (1946) as 0.05 mM. According to ORLANDO and NEILANDS (1982), maximum growth rate of L. gibba and L. minor was obtained with an iron supply of 0.02 mM (added as ferrichrome or ferric EDTA). The maximum iron content was reached at 10 mM in S. polyrrhiza (EYSTER 1966). Already at 1 mM Fe, S. polyrrhiza showed a slow growth and light green fronds. SCHREINEMAKERS (1984) obtained an inhibition of S. polyrrhiza with 0.2 mM Fe and nearly a stop of growth at 0.4 mM. DOCAUER (1983) measured the Fe content of ponds with different Lemnaceae cover. He concluded a different need of Fe for Lemnaceae species. According to his observations, L. turionifera and L. minor require more iron than S. polyrrhiza and W. columbiana. EYSTER (1966) estimated the amount of Fe required to produce one gram dry weight of S. polyrrhiza as 0.017 mg.

A flower-promoting effect of iron in L. aequinoctialis was reported by NISHIOKA et al. 1986. For further effects of iron on flowering see chapter 2.4.3.1.2.

2.3.3.4.7.5. Manganese (Mn)

The essentiality of manganese for Lemnaceae was already proved by HOPKINS (1931) (for L. minor), MCHARGUE and COLFUE (1932) (for S. polyrrhiza), CLARK (1933) (for S. polyrrhiza), and SAEGER (1933a) (for S. polyrrhiza). S. polyrrhiza needs at least 10^{-5} mM Mn. At this concentration, the fronds were small, rich in anthocyanin and clustered. Shortage of Mn depresses root growth in S. polyrrhiza (SAEGER 1933a) and L. minor (HOPKINS 1931). Maximum growth was attained in S. polyrrhiza between 10^{-3} and 0.06 mM Mn (EYSTER 1966) and between 0.009 and 0.047 mM (CLARK 1933). L. minor requires 10^{-3} mM Mn (STEINBERG 1946), L. minuscula 10^{-2} mM Mn, (JUNGnickel and GEBHARD 1986) for good growth. Much lower key concentrations were measured by YOSHIMURA (1941) for S. polyrrhiza, L. aequinoctialis and L. valdiviana: 5×10^{-7} mM Mn was the minimal amount for growth and 10^{-5} mM was needed for optimal growth. The maximum concentration to still allow growth for S. polyrrhiza is given by EYSTER (1966) with 0.3 mM Mn. At 3 mM Mn the plants stopped growth, were very small, clumped, and pale green. L. minuscula showed toxic effects at 1 mM Mn (and 50% of the maximum growth rate, JUNGnickel and GEBHARD

1986). According to NASU and KUGIMOTO (1981) and NASU (1983), Mn concentrations of 0.02 mM inhibit the growth of L. aequinoctialis to 50% (Bonner-Devirian medium). Reduced growth (50% of the maximum growth rate) was obtained by JUNGnickel (1984) at Mn concentrations of 0.1 mM for L. aequinoctialis, at 1 mM for S. polyrrhiza, W. arthiza, and L. minuscula, the latter species developing the highest tolerance against Mn. The root growth of L. gibba is inhibited at 0.2 mM Mn and stopped completely at 1 mM (LIEBERT 1986a).

EYSTER (1966) estimated the minimal amount of Mn required to form one gram dry weight of S. polyrrhiza as 0.0033 mg. The Mn requirements are higher in darkness than in the light (EYSTER 1966).

Under mixotrophic conditions using nitrate as sole nitrogen source, the minimum Mn concentration for good growth of L. minuscula was 10 times lower (10^{-3} mM) than in autotrophic cultures. The minimum Mn requirement per frond amounted to 1 pg compared with 50 pg in autotrophic cultures (JUNGnickel and GEBHARD 1986).

Mn⁺⁺ raises the activity of phenylalanine ammonia-lyase similarly as a reducing agent (GORDON 1977). SCHREINEMAKERS (1984) and SCHREINEMAKERS and DORHOUT (1985) point to the possible existence of a coupled uptake mechanism for Mn and Mg in S. polyrrhiza. For activities of Mn on flowering see chapter 2.4.3.1.2.

2.3.3.4.7.6. Zinc (Zn)

THIMANN and EDMONDSON (1949) proved the indispensability of Zn for S. punctata. The minimal Zn concentration to achieve good growth of L. minor is 0.0006 mM (STEINBERG 1946). HUTCHINSON and CZYRSKA (1975) noted a growth stimulation for L. minor at Zn concentrations of 0.0008 and 0.0016 mM. The lowest Zn concentration to allow a slow growth of S. polyrrhiza was 0.002 mM (EYSTER 1966). Best growth was attained for S. polyrrhiza with 0.02-0.2 mM Zn (EYSTER 1966), for L. minor with 0.002 mM Zn (HUTCHINSON and CZYRSKA 1975), and for L. obscura (named as L. minor) with 0.1 mM (STANLEY and MEADWELL 1976a). With 2 mM Zn the growth rate of S. polyrrhiza is slowed down, and with 20 mM the fronds died (EYSTER 1966). SAROSIEK et al. (1982) observed toxic symptoms in L. minor at concentrations of 0.1 mM (they used tap water as medium). Sixteen mM Zn resulted in the death of L. obscura (STANLEY and MEADWELL 1976a). Fronds of L. aequinoctialis produced brown spots at Zn concentrations of 3 mM

and died at 8 mM (NASU and KUGIMOTO 1981, NASU 1983). VAN DER WERFF and PRUYT 1982) observed no toxic effect of Zn concentrations up to 0.5 mM in S. polyrhiza and L. gibba. An inhibiting effect of 0.1 mM Zn and higher on the root growth of L. gibba was reported by LIEBERT (1986a). Root growth stopped completely with 1 mM Zn.

According to EYSTER (1966), the minimal zinc requirement for the production of one gram dry weight of S. polyrhiza amounts to 0.26 mg. Zinc deficiency (in contrast to copper deficiency) lowers the superoxide dismutase activity in L. gibba (VAUGHAN et al. 1982). SCHREINEMAKERS (1984) and SCHREINEMAKERS and DORHOUT (1985) suggest a possible coupled uptake mechanism for Zn and Ca in S. polyrhiza. The toxic effect of Zn was neutralized by raising the Mg content (MARTIN 1955).

2.3.3.4.7.7. Copper (Cu)

The evidence of the indispensability of copper could not be demonstrated by SAEGER (1937), STEINBERG (1946), THIMANN and EDMONDSON (1949) and EYSTER (1966). EYSTER (1966) showed that many chemicals (e.g. KH_2PO_4 , $\text{MgSO}_4 \cdot 7 \text{H}_2\text{O}$, $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, H_3BO_3) and even distilled water contain traces of Cu. Therefore the proof of Cu requirement is extremely difficult. Nevertheless, most authors (e.g. SAEGER 1937) think that copper is an obligatory element for Lemnaceae. THIMANN and EDMONDSON (1949) suppose that a copper containing enzyme participates in the formation of anthocyanin in S. punctata. A growth stimulation in S. polyrhiza was observed by EYSTER (1966) with 0.001 mM Cu, and by HUTCHINSON and CZYRSKA (1975) in L. minor with 0.0001 mM Cu. The latter authors report on growth inhibition already with 0.0008 mM Cu. Growth reduction was recorded in L. minor with 0.06 mM (FILBIN and HOUGH 1979), and in L. aequinoctialis with 0.0015 mM Cu (NASU and KUGIMOTO 1981, NASU 1983). VAN DER WERFF and PRUIT (1982) did not note any toxicity of Cu concentrations up to 0.5 mM in S. polyrhiza and L. gibba. Lethal doses are shown in S. polyrhiza as 0.01 mM (EYSTER 1966), in L. obscura (named as L. minor) as 2 mM (STANLEY and MADEWELL 1976a), and in L. minor as 0.8-1 mM (SAROSIEK and WOZAKOWSKA-NATKANIEC 1980, SAROSIEK et al. 1982), as 0.06 mM (FILBIN and HOUGH 1979) and as 0.005 mM (HUTCHINSON and CZYRSKA 1975). MATTOO et al. (1986a) observed damages of intracellular membranes in S. punctata by copper at concentrations of 0.02 mM and higher. The strikingly different results reflect the fact that the composition of

the nutrient solution, especially the pH and the presence of a chelating agent, has a great influence on the effect of copper and other trace elements. Root growth of L. gibba is inhibited at Cu concentrations of 0.01 mM or higher and is stopped completely at 0.1 mM (LIEBERT 1986a). The effect of high copper concentrations results in the following characteristics (FILBIN and HOUGH 1979): reduction of chlorophyll content, reduction of CO_2 uptake, reduction of sucrose uptake in the dark, and elevation of the respiration rate in the light (dark respiration was not changed).

EDTA in a concentration of 0.03 mM prevented the absorption of Cu (0.005-0.01 mM). If ammonium was added to the solution, the Cu uptake was also greatly suppressed. However, the absorption of Cu was not significantly affected by the initial pH between 3.6 and 5.1 (NASU et al. 1983). A relatively great amount of Cu (0.0005-0.001 mM) in the nutrient solution induces flowering of the short-day plant L. aequinoctialis in both long and short-day regimes, whereas the long-day plant L. gibba loses its ability to flower (HILLMAN 1962, 1965). HILLMAN assumes an effect of Cu on the photoperiodic sensitivity rather than directly on flowering. Also TANAKA and CLELAND (1980) showed that flowering of L. gibba is inhibited by copper under continuous light. This inhibition is largely abolished by salicylic acid (SA) or by EDTA (TANAKA et al. 1982a). Unlike EDTA and ammonium, SA does not affect the absorption of Cu. In L. aequinoctialis, SA multiplied the flowering effect of Cu. The absorption of Cu is not influenced by cadmium, but the flower-inducing activity on L. aequinoctialis is inhibited (NASU et al. 1984). At pH 4.1 0.005 mM Cu results in an unchanged chlorophyll content of L. aequinoctialis, a somewhat reduced growth rate and a substantial induction of long-day flowering. By contrast, at pH 5.1 the same copper concentration causes a reduction in the chlorophyll content and a strong inhibition of growth, but it does not induce long-day flowering (TANAKA et al. 1982b). The flower-inducing activity of Cu is assumed to be the consequence of the suppression of nitrate reductase activity (TANAKA et al. 1986a). KHURANA and MAHESHWARI (1983c) measured a significantly higher copper level in plants of L. aequinoctialis treated with the chelating agent 8-hydroxychinoline for flowering. Ethylene production in S. punctata is markedly stimulated by CuSO_4 (MATTOO and CONLON 1985). For further details on the flowering effect of copper see chapter 2.4.3.1.3. The addition of nickel increases the copper uptake in L. minor (HUTCHIN-

SON and CZYRSKA 1975). Coupled uptake mechanisms for copper and iron are assumed by SCHREINEMAKERS (1984) and SCHREINEMAKERS and DORHOUT (1985). ROGERS et al. (1978) showed that copper acts synergistically with zinc and mercury.

CuSO₄ at concentrations of 0.006 and 0.06 mM greatly reduces the growth of Lemnaceae if sprayed as a herbicide. The chlorophyll content of the plants is lowered; the nitrogen content decreases from 4% of the dry weight to 0.3-1% (SINHA and SAHAI 1975).

2.3.3.4.7.8. Cobalt (Co) and nickel (Ni)

EYSTER (1966) was not able to prove any stimulating effect of cobalt. Co was toxic to S. polyrrhiza at concentrations of 10 mM. POLAR (1984 in lit.) reported a concentration lethal to a L. minor clone of 0.05 mM Co; another clone showed only a reduction in growth rate of 60% at the same concentration. The uptake of cobalt by L. gibba is enhanced at a low pH (pH was studied between 5 and 9) (EL SHINAWY and ABDEL-WALIK 1980).

For nickel, a stimulating effect was reported by GORDON et al. (1978). At a Ni concentration of 0.05 mM, the growth rate of L. aequinoctialis was doubled if urea was used as a sole nitrogen source. The curve of respiration (CO₂ output) shows a double peak daily if Ni is added to the solution whereas without Ni a single peak occurs. According to HUTCHINSON and CZYRSKA (1975), an addition of 0.00017, 0.00085, and 0.0017 mM stimulated the growth of L. minor. At concentrations of 0.005 and 0.008 mM Ni, growth is inhibited, and at 0.017 mM Ni, the fronds die. Ni and Cu have a synergistic effect. If combined they are lethal for L. minor at concentrations at which the single element is not.

2.3.3.4.7.9. Chromium (Cr)

No distinct stimulating effect of chromium is known. However, Lemnaceae are relatively tolerant to Cr. HUFFMANN and ALLAWAY (1973) did not note any influence of Cr at concentrations of 0.0005 mM on the growth rate of Lemna sp. Inhibition of growth was observed by MANGI et al. (1978) in S. polyrrhiza at concentrations of 0.02 mM Cr and in L. minor at 0.0002 mM. Reduced growth of L. minor determined by the frond area, lower chloroplast volume, and decreased content of chlorophyll a and b was measured by BASYNSKI et al. (1981) in solutions with 0.001 to 1 mM Cr; at 10

mM Cr growth was practically stopped. Also CORRADI et al. (1987) observed the loss of grana in chloroplasts of L. minor with 0.02 mM Cr. Chromium affects especially the photosystem II. For L. aequinoctialis, NASU and KUGIMOTO (1981) found growth inhibition at Cr concentrations of 0.02 mM and higher. SHIROISHI and SHIMIZU (1984) observed a slight stimulation of the growth of L. aequinoctialis at concentrations up to 0.002 mM Cr and an inhibition from 0.004 mM upwards. Mortality of L. aequinoctialis was reported by CLARK et al. (1981) at Cr concentrations of 0.005 mM and higher. Concentrations of 0.4 and 1 mM Cr are lethal to L. minor (SAROSIEK and WOZAKOWSKA-NATKANIEC 1980).

STAVES (1980) and STAVES and KNAUS (1985) studied the Cr uptake and the short-time effect of Cr on growth of S. polyrrhiza, S. punctata, and L. gibba using the radiotracer ⁵¹Cr. At Cr levels higher than 0.02 mM growth was greatly inhibited, S. polyrrhiza showing the least tolerance and S. punctata the greatest tolerance towards Cr. At lower concentrations than 0.02 mM no negative effect of Cr has been observed. Between 0.002 mM and 0.4 mM, the uptake of Cr was directly related to ambient Cr concentrations.

2.3.3.4.7.10. Lead (Pb) and cadmium (Cd)

Neither lead nor cadmium stimulate the growth of Lemnaceae.

Lead concentrations of 1.5 to 2.5 mM are toxic to L. minor (SAROSIEK and WOZAKOWSKA-NATKANIEC 1980, SAROSIEK et al. 1982). Longest roots were observed with 0.05 mM Pb (longer than at lower or at higher concentrations). Different clones of L. minor showed somewhat different response to Pb. According to FIUSELLO (1973), the chlorophyll content is reduced at 0.1 and 1 mM Pb. The toxicity of Pb is higher if Pb is added as acetate than as nitrate (SCHWABEL and ANDERSON 1982). S. polyrrhiza (and Eichhornia) have a higher capacity to accumulate Pb than terrestrial species (the Pb given as Pb-EDTA). Under the investigated concentrations of up to 0.5 mM Pb no toxicity was evident.

The uptake of Pb by L. trisulca was studied by EVERARD and DENNY (1985). The absorption is very fast. Within 10 minutes, L. trisulca takes up 0.1 mg Pb per gram dry weight from a solution of 0.005 mM Pb. The uptake is followed by a small net efflux of Pb.

Cadmium inhibits growth of L. minor at concentrations higher than 0.0001 mM; it is lethal at 0.005 mM (HUTCHINSON and CZYRSKA 1975). ORNES (1979)

observed growth inhibition of S. polyrrhiza at concentrations of 0.0004 mM. The fronds did not die if temporarily grown with 0.07 mM Cd in contrast to other water plants such as Azolla, Salvinia, and Ceratophyllum. The root growth of L. gibba was reduced at 0.001 mM Cd and stopped at 0.2 mM (LIBERT 1986a). SETO et al. (1979) report on chlorotic fronds of L. gibba at Cd concentrations of 0.001 and 0.01 mM in a diluted nutrient solution. POLAR and KUECKEZEZAR (1986) mention growth inhibition of L. gibba with 0.01 mM Cd. Differently, VAN DER WERFF and PRUIT (1982) did not find any toxicity symptoms in S. polyrrhiza and L. gibba at Cd concentrations of up to 0.5 mM. The toxicity of cadmium is much dependent on the composition of the nutrient solution and on the light regime. It is much more toxic in deionized water than in river water (WANG 1986b). Further results on toxicity and bioaccumulation of Cd are reported by CHARPENTIER et al. (1987).

In the Hillman M solution (modified Hoagland solution) and in the solution of Bonner-Devirian, 0.0001 mM Cd result in growth inhibition of L. aequinoctialis. In 1/2 Hutner solution, the same effect was achieved at 100 times higher Cd concentrations (NASU and KUGIMOTO 1981, NASU 1983). To prevent the absorption of 0.005 to 0.01 mM Cd, a supply of 0.4 mM EDTA is necessary. Ammonium suppresses Cd absorption only slightly. The absorption of Cd increases with higher initial pH (between 3.6 and 5.1) (NASU and KUGIMOTO 1981, NASU et al. 1983). The uptake of Cd is reduced considerably with increasing concentrations of Cu in the culture solution (NASU et al. 1984). Zn increases the toxicity of Cd (HUTCHINSON and CZYRSKA 1975).

The uptake of Cd by S. polyrrhiza is proportional to the Cd concentration given in the solution. If the Hoagland solution is diluted to 10%, the Cd is absorbed faster than from 50% Hoagland. Afterwards, a release of up to 90% of the Cd absorbed is noted in 10% Hoagland, contrary to the situation in the 50% Hoagland (ORNES 1979). The iron content of L. gibba fronds increases with increasing concentrations of Cd in the solution, the manganese and zinc content decrease (POLAR and KUECKEZEZAR 1986).

SUN et al. (1985, paper not seen) report on physiological changes and injury mechanisms in Lemnaceae treated with Cd. SCHREINEMAKERS (1986) studied the Cd absorption and Cd compartmentation in W. gladiata. The Cd concentration in the cytoplasm reached about 20 mM, 240 minutes after adding 0.4 mM Cd to the solution. It amounted to 12 mM at a concentra-

tion of 0.04 mM Cd and to 3 mM at a concentration of 0.001 mM Cd.

2.3.3.4.7.11. Arsenic (As), selenium (Se) and vanadium (V)

No stimulation was observed by the addition of As, Se, or V.

L. obscura (named as L. minor) is able to grow at arsenic concentrations of 0.015 mM (as NaAsO_2) or of 0.2 mM (as Na_2AsO_4) without great damage (STANLEY and MEADWELL 1976a). The growth rate of L. aequinoctialis is inhibited (50%) by 0.01 mM As (NASU and KUGIMOTO 1981, NASU 1983, NASU et al. 1983) and 0.007 to 0.008 mM As reduces the growth rate of L. minor to 50% (SZABADOS et al. 1983). NaAsO_2 is toxic to Lemnaceae at concentrations of 0.01 mM if sprayed as a herbicide (SINHA and SAHAI 1975).

Arsenic is methylated by S. polyrrhiza, L. minuscula, and W. columbiana, and later, arsoniumphospholipids are formed (BENSON et al. 1981, NISSEN and BENSON 1982). Similarly to other aquatic plants but differently to terrestrial plants, Lemnaceae have evolved mechanisms for rapid detoxification of arsenate, arsenite, and other toxic arsenic substances. According to OERENCIC et al. (1982), NaAsO_3 does not inhibit chlorophyll synthesis of L. gibba. In this species, however, flowering is inhibited by 0.1 to 0.5 mM disodium arsenate (KANDELER 1968, OOTA 1969a). ULLRICH-EBERIUS and NOVACKY (1984) suggest that the arsenate influx proceeds via the P carrier.

Selenium in form of H_2SeO_3 is absorbed by L. obscura (named as L. minor) up to a concentration of 0.03 mM without great damage (STANLEY and MEADWELL 1976a). Selenium is a competitor to sulfur. Therefore, it is especially toxic in solutions which lack sulfur (DATKO et al. 1980a). Replacement of the sulfur atom in amino acids by selenium atoms had a toxic effect on L. minor (GULATI et al. 1981). In a solution with equivalent amounts of selenite and selenate, S. punctata takes up three times as much selenite. Colloidal Se is also absorbed. In the soluble cell fraction, Se is present as selenocystine, in the protein as selenomethionine (BUTLER and PETERSON 1967).

Vanadium at a concentration of 0.2 mM inhibits growth of L. minor (PIIS-PANEN and LAHDESMAKI 1983). The vanadate/ H^+ cotransport proceeds contrary to arsenate via a carrier which is independent of the P carrier. Vanadate is neither a strong nor a specific inhibitor of the plasmalemma H^+ -ATPase in vivo for L. gibba as supposed for other plants (ULLRICH-EBERIUS and NOVACKY 1984).

2.3.3.4.7.12. Silver (Ag) and mercury (Hg)

Silver inhibits the growth of L. aequinoctialis at a concentration of 0.0001 mM (NASU and KUGIMOTO 1981) and of L. minor at 0.0002 mM (HUTCHINSON and CZYRSKA 1975). The uptake of Ag is not detectable. TAKIMOTO and TANAKA (1973), TAKIMOTO (1981) and NASU and KUGIMOTO (1981) describe a flower-promoting effect of silver in L. aequinoctialis at 0.0001 and 0.01 mM Ag (especially at the low pH of 4.1). This effect is supposed to occur due to the inhibition of the activity of SH groups by Ag.

Mercury suppresses growth of Lemnaceae at a concentration of 0.002 mM Hg (CHANG et al. 1977). According to FERNANDEZ et al. (1983), L. aequinoctialis shows a reduced growth rate and brown spots on the frond surface with 0.02 mM Hg; the plants die with 0.03 mM Hg. Hg has a similar effect on flowering of L. aequinoctialis as Cu (HILLMAN 1962, TAKIMOTO 1981, BEPPU and TAKIMOTO 1983). HgCl_2 at concentrations higher than 0.01 mM depolarizes the membranes of L. gibba markedly and irreversibly. It was concluded that Hg^{++} binds irreversibly to SH groups of the H^+ /sugar and H^+ /amino acid transport carriers and inhibits these carriers (SCHWEIGER et al. 1984). Further studies on the effect of HgCl_2 on L. gibba suggest the following sequence of sensitivity of physiological functions: amino acid and glucose transport and probably Na^+ and K^+ proton exchange carriers at the plasmalemma > mechanisms establishing a metabolism-dependent electrical potential at the plasmalemma = passive Na^+/K^+ permeabilities at the plasmalemma > respiration and photosynthesis (LOOS and LUETGE 1984).

2.3.3.4.7.13. Germanium (Ge) and silicone (Si)

Germanium inhibits growth of L. minor at concentrations of 0.3 mM GeO_3 . W. arnhiza is able to tolerate somewhat higher concentrations. The inhibition is partly neutralized if silicic acid is added to the solution (WERNER 1967). The author supposes that germanic acid disturbs the balance of silicic acid. However, it has never been proved that silicone is an essential element for Lemnaceae. LOOMIS (in lit. 1987) tried to substitute germanic acid for boric acid in the nutrient solution. He succeeded in culturing L. minor for 6 months in 0.05 and 0.1 mM GeO_2 . The plants formed rootless colonies and did not separate normally.

2.3.3.4.7.14. Other minerals

The uptake and accumulation of further minerals is dealt with in chapter 3.5.3.

2.3.3.5. Organic substances

2.3.3.5.1. General remarks

Lemnaceae are able to take up water soluble organic substances besides the inorganic chemical compounds. Four groups of organic substances can be distinguished according to their mode of action in the metabolic processes:

- 1) Energy suppliers and building material: substances which are used as a basis for respiration or which are incorporated in the plant body as reserve substances or as elements of the plant organism.
- 2) Chelating agents: organic substances which form complex compounds with metallic ions enabling a better uptake of the ions and a detoxification of some metallic elements.
- 3) Growth factors: vitamins or hormone-like substances which regulate different metabolic processes; they act in very small doses.
- 4) Toxic substances: chemical compounds which disturb or interrupt certain metabolic processes, or which damage parts of the cells (e.g. pesticides, detergents).

Some of the organic compounds have combined effects. Members of the groups 1) and 3) may also act as chelating agents. Some growth factors behave as toxins in higher doses, and some toxins behave as growth factors in small doses.

The uptake of organic substances in L. aequinoctialis was studied by DAYKO and MUDD (1985, 1986). The authors detected 8 distinct uptake and transport systems for organic substances.

2.3.3.5.2. Energy suppliers and building material

2.3.3.5.2.1. Carbohydrates

Lemnaceae use carbohydrates as energy suppliers under non-saturating light conditions (cf. HOPKINS 1931, STEINBERG 1941, 1946, GORHAM 1950,

LANDOLT 1957, HILLMAN 1961a, ULLRICH-EBERTUS et al. 1978; cf. also chapter 2.3.5.4). The best effect is realized with glucose, fructose, or sucrose, and to a lesser degree with maltose (YOSHIMURA 1944). Sucrose is changed into invert sugar (glucose + fructose) by autoclaving. The following carbon sources are not usable by *Lemnaceae*: ethanol, glycerol, mannitol, inuline (YOSHIMURA 1944), starch, lactose, arabinose, ribose, tartrate, succinate, acetate (HILLMAN 1954), citrate, malonic acid, oxalic acid, glucuronic acid, and glutaric acid (CHANG et al. 1977). For *L. gibba* 0.1% galactose proved to be toxic. The effect is neutralized if glucose is added (DECOCK et al. 1979). According to HUBALD and AUGSTEN (1977a), galactose, ribose, and xylose inhibit the growth of *L. gibba*. Sorbose has a similar effect on the growth of *L. minor* as galactose (STROTHER 1981). Five mM sorbose results in complete bleaching of the cultures. The effect can be partly overcome by fructose.

It is somehow unclear if glucose or fructose are more favourable to growth of *Lemnaceae*. The effect is probably dependent on the culture conditions or on the species. THIMANN et al. (1951) achieved better growth in *S. punctata* with fructose than with glucose. The same was




Fig. 2.10. Growth rate of *Spirodela polyrrhiza* in relation to sucrose concentration at 26°C and light duration of 18 hours per day at 1000 lux (from LANDOLT 1957)

observed by HENSEN (1954) in *S. polyrrhiza*. On the other hand, HUBALD and AUGSTEN (1977a) report better growth of *L. gibba* with glucose than with fructose. DATKO et al. (1980a) mention a similar growth rate in *L. aequinoctialis* with either glucose or sucrose at the same concentrations.

The best concentration of sucrose for *Lemnaceae* growth is between 0.5 and 2.5% (GORHAM 1950, LANDOLT 1957, ROMBACH 1976) (fig. 2.10). According to DATKO et al. (1980a), the growth rate of *L. aequinoctialis* still rises slightly between 0.5 and 1% sucrose.

All species of *Lemnaceae* which are kept under suboptimal light conditions grow faster after adding sucrose. The growth stimulation is limited when the maximum growth rate of the species under optimal light conditions and continuous light is reached. At saturating light intensities, sucrose inhibits growth (LANDOLT 1957). The growth rate of *Lemnaea* in solution with sucrose at low light intensities consists additively of the growth rate in darkness (with sucrose) and of the growth




Fig. 2.11. Growth rate of *Spirodela polyrrhiza* in the dark with 1% sucrose (2), at 1000 lux without sucrose (1) and at 1000 lux with 1% sucrose (3). --* marks the added values of curve (1) and (2) which corresponds roughly to curve (3) (from LANDOLT 1957)

rate at the same low light intensity in solutions without sucrose (LANDOLT 1957) (fig. 2.11). Species which do not grow on the surface of the water but submerged (L. trisulca, L. valdiviana, Wolffiella species) are dependent upon sugar in the solution if it is not supplemented with a carbon dioxide source (aeration with CO₂, addition of carbonate or hydrocarbonate) (cf. chapters 2.3.2.2 and 2.3.3.4.5). MONSELISE et al. (1984) investigated a mutant of L. aequinoctialis (6746) incapable of photosynthesis. It was dependent on the supply of sucrose for its growth and showed unusual structures of mitochondria and chloroplasts.

The size of the frond of L. minor decreases with raising of the sucrose concentration from 1 to 3% to about half (ROMBACH 1976). The disconnection of the daughter fronds from the mother fronds is impeded by sucrose; therefore, large colonies develop (e.g. YOSHIMURA 1944, HILLMAN 1954, own observations). The effect of sugars on flowering is presented in chapter 2.4.3.1.4. Fructose and sucrose stimulate the formation of anthocyanin in S. punctata; glucose does this only slightly (THIMANN et al. 1951). GROB and RUFENER (1969) report that 1% glucose in the light changes the chloroplasts of S. punctata. After 25 days, chlorophyll is no longer present; the chloroplasts are filled with starch, the thylakoids have entirely disappeared. Bleached fronds brought into a new glucose-free medium turn green again after a few days. This phenomenon could not be observed in our cultures with sucrose. Glucose is supposed to stimulate the nitrate assimilation (GROB et al. 1973). The content of IAA in S. punctata is enhanced by sucrose (WITZTUM et al. 1978). Sucrose was found to reduce glutamate dehydrogenase activity in L. aequinoctialis; glucose on the other hand increased the activity (DUKE 1975, DUKE and KOUKKARI 1977).

Sucrose uptake was investigated by WOHLER (1966). It amounted to about the same rate in the dark as well as at different light intensities. The uptake rate was different for different species. The following maximum rates were measured in mg sucrose per g dry weight and hour: 10 for S. polyrhiza, 20 for L. minor, 40 for L. trisulca, and 65 for W. gladiata. For hexoses (glucose and fructose), the absorption is energy-dependent, at least for short times (ULLRICH-EBERIUS et al. 1978). The uptake of hexoses was 1 1/2 to 3 times higher in the light than in the dark. DAYKO and MUDD (1985) studied the uptake of aldohexoses (D-glucose, D-galactose, D-mannose) in L. aequinoctialis. Fastest uptake was observed for glucose. Galactose and mannose were absorbed nearly as fast. Fructose,

sucrose, arabinose, ribose, and glycerol showed much lower uptake rates. In nature, sugars might be found in lake water (VALLENTYNE and WHITTAKER 1956, SAUNDERS 1957, HARTMAN 1960, WALSH 1965a,b). Heterotrophic growth in nature is assumed for submerged species such as W. neotropica, W. lingulata, W. oblonga, L. valdiviana. In South America, these species could be observed in the dark under mats of Salvinia, Azolla, and Scirpus cubensis (LANDOLT and ZARZYCKI in prep.).

2.3.3.5.2.2. Amino acids

It is presumed that the addition of amino acids to a well balanced inorganic nutrient solution supplied with a chelating agent does not stimulate growth of Lemnaceae. Some amino acids may inhibit growth because they interfere with the synthesis of other amino acids and metabolic processes. In unfavourable nutrient solutions, especially at low nitrogen concentrations, without a chelating agent or at low light intensities, some amino acids might serve as a nitrogen source, as a chelating agent or merely as an amino acid source. Some of the contradictory results listed below can be explained by not always optimal growth conditions. Mechanisms of uptake of amino acids are described more extensively in chapter 2.5.2.3.1.

If inorganic nitrogen sources are lacking, L. minor is able to use a mixture of glutamic acid, aspartic acid, and arginine as well as casamino acids. Other amino acids such as alanine, valine, methionine, and leucine inhibit growth (JOY 1969a). According to FELLER and ERISMANN (1976), the following amino acids reduce the growth rate of L. minor if added singly to the nutrient solution: arginine, alanine, glycine, asparagine, aspartic acid, glutamine, glutamic acid, ornithine, and citrulline. The inhibiting effect is greater with NO₃⁻ than with NH₄⁺ as nitrogen source. Arginine enlarges the size of the frond of L. minor, promotes the stripes and inhibits the separation of the daughter fronds. Contrary to the results of FELLER and ERISMANN (1976), HUBALD and AUGSTEN (1977a) observed a growth stimulation of L. gibba with the addition of 10 mM L-arginine and L-asparagine. (HUBALD and AUGSTEN used a nutrient solution without EDTA, and a relatively low light intensity of 900 lux). Neither stimulation nor inhibition was found with the following amino acids: glycine, DL-alanine, D-valine, DL-serine, DL-methionine, L-tyrosine, L-histidine, L-proline, and L-tryptophane. NICKELL (1956)

centration of 0.005 mM. Simultaneous application of canavanine and canavanine caused an additive reduction of the growth rate. Ureidohomoserine interacted with canavanine or canavanine. It increased the additive growth reduction caused by canavanine plus canavanine (ROSENTHAL et al. 1975, 1976). Already NICKELL (1956) reported a toxic effect of canavanine on L. minor which could be neutralized by arginine and citrulline. MONSE-LISE et al. (1986) were able to obtain a partial recovery of plastids in a mutant of L. aquinoctialis which was incapable of ammonium uptake and photosynthesis, by addition of L-methionine and L-cystine.

NAKASHIMA (1964) observed inhibition of flowering of L. gibba by all 20 investigated amino acids (at concentrations between 0.1 and 1 mM). Thirteen of these amino acids did not inhibit growth at the same concentration. The addition of the amino acids glutamic acid, leucine, and serine (at concentrations of 0.001 mM) enhance the flowering percentage of S. polyrrhiza and L. minor under long-day conditions. Valine, tyrosine, and proline stimulated flowering under short-day conditions; cystine and some other amino acids proved to be inhibitory (KRAJNCIC and DEVIDE 1982b). FERGUSON (1970) noted, in S. punctata, a faster uptake of asparagine and glutamine than of nitrate; aspartate and glutamate were taken up at about the same rate as nitrate. NEWTON (1972a) indicates that DL-leucine and uridine are actively transported across the plasmalemma in the roots of L. minor. The uptake of glycine in L. gibba was studied by FISCHER and LUETTGE (1980). The uptake of neutral L-amino acids in L. gibba was shown to be a H^+ cotransport mechanism driven by a proton-electrochemical gradient at the plasmalemma (JUNG and LUETTGE 1980). Similar results were achieved with acidic amino acids (L-alanine, L-serine, L-aspartate) by JUNG et al. (1982). DATKO and MUDD (1985) investigated the uptake of amino acids and other organic compounds in L. aquinoctialis. They identified 8 discrete transport systems two of which for amino acids: a) neutral amino acids (e.g. L-leucine), b) basic amino acids (e.g. L-arginine), c) purine bases (e.g. adenine), d) choline, e) ethanolamine, f) tyramine, g) urea, and h) aldohexoses.

who also worked with a nutrient solution without EDTA reported growth stimulation of L. minor with the amino acids arginine, ornithine, citrulline, and isoleucine. Similar stimulation in S. polyrrhiza with alanine, glutamic acid, and tryptophane was achieved by SMIRNOVA (1975b, 1981, also in solutions without EDTA). The addition of amino acids resulted in a higher dry weight per frond. GULATI et al. (1981) investigated the influence of 55 amino acids not occurring in natural proteins on the growth of L. minor. A replacement of S atoms by Se atoms or certain substitutions by F atoms resulted in toxic effects. Additional methyl and phenyl groups did not change this effect. The authors divided the amino acids studied into 5 groups, according to their toxicity on Lemnaceae.

Alanine, aspartic acid, glycine, and serine inhibit the activity of glutamine synthetase in L. minor, but not that of glutamate dehydrogenase (STEWART and RHODES 1977a). The addition of a single amino acid such as L-valine, L-isoleucine and L-leucine, inhibits the growth of S. polyrrhiza. A combined addition of L-valine and L-isoleucine shows no inhibition, contrary to other combinations (BORSTLAP 1970). Apparently, valine prevents the synthesis of isoleucine, isoleucine that of leucine, and leucine that of valine and isoleucine through blocking of enzymatic processes of synthesis (BORSTLAP 1972, 1977b). The uptake of L-glutamic acid in S. polyrrhiza declined between pH 4.0 and 6.0 and that of L-leucine between pH 4.0 and 8.0 whereas L-lysine uptake was optimal at pH 6 (BORSTLAP et al. 1986). The authors detected two saturable and one non-saturable uptake systems for amino acids. WONG and DENNIS (1973) reported that threonine, homoserine, isoleucine, and to a lesser extent, lysine inhibit growth of L. minor at a concentration of 0.25 mM. Methionine was lethal at the same concentration; so was a combination of lysine and threonine. It is supposed that lysine inhibits the serine hydroxymethyl transferase and in this way the synthesis of methionine (WONG and COSINS 1976). BORSTLAP and VERNOOY-GERRITSEN (1985) studied the influence of leucine and valine, added to the solution, on the leucine and valine synthesis in S. polyrrhiza. If the leucine content increases, leucine synthesis is markedly reduced, valine synthesis only slightly, and isoleucine synthesis apparently not at all. A higher valine content strongly affected the valine synthesis, and the leucine synthesis lesser but not the isoleucine synthesis. The amino acids of the canavanine-urea cycle (canavanine, canaline, ureidohomoserine) are toxic to L. minor at a con-

2.3.3.5.3. Chelating agents

2.3.3.5.3.1. General remarks

In many older publications, a growth stimulation was reported in cultures supplied with organic substances. BOTTOMLEY (1920b), MOCKERIDGE (1920), and CLARK (1926) observed better growth in Lemnaceae cultures to which extracts of soil, manure, or peat were added. STELING (1937) prevented chlorosis of L. minor by supplying Fe humate to the nutrient solution. In a similar way, TATKOWSKA and KOBYLANSKA (1978) promoted growth rate, dry weight, chlorophyll, phosphorus and iron content in S. polyrrhiza by means of adding Fe humates to the culture medium. All these effects were achieved in solutions with no other chelating agent. Fe humate is also able to neutralize the toxic effect of the detergent DBSS on S. polyrrhiza diminishing the ratio of P/Fe (GUMINSKI et al. 1978). NICKELL and FINLAY (1954) achieved a growth stimulation of L. minor with addition of antibiotics which might be attributed to chelating properties of these chemicals. Also, a supply with citrate (BITCOVER and STELING 1951) and kojic acid (YOKOTA and SHIMADA 1958) resulted in higher growth rates of S. polyrrhiza if the solutions were void of a chelating agent. OLSEN supposed, already in 1930, that humus substances promote the availability of Fe to the plants in nutrient solution with a high pH by forming a complex salt. Humic acid otherwise stimulates the production of superoxide radicals generated in the xanthine - xanthine oxidase system. The activity of superoxide dismutase in L. gibba is also enhanced (VAUGHAN and ORD 1982). Oxalate promotes the growth of L. gibba and raises the uptake of Ca, according to DEKOCK et al. (1973). CLARK and ROLLER (1931), SIEGER (1925, 1930), WOLFE (1926), ASHBY (1929b), CLARK (1932), and many other authors proved clearly that good Lemnacean growth is possible in aseptic cultures without addition of organic substances. Today, most authors agree that many of the growth promoting organic substances are effective through formation of chelates with Fe and other metallic ions (e.g. Mn, Zn, and partly also Ca and Mg). If these ions are bound in a complex way, they do not precipitate and are available to the plants. On the other hand, the toxic effect of too high concentrations of metallic ions are moderated if complex salts are formed. Besides citrate and humate, tartrate (in form of Fe tartrate) was frequently used as a chelating agent (e.g. in the Hoagland

solution). However, EDTA is more wide-spread in culture solutions as a chelating agent today. EDDHA and SA which are discussed here under the chelating agents have many effects similar to hormones (e.g. cytokinins) and are used in cultivation of Lemnaceae only for special purposes (e.g. to induce flowering).

EYSTER (1966) compared the effect of EDTA, sesquestrene, tartrate, citrate, and α , α , α -tris-(hydroxymethyl)-methylamine (tris). Best growth was clearly achieved with EDTA.

2.3.3.5.3.2. Ethylenediaminetetraacetic acid (EDTA)

The optimal concentration of EDTA for growth of S. polyrrhiza was found to be 0.6-2.0 mM if Fe was added in concentrations of 0.05-2.0 mM (fig. 2.12). According to EYSTER (1966), EDTA is destroyed in culture solutions exposed to the light and loses most of its effect after 8 days. Aeration also reduces the activity of EDTA. The destructive action of light on EDTA is independent of the presence or absence of Lemnaceae in the solution. In certain nutrient solutions (e.g. Knop), the activity of




Fig. 2.12. The effect of EDTA on the growth of Spirodela polyrrhiza (from EYSTER 1966)

EDTA is already reduced by autoclaving. It is not clear from our experience why there is no reduction of the growth rate after 8 days cultivation when the EDTA should be destroyed. Under long-day conditions EDTA promotes length of the root, size of the frond and gibbosity of *L. gibba*, it also intensifies the green colour and stimulates flowering (HILLMAN 1961b). *L. gibba* which does not flower without EDTA becomes a long-day plant in solutions with EDTA, and the day-length indifferent *L. aequinoctialis* a short-day plant. This effect is probably due to a preventing action of Cu ions (HILLMAN 1961e, 1962). The flowering effect of old cultures on *L. aequinoctialis* even under long-day conditions described by LANDOLT (1957) and many other authors may be attributed to the destruction of EDTA by light.

According to BOLGIANO (1979) EDTA removes Ca^{++} ions from the solution and therefore, may be toxic to Lemnaceae in solutions with a low Ca content. The author observed an adaptation of the fronds after some time which he ascribes to a possible modification of the enzyme system. It seems more probable that the "adaptation" is due to the destruction of the EDTA after the first few days.

2.3.3.5.3.3. EDDHA, SA, ASA, BEA, 8-HQ

EDDHA and SA have similar effects on *L. gibba* (PIETERSE and MUELLER 1977) and *S. punctata* (SCHARFETTER et al. 1978). In *L. gibba*, both substances initiate gibbosity (10 and 1 ppm, resp.) and flowering under short-day conditions. The stimulation of flowering and the enhancement of the dry weight by 0.01 mM SA was already observed by BHALLA and SARHARVAL (1975). At higher concentrations of SA, flowering and dry weight are reduced. In *S. punctata*, EDDHA and SA cause a reduction of the size of the frond, a prolongation of the root, a diminution of the number of roots from 2-4 to 1, more intercellular air spaces, more prominent papules, and a delayed separation of daughter fronds; also more chlorophyll and more anthocyanins are formed, and flowering stimulated. The flowers of *L. aequinoctialis* are induced by SA and aspirin under non-flowering conditions (KHURANA and MAHESHWARI 1978). Also the short-day plant *W. microscopica* flowers under long-day conditions if either EDDHA (MAHESHWARI and SET 1966a), SA, BEA, aspirin, or salicylaldehyde are added to the solution (KHURANA and MAHESHWARI 1983a). EDDHA, BEA and SA also initiate the flowering of the day-neutral *S. polyrhiza*. The fronds

of this species become gibbous and the chlorophyll and anthocyanin content is enhanced (KHURANA and MAHESHWARI 1980, 1986b). LANDOLT (unpubl. results) was able to get the following species to flower in solutions with EDDHA: *S. intermedia*, *S. polyrhiza*, *S. punctata*, *L. gibba*, *L. dispersa*, *L. minor*, *L. obscura*, *L. japonica*, *L. trisulca*, *L. valdiviana*, *W. Weltschii*, *W. brasiliensis*, *W. australiana*, and *W. columbiana*. TAMOT et al. (1987) and TAMOT and MAHESHWARI (1987) were able to induce flowering of *W. hyalina* under short-day conditions if 10^{-2} mM SA was provided. EDTA, EDDHA, 8-HQ, and zeatin had no effect. The flowering effect of SA only occurs if ammonium, copper, or iron are present in the solution (DEKOCK et al. 1974). More details on the flowering effect of EDDHA and SA can be looked up in chapter 2.4.3.1.2.

SA inhibits the growth rate of *L. obscura* at concentrations of 0.1 mM or higher. Anthocyanin formation is stimulated at 0.0005 mM and inhibited at 0.05 mM and higher (LEATHER and EINHELLIG 1985). The gibbosity effect of EDDHA and SA on *L. gibba* is reversed by β -naphthol (PIETERSE 1978b) and by breakdown products of GA_3 (formed during autoclaving) but not by GA_3 itself (PIETERSE 1976, 1982). In *S. polyrhiza* (DE LANGE and REYER 1982) and in *S. intermedia* (own observations), no gibbosity is induced under the same conditions as for *L. gibba*. However, this seems to be a question of the right concentration since KHURANA and MAHESHWARI (1980) obtained gibbous *S. polyrhiza* with EDDHA. At 0.1 mM SA, the net photosynthesis is completely inhibited in *L. minor* (MARTI et al. 1986). Probably, EDDHA and SA activate the IAA oxidase by delivering of Mn and in this way lower the auxin level of the plants. NAA shows antagonistic effects on EDDHA and SA (SCHARFETTER et al. 1978). CLELAND et al. (1982) suppose that EDDHA is degraded to SA-like fragments within the plants; this explains the similar effect of both substances. SCHARFETTER et al. (1986) assume that EDDHA decreases the endogenous level of free auxins (and, possibly, the endogenous cytokinin level) thus changing the sensitivity to endogenous ethylene. Flowering, production and development of the number of roots, length of the roots, and the size of frond colonies in *S. punctata* seem to need a certain intermediate ethylene level which can be suppressed by AVG, and AOA or increased by ACC and etrel. Benzoic acid (BEA) and BEA derivatives have a flowering effect on *L. minor* and *L. aequinoctialis* (KAIHARA et al. 1981, TAKIMOTO 1981, WATANABE and TAKIMOTO 1979, WATANABE et al. 1981, 1983, TAKIMOTO and KAIHARA 1986). Two clones of *L. aequinoctialis* flowered readily on application

of BEA, GA, IAA, ABA, and lutein neutralized the effect. The addition of trans-zeatin on the other hand had a synergistic effect (FUJIOKA et al. 1983b, 1985, 1986d). Cytokinins in general improve the activity of BEA but not of SA. SA is more effective in *L. gibba* than BEA, but less effective in *L. aequinoctialis* (CLELAND 1974c, WATANABE and TAKIMOTO 1979). Similar effects of SA, ASA, and BEA on the long-day plant *L. gibba* are described by CLELAND and TANAKA (1979), CLELAND and BEN-TAL (1982) and FUJIOKA et al. (1985). An addition of BEA intensifies the flower-inducing activity of vitamin K in *L. aequinoctialis* (KAIHARA and TAKIMOTO 1985a, TAKIMOTO and KAIHARA 1986). Chlorination of BEA suppresses the uptake rate in *L. minor* (KENNEY-WALLACE and BLACKMAN 1972). KHURANA and MAHESHWARI (1983c, 1984) used 8-hydroxyquinoline (8-HQ), a well known copper chelating agent, to induce flowering in *L. aequinoctialis* and *W. microscopica* (KHURANA et al. 1986) under non-flowering conditions. The endogenous levels of copper and iron are significantly higher in the plants treated with 8-HQ.

SCHARFETTER et al. (1986) showed that the antiauxin PCIB can mimic EDDHA effects in *S. punctata* to a high degree, causing only one root per frond, large colonies of fronds, good visibility of air chambers and pronounced papillae, but not flowering.

2.3.3.5.4. Vitamins

In cultures with optimal light intensity, Lemnaceae form vitamins endogenously. Therefore, an addition of vitamins has no stimulating effect. CLARK et al. (1938) demonstrated the presence of vitamins A, B₁, and C in *S. polyrrhiza* and NAKAMURA (1960) the vitamins A, B₂, B₆, and C in *W. globosa* (named as *W. arizhiza*). Further references are put together in chapter 1.2.10. YONE and TOMIZAWA (1952) reported growth promotion of *L. aequinoctialis* by purified liver fraction which was not due to folic acid nor to vitamin B₁₂ (this compound stimulates growth only slightly). It must be assumed that this unidentified growth promotion was only possible under the not very favourable conditions applied. However, in darkness, some of the Lemnaceae species need vitamins from the B group (especially B₁) (cf. GORHAM 1950, ONO 1952a, LANDOLT 1957, HILLMAN 1961a, ROMBACH 1974a,b, 1976, LEUCHTMANN 1979). For more details see chapter 2.3.5.4 on heterotrophic growth. The thiazole component of thiamine substitutes in *L. minor* for the thiamine (B₁) (ROMBACH 1976).

Vitamin C (0.1 mM) promotes flowering of *L. aequinoctialis* under continuous light (KÄNDELER 1971a). BARBER and CALDWELL (1976) studied the effect of vitamin C (ascorbic acid) on *L. gibba*. They observed an influence of vitamin C on the RNA synthesis (2 H-uridine incorporation) and on flowering. Possibly, vitamin C acts as a regulator of the gene activity (especially in relation to flowering). According to FERGUSON and KNYPL (1974), ascorbate can induce high levels of nitrate reductase activity in *S. punctata*.

A flower inducing activity of vitamin K (K₁, K₃, K₅) in *L. aequinoctialis* was observed by KAIHARA and TAKIMOTO (1985a). K₅ was most efficient in inducing flowering at 0.001 mM. The activity is higher at a pH of 4 than of 5. The effect was intensified by the addition of 0.0005-0.001 mM BEA, but not of BA. Copper as CuSO₄ and ferricyanide (K₃Fe(CN)₆) worked synergistically with vitamin K₅. Dicoumarol and 4-hydroxycoumarin (but not coumarin), known as antagonists of vitamin K, promoted flower-inducing activity of vitamin K₅ and induced flowering in *L. aequinoctialis* in a similar way to BEA.

Folic acid (vitamin B₂ group) shows antagonistic activities against high cytokinin concentrations which inhibit growth of Lemna. It is assumed that folic acid accelerates the photochemical degradation of cytokinins (VAN EYCK 1963, BEZEMER-SYBRANDY et al. 1968, BEZEMER-SYBRANDY 1969). Also ROMBACH (1976) observed a stimulation of the growth rate of *L. minor* in solutions with 0.001 mM folic acid. FROMM and O'DONNELL (1952) succeeded in enhancing the growth rate of *L. minor* with PABA, which is an intermediate in the biosynthesis of folic acid. PABA is able to neutralize the inhibiting effect of sulfanil amide but not of ammonium sulfate, sulfaguanidine and sulfathiazole (FROMM and O'DONNELL 1953, 1955). Nicotinic acid (vitamin B₃) extracted from culture solutions with flowering *L. gibba* and *L. aequinoctialis* showed flower-inducing activity on the same species (FUJIOKA et al. 1986a). An addition of IAA, GA₃, or ABA inhibits the flower-inducing effect on *L. aequinoctialis* whereas zeatin leads to a further stimulation of flowering.

The minimum concentration of nicotinic acid to induce flowering of *L. aequinoctialis* is 10 to 1000 times higher (0.01 mM) than of SA, BEA, and EDDHA (TAKIMOTO and KAIHARA 1986). In *L. gibba*, the flowering effect of nicotinic acid is inhibited by all four hormones mentioned (FUJIOKA et al. 1986b).

- 13 'The product number for 5g of Ninhydrin AnalaR is 10132 2C.
- 23 Product 10380 is cyclohexanone **NOT** cyclohexane.
- 36-39 The third column heading should read 'molecular formula **NOT** empirical formula'.
- 40 Lead grain is soluble in H_2SO_4 and HNO_3 .
- 41 The second column in the first table should be headed 'cm³ of 200mmol l⁻¹ AnalaR KCl'.
- 43 The final word in the first line of the second paragraph should be 100 cm³.
- 51 The symbol for Actinium is Ac.
- 52 The symbol for Ruthenium is Ru.
- 54 Relative humidities are expressed as a %.
- 57 The abbreviation for Molar is mol l⁻¹ **NOT** M.
- 57 The definition for volume percent is
- $$\frac{\text{volume of Solute} \times 100}{\text{volume of Solution}}$$

2.3.3.5.5. Hormones and hormone-like substances

2.3.3.5.5.1. Auxins, auxin antagonists and effectors of auxin metabolism (BZ, 2,4-D, IAA, IBA, NAA, PAA, PCIB, PPA, TIBA)

Growth rate. The auxins regulate mainly the extension of the cells. At low concentrations, there is probably no distinct effect on multiplication of cells and on growth rate (GORHAM 1941 with IAA, IBA, and NAA in L. minor; BLACKMAN and ROBERTSON-CUNNINGHAME 1954 with IAA in L. minor; CLARK and FRAHM 1940a,b with IAA, PAA, and PPA in L. minor). ONO (1952a) reported on a stimulation of the growth rate of S. polyrrhiza and Lemna sp. with IAA and NAA at concentrations of 10^{-4} to 10^{-2} mM. However, he had a very incomplete nutrient solution (no Fe, no trace elements), his results are therefore not representative. The same is true for the results of the following authors which did not work with very balanced nutrient solutions either. CLARK and FRAHM (1940a,b) stimulated the growth rate of L. minor with intermittent addition of IAA, PAA, and PPA. According to CHANG et al. (1977, 1978), IAA, IBA, and NAA (0.01-0.025 ppm) improved the growth rate of L. aequinoctialis, but not of S. polyrrhiza and W. globosa (named as W. arhiza). The growth was only measured for 5 days. Possibly, the increase of fronds was achieved by faster elongation of the cells which makes the daughter fronds visible earlier. TIBA at concentrations of 10^{-2} to 10^{-1} mM inhibited the growth rate of L. gibba (OOTA 1965).

Size of frond. The frond is enlarged by BZ (cells larger, but not greater in quantity (HILLMAN 1954; 1955)). The lengths of fronds and stalks of L. trisulca are increased by the addition of IAA to the solution (BATA and NESKOVIC 1982a). In S. intermedia the frond size is reduced with increasing concentrations of IAA and with high concentrations of TIBA (1 mg/l), but enlarged with low concentrations (0.1 mg/l) of TIBA (KLICH et al. 1987b).

Length of roots. Low concentrations of IAA and NAA (10^{-4} to 10^{-3} mM) result in longer roots of S. polyrrhiza and Lemna spp., higher concentrations in shorter roots (ONO 1952). Similar results were obtained in L. minor by BLACKMAN and ROBERTSON-CUNNINGHAME (1954) with IAA, by HILLMAN (1954) with IAA and NAA, and by CLARK and FRAHM (1940a, b) with IAA, PAA, and PPA. The roots of L. trisulca become shorter if IAA is added at concentrations of 0.001, 0.01, 0.1, and 1 ppm (BATA and NESKOVIC 1982a).

Root elongation of L. minor is completely inhibited by 3.4 mM BZ (HILLMAN 1955). TILLBERG and ELIASSON (1976) report on a root elongation in L. gibba with 10^{-6} mM IAA and an inhibition with 10^{-4} mM and higher. IAA reduces root length but not root number of S. intermedia with increasing concentrations (KLICH et al. 1987b).

Stipes of the daughter fronds. The stipe of L. minor becomes elongated by auxine-like substances: CLARK and FRAHM (1940a, b) with IAA, PAA, and PPA, BLACKMAN and ROBERTSON-CUNNINGHAME (1954) with IAA, and HILLMAN (1954, 1955) with IAA, NAA, BZ, and PCIB (the last only at high concentrations). The stalk of L. trisulca is also elongated by IAA (BATA and NESKOVIC 1982a).

Formation of accessory nerves. The addition of IAA, NAA, and TIBA results in the formation of accessory lateral nerves in L. minor (SARGENT 1957).

Closing of stomata. TIBA (1 ppm) is able to close the stomata of S. intermedia completely. The closing is coupled with a characteristic lower potassium content and a lower starch content within the stomata cells (KLICH 1986).

Development of daughter fronds. If TIBA is added to the culture solution, the daughter fronds of L. minor develop simultaneously on both sides of the mother fronds. It can happen that the first daughter frond appears on the side where normally the second one is formed. Additional IAA and NAA strengthens the effect at medium concentrations of TIBA, but lowers it at low and high concentrations (WANGERMANN and LACEY 1953, SARGENT 1957). According to WITZTUM (1966), TIBA prevents the formation of axillary buds in Lemna.

Flowering. The flowering of L. aequinoctialis (GUPTA and MAHESHWARI 1970, TAKIMOTO 1981) and of L. gibba (OOTA 1965, TAKIMOTO 1981, FUJIOKA et al. 1983b) is inhibited by auxin-like substances. According to OOTA and TSUDZUKI (1971), IAA hampers the flowering of L. gibba at concentrations of 10^{-3} mM and stimulates it at 10^{-6} mM.

Protein content. MACIEJEWSKA-POTAPCZYK et al. (1976) observed a 39% increase of the protein content in solutions with 5×10^{-6} mM IAA. Uptake of the substances and mode of action. BLACKMAN and SARGENT (1959) and BLACKMAN et al. (1959) observed a constant absorption of PAA by L. minor whereas 2,4-D and TIBA are released again after the first two hours. PAA and TIBA are more easily taken up at low pH than at high pH. On the other hand, the effect of BZ in L. minor is greater at pH 6.2

than at pH 4.2 (HILLMAN 1954, 1955); the cell elongation effect is not affected by purines, but neutralized by PCIB. The higher the chlorination of PAA, the faster is the uptake (KENNEY-WALLACE and BLACKMAN 1972). The uptake and the hydrolysis of IAA conjugates in *L. gibba* have been studied by SLOVIN et al. (1984). IAA and IBA substantially increase the inhibitory effect of ABA on growth of *L. gibba*. NAA and 2,4-D do not show any synergistic effect. PCIB used as an antiauxin did not counteract the synergistic action of IAA and ABA. Apparently, the enhancing effect of IAA on the ABA action is not a typical auxin effect (TILLBERG and ELIASSON 1976). A chelating effect of IAA for Cu and Fe ions has been observed by OOTA and TSUDZUKI (1971).

2.3.3.5.5.2. Cytokinins (adenine, BA, iP, kinetin, purine, thiokinetin, zeatin)

Growth rate and weight. Cytokinins (at low concentrations) improve the growth rate and enhance the dry weight of *Lemnaceae* at low light intensities (e.g. 900 lux) and in darkness but have no marked effect at high light intensities (see also chapter 2.3.5.4). It is assumed that under optimal conditions a sufficient amount of cytokinins is formed by the plants themselves to perform maximal cell multiplication. However under suboptimal conditions, the cytokinin level in the plants is lowered. In this situation, the addition of cytokinin has an accelerating effect on cell division (TREWAWAS 1979). ROMBACH (1976) achieved a stimulation of the growth rate in *L. minor* to about 170% by kinetin. The stimulation disappears at light intensities above $200 \mu\text{W cm}^{-2}\text{s}^{-1}$. TILLBERG et al. (1979) enhanced the growth rate to about 110% and the fresh weight to 120-140% with 10^{-6} mM BA and at light intensities of $132 \mu\text{W cm}^{-2}\text{s}^{-1}$. The effects of cytokinins on *Lemnaceae* are surveyed in table 2.18. The effective concentration is dependent on species, form of cytokinin, and culture conditions (e.g. light intensity, day-length, see OOTA 1966 and AL-SHALAN and KANDELER 1978). In general, concentrations of cytokinins higher than 10^{-2} mM are lethal for *Lemnaceae* (PERLMAN and SEMAR 1967). **Size and form of frond.** The fronds of *Lemnaceae* are generally enlarged (even at high light intensities) when cytokinins are added to the culture solution (ROMBACH 1961, 1976, OOTA 1965, LETHAM 1967, MAHESHWARI and VENKATARAMAN 1966, BEZEMER-SYBRANDY 1969, VENKATARAMAN et al. 1970, SCHWEBEL 1973, LE PABIC 1976a, TILLBERG et al. 1979, KLICH et al.

1987c). In *L. minor*, the cell is also enlarged by adenine (WANGERMANN and LACEY 1953). In *L. trisulca*, the frond is not enlarged significantly, but the stalk is elongated (BATA and NESKOVIC 1982a). The fronds of *S. punctata* become vaulted after the addition of zeatin (KERN and NAEP-ROTH 1975).

Table 2.18. Influence of cytokinins on growth rates of *Lemnaceae*

Kind of cytokinin	concentration in mM		species	author
	stimulation	inhibition lethal		
kinetin	10^{-3}		<i>S. punctata</i>	13
	10^{-3}		<i>L. gibba</i>	14
	10^{-2} - 10^{-3}		<i>L. minor</i>	8
	10^{-3}	10^{-2}	<i>L. minor</i>	16
	10^{-2} - 10^{-4}		<i>L. minor</i>	3
	10^{-5} - 10^{-4}		<i>L. minor</i>	6
	10^{-4}	10^{-2}	<i>L. minor</i>	7
	10^{-2} - 10^{-4}	10^{-3}	<i>L. minor</i>	11
	10^{-3} - 10^{-4}		<i>L. trisulca</i>	2
	10^{-2}		<i>L. aequinoctialis</i>	4
thiokinetin	10^{-3} - 10^{-4}		<i>W. columbiana</i>	9
	10^{-5} - 10^{-4}		<i>W. globosa</i>	4
	10^{-2} - 10^{-5}		<i>L. minor</i>	6
	10^{-3} - 10^{-4}	10^{-1} - 10^{-2}	<i>S. polyrrhiza</i>	12
	10^{-3}		<i>S. punctata</i>	13
	10^{-3}		<i>L. gibba</i>	5
	10^{-2} - 10^{-4}		<i>L. minor</i>	17,18
	10^{-3}	10^{-1}	<i>L. minor</i>	3
	10^{-3} - 10^{-4}		<i>L. minor</i>	18
	10^{-3} - 10^{-4}		<i>L. minor</i>	1
iP	10^{-5} - 10^{-7}	10^{-3} - 10^{-4}	<i>L. aequinoctialis</i>	5
	10^{-3}		<i>L. aequinoctialis</i>	15
	10^{-4}		<i>L. aequinoctialis</i>	10
	10^{-4}		<i>L. minor</i>	18
zeatin			<i>S. punctata</i>	13

References

- 1 ALBANELL et al. 1985
- 2 BATA and NESKOVIC 1982a
- 3 BEZEMER-SYBRANDY 1969
- 4 CHANG et al. 1977
- 5 CURTIS 1971
- 6 DEYSSON 1959
- 7 ERISMANN and WEGNER 1967
- 8 HILLMAN 1957
- 9 HILLMAN 1960a
- 10 KOUKKARI and DUKE 1973
- 11 KRAINCIC 1982
- 12 LE PABIC 1967a, b
- 13 LETHAM 1967
- 14 OOTA 1965, 1966
- 15 PERLMAN and SEMAR 1967
- 16 ROMBACH 1961, 1976
- 17 SCHWEBEL 1973
- 18 TASSERON-DE JONG and VELDSIPA 1971a,b

Length of roots. The root growth is inhibited in L. gibba by BA and kinetin (BEZEMER-SYBRANDY 1969), and in L. trisulca by kinetin (BATA and NESKOVIC 1982a). However, BA increases the length of the root of L. gibba at low concentrations (10^{-5} to 10^{-4} mM). Only at higher concentrations (10^{-3} to 10^{-1} mM) there is an inhibiting effect (LIEBERT 1980a). In S. intermedia, the number and length of roots are not affected by BA up to a concentration of 0.1 mg/l (KLICH et al. 1987c). Development and disconnection of daughter fronds. Kinetin (5×10^{-1} mM) and still more effectively BA (10^{-2} mM) stimulate the development of the second daughter frond neutralizing the inhibition by the first daughter frond (GUERN 1965). BA inhibits the disconnection of daughter fronds of L. minor (SCHWEBEL 1973).

Turion formation. Turion formation is completely inhibited at BA concentrations of 10^{-2} mM (JUNGnickel and AUGSTEN 1986). Turion germination. Cytokinins (IP, BA) are able to overcome turion dormancy in S. polyrrhiza and stimulate germination (MALEX and COSSINS 1983).

Flowering. BA inhibits the flowering of L. gibba but not of L. aequinoctialis (CURTIS 1971). Kinetin at high concentrations impedes flowering of L. gibba, too (similar to IAA); at low concentrations (10^{-6} mM) the flowering is stimulated. The flowering of L. aequinoctialis is promoted by cytokinins under short-day conditions, but the critical dark period required is not altered (KHURANA and MAHESHWARI 1983b). The flowering of L. aequinoctialis LP₆ a photoperiodically neutral clone is stimulated by cytokinins in the simultaneous presence of a high iron level (0.1 mM) (KHURANA and MAHESHWARI 1986d). The authors assume that cytokinins might stimulate the effect of chelating agents. Cytokinins induce the flowering of the short-day plant W. microscopica (MAHESHWARI and VENKATARAMAN 1986, VENKATARAMAN et al. 1970). Flowering of S. polyrrhiza and L. minor is enhanced under flowering conditions by kinetin at concentrations of 0.025- 0.05 ppm and 0.05-0.1 ppm, respectively; kinetin stimulates apical floral induction as well as the postinductive stages in the floral development. 0.2 ppm of kinetin inhibit flowering of both species (KRAJNCIC 1982). W. arhiza, too, achieves higher percentage of flowering under flowering conditions (preculture under long-day, culture under short-day conditions) if BA, zeatin, or kinetin is added to the culture solution. Flowering under short-day conditions (without long-day preculture) is induced by BA, zeatin and kinetin; flowering under long-day

conditions is achieved only by application of zeatin (KRAJNCIC 1983). When isolated flower primordia are cultivated in vitro, L. aequinoctialis has a tendency to form more readily the male organs, L. gibba the female ones. Kinetin is able to neutralize this unilateral tendency (HUEGEL 1976c).

Senescence. Senescence in L. gibba is delayed by cytokinins (PARK et al. 1987).

Content of starch and sugars. The starch content of L. minor fronds rises by 200% 4 days after application of kinetin, the content of soluble sugars by 40% (FANKHAUSER and ERISMANN 1969a). This causes the chloroplasts to burst after 5 days (ERISMANN and WEGNER 1967, ERISMANN and FANKHAUSER 1967). According to FANKHAUSER and ERISMANN (1969a) kinetin does not directly influence the sugar metabolism but rather the protein synthesis. Due to a general growth inhibition coupled with a relatively intact photosynthesis, the carbohydrates are stored. Also BA results in starch accumulation in L. minor. The starch content rises from 10% to 25% of the dry weight in a medium with 0.1 mM BA (TASSERON-DE JONG and VELDSTRA 1971a). KLICH et al. (1987c) did not observe any change in the starch content of the cells of S. intermedia after addition of kinetin or BA.

Protein metabolism. Cytokinins enhance the protein content of Lemnaea (ERISMANN and FANKHAUSER 1967, KRZECZOWSKA and ZIMNA 1972, MACIEJEWSKA-POTAPCZYK et al. 1976, LE PABIC 1976a). VAUGHAN et al. (1983) report of a decrease of protein content of L. gibba in solutions with 4 and 40×10^{-4} mM BA. The protein synthesis is accelerated and the decomposition reduced by cytokinins (TREWAWAS 1972a,b). Kinetin is able to enhance the RNA content of L. minor for a short time (ERISMANN and FANKHAUSER 1967); the synthesis as well as the decomposition of RNA are stimulated (TREWAWAS 1970). McCOMBS and RALPH (1972a,b) were able to demonstrate protein synthesis of S. punctata in darkness when kinetin was added to the solution. The formation of DNA and RNA was accelerated. According to BEZEMER-SYBRANDY and VELDSTRA (1971a,b), BA is incorporated in nucleotides of L. minor; with alkaline hydrolysates cytokinins are established in the t-RNA. Kinetin inhibits the degradation rates of mRNAs encoding two major chloroplast proteins in L. gibba (TOBIN and TURKALY 1982). A cytokinin-modulated regulation of mRNA levels for the chlorophyll a/b protein in L. gibba is reported by FLORES and TOBIN (1985). Lipid composition. BERUBE et al. (1982) observed distinct modifications in the lipid composition of L. minor after application of BA: greater

proportions of phospholipids as compared with galactolipids, greater α -linolenic acid content of total neutral lipids. The author supposes that the cell membranes (particularly those of the chloroplasts) and their functions may be altered by high doses of cytokinins. LE PABIC (1980) noted a greater content of total phospholipids after application of 10^{-3} mM BA. The proportion of unsaturated polar lipid fatty acids increased slightly.

Content of chlorophylls, carotenoids and flavonoids. The chlorophyll and carotenoid content in S. polyrrhiza is lowered by 10^{-2} mM BA and slightly raised by 10^{-4} - 10^{-3} mM BA (LE PABIC 1976a,b). The anthocyanin content of S. polyrrhiza is also reduced at kinetin concentrations of 5×10^{-1} mM and at BA concentrations of 10^{-2} mM. ERISMANN and WEGNER (1967) observed a reduction of the chlorophyll content of L. minor in solutions with kinetin. Cyclic adenosine monophosphate (cAMP) treatment inhibits chloroplast replication of W. arrhiza but results in an increased exponential chloroplast growth rate: 260% of that of the kinetin control treatment; whereas cyclic guanosine monophosphate (cGMP) results in a rate of that of the kinetin treatment (HONDA 1983a,b). The flavonoid content of S. intermedia was enhanced by kinetin (10^{-3} mM) in the light but not in the dark (McCURE 1973).

Other effects and mode of action. The growth stimulation caused by BA results in a lower content of calcium in S. polyrrhiza; the content is enhanced again by the addition of ABA (DEKOCK and HALL 1981). Cytokinins generally counterbalance the effects of ABA. BA is able to neutralize the growth inhibiting influence of ABA (VAN OVERBEEK and MASON 1968). Kinetin prevents the turion formation of S. polyrrhiza stimulated by ABA (STEWART 1969). Adenine is able to neutralize the growth stimulating as well as the inhibiting effect of cytokinins (GUERN 1965, PERLMAN and SEMAR 1967). According to BEZEMER-SYBRANDY (1969), adenine and adenosine prevent only the uptake of cytokinins but do not succeed in suppressing the growth stimulating effects. BA has no influence on the pectinase activity in L. minor (SCHWEBEL 1973). However, it decreases the activity of superoxide dismutase and the content of zinc in L. gibba at concentrations of 4×10^{-4} mM (VAUGHAN et al. 1983). Kinetin has no effect on the photosynthesis of L. minor (ROMBACH 1976). Addition of cytokinins reduces the longevity of L. minor in darkness (TASSERON-DE JONG and VELDSTRA 1971b). According to VAN EYCK and VELDSTRA (1966), the kinetin-type activity on the growth of L. minor seems to be restricted to adenine derivatives.

2.3.3.5.5.3. Gibberellins (GA)

Gibberellic acid (GA_3) is transformed to alloligibberic acid in the nutrient solution after autoclaving. This latter substance is more effective than GA_3 in inhibiting flowering of L. aequinoctialis. In a similar way, 1,3-deoxy alloligibberic acid originates from GA_3 (PRYCE 1973a).

Growth rate and dry weight. The growth rate of L. gibba (PIETERSE 1974b, 1975a), of L. minor (LOOS 1962), and of L. aequinoctialis (HILLMAN 1960b, PRYCE 1973a, CHANG et al. 1977) is raised by GA, that of S. intermedia lowered (at a concentration of 10^{-2} mM) (McCURE 1973) and that of L. trisulca remains nearly unchanged but the dry weight per frond is lowered (BATA and NESKOVIC 1982). CCC as a blocker of gibberellin synthesis inhibits growth of L. minor at concentrations of 10^{-3} mM and is lethal at 1 mM (SUPNIEWSKA 1963).

Size of frond. The size of the fronds of L. minor (LOOS 1962), of L. aequinoctialis (HILLMAN 1960b, PRYCE 1973b), and of L. trisulca (BATA and NESKOVIC 1982a) is reduced by GA; however, the length of the stalk in L. trisulca is not altered.

Gibbosity and buoyancy. The gibbosity of L. gibba fronds is lowered drastically after addition of GA to the nutrient solution (PIETERSE 1975a). Correspondingly, addition of CCC results in thicker fronds of L. minor (SUPNIEWSKA 1963). No effect on gibbosity was observed in S. intermedia at concentrations of 0.1 mg GA_3 /l. But at concentrations of 1 and 3 mg/l, the ability of the frond to float was affected by allowing the entrance of water into the air spaces (KLICH et al. 1985). The percentage of submersed fronds increased up to 100. An indirect action of GA on the epidermal structure affecting water permeability is not excluded by the authors.

Epidermis and stomata. GA (1 or 3 mg/l) increases the number of stomata and decreases the number of epidermis cells per unit area in S. intermedia (KLICH et al. 1987d).

Length of roots. The roots of L. minor (LOOS 1962) and of L. trisulca (BATA and NESKOVIC 1982a) are shorter in solutions with GA compared with controls.

Development of daughter fronds and connecting stipes. GA stimulates the growth of the second daughter frond of S. polyrrhiza and L. trisulca; the growth of the stipes is promoted by GA (GUERN 1965). SUPNIEWSKA (1963) reported on inhibited disconnection of daughter fronds in L. minor by the action of CCC.

Flowering. The flowering of *L. aequinoctialis* is inhibited by GA (HILLMAN 1960b, PRYCE 1973b). GA promotes the formation of male organs in explanted flower primordia of *L. gibba* and *L. aequinoctialis* (HUEGEL 1976a,b,c). Flowering of *L. gibba* is stimulated by GA at concentrations of 10^{-2} mM and prevented at 10^{-1} mM (OOTTA and TSUDZUKI 1971). According to PIETERSE (1974b), allogibberic acid inhibits flowering of *L. gibba* at concentrations higher than 10^{-3} mM (under continuous light). HUEGEL and KANDELER (1974) suppose that GA is produced in the meristem of the long-day plant *L. gibba* at lower concentrations than in the short-day plant *L. aequinoctialis*. The short-day plant is believed to contain more GA in relation to the counteracting ethylene. The amount of GA which is taken up from the solution and transported to the meristem is regulated by light through the phytochrome system (HARTUNG and KANDELER 1976). The synthetic compound hexahydrofluorene-9-carboxylic acid has the same effect on flowering of *L. aequinoctialis* as allogibberic acid (PRYCE 1974, 1978). Chapter 2.4.3.1 deals with further information on flowering.

Content of chlorophylls, carotenoids, flavonoids, and fatty acids. The decomposition of chlorophyll in old fronds of *L. trisulca* is prevented by GA. This results in a prolongation of the green colour of old fronds for 2-3 weeks compared with the fronds of controls (BATA and NESKOVIC 1974). Chlorophyll and carotenoid content is reduced at 10^{-2} mM gibberellic acid concentration (D'HARLINGUE 1976). SUPNIEWSKA (1963) reported higher chlorophyll content in *L. minor* with the addition of CCC to the solution. GA₃ at concentrations of 10^{-2} mM inhibits the formation of flavonoids of *S. intermedia* in the light but not in the dark (McCLURE 1973). The total fatty acid content of *S. polyrhiza* is reduced, the sterole content increased in 0.1 mM gibberellic acid (D'HARLINGUE 1976). GA uptake and transport. The uptake of gibberellin in *L. aequinoctialis* and its accumulation and distribution was studied by AL-SHALAN and KANDELER (1979), in *L. gibba* by KANDELER and HARTUNG (1976). The rate of GA transport within the plant amounts to about $2.5-3.0 \text{ mm h}^{-1}$ (HARTUNG and KANDELER 1976).

2.3.3.5.5.4. Abscissic acid (ABA)

Growth rate. ABA inhibits growth processes. The cell elongation of *L. gibba* is hampered by ABA concentrations of 10^{-4} mM (CHEN and PARK 1976). At higher concentrations, cell division as well as cell elongation in

L. minor is prevented (NEWTON 1972b, 1977). According to CURTIS (1971), growth reduction of *L. gibba* and *L. aequinoctialis* can be observed at ABA concentrations higher than 10^{-6} mM. Evidently, the effect of ABA is dependent on culture conditions. Further growth inhibitions are reported by KLICH et al. (1987a) in *S. intermedia*, by STEWART (1969) in *S. polyrhiza*, VAN STADEN and BORNHANN (1969) in *S. punctata*, CHEN and PARK (1976), LIEBERT (1977), AL-SHALAN and KANDELER (1978), DEKOCK et al. (1978), TILLBERG et al. (1979), and DEKOCK and HALL (1981) in *L. gibba*, VAN OVERBEEK and MASON (1968), SCHWEBEL (1973), MCLAREN and SMITH (1976), NEWTON (1977), and ALBANELL et al. (1985) in *L. minor*, CURTIS (1971) in *L. aequinoctialis*, and PIETERSE (1972) in *W. gladiata*. Growth promotion has been observed at low ABA concentrations: at 10^{-7} to 10^{-5} mM in *S. polyrhiza* (MCWHIA and JACKSON 1976, SMART and TREWAVAS 1983a), at 10^{-6} mM in *S. punctata* (VAN STADEN and BORNHANN 1969), and at 10^{-7} to 10^{-4} mM in *L. gibba* (CHEN and PARK 1976, AL-SHALAN and KANDELER 1978).

Size of frond, thickness, and colour. The frond of *L. minor* is smaller in solutions with ABA concentrations of 10^{-2} mM than in controls, and the colour is more intensely green (NEWTON 1972b, SCHWEBEL 1973). The size of the frond of *L. gibba* only amounts to 53-65% of the control in solutions with 10^{-4} mM ABA (TILLBERG et al. 1979). The layers of air spaces in *L. minor* are reduced by ABA (SEVERI and BARONI FORNASIERO 1983a). KLICH et al. (1987a) did not see any change of aerenchymatic tissue of *S. intermedia* after addition of ABA. However, the frond size became smaller at concentrations of 10^{-3} mM or higher.

Length of root. ABA inhibits the growth of the root of *L. minor*. Both, maximum length and number of the cells are reduced (NEWTON 1972b, 1974a, 1977). In a similar way growth of *L. gibba* roots is hampered at ABA concentrations of 10^{-5} mM and higher (LIEBERT 1980a). In *S. intermedia*, the root length is reduced by ABA but the root number is not affected (KLICH et al. 1987a).

Development and disconnection of daughter fronds. ABA treatment results in the production of fewer daughter fronds in *L. minor* (SCHWEBEL 1973) and in *L. gibba* (KANG and CLELAND 1985). The disconnection of daughter fronds of *L. minor* is induced by ABA (SCHWEBEL 1973, OSTROM-SCHWEBEL 1979). The area of disconnection in the stipe of *L. minor* was studied by OSTROM-SCHWEBEL (1979) in detail. Contrary to these results, NEWTON (1972b, 1977) observed a retardation of the disconnection of *L. minor* in

solutions containing ABA; the fronds remained in clusters. However, his concentrations of ABA were nearly three times stronger (2 ppm) than those of SCHWEBEL (1973).

ion formation. Turion formation of S. polyrhiza is stimulated by ABA concentrations of 10^{-3} to 10^{-5} mM (PERRY and BYRNE 1969, STEWART 1969, SAKS et al. 1975, 1980, SMART and TREWAVAS 1983a). The same is true for the formation of resting fronds of L. minor (VAN OVERBEKE et al. 1968, SEVERI and BARONI FORNASIERO 1983a), of L. perpusilla and of aequinoctialis (KANDELER and HUEGEL 1974a). PERRY and BYRNE (1969) report a mutant of S. polyrhiza which showed neither growth inhibition or turion formation in response to ABA. The ABA induced turions of S. polyrhiza are characterized by reduced Ca^{++} exchange across the tonoplast and low vacuolar Ca^{++} content compared with vegetative fronds. In addition, the turions exhibit a higher plasmalemma flux with a correspondingly high Ca^{++} concentration in the cytoplasm. The accumulation of Ca^{++} and Cl^{-} is much lower in the cytoplasm of the ABA induced turion than the vegetative frond (SMART and TREWAVAS 1984b). For further information on turion formation see chapter 2.4.2.

flowering. ABA (at concentrations of 0.1-1 µg/l) significantly promoted flowering of the long-day plant L. minor under long-day conditions. At higher concentrations (10-50 µg/l), it inhibited or suppressed the flowering of S. polyrhiza, L. gibba, L. minor, and W. arhriza. ABA (0.1-1 µg/l) in combination with EDDHA (7 mg/l) had a synergistic effect on flowering of L. minor and S. polyrhiza (KRAJNCIC 1985). At concentrations of 0.1 µg/l, ABA enhances flowering of L. minor (especially the initial stages of apical flower induction), at concentrations of 10 µg/l and higher, it inhibits flowering (KRAJNCIC and BAVEC 1986). Increasing concentrations of ABA reduced stamen development in explanted flower buds of L. gibba and pistil development in such explants of L. aequinoctialis (HUEGEL 1976c). ABA counteracts the flowering effect of (FUJIOKA et al. 1983b).

Senescence of L. gibba is prematurely induced by ABA (PARK et al. 1987). The starch and sugar content of Lemnaceae enhanced in solutions with ABA (VAN STADEN and BORNWANN 1969, McLAREN and SMITH 1976, 1977, DEKOCK et al. 1978, TILLBERG et al. 1979, KLICH et al. 1987a). The uptake of glucose and glycine is stimulated by ABA (HARRINGTON et al. 1980). SEVERI and BARONI FORNASIERO (1983a) observed plas-

tids with especially large starch grains in L. minor growing on solutions containing ABA. THORNSTEINSSON et al. (1985) recorded increasing amounts of free sugars in L. gibba in solutions with higher ABA content. The effect of ABA was in part counteracted by BA.

Mineral composition. ABA changes the mineral composition in the frond of L. gibba; it lowers the P/Fe ratio from 33 to 6 and the K/Ca ratio from 3 to 0.6 (DEKOCK et al. 1978). ABA (10^{-3} mM) lowers the K content in L. gibba from 60 mg/g dry weight to 20 mg (LIEBERT 1977).

Chlorophyll and carotenoid content. ABA reduces the chlorophyll content of L. minor and enhances the carotenoid content (SEVERI and BARONI FORNASIERO 1983a). ALBANELL et al. (1985) observed a higher chlorophyll content in fronds of L. minor at concentrations of 10^{-3} mM ABA.

Photosynthesis and respiration. The CO_2 fixation in L. minor is inhibited by ABA; a direct effect on the transport of electrons of the photosystem II is supposed (BAUER et al. 1976). Also TILLBERG et al. (1981) noted an inhibition of the CO_2 uptake in L. gibba after addition of ABA to the solution. The activity of photosynthetic enzymes in L. minor is, however, enhanced by the action of ABA, according to SANKHLA and HUBER (1979). Respiration in the dark was stimulated within the first 2 to 7 days. Unlike other phanerogams, Lemna does not close the stomata after ABA application (TILLBERG et al. 1981).

Other life processes and mode of action. DEKOCK et al. (1978) observed a rise of enzyme activities of phosphatase, peroxidase, amylase and invertase in L. gibba after application of ABA. The RNA content is, according to this author, higher. On the other hand, NEWTON (1972b, 1974a) reports an inhibition of the RNA synthesis of L. minor in solutions containing ABA. Also TREWAVAS (1970) noted a markedly reduced synthesis rate of ribosomal RNA but unaltered degradative rates in L. minor after adding ABA. According to ANDRES and SMITH (1976), the content of free amino acids in fronds of S. polyrhiza, L. minor, and L. aequinoctialis is raised by ABA; the same is true for the release of amino acids to the culture solution. The authors assume that ABA increases the permeability of the cell membrane. ABA does not influence the ability of S. polyrhiza to induce nitrate reductase (STEWART and SMITH 1972). The cellulase activity of L. minor becomes enhanced by 3×10^{-3} mM ABA whereas pectinase activity is not significantly affected (SCHWEBEL 1973). VAUGHAN et al. (1983) recorded an increase of superoxide dismutase and zinc, copper, and protein content in L. gibba at ABA concentrations of 4 to 40x

10^{-4} mM. DNA synthesis in the developing turions of S. polyrrhiza which have been induced by ABA is inhibited by ABA and followed by a repression of protein synthesis while RNA synthesis was not altered (SMART and TREWAVAS 1984a).

The effects of ABA are partly neutralized by BA and other cytokinins but not by gibberellins (VAN OVERBEEK and MASON 1968, VAN STADEN and BORNHANN 1970b, DEKOCK et al. 1978, ALBANELL et al. 1985). The turion stimulation by ABA is also suspended by kinetin (SAKS et al. 1975).

2.3.3.5.5.5. Ethylene and ethylene releasing substances (ethrel, ACC)

The effect of ethylene on Lemnaceae is partly similar to that of GA and partly to that of ABA. The effect of ethylene is reversed by autoclaved IA_3 (PIETERSE 1976, 1982).

Growth rate. At concentrations of 4×10^{-2} mM, ethylene promotes the growth rate of S. polyrrhiza by 50%; at 4×10^{-1} mM it inhibits the growth rate (NEGBI et al. 1972).

Size of frond and gibbosity. An enlarging effect on fronds of L. gibba in a mould-infected culture was explained by WETSTEYN et al. (1984) by the assumed production of ethylene by the mould. Gibbosity of L. gibba is caused by ethylene at concentrations of 10^{-3} mM and higher. Similar concentrations could be measured in the field at places where gibbous L. gibba occurred; at places where flat L. gibba was present, the ethylene concentration was significantly lower (ELZENGA et al. 1980). Chloroethanephosphonic acid (ethrel) which was found to release ethylene within the Lemna fronds has the same effect on gibbosity as ethylene (PIETERSE 1976).

Nerves and disconnection of the daughter fronds. Ethylene (0.1%) was found to induce the 4th and 5th nerve in L. minor (SARGENT and WANGERMAN 1959). The disconnection of daughter fronds of S. polyrrhiza is accelerated by ethylene (NEGBI et al. 1972).

Flowering. According to PIETERSE (1982), ethylene does not induce or enhance flowering in L. gibba, but SCHARFETTER et al. (1984, 1986) report a role of ethylene in flowering of Lemnaceae. Investigations with ACC, a direct precursor of ethylene, showed a stimulation of flowering in the long-day plant L. gibba at 10^{-5} mM. At 3×10^{-4} mM ACC was flower-inhibiting. The flowering of the long-day plant S. punctata was prevented by 10^{-6} to 10^{-4} mM ACC. In contrast, flowering in the short-day plant L.

aequinoctialis was promoted by 10^{-4} to 10^{-3} mM ACC under long-day conditions (SCHARFETTER et al. 1987). Ethrel promotes the development of female flower parts in explanted flower primordia of L. gibba and L. aequinoctialis (HUBGEL 1976b). For more details on effects of ethylene on flowering of Lemnaceae see chapter 2.4.3.1.

Senescence. Ethylene induces premature senescence in L. gibba (PARK et al. 1987).

2.3.3.5.6. Toxic substances and phytochemical substances not classified

2.3.3.5.6.1. General remarks

There is an enormous literature on toxic effects of different chemicals on growth and behaviour of Lemnaceae which is not easy to survey. A rather incidental selection is presented here. The reasons for the extensive work are as follows:

- 1) Lemnaceae are suitable test plants for checking the toxicity to organisms of new chemicals (e.g. pesticides, detergents) (see chapter 3.6)
 - 2) In some regions it is necessary to keep mass development of Lemnaceae under control (duckweeds are obstacles for shipping, bathing; they may deteriorate the quality of fish ponds etc.).
- Some of the investigated substances behave as hormone-like factors at lower concentrations and are therefore dealt with in chapter 2.3.3.5.5. A few substances toxic for Lemnaceae are put together in tables 2.19 and 2.20.

2.3.3.5.6.2. Herbicides

In general, Lemnaceae are very sensitive to herbicides. They are often used as test organisms to detect the presence of herbicides in water: e.g. L. minor for prometryn and diuron (DECLERE and DE CAT 1977), S. polyrrhiza for ametryn and cyanatryn (SUTANTO et al. 1976) and for simazine, linuron, bromacil, fluorodifen, paraquat, and MCPA (DAMANAKIS 1970, 1972) (see also chapter 3.6).

Heterocyclic compounds (e.g. 6-methylpurin), urea derivatives, and tertiary ammonium compounds belong to the most toxic substances for Lemnaceae. They still affect the growth rate of Lemnaceae at 10^{-4} to 10^{-5} mM.

Table 2.19. Effects of some herbicides on Lemnaceae

	Concentration in mM inhibiting	lethal or inhibiting >50%	species	author
<u>Inorganic compounds</u>				
Ammonium sulfamate		1	d	7
Ammonium thiocyanate		1	d	7
Sodium chlorate		5	d	7
<u>Phenol derivatives</u>				
Nitrofen	10 ⁻³	10 ⁻³	c	33
Fluorodifen	10 ⁻⁴	10 ⁻³	a	5
<u>Urea derivatives</u>				
Diuron, DCNU	10 ⁻⁴ -10 ⁻³		c	14
	10 ⁻⁵	10 ⁻³	e	24
	10 ⁻⁵	10 ⁻³	m	18
	10 ⁻⁵	10 ⁻³	m	18
	10 ⁻⁴		a	5
		10 ⁻⁴	h	35,36
		10 ⁻⁴	h	35,36
<u>Fluometuron</u>				
Linuron	10 ⁻³		d	6
Chlorobromuron	10 ⁻³	10 ⁻²	a	31
Diphenoxystyrene	10 ⁻³	10 ⁻²	d	1,2,7,27,30
2,5-Dimethyldodecanoic acid	10 ⁻³		c	22
Aliphatic carbonic acids and nitriles	10 ⁻³		g	34
Chloroflurenol	10 ⁻⁴		a	4,5
2,4-D	10 ⁻⁴		d	19,20
MCPA	10 ⁻⁴		d	10
<u>Amines</u>				
Ethylfluralin (10 ⁻⁵ mM stimulating)	10 ⁻⁵	10 ⁻⁴	d	10
<u>Carbonic acid amides</u>				
Alachlor	10 ⁻⁵	10 ⁻⁴	a	0,4,5
Quaternary ammonium compounds	10 ⁻⁵	10 ⁻⁴	a	0,4,5
Paraquat	10 ⁻⁴	10 ⁻³	d	20
<u>Heterocyclic compounds</u>				
Fluridone	10 ⁻⁴	10 ⁻³	h	35,36
Lenacil	10 ⁻⁴	10 ⁻³	a	5
Bromacil	10 ⁻⁴	10 ⁻³	a	5
Sisazone	10 ⁻⁴	10 ⁻²	d,k	15
Atrazine (10 ⁻⁴ mM stimulating)	10 ⁻³ -10 ⁻²	10 ⁻¹	a,c,m	16,17,18
CPS (10 ⁻⁴ , 10 ⁻⁴ mM stimulating)	10 ⁻¹		a,c	26
CTS	10 ⁻²		a	26
	10 ⁻⁴	10 ⁻³	c	26
	10 ⁻⁵	10 ⁻³	m	18
	10 ⁻⁵	10 ⁻³	d	20
	10 ⁻⁵	10 ⁻²	m	18
	10 ⁻⁵	10 ⁻²	d	21
	10 ⁻⁵	10 ⁻²	d	21
<u>Thiouracil, TU</u>				

Table 2.19 (continued)

	Concentration in mM inhibiting	lethal or inhibiting >50%	species	author
Purin		10 ⁻²	e	23
6-Chloropurin		10 ⁻²	e	23
6-Methylpurin, MP	10 ⁻⁸	10 ⁻⁵	e	23
		10 ⁻³	c	32
<u>Organic compounds containing phosphorus</u>				
Glyphosate, Phosphonomethylglycine		10 ⁻⁴ -10 ⁻²	c	11
SC-0224	10 ⁻²		d	9,25
9-Amino-9-fluorenephosphonic acid	10 ⁻²		c	6a
9-Amino-9-fluorenephosphine oxides	10 ⁻²		b	3
<u>Organic compounds containing arsenic</u>				
MSMA	10 ⁻¹		m	8,13
				18

Species

- a Spirodela polyrrhiza e L. aquinoctialis i W. columbiana
b S. punctata f Wolffia gladiata k Wolffia sp.
c Lemna gibba g Wolffia brasiliensis l Lemna sp.
d L. minor h W. arrhiza m duckweeds

References

- 0 BLACKBURN and WELDON 1965 18 LIU and CEDENO-MALDONADO 1979
1 BLACKMAN 1952 19 LOCKHART and BLOUW 1979
2 BLACKMAN and 20 LOCKHART et al. 1983
ROBERTSON-CUNNINGHAM 1954 21 NICKELL 1962
3 CZERWINSKI et al. 1982 22 OOTA 1966
4 DAMANAKIS 1970 23 PERLMAN and SEMAR 1967
5 DAMANAKIS 1972 24 FOSNER et al. 1977
6a FOY and COOLEY 1987 25 PRASAD 1984
6 ENTZEROOTH et al. 1985 26 RASOL and RENDIC 1977
7 FROMM 1951 27 SHAMBHU et al. 1976
8 GANCARZ et al. 1985 28 SIMON and BLACKMAN 1953
9 HARTMAN and MARTIN 1984 29 STEPHENSON and KANE 1984
10 HARTMAN and MARTIN 1985 30 SUPNIEWSKA 1963
11 HONGLAND and PAUL 1978b 31 TREICHEL 1974a
12 JAWORSKI 1972 32 TROGICH and ULLRICH 1984
13 KAFARSKI et al. 1985 33 WEJNAR and TALS 1983
14 KANDELER 1969b 34 WORTHLEY and SCHOTT 1971
15 KARELS and LEMBI 1973 35 ZAWADZKI 1974
16 KRSNIK-RASOL and RENDIC 1975 36 ZAWADZKI and REJMAN 1978
17 KRSNIK-RASOL and RENDIC 1977

The following classification follows the monographic treatment of herbicides by WEGLER (1977).

Inorganic compounds

Ammonium sulfamate, ammonium thiocyanate, and sodium chlorate were tested by FROMM (1951) on L. minor. They proved to be moderately toxic (lethal at c. 1 mM). Copper sulfate, sodium arsenite, ammonium thiocyanate, and ammonium 10% were used as foliar spray solutions on S. polyrrhiza and showed lethal effects at the same concentrations (SAHAI et al. 1980).

Phenol derivatives

Of 36 tested nitrophenols 12 impeded the formation of chloroplasts in Lemnaceae (PRICE and WAIN 1976). Whereas phenol only inhibits L. minor at concentrations of 5 mM, 2,4-dinitrophenol does already at 10^{-2} mM (BLACKMAN et al. 1954). The biological effect of substituted phenols was enhanced if alkyl side chains were lengthened or the number of substituted chlorine atoms or alkyl groups in the ring were increased. The substitution on the para position was more toxic than substitution on the ortho position (BLACKMAN et al. 1954, SIMON and BLACKMAN 1953).

Nitrofen at concentrations of 0.1 ppm (c. 5×10^{-4} mM) inhibits growth rate and root growth of L. gibba and lowers the content of pigments, chlorophylls a and b, β -carotene, lutein, violaxanthin, and neoxanthin (WEJNAR and TAIS 1983). It destabilizes the chloroplast structure: formation of cisterns within the stroma, widening of the thylakoid interiors, and disappearance of the grana structure (at nitrofen concentrations of 2×10^{-4} , 5×10^{-4} , 5×10^{-3} mM) (WEJNAR and MICHEL 1983).

Pentachlorophenol (PCP) inhibits the photosynthetic O_2 production of L. minor and decreases its chlorophyll content. There is no pronounced effect on O_2 consumption during respiration in the dark. The influence of the toxin on many enzymes was studied (HUBER et al. 1982). Of 11 aquatic animal and plant species tested, L. minor was the least sensitive to PCP with no apparent effects up to concentrations as high as 1.4 mg/l (HEDTKE et al. 1986). Another study with pentachlorophenol was made by YUI and KOIKE (1955) and YUI (1958).

Carbonic acid and thiocarbonic acid derivatives

CIPC in concentrations up to 10 mg/l has no effect on the photosynthesis of S. polyrrhiza (BIELECKI and SKRABKA 1976).

Urea derivatives

Diuron (DCMU), a photosynthetic inhibitor inhibits growth and stimulates

flowering in L. gibba (KANDELER 1969b). This effect is not due to altered sizes of ATP pools (GOWER and POSNER 1979). DCMU also inhibits growth of L. aequinoctialis at concentrations of 10^{-5} mM and enables flowering under long-day conditions. A mutant type of strain 6746 is stimulated by concentrations up to 10^{-3} mM DCMU (POSNER et al. 1977). The degradation of the 32 kDa protein in S. punctata is inhibited by DCMU (MATTOO et al. 1986b). Diuron and fluometuron stimulate the activity of the nitrate reductase in S. polyrrhiza at concentrations of about 10^{-5} mM. The activity is lowered at 10^{-3} mM (diuron) or 10^{-2} mM (fluometuron) (LIU and CEDENO-MALDONADO 1979).

Araliphatic carbonic acids and nitriles

Morphactins as chlorofluoreneol lower growth rate, length of root, rate of respiration, and photosynthesis of L. gibba. They enhance the content of starch and chlorophyll as well as the level of ATP and the activity of amylase and phosphorylase (TREICHLER 1974a,b). KAFARSKI et al. (1985) investigated the toxicity of phosphonic analogues of morphactins in S. punctata. The glycylglycyl and threonyl peptides showed great activity in reducing dry weight.

2,4-D has a relatively low activity against Lemnaceae. The activity is dependent on the pH of the solution: at 4.6 the effect on L. minor is ten times greater than at 6.1 (BLACKMAN and ROBERTSON-CUNNINGHAME 1953). FERNANDEZ et al. (1972) report a similar pH dependence of the 2,4-D effect on S. intermedia (compare chapter 2.5.2.3.2). At higher light intensities, the effect of low concentrations of 2,4-D is greater than at lower ones. IAA partly neutralizes the inhibiting influence of 2,4-D on L. minor. At high concentrations of 2,4-D, the light intensity is of no significance for the toxic effect. However, a pretreatment with low light intensity enhances the effect (BLACKMAN and ROBERTSON-CUNNINGHAME 1954). 2,4-D is absorbed by S. intermedia mainly by the roots, according to FERNANDEZ et al. (1972). If polystyrene is used as a carrier substance, 2,4-D shows a long-time effect in water. To suppress growth of L. minor, it is necessary that the concentration of 2,4-D does not sink below 10^{-1} mM (SHAMBHU et al. 1976).

2,4-D and CIPC have no effect on the photosynthesis of S. polyrrhiza at concentrations up to 10 mg/l (BIELECKI and SKRABKA 1976). TITOVA (1978a) reports the accumulation of 2,4-D in Lemnaceae. TCDD, a very poisonous impurity of the herbicide 2,4,5-T is accumulated 1000 to 5000 times by L. minor (ISENSE and JONES 1975a) (see also chapter 3.5.3).

2,4,5-T and 2,4-D have been investigated by SAHAI et al. (1980) applied as a foliar spray on S. polyrrhiza at concentrations of 100 ppm and higher. In a similar way, KHARE (1977, 1979) spread MCPA, 2,4-D, and 2,4,5-T on the surface of S. polyrrhiza fronds. Necrosis showed up at concentrations of 1 to 10^{-1} mM. Control of duckweeds in fish ponds was accomplished successfully by spraying 0.7 g/ha endothal and 0.09 g/ha triad which together proved very toxic to Lemnaeae. The herbicide mecoprop killed S. punctata within 10 to 20 hours in sublethal doses of 10^{-1} mM if 0.5-1.5 $\times 10^{-2}$ mM of derivatives of 9-amino-9-fluorenephosphonic acid were added simultaneously (CZERWINSKI et al. 1982).

Quaternary ammonium compounds

Paraquat is very toxic to S. polyrrhiza (DAMANKIS 1970, 1972). A similar effect on Lemnaeae is caused by diquat (FUNDERBURN and LAWRENCE 1963, BIRMINGHAM and COLMAN 1983). Diquat and paraquat give rise to chlorosis in S. polyrrhiza, L. minor, W. gladiata, and W. columbiana; most sensitive is S. polyrrhiza, least W. columbiana. After seven days treatment with 0.01 ppm diquat (in brackets paraquat) the following percentage of chlorosis can be observed: 95% (80%) in S. polyrrhiza, 85% (70%) in L. minor, 75% (65%) in W. gladiata and 5% (5%) in W. arhiza. The activity of the herbicides rises with light intensity; it is highest in red light and lowest in green light (BLACKBURN and WELDON 1965). Diquat at 1.1-3.4 kg/ha gave the best long-term control of S. polyrrhiza in Florida (better than glyphosate or endothal) (THAYER and HALLER 1985).

Orellanine, a toxic substance produced by the basidiomycete Cortinarius speciosissimus has a similar structure as diquat and paraquat. It destroys the photosynthetic pigments and the chloroplasts of L. minor in the same way as the two herbicides (HOLLAND 1983). Photosynthetic activity of L. minor is efficiently inhibited by orrellanine at a concentration of 0.4 mM (RICHARD et al. 1987).

The change in cell membrane permeability of L. minor was the criterion used by O'BRIEN and PRENDEVILLE (1978, 1979) to detect the presence of paraquat and diquat in water. The minimum herbicide concentration that could be detected ranged from 0.00017 to 0.00018 ppm (ca. 10^{-7} to 10^{-6} mM). Diquat remains toxic to Lemnaeae for many weeks after treatment if it is absorbed by clay minerals of the aquatic sediment (BIRMINGHAM and COLMAN 1983). A mixture of diquat and cutrine plus was most effective against Lemnaeae for application over wide areas (HARGROVE 1976).

Heterocyclic compounds

NICKELL (1962) investigated the effect of 87 pyrimidines on L. minor, 7 of which (including 2,4-dichloropyrimidine and thiouracil) proved to be strongly inhibiting at concentrations of 10^{-2} mM. Eight compounds had a stimulating effect at the same concentration. Naturally occurring cytosins did not show any effect on L. minor. UMEMURA and OOTA (1965a) studied the effect of FDU, FU, TU, and other antimetabolites and of corresponding metabolites (thymidine, orotic acid, uracil, guanine, methionine) on growth and flowering of L. gibba. All antimetabolites reduced flowering. RIMON and GALUN (1967) also investigated the impact of pyrimidine base analogues on growth of S. punctata. FDU stopped cell division at 4×10^{-4} mM and higher. This effect could be multiplied by the addition of 4×10^{-2} mM thymidine. Neither uridine nor uracil had an antitotal effect. FU at 10^{-2} mM affected cell division, frond elongation, and differentiation. No counteraction by thymidine or uracil was visible. TU inhibited differentiation of frond tissue rather specifically. Uracil and uridine counteracted the effect. Atrazine at sublethal doses raises the content of water, chlorophyll, soluble protein, nitrogen, and DNA of L. minor and lowers its photosynthesis. At low concentrations, however, the photosynthesis is stimulated (BEAUMONT et al. 1976, 1978), and the total lipid content enhanced (GRENIER et al. 1979). Phosphatidyl glycerol content is increased and the content of phosphatidyl ethanolamine and of phosphatidyl inositol decrease in the presence of atrazine (GRENIER et al. 1982). Also, sublethal doses of atrazine stimulate the desaturation of fatty acids and the lipid metabolism of the chloroplasts (increase of the number of grana per chloroplast) (GRENIER and BEAUMONT 1983). MATTOO et al. (1984b) noted a higher linolenic/linoleic acid ratio in the thylakoid membrane lipids of S. punctata under sublethal concentrations of atrazine. Fourteen days after application of atrazine, the chlorophyll content is greatly reduced and fronds become chlorotic (GUNKEL 1983). Atrazine inhibits the 32 kDa protein degradation in S. punctata (MATTOO et al. 1986b). GRESSEL (1982a,b) reports on the possible triazine interaction with a 32000 M_r thylakoid protein in S. punctata. Triazine inhibits photosynthesis by binding to thylakoids. Simazine inhibits the photosynthesis rate of L. minor at concentrations of 1 ppm (SUTTON et al. 1969). The substance was used as a foliar spray solution (at concentrations of 100 ppm and higher) on S. polyrrhiza by SAHAI et al. (1980). Ametryn and prometryn stimulate the activity of the

nitrate reductase in *S. polyrrhiza* at concentrations of about 10^{-8} mM. The activity decreases at concentrations of 10^{-3} mM. BAHADIR and PFISTER (1985) report a sufficient checking of *L. minor* in a simulated flow system by 20 ppb (c. 10^{-4} mM) terbuthryn. Terbuthryn at concentrations of 10^{-3} mM is destroyed faster in ponds with Lemnaceae and *Typha* sp. than in ponds without water plants, the half-life period being 21 days instead of 30 days (MUIR et al. 1981). In channels sprayed with terbuthryn or cyanatryn, Lemnaceae are able to thrive due to lack of competition (MURPHY et al. 1981). Of all herbicides investigated, 6-methylparin was most toxic. PERLMAN and SENAR (1967) observed an inhibiting effect on *L. aequinoctialis* at concentrations as low as 10^{-8} mM. At concentrations of 10^{-3} mM it completely inhibits the growth rate of *L. gibba*, and reduces chlorophyll and protein content and the activity of nitrate reductase (TROGTSCH and ULLRICH 1984).

Organic compounds containing phosphorus

Glyphosate is not very effective against *L. minor* and *W. arnhiza* according to KARELS and LEMBI (1973). They had to apply about 1 mM to get toxic effects. However, HARTMAN and MARTIN (1984) noted that glyphosate is lethal to *L. minor* at concentrations of about 10^{-4} mM. If suspended betonite clay was added to the solution, glyphosate was not toxic at the same concentration due to unavailability of the herbicide bound to clay particles. N-phosphonomethylglycine appears to inhibit the aromatic amino acid biosynthetic pathway in *L. gibba* (JAWORSKI 1972).

Glyphosine, a plant growth regulator which causes chlorosis in maize and increases sucrose levels in sugar cane, inhibits the growth rate of *L. gibba* at 1 mM and stops it at 10 mM. In solution with 5×10^{-1} mM glyphosine, newly formed fronds are white or pale yellow and contain less than 0.4% of the chlorophyll of a green frond (SLOVIN and TOBIN 1980, 1982). Lemnaceae often concentrate herbicides within their fronds.

Further organic compounds used as herbicides

Hydrothol-47 and TD-1929 show a toxic effect on *Wolffia* sp. but not on *Lemna* sp. at concentrations of 1-3 ppm (c. 10^{-2} mM) (KARELS and LEMBI 1973).

Cyanoacrylate herbicides are potent inhibitors of the photosynthetic electron flow in isolated chloroplast systems. They are highly toxic to *L. minor* and other water plants (HUPPATZ and PHILLIPS 1981).

SAHAI et al. (1980) tested benzene, kerosene, and xylene as a foliar spray on *S. polyrrhiza*. Toxic effects showed up at c. 1 mM and higher.

Table 2.20. Effects of some algicides, fungicides, insecticides, and other toxic substances on Lemnaceae

Substances	concentrations in mM inhibiting	lethal or inhibiting > 50%	species	author
Algicides				
HOE 2977		10^{-2}	d	2
PH 40:62		10^{-3}	d	10
	5×10^{-3}	5×10^{-2}	b	15
Fungicides				
QAC		10^{-2}	c	1
Cuso		10^{-2}	c	1
Insecticides				
2,4,6-ICP	10^{-4}	10^{-1}		
Metacil		10^{-2}	d	5
Methyl parathion	$> 10^{-2}$		d	11
DDT	10^{-1}		f	16
Detergents				
DBST, ENF (10^{-4} mM stimulating)		10^{-2}	a	13
"Super Surf"		10^{-1}	a	7
Bleaching substances				
Phenol, Catechol	10^{-2}	10^{-2}	e	6
	10^{-2}	10^{-1}	d	12
Guaiacol	10^{-2}	10^{-1}	e	6
9,10 Epoxystearic acid	10^{-1}	1	e	6
Disinfectants				
o-Cresol	10^{-2}	10^{-1}	c	1
CS		10^{-1}	b, e, f	8
Other toxic substances				
Patulin	10^{-5}	10^{-2}	c	9
Aflatoxin	10^{-2}		d	3
Hydroquinone			d	12
Glycolate, Glyoxylate (0.5-3 mM stimulating)	5×10^{-2}	10^{-1}	c	14
DMPP, DMNH	10^{-2}		b	4

Species
a *Spirodela polyrrhiza* c *Lemna gibba* e *L. aequinoctialis*
b *S. punctata* d *L. minor* f *Wolffia brasiliensis*

References

- 1 DAVIS 1981
- 2 HARTZ et al. 1972
- 3 JACQUET and BOUTIBONNES 1970
- 4 KNYPL and OSWIECINSKA 1986
- 5 PRASAD 1981
- 6 ROWE et al. 1982
- 7 SAHAI et al. 1977a, b
- 8 SCHOTT and WORTHLEY 1973
- 9 SLOVIN et al. 1984
- 10 SPENCER-JONES 1981
- 11 STEPHENSON and KANE 1984
- 12 STOM and ROTH 1981
- 13 TATKOWSKA and TOPOROWSKA 1978
- 14 TILLBERG 1980
- 15 WALKER and EVANS 1978
- 16 WORTHLEY and SCHOTT 1971

The herbicides arealine, atrataf, tafazine, tafapon, and amitrole were spread on the frond surface of S. polyrrhiza by KHARE (1977, 1979). They produced necrosis at concentrations of 1 to 10^{-1} mM.

2.3.3.5.6.3. Algicides and fungicides

The algicides investigated so far (2-dichloroacetamido-3-chloro-1,4-naphthaquinon; quaternary ammonium compounds; PH 40:62) mostly have a stronger effect on Lemnaceae than on water plants rooted in the soil (HARTZ et al. 1972, WALKER and EVANS 1978, SPENCER-JONES 1981). Fungicides such as 2,4,6-TCP are rather toxic to L. gibba (DAVIS 1981). Cyanobacterin, a natural algicide controlled growth of Lemna if added to the medium (GLEASON and CASE 1986; see also KIRPENKO 1986).

2.3.3.5.6.4. Insecticides and other pesticides

Insecticides spread in nature are often concentrated in Lemnaceae. S. polyrrhiza and L. minor are able to accumulate up to 2 mg DDT per kg fresh weight and up to 1.2 mg HCHC (VROCHINSKIY et al. 1970). The toxin becomes dissolved in the lipids of the cell membrane and is excreted on the inner side of the cell. The dead fronds store DDT at the bottom of the water (VROCHINSKIY et al. 1971). Of 10 pesticides studied, DDT and 2,2',4,4',5,5' hexachlorobiphenyl become most enriched in L. minor. The concentration of the chemicals was more than 1000 times higher than in the water (LOCKHART et al. 1983). Polychlorinated biphenyls (PCB) only slightly soluble in water, are greatly accumulated by Lemnaceae. Twenty-one days after application, the plants contained 0.023 mg/g dry weight, 70 times more than Scirpus, and 6 times more than Eichhornia (ENGLAND and KAIGATE 1981). Five ppm atrachlor 1242 a polychlorinated biphenyl produces a characteristic chlorosis pattern in S. punctata; chlorophyll content is reduced to 56% and content of nucleic acid to 73% (desoxyribonucleic acid content remains unchanged) (MAHANTY and MCWHIA 1976). Fenitrothion is concentrated 6 times by L. minor in an aquatic ecosystem but is decomposed within a short time (MOODY et al. 1978). MALIS and MUIR (1984) measured a 700fold concentration (up to 23 ppm) of fenitrothion in Lemnaceae (dry weight) five days after application. The amount is reduced to about 10% 77 days later. Dowanol TPM (tripropylene glycol-methylether) is in use as an adjuvant in fenitrothion spray formula-

tions. At 5×10^{-3} mM it has a specific deleterious effect on the photosynthetic activity of L. minor (WEINBERGER and CAUX 1985). The insecticide matacil containing nonylphenol proved to be lethal to L. minor at concentrations of 10^{-2} mM. Probably the membranes of chloroplasts are destroyed causing the chlorophyll to emerge (PRASAD 1981). The other components of matacil (aminocarb, diluent oil) are not very effective (WEINBERGER and IYENGAR 1983).

Carbofuran does not show any toxicity to L. minor in concentrations up to 10 mg/l (HARTMAN and MARTIN 1985). The evidence of alkylbenzene sulfonate and ammonium carbamate in water was established by WARD et al. (1981) using S. polyrrhiza and L. minor as test organisms.

Permethrine is also readily absorbed by Lemnaceae, but is not persistent (RAMN et al. 1982).

Carbaryl and 3,5-xylyl methylcarbamate (XMC) is accumulated up to 4000-fold in Lemnaceae (on a dry weight basis) compared to 100-500fold in snails, fish and crustaceae (KANAZAWA et al. 1975).

The fluorinated derivatives of SA which are used as molluscicides damage L. minor whereas other higher plants are not affected (WILLOWITZER et al. 1972).

Sodium fluoracetate, a pesticide against rabbits and opossum inhibits the growth of S. polyrrhiza, S. punctata, and L. minor at concentrations of 10^{-2} mM (BONG et al. 1980).

Dinitro-ortho-cresol (DNOC) reduces the chlorophyll and carotenoid content in L. minor at a concentration of 10 ppm (KRIVOKAPIC and ERIC 1984).

2.3.3.5.6.5. Detergents

A stimulating effect of the detergents DBST and ENF on Lemnaceae at low concentrations is explained by TATKOWSKA and TOPOROWSKA (1978) with a chelating action. At higher concentrations (10^{-2} to 10^{-1} mM), the detergents damage the fronds. An inhibition of nitrite reductase activity and consequently an accumulation of NO_2^- and a lowering of the protein content in S. polyrrhiza by 5 ppm DBST and ENF was observed. DBST also lowered the nitrate reductase activity and the dry weight of the fronds (BUCZEK 1984b). Sodium humate counteracts the effect of DBST in S. polyrrhiza (GUMINSKI et al. 1978).

GODZIK (1964) investigated the effect of the detergents dibutyl-naphtha-

lene sodiumsulfonate and alkylbenzene sodiumsulfonate on the O_2 uptake of L. minor. "Super Surf" which contains alkylbenzene sulfonic acid shows a lethal effect on S. polyrhiza at concentrations of 50 ppm and higher. At lower concentrations, a pronounced inhibition is coupled with a lower chlorophyll content (SAHAI et al. 1977a,b). L. gibba still grows relatively well at concentrations of 15 ppm tetrapropylene benzolsulfonate (AGAMI et al. 1976). KICKUTH and TITZLER (1974) investigated the effect of p-toluol sulfonic acid on different water and marsh plants. At concentrations of 1 mM, the substance inhibits L. minor. Compared to other water plants, L. minor shows a medium tolerance towards the detergent. The effect is believed to be caused mainly by a lowering of the pH in the solution. OERENCIK et al. (1982) observed a slightly lower chlorophyll content in L. gibba growing in waters polluted by detergents. POLAR (pers. comm. 1987) detected different tolerances to detergents between different clones of L. minor.

2.3.3.5.6.6. Further toxins and other plant active substances

All substances tested on Lemnaceae and not incorporated in one of the chapters from 2.3.3.5.3 to 2.3.3.5.6 are placed here. Sometimes it is difficult to classify a substance or group of substances correctly. Therefore, not all readers may agree with the placing of the different compounds. Sometimes, especially if the author worked with many different substances, a single substance may occur at more than one place. The effect of many toxic substances was tested on Lemnaceae: chlorate, bromate, iodate (HESSENLAND and FROMM 1932, HESSENLAND et al. 1933), methoxone (KAR 1947), substituted acryl acids (FROMM 1948, 1949, 1955), maleic hydrazide (BERTOSI 1950), sulfonamides (FROMM and O'DONNELL 1951, FROMM and PACE 1957), methylcoumarines (CIPFERRI and CIPFERRI 1947). 2,4-Dinitrophenol, sodium azide, sodium arsenite (at concentrations of 10 ppm) reduced the chlorophyll content of L. gibba only slightly but did not inhibit chlorophyll synthesis, according to OERENCIK et al. (1982). A six hour application of 1 mM sodium azide changes the phase of potassium uptake rhythm but leaves the potassium uptake itself intact (KONDO 1983a). 0.1 mM NaN_3 prohibits the CO_2 uptake in light. HILLMAN (1954) studied the effect of cyanide, azide, phenylthiourea, 2,4-dinitrophenol, and iodineacetate on the respiration of L. minor. He observed a pronounced inhibition by all substances. Flowering of L. gibba under short-

day condition is induced by 2×10^{-3} mM $K_3Fe(CN)_6$ (TANAKA and CLERLAND 1978, 1980). Cyanide induces flowering in L. aequinoctialis (TAKIMOTO 1981). It is supposed that the release of HCN from ferricyanide and ferrocyanide is responsible for the flower induction of these agents (TANAKA et al. 1983). Cyanide depolarizes the electrical membrane potential of L. gibba to the energy independent diffusion potential in the dark (ULLRICH-EBERIUS et al. 1983). A 6 hour pulse of 3 mM cyanide or 0.01 mM DCMU shifted the potassium uptake rhythm in L. gibba and considerably reduced the CO_2 uptake in the light and the CO_2 release in the dark. In addition, the potassium uptake activity was greatly inhibited (KONDO 1983a).

Nalidixic acid and other inhibitors of the protein and nucleic acid synthesis (e.g. chloramphenicol, cycloheximide, tetracycline, streptomycin, kanamycin, fluorodeoxyuridine, azaguanine, azaracil, aminotriazole) have a bleaching effect on S. polyrhiza and prevent chlorophyll synthesis (FRICK 1972, FRICK and JONES 1975). SCHER and AARONSON (1958) report the reversible bleaching effect of streptomycin on S. punctata. Streptomycin was found to be effective in mutagenesis of L. gibba tissue culture (SLOVIN and COHEN 1986). NICKELL and FINLAY (1954) studied the effect of different antibiotics (including penicillin G and oxytetracycline), they noted a stimulating effect on L. minor. HILLMAN (1961a) supposes that the stimulation is caused by a chelating effect due to the unbalanced composition of the nutrient solution. Some mycotoxins are much more toxic. Orellanine which has a similar structure as diquat and paraquat has already been mentioned (HOILAND 1983). Patulin is toxic to L. gibba at concentrations of 10^{-2} mM (SLOVIN et al. 1984). The aflatoxin flavacoumarin inhibits the chlorophyll synthesis at 10^{-2} mM (JACQUET et al. 1971). Fusicoccin increases the membrane potential of L. gibba. The uptake of glucose and glycine was significantly enhanced (BOEGER et al. 1980). The toxin hyperpolarizes the membrane of energy deficient L. gibba considerably (LUETTGE et al. 1981). Tentoxin, a toxin from Alternaria alternata, caused an inhibition of photoautotrophic growth of L. gibba and a decrease of transmembrane potential in the light (BOEGER and NOVACKY 1981). ROWE et al. (1982) report the effect of bleaching substances (phenols, catechols, guaiacols, p-cyrenes on L. aequinoctialis. The higher chlorinated a substance the more toxic it was. Leucocyanidin, procyanidin, propelargonidin, proanthocyanidin, epicate-

chin, epigallocatechin promote the growth rate of L. aequinoctialis, increase the RNA and protein content; the soluble nitrogen content is lowered and the DNA content not influenced by these substances (RAO and RAO 1984). Chlorophyll a and b content is enhanced considerably after addition of 0.1 and 0.5 ppm of the first four substances (RAO and RAO 1985a). According to LEATHER and EINHELLIG (1985) and EINHELLIG et al. (1985), the allelochemicals catechin, esculetin, chlorogenic acid, scopoletin, and vanillic acid inhibit growth of L. minor in concentrations of 1 mM, 0.5 mM, 0.25 mM, 0.5 mM, and 0.5 mM, respectively. Catechin stimulated growth of L. minor at concentrations of 0.05 and 0.1 mM. Tannic acid induces flowering of L. aequinoctialis under non-inductive photoperiods (KHURANA and MAHESHWARI 1986a). Mangiferin, a phenolic growth inhibitor (xanthone group) inhibits the growth of L. aequinoctialis at concentrations of 0.05 ppm (RAO and RAO 1985b). P-Commaric acid and P-ansaic acid are inhibiting growth of L. minor at 1 mM and 0.1 mM, resp. (EINHELLIG et al. 1985). Uptake and toxicity of coumarin and ethyl-N-phenylcarbamate in S. polyrrhiza are influenced by the pH (GREENHAM 1986). Sodiumphytate at 0.1 mM slightly inhibits the growth rate of L. minor and results in longer roots. The root elongation is much more pronounced if distilled water is used instead of Hoagland solution (BOCCHI 1981, 1982). Methionine sulfoximine reduces the photosynthesis of L. gibba (JOHANSSON and LARSSON 1985). Halogenated analogues of IAA proved to be very toxic to L. gibba (BALDI et al. 1984).

The antifreeze ethylene glycol and the dl(2-ethylhexyl) phthalate which is used in vacuum pumps are not very toxic to Lemnaceae. A lethal effect was observed at concentrations of 10 M and higher (DAVIS 1981). Polyethylene glycol reduces growth rate, and the content of protein, liquid chlorophyll and carotenoid in S. polyrrhiza (LECHEVALLIER 1977b). Croton aldehyde, a waste product of chemical factories, inhibits respiration activity of L. minor (GODZIK 1964). Caprolactam as a monomer for Nylon-6 is scarcely toxic to L. minor. It reduces the growth rate at 1% concentration and inhibits it completely at 10% (LOEWENGART 1984). STOW and ROTH (1981) investigated the influence of the following substances (in brackets the concentrations at which the growth rate was inhibited to 50% (in mM): phenol (1.8), resorcinol (1.5), dimethyl hydroquinone (0.75), dimethyl pyrocatechol (10), p-benzoquinone (0.2), hydroquinone (0.07), pyrocatechol (0.12). It was assumed that most of

these substances affect cell proteins unspecifically. CS, diazinon, malathion, dieldrin, aldrin, and sevin were tested by WORTHLEY and SCHOTT (1971) and found moderately toxic to W. brasiliensis (inhibition of growth at concentrations of 10^{-2} to 10^{-1} mM). Of three different species of Lemnaceae tested with CS, L. aequinoctialis proved to be most sensitive (more so than S. punctata and W. brasiliensis) (SCHOTT and WORTHLEY 1973). Ethanol inhibits growth rate and anthocyanin production of L. obscura at concentrations of 0.13% and higher (LEATHER and EINHELLIG 1985). 50% growth inhibition of L. aequinoctialis was achieved with the following concentrations of lower fatty acids: 0.2 mM of propionic or butyric acid, and 3 mM of acetic acid (TACHIMOTO et al. 1986). Puffs of tobacco smoke and some constituents of it (dimethylnitrosamine, methylbenzo(a)pyrene) inhibit both growth and flowering of L. gibba. Tobacco smoke condensate, polynuclear aromatic hydrocarbon fraction of tobacco smoke and benzo(a)anthracene have a marked inhibitory effect on flowering but little influence on growth rate (SABHARWAL and BHALLA 1973). The aqueous extracts of fuel oil, crude oil and especially raw coal distillate reduce the growth rate of L. gibba, L. minor and L. aequinoctialis (KING and COOLEY 1985).

SHIMOMURA et al. (1981) tested the influence of 108 crude drugs and of 10 species of medicinal plants on the growth of L. aequinoctialis and other plants. The following drugs had a stimulatory effect on L. aequinoctialis at concentrations of 2 o/oo: Achyranthis radix, Angelicae dahuricae radix, Asiaticae radix, Pruni cortex, Aucubae japonicae foliae, Casuarina (stem). The following drugs showed marked inhibiting effect at the same concentration: Chamomillae flos, Alismatis rhizoma, Astragali radix, Cinnamomi cortex, Coptidis rhizoma (was lethal), Corydalis tuber, Galla rhois (was lethal), Glycyrrhizae radix, Ipecacuanhae radix, Myrica cortex, Phellodendri cortex, Polygalae radix, Rhei rhizoma (was lethal), Senegae radix, Ginseng radix, Granati cortex, Lonicerae flos, Magnoliae cortex, Menthae herba, Pyrethri flos, Schizonepetae herba, Scutellariae radix, Sinapis semen, Araliae cordatae foliae, Chloranthi glabri foliae, Heracleum lanatum var. nipponicum (root).

Cell wall fragments of different plants, e.g. from sycamore (Acer pseudoplatanus) and soybean (Glycine max) proved to be inhibitory on flowering of L. gibba but stimulated the growth rate (MCNEIL et al. 1984, GOLDIN et al. 1984). The authors assume that plant cell wall fragments participate in the control of growth and development in plants.

2.3.4. Influence of temperature

2.3.4.1. Dependence on temperature of characteristics and metabolism

The growth rate of *Lemnaceae* shows a clear relationship to the temperature with minimum, optimum, and maximum temperature (fig. 2.11). The cardinal points of the growth rate curve are partly dependent on other factors, especially light intensity (fig. 2.13) (see also chapter 2.3.5.1.1.2). The growth rate rises up to 25°C to 31°C depending on the species (see chapter 2.3.4.3 and fig. 2.14), and falls down rather abruptly at higher temperatures. However, the dry weight per frond is much lower at optimum temperatures (fig. 2.15; ASHBY and OXLEY 1935). LANDOLT (1950) also observed a much lower dry weight per frond in *L. minor* at higher temperatures (20-32°C) than at lower ones (14°C). The area




Fig. 2.13. Growth rates of *Lemna minor* at different temperatures and different light intensities (from ASHBY and OXLEY 1935)

per dry weight in *L. minor* rises from 12.5°C to 27.5°C to the threefold value (HODGSON 1970). The area itself is greatest at 20°C (5.7 mm²) and smaller at 10°C (4.4 mm²), 26°C (4.0 mm²) and 32°C (2.6 mm²). In solutions with sugar, the maximal area is somewhat shifted to higher temperatures (LANDOLT 1957). The frond length of *W. microscopica* is greater (1.1 mm) at 22°C and short-day conditions than at 27°C and long-day conditions (0.8 mm) (RIMON and GALUN 1968b).

The growth rate of *Lemnaceae* is the result of many temperature-dependent chemical processes with intricate interactions: nutrient uptake, nutrient assimilation and transport, photosynthesis, respiration, and many other processes which are each composed of a series of enzymatic activities. Physical processes as diffusion of elements, water flow (transpiration) and permeability are also important temperature-dependent events. Photosynthesis and respiration, two dominating metabolic processes, do not have the same temperature curves. The respiration rate of




Fig. 2.14. Growth rate of different Lemnaceae species in relation to temperature (from DOCAUER 1983)

S. polyrrhiza rises to 45°C whereas the photosynthesis rate reaches its peak at 30°C (CZOPKE 1967). HODGSON (1970) noted that as a result the rate of net assimilation in L. minor only slightly rises from 12.5°C to 17.5°C and falls to 2/3 of the maximum value at 27.5°C. However, the growth rate is higher at 27.5°C than at 22.5°C. According to ASHBY and OXLEY (1935), the net assimilation rate (mg CO₂ per area) for the same species is higher at 21°C than at 18°C and 24°C (fig. 2.16). The relation between net assimilation rate and growth rate is shown in fig. 2.17. From a certain assimilation rate upwards, ranging between c. 1 and 6 mg CO₂ per dm² and hour (higher values at higher temperatures), starch is formed in the frond (ASHBY and OXLEY 1935).

WEDGE and BURRIS (1979, 1982) measured the photosynthetic O₂ evolution and the ¹⁴CO₂ fixation. The maximum O₂ evolution occurred in L. minor at




Fig. 2.15. Mean dry weight per frond of Lemna minor at different temperatures and different light intensities (from ASHBY and OXLEY 1935)

30°C and in S. punctata at 35°C. The maximum CO₂ fixation was achieved in L. minor between 20 and 30°C and in S. punctata at 30°C. FUHRER and ERISMANN (1984) observed, in L. minor cultivated in a solution with nitrate, a lower C net fixation and a lower oxidation/carboxylation ratio at 15°C compared with 25°C. If ammonium replaces nitrate, only net fixation is reduced at the lower temperature. It seems that the photosynthesis rate (and the growth rate) varies not only with light intensity but also with the method used and with the time of adaptation of the fronds to the experimental conditions.

Certain temperatures induce or keep physiological processes going. Temperature plays an important role for flower induction (see chapter 2.4.3.1.6). Low night temperatures are known to induce turion formation in S. polyrrhiza (JACOBS 1947). The relation between constant temperature and turion formation in S. polyrrhiza and L. turionifera is presented in fig. 2.18. The turion germination of the same species is also




Fig. 2.16. Mean net assimilation rates of Lemna minor in mg CO₂ dm⁻² h⁻¹ plotted against temperature (from ASHBY and OXLEY 1935)

dependent on temperature (fig. 2.19) (DOCAUER 1983). Chilling was found to be effective in breaking turion dormancy (JACOBS 1947, HENSSEN 1954, PERRY 1968, NEWTON et al. 1978) as well as seed dormancy in L. perpusilla (KANDELER 1975). Germination of seeds of L. gibba takes place faster at higher temperatures than at lower temperatures, continuously from 17°C to 33°C (REJMANKOVA 1976). The germination of turions and seeds is also dealt with in chapters 2.4.2.2. and 2.4.2.3. The roots of S. polyrrhiza are thicker and stiffer at 23°C than at 28°C (DAVIDSON and SIMON 1981a,b).




Fig. 2.17. Relation between growth rate and assimilation rate of Lemna minor at different temperatures. The arrows indicate the assimilation levels at which starch appears in the fronds (from ASHBY and OXLEY 1935)

2.3.4.2. Influence of constant and varying temperatures

For the growth rate of Lemnaceae, it is not important, whether temperatures between 15°C and 25°C at medium light intensities are kept constant or whether they vary around an average value. LANDOLT (1957) did not find very pronounced differences between growth rates of several Lemnaceae species kept at a constant temperature of 22°C or alternating at 26°C (16 hours) and 14°C (8 hours). Under the alternating conditions the growth rate was slightly higher in L. turionifera and slightly lower in L. gibba, L. minor, and L. minuscula. Also SHARITZ and LUVALL (1978)




Fig. 2.18. Turion formation of Spirodela polyrrhiza and Lemna turionifera in relation to temperature (from DOCAUER 1983)

did not find different growth rates in S. punctata at constant temperatures of 17.5°C and at regularly or acyclicly changing temperatures between 25°C and 10°C with the same average. However, it is possible that in nature the daily change of temperature has some influence if the extreme temperature values lie near the minimum or maximum temperature of the species. At low temperatures, the alternation might have a positive effect assuming that no resting stage is induced. It is also conceivable that varying temperatures might enable a better gas exchange in the culture. FERNANDEZ and MUJICA (1973) observed a greater number of stomata per number of epidermis cells and in addition bigger stomata in S. in-terminata cultivated at the same day temperatures but at cooler night temperatures than at constant temperature. It has never been studied if varying temperatures have a positive or negative effect on flowering of Lemnaceae.

Some differences in values of growth rates at the same temperatures indicated by different authors might be due to the difficulty of measuring




Fig. 2.19. Germination of turions in relation to temperature (from DOCAUER 1983)

the physiologically active temperature in Lemnaceae. The temperature in cultures with different day-lengths can only remain constant if the solution is temperature controlled. In experiments with air controlled temperatures, the temperature in the solution is warmed up during day-time by 1 to 4°C, according to the light intensity and the light source used. In addition, the temperature of the frond is again 1 to 3°C higher than the temperature of the surrounding water during irradiation. CASPERSON (1956) measured, at 5000 lux, in open cultures of S. polyrrhiza and at an air and water temperature of 18°C a 2.7°C higher temperature at the upper surface and a 1.4°C higher temperature at the lower surface of the frond. For thin or partly submerged fronds, the difference is probably negligible.

2.3.4.3. Optimum temperature

LANDOLT (1957) noted for all tested species the optimum temperature for growth rate in solutions without sugar and at light intensities between 1000 and 9000 lux within the temperature range of 20°C to 30°C. In solution with sucrose, the optimum temperature was somewhat higher. For S. polyrrhiza, JACOBS (1947) reported an optimum temperature of 25°C in solution without sucrose. ASHBY and OXLEY (1935) found an optimum temperature of 29°C for L. minor at light intensities between 800 and 16000 lux (fig. 2.13). It is possible that the lower optimum temperatures of the first authors who worked in contrast to ASHBY and OXLEY with closed and not aerated culture vessels are due to a shortage of CO₂ at higher temperatures. Differences in growth rates for different species and clones in solution with sucrose and at 2500 lux were presented by LANDOLT (1957). The growth rates were measured at 21°C, 26°C, and 30°C. Three categories could be distinguished:

- 1) Clones with a rather low optimum (growth rate best at 26°C, and higher at 21°C than at 30°C): S. punctata (1 clone), L. trisulca (2 clones), L. gibba (4 clones), L. minuscula (5 clones)
- 2) Clones with a medium optimum (growth rate best at 26°C, and higher at 30°C than at 21°C): L. minor (7 clones), L. turionifera (3 clones), L. minuscula (4 clones), W. arrhiza (1 clone), W. columbiana (1 clone).
- 3) Clones with a high optimum (growth rate best at 30°C): S. polyrrhiza (3 clones), L. aquinoctialis (3 clones), W. globosa (2 clones).

BOSS et al. (1963b) measured optimum temperatures for all 6 species tested (S. polyrrhiza, S. punctata, L. gibba, L. minor, L. aequinoctialis, L. minuscula) at about a temperature between 27°C and 29°C. The optimum temperature for L. gibba was noted by REJMANKOVA et al. (1986) with 28°C. DOCAUER (1983) gives optimum temperatures of 29°C for L. minor and W. borealis, and 31°C for S. polyrrhiza and W. columbiana (fig. 2.14).

In general, species with a wide-spread tropical distribution show a high optimum temperature, whereas species with main distribution in cooler and more oceanic areas have a lower optimum temperature.

2.3.4.4. Minimum temperature

2.3.4.4.1. Minimum temperature for survival

Even tropical species are able to tolerate temperatures of 0°C or slightly lower for short periods. GODZIEMBA-CZYŻ (1970) reported that fronds and turions of W. arrhiza can survive temperatures of -2°C for 8 to 10 days, and of 1°C for 10 to 20 days. Turions of S. polyrrhiza died if exposed to -12°C for 2 days, but remained alive at -8°C for 24 hours and at -4°C for at least 3 months (JACOBS 1947). At about +4°C they survive for at least 4 years (KRONBERGER, personal comm.). Turions of the mainly tropical species W. globosa were able to survive temperatures of 5°C in the dark for 2 months, which means that this temperature might approach the minimum temperature for turion germination but not for frond growth which is much higher (see below) (LANDOLT 1957).

To stand cold seasons, Lemnaceae species often sink into deeper layers of water where the temperatures never fall much below freezing point. Fronds of L. minor can stay alive even when enclosed in ice for a longer time (KRONBERGER, personal comm.). Some of the Lemnaceae species can survive cold winters in form of seeds (e.g. L. gibba, L. perpusilla, L. aequinoctialis), others in form of true turions (e.g. S. polyrrhiza, L. turionifera, many Wolffia species) or in form of resting fronds (e.g. L. gibba, L. perpusilla, L. aequinoctialis, W. gladiata), and still others sink as whole frond groups to deeper layers (e.g. L. minor, L. trisulca). For more details, see volume I, chapters 2.5 and 4.4.2 (LANDOLT 1986). In this way, Lemnaceae can survive in regions with winter temperatures far below the minimum temperature for growth.

2.3.4.4.2. Minimum temperature for growth

Minimum temperatures which just enable a very slow permanent growth rate were studied by LANDOLT (1957) for several species in dark cultures. 5 categories of behaviour towards low temperature could be distinguished:

- 1) Minimum temperature distinctly lower than 8°C (the growth rate at 8°C amounts to at least 10% of the maximum growth rate): L. gibba (6 clones), L. minor (3 clones), L. trisulca (3 clones).
- 2) Minimum temperature around 8°C (growth is visible at 8°C but much slower than 10% of the maximum growth rate): S. punctata (2 clones), L. minor (5 clones), L. turionifera (5 clones), L. trisulca (1 clone), L. minuscula (8 clones).
- 3) Minimum temperature between 8°C and 13.5°C (not growing at 8°C; growth rate at 13.5°C at least 10% of the maximum growth rate): S. polyrrhiza (1 clone), L. turionifera (1 clone), L. aequinoctialis (1 clone), L. valdiviana (1 clone), L. minuscula (2 clones), W. gladiata (1 clone).
- 4) Minimum temperature between 13.5°C and 16.5°C (not growing at 13.5°C; growth rate at 16.5°C at least 10% of the maximum growth rate): S. polyrrhiza (2 clones), L. aequinoctialis (4 clones), W. arrhiza (1 clone), W. columbiana (1 clone), W. globosa (1 clone).
- 5) Minimum temperature between 16.5°C and 20°C (not growing at 16.5°C; growth rate at 20°C at least 10% of the maximum growth rate): S. polyrrhiza (1 clone), L. aequinoctialis (1 clone).

DOCAUER (1983) estimated the minimum temperatures for a clone of: L. minor at 5°C, L. turionifera at 9°C, S. polyrrhiza, W. borealis, and W. columbiana at 12.5-13.5°C. This is more or less in accordance with the results of LANDOLT (1957).

The results explain why the species with high minimum temperature are found in nature only in regions with very warm growth periods whereas species with lower minimum temperatures are also occurring in regions with cool summers.

2.3.4.5. Maximum temperature

2.3.4.5.1. Short-term maximum temperature

The turions of *S. polyrrhiza* are able to stand 24 hours at 50°C and one week at 45°C (JACOBS 1947). STANLEY and HEADWELL (1976b) studied the short-term tolerances of *L. obscura* (named as *L. minor*) at temperatures between 40°C and 60°C. The temperature of 50% inhibition of growth rate and the temperature of 50% lethality are very near together. Temperatures above 60°C are not tolerated at all, even not for very short periods. The 50% lethality is achieved at 55°C after 30 seconds, at 50°C after 5 minutes and at 42°C after 2 hours. The heat tolerances are somewhat enhanced by the addition of calcium (up to at least 3 mM). Also the nitrogen content has an influence on the temperature sensitivity. The optimal nitrogen concentration for growth at 0.7 mM enabled the highest tolerance. In the light, the fronds survived for a longer time at 45°C than in darkness, but at 55°C, there was scarcely a difference.

2.3.4.5.2. Long-term maximum temperature

Long-term temperature limits are not the consequence of plasma damages by temperature but probably due to an unbalance of metabolic processes at this temperature. Thus, the photosynthesis rate is lowered in *S. polyrrhiza* at temperatures above 30°C whereas the respiration rate is at this temperature still rising (CZOPEK 1967) resulting in a negative net assimilation rate and finally in starvation. Also other processes must be involved in the long-term effect at high temperatures because the fronds die in solution with sucrose nearly at the same temperature as without sucrose irrespective of cultivation in the light or in the dark. For *S. punctata* it is known that it loses the ability of synthesizing chlorophyll and evolving oxygen at temperatures around 37°C. This effect was only observed in the light but not in the dark (PORATH and BEN-SHAUL 1971). SCHER and AARONSON (1958) noted a reversible bleaching in *S. punctata* already at 34°C. STEPHENSON et al. (1980) also report of a growth stop of *S. punctata* at 35°C.

Long-term maximum temperatures were established by LANDOLT (1957). 5 categories can be distinguished:

- 1) Maximum temperature below 30°C: *L. minor* (1 clone), *L. trisulca* (3 clones), *L. minuscula* (1 clone).
- 2) Maximum temperature between 30°C and 32°C: *L. gibba* (2 clones), *L. minor* (5 clones), *L. turionifera* (2 clones), *L. trisulca* (1 clone), *L. minuscula* (8 clones), *W. arrhiza* (1 clone), *W. gladiata* (1 clone).
- 3) Maximum temperature between 32°C and 33°C: *L. gibba* (2 clones), *L. turionifera* (2 clones), *L. minuscula* (1 clone), *L. valdiviana* (1 clone), *W. columbiana* (1 clone).
- 4) Maximum temperature between 33°C and 34°C: *S. punctata* (1 clone), *L. gibba* (1 clone), *L. minor* (1 clone), *L. aequinoctialis* (3 clones), *W. globosa* (1 clone).
- 5) Maximum temperature higher than 34°C: *S. polyrrhiza* (3 clones).

DOCAUER (1983) obtained similar results for the upper temperature limit of the following species (only one clone measured): 32.5°C for *L. minor*, 34°C for *L. turionifera*, 35°C for *W. borealis*, 36°C for *W. columbiana*, and 38°C for *S. polyrrhiza*. DOCAUER measured the temperature at which the species cease growth, LANDOLT the temperature at which the species still grows slowly. This might explain the higher temperatures of DOCAUER. Though different clones of the same species do not behave exactly in the same way, some species tolerate higher temperatures on the average (e.g. *S. polyrrhiza*) and others only lower temperatures (e.g. *L. trisulca*). In general, species with higher tolerance are distributed in warmer or more continental areas whereas species with a low tolerance are to be found in rather cool areas. However the differences are not so pronounced as one would suppose from the distribution of the species in different climate zones.

2.3.5. Influence of light and other radiations

2.3.5.1. Light (including UV radiation)

2.3.5.1.1. General remarks

The solar radiation within the visible range is absorbed by the Lemna-case through different pigment systems. It is used for energy supply as well as for regulation of different life processes.

The measurements of light energy are not always comparable. In the older literature, the light intensity was measured by photo cells in lux or foot-candles (English speaking countries). In the present work the foot-candle was transferred to lux approximately by the factor 10. In more recent publications, the intensity is either described in energy units (Joule or erg) (mostly restricted to the visible range of the radiation spectrum), or in units of the photon flux density which is expressed in Einstein, or mmol per squaremeter and second. 1 erg corresponds to 10^{-7} Joule (J). 1 Einstein is identical with $1 \text{ mmol m}^{-2} \text{ s}^{-1}$. The energy content of a unit of photon flux density ($1 \text{ mol m}^{-2} \text{ s}^{-1}$) is different at different wavelengths. It amounts to $800 \text{ kJ m}^{-2} \text{ s}^{-1}$ at 150 nm and $170 \text{ kJ m}^{-2} \text{ s}^{-1}$ at 700 nm. Just to give an approximate idea of the correspondence of the different measurement systems, the following example may be given. A light intensity of 20000 lux might correspond to a total energy content of $100 \text{ J m}^{-2} \text{ s}^{-1}$ (white fluorescent light) and roughly to about $1 \text{ mmol m}^{-2} \text{ s}^{-1}$ at a wavelength of c. 500 nm. However, this relation is dependent on the type of light source and may vary greatly.

2.3.5.1.2. Light as source of energy in photosynthesis

Relatively high light intensities are necessary for photosynthesis, in contrast to other light-dependent metabolic processes. The chlorophyll system is responsible for the energy transfer. The maximum solar energy conversion efficiency in S. punctata was measured with 4.9% of the photosynthetically active radiation (PAR) (including UV light) (MESTAYER et al. 1984). The response of Lemna to light intensity is dependent on species, clone, temperature, as well as on nutrient and CO_2 supply. Also, the light quality is important (EICHORN 1983; fig. 2.20).

2.3.5.1.2.1. Light saturation, minimal and maximal light intensity

At low temperatures, the light saturation is lower than at high temperatures (see chapter 2.3.5.1.2.2). Light saturation is generally lower for growth rate than for photosynthesis rate. Therefore, the dry weight still rises at intensities higher than light saturation for growth rate. In our own experiments (LANDOLT 1957 and unpubl.) the saturation for growth rate was reached at intensities between 5000 and 15000 lux depending on species and clones (see chapter 2.3.5.1.2.4) (continuous light). BOSS et al. (1963b) observed light saturation for growth rate (but only 12 hours day-length) in L. aequinoctialis (2 clones), S. polyrrhiza (1 clone), S. punctata (1 clone), L. obscura (named as L. minor, 1 clone) and L. minuscula (1 clone) at c. 17000 lux (27-29°C). L. gibba reached saturation at still higher light intensities. MESTAYER et al. (1984) noted c. $50 \text{ J m}^{-2} \text{ s}^{-1}$ at light saturation for growth rate of S. punctata. DOCAUER (1983) measured the half saturation constants K_L




Fig. 2.20. Relation between light intensity and growth rate in Lemna arrhiza chemostat cultures (from EICHORN 1983)

for growth rate at 28°C of one clone each of 5 species (see chapter 2.3.5.1.2.4) between 0.024 mmol m⁻² s⁻¹ (*S. polyrrhiza*) and 0.086 mmol m⁻² s⁻¹ (*W. borealis*). The light saturation is reached at roughly four times higher light intensities than K_L . In the following references, the light saturation for photosynthesis was determined: CZOPEK (1967) 91 J m⁻² s⁻¹ for normal fronds and 32 J m⁻² s⁻¹ for turions of *S. polyrrhiza*; GAPONENKO and STAZHETSKYI (1969) 20 J m⁻² s⁻¹ or 5000 lux for *S. polyrrhiza*; LINDEMAN (1951, 1972) 100 J m⁻² s⁻¹ for *L. minor* (enrichment of the CO₂ content of the air); INGEMARSSON et al. (1984) 1.1 mmol m⁻² s⁻¹ for *L. gibba*; ANDERSEN (1983) 0.5-0.7 mmol m⁻² s⁻¹ for *L. gibba*; WEDGE and BURRIS (1982) 0.3-0.6 mmol m⁻² s⁻¹ for *L. minor* and 0.6-1.2 mmol m⁻² s⁻¹ for *S. punctata*; FILBIN and HOUGH (1985) c. 0.35 mmol m⁻² s⁻¹ for *L. minor*.

Long-term compensation point for *L. minor* is near 2-3% of full sunlight (WILKINSON 1963). The compensation point of normal fronds of *S. poly-*




Fig. 2.21. Growth rates of *Lemna minor* at different light intensities and different temperatures (from ASHBY and OXLEY 1935)

rrhiza was noted by CZOPEK (1967) at 15 J m⁻² s⁻¹ and for turions at 11-13 J m⁻² s⁻¹. ASHBY and OXLEY (1935) measured a distinct photosynthesis rate for *L. minor* already at 500 lux (lower intensities not investigated). ULLRICH-EBERIUS et al. (1978) state that photosynthetic O₂ evolution in *L. gibba* commences at 100 lux. An occurrence of *L. trisulca* (together with other water plants) was found by PIP and SIMMONS (1986) in Shoal Lake (Canada) at depths of 12-14 m. At these depths plants received an estimated 0.5-1% of surface light. According to INGEMARSSON et al. (1984a), the light compensation point measured as minimal needed photon flux density lies for *L. gibba* at c. 0.05 mmol m⁻² s⁻¹. It is not affected by nitrogen limitation.

The relation between light intensity and growth rate in *Wolffia arrhiza* chemostat cultures is shown in fig. 2.20. At low light intensities (see also figs. 2.22 and 2.23) the rates of growth and photosynthesis rise proportionally to the light intensity. At higher intensities, they




Fig. 2.22. Growth rate of different species in relation to light intensity (from DOCAUER 1983)

1939, BORNKAMM 1970b, ROMBACH 1976). L. gibba also has longer roots at higher light intensities (ANDERSEN 1983). The nitrate uptake of W. arrhiza rises with increasing light intensities up to $87 \text{ J m}^{-2} \text{ s}^{-1}$ (SWADER et al. 1975). It decreases again above $174 \text{ J m}^{-2} \text{ s}^{-1}$. At low light intensities, L. trisulca forms only one daughter frond on one side, consequently spiral colonies develop (GOEBEL 1921). HILLMAN (1961a) supposes that the low sugar content in the fronds regulates the limited production of daughter fronds. In nutrient solutions with sucrose, L. trisulca is able to form daughter fronds on both sides even in darkness.

It is not quite clear if there is a maximum light intensity above which Lemnaeae fronds are damaged. Data in the literature are controversial. WHITE (1936c) reports that L. minor shows injuries at light intensities above 15000 lux . Fig. 2.20 demonstrates that W. arrhiza does not grow at light intensities above $6 \text{ to } 9 \text{ J m}^{-2} \text{ s}^{-1}$. This value is extremely low (at least 10 times lower than in other publications) and is probably caused




Fig. 2.24. Dry weight per frond in relation to light intensity (from DOCAUER 1983)

approach an optimum asymptotically (ASHBY and OXLEY 1935, WHITE 1937a, MIYATA 1971b, DOCAUER 1983) (figs. 2.21, 2.22). The dry weight increases linearly with the light intensity up to $5000 \text{ to } 9000 \text{ lux}$ depending on temperature (ASHBY and OXLEY 1935, DOCAUER 1983; figs. 2.23 and 2.24). The size of the frond is also mostly higher at high light intensities than at lower ones (KRZECZOWSKA et al. 1975, ROMBACH 1976 for L. minor, ANDERSEN 1983 for L. gibba). ASHBY (1929a), HICKS (1934) and WHITE (1936c) did not find a clear relation between light intensity and area of the frond. The following characteristics of L. minor become greater with increasing light intensity: root length, respiration rate per area (contrary to the respiration rate per dry weight), contents of protein, chlorophyll, and carotenoid (WHITE and TEMPLEMAN 1937, WHITE 1937b,




Fig. 2.23. Mean dry weight per frond of Lemna minor at different light intensities and different temperatures (from ASHBY and OXLEY 1935)

by the submerged state of plants in chemostat culture. MESTAYER et al. (1984) measured lower growth rates at levels of PAR above $80 \text{ J m}^{-2} \text{ s}^{-1}$ than between 50 and 80. According to ZURZYCKI (1957a) photosynthesis of *L. trisulca* does not function any longer at intensities above $500 \text{ J m}^{-2} \text{ s}^{-1}$. Photosynthetic O_2 evolution of *S. punctata* and *L. minor* was photoinhibited above a photon flux density of $1.2 \text{ mmol m}^{-2} \text{ s}^{-1}$ (WEDGE and BURRIS 1982). OEGREN et al. (1984) and OEGREN and OEQUIST (1984a,b) observed photoinhibition of photosynthesis in *L. gibba* at 0.65 and $1.75 \text{ mmol m}^{-2} \text{ s}^{-1}$, subsequently isolated chloroplasts showed pronounced reduction in the capacity of whole chain electron transport. LANDOLT (1957 and unpubl.) did not see any damage in fronds of all investigated species cultivated at light intensities up to 33000 lux (higher intensities could not be achieved). In nature, much higher light intensities are measured on sunny days (up to 120000 lux). Possibly the relatively low upper limits of light intensities recorded are caused by the experimental conditions. As in other phanerogams, Lemnaceae form different fronds under shady and sunny conditions (LINDEMAN 1973, 1979). Therefore any measurements at high light intensities to determine a long lasting photoinhibition can only be made after having exposed the cultures to the new conditions for at least five weeks.

2.3.5.1.2.2. Light effect in relation to temperature

Figs. 2.25 and 2.21 show the relationship between growth rate and light intensity at different temperatures. According to LANDOLT (1957), light saturation was not reached in the two clones of *L. gibba* at 9000 lux if temperatures surpassed 20°C . At 18°C the maximum growth rate was achieved at 9000 lux ; at 10°C 2500 lux and at 5°C even 1000 lux were sufficient for maximum growth rate. Further studies revealed that all investigated species and clones reached their maximum growth rate at lower temperatures at a much lower light intensity.

OEGREN et al. (1984) and OEGREN and OEQUIST (1984a,b) demonstrated the dependence on temperature of photoinhibition. *L. gibba* was exposed for two hours to various combinations of photon flux densities and temperatures. The extent of photoinhibition was assayed by measuring the net CO_2 uptake before and after photoinhibitory treatment. At a constant photon flux density of $0.65 \text{ mmol m}^{-2} \text{ s}^{-1}$ the extent of photoinhibition increased with decreasing temperature. About 60% photoinhibition of the




Fig. 2.25. Growth rate of *Lemna gibba* (no. 6566 and 6745) at three light intensities in solutions with and without sucrose in relation to temperature (from LANDOLT 1957, completed)

quantum yield for CO_2 uptake could be obtained either by a high photon flux density of $1.75 \text{ mmol m}^{-2} \text{ s}^{-1}$ at 25°C or by a moderate photon flux density of $0.65 \text{ mmol m}^{-2} \text{ s}^{-1}$ at 3°C .

It is supposed that the light compensation point is similarly dependent on the temperature. This means that the lower the temperature, the lower is the compensation point.

2.3.5.1.2.3. Light effect in relation to nutrient supply

ASHBY (1929a) noted light saturation of a clone of *L. minor* already at

1500 lux (at 25°C). This must be explained by the unfavourable nutrient solution. According to WHITE (1937a,b, 1938, 1939) and WHITE and TEMPLEMAN (1937), the optimal nitrogen and potassium content for L. minor is higher at higher light intensities, e.g. optimal K content at 600 lux is 0.05 mM, and at 3000 lux 5 mM. This is in accordance with the results of SWADER et al. (1975) who showed that the nitrate uptake of W. arhriza rises with higher light intensities. LINDEMAN (1951, 1952) observed that lack of P inhibits the photosynthesis near light saturation. The inhibition was stopped within a few hours after adding PO_4^{3-} to the solution.

2.3.5.1.2.4. Differences in the light effect between different species and clones

HICKS (1934) observed different light saturations (9000 lux and 15000 lux) for two clones of L. minor. WHITE (1936c) also found different responses of different clones of L. minor to light intensities. L. trisulca is able to grow at lower light intensities than L. gibba and L. minor (AMBROSE 1978). DOCAUER (1983) measured the following light saturations for 5 different species in $mmol\ m^{-2}\ s^{-1}$: S. polyrrhiza (0.024), L. minor

Table 2.21. Growth rate of different species of Lemnaceae at two light intensities (2000 lux and 9000 lux continuous light) in percent of the maximum growth rate at 24°C (after LANDOLT 1957)

Species (number of clones)	light intensity in lux	
	2000	9000
<u>S. polyrrhiza</u> (5)	50-55	91-96
<u>S. punctata</u> (1)	58	95
<u>L. gibba</u> (6)	55-63	83-96
<u>L. turionifera</u> (5)	48-63	91-99
<u>L. minor</u> (7)	57-63	88-98
<u>L. aequinoctialis</u> (4)	59-75	88-100
<u>L. valdiviana</u> (1)	70	101
<u>L. minusculea</u> (9)	66-86	96-105
<u>W. lingulata</u> (1)	32	53
<u>W. gladiata</u> (1)	43	68
<u>W. arhriza</u> (1)	56	99
<u>W. columbiana</u> (1)	48	60
<u>W. globosa</u> (1)	64	95

(0.035), L. turionifera (0.047), W. columbiana (0.047), and W. borealis (0.086). Table 2.21 lists differences in the behaviour of different species and clones at two light intensities (from LANDOLT 1957). In general, the differences within a species are nearly as great as between the species. The relatively low percentage of maximum growth rates of W. lingulata, W. gladiata, and W. columbiana is remarkable. According to the results of DOCAUER (1983), W. borealis has a still lower percentage. Clones (and species) with a low percentage of the maximum growth rate at 9000 lux have a high light intensity saturation and therefore grow in open not shaded places or they grow mixotrophically. On the other hand L. minusculea or clones of L. aequinoctialis with a high percentage of the maximum growth rate at 2000 lux are able to grow in shady areas. From preliminary indications, L. trisulca also belongs to this group.

2.3.5.1.2.5. Relation between light effect and the age of the fronds

When older, fronds of S. polyrrhiza grow more slowly (at $14\ J\ m^{-2}\ s^{-1}$). A short illumination of 100, 250, or $500\ J\ m^{-2}\ s^{-1}$ enhances the photosynthetic rate in parallel to the light intensity and the age of the fronds (GAPONENKO and STAZHETSKII 1969). The mechanism of this effect is not sufficiently known.

2.3.5.1.2.6. Indirect effects of light intensity

YOSHIMURA (1952) detected a higher NH_4^+ uptake in the light than in the dark for S. polyrrhiza and L. valdiviana whereas the NO_3^- uptake was slightly inhibited in the light. HILLMAN (1961a) supposes that this effect is due to the higher partial O_2 pressure during photosynthesis. BLACKWOOD and LEAVER (1977) showed that in L. minor light stimulates amino acid incorporation specifically into three proteins. LIEBERT (1986a) reports a higher content of Ca, Mg, and K in L. gibba if cultivated in the light compared with in the dark.

The plasma adhesion to the cell wall of parenchyma cells containing chlorophyll of L. minor roots is lower in darkness than in the light. It is assumed that this is the result of the higher CO_2 level in darkness and the subsequently lowered pH (SCHAEFER 1956).

2.3.5.1.2.7. Duration of light

The growth rates of Lemnaceae rise proportionally to the length of day at least at suboptimal light intensities (figs. 2.26 and 2.27). They are highest under continuous light (ASHBY 1929a, LANDOLT 1957). Near light saturation, the increase is no longer linear. EYSTER (1966) did not observe a distinct rise of growth rate in *S. polyrrhiza* above 16 hours light duration at 25°C and 6000 lux. Growth rate under alternating periods of 2 hours does not differ significantly from the growth rate under a 12 hour period (ASHBY 1929a).

Some authors report damaged fronds under continuous light conditions, e.g. CLARK (1925) in *S. polyrrhiza*, at light intensities of 4000 and 9000 lux, and WHITE (1936c) in *L. minor*, at light intensities of 15000 lux. Other authors did not note any injuries to fronds under continuous light and at different intensities (e.g. LANDOLT 1957, EYSTER 1966).




Fig. 2.26. Growth rate of *Lemna gibba* (no. 6566) and in relation to the length of day in solution with sucrose (S) and without sucrose (O) (from LANDOLT 1957)

Therefore, it is possible that the damaging effect must be attributed to other factors than lack of dark periods, e.g. too high temperatures, unfavourable nutrient conditions, too sudden change from conditions of low light intensity to high intensity.

Compared with continuous light, a day of 12 hours results in smaller fronds, less stomata per frond area, and smaller stomata in *S. intermedia* (FERNANDEZ and MUJICA 1973). The life-span of *L. minor* is not influenced by the length of day (WANGERMANN 1952). Differently, OSTROW and DIJKMAN (1969) observed a pronounced reduction of the life-span of *S. punctata* fronds with prolongation of the day.

HUBALD and AUGSTEN (1979) showed that alternating light and dark conditions enhance the activity of nitrate reductase and the content of soluble protein in *W. arrhiza* compared with continuous light conditions.

For other effects of length of day (especially on flowering), see chapters 2.3.5.1.2 and 2.4.3.1.5.




Fig. 2.27. Growth rate of *Wolffia arrhiza* (no. 6862) in relation to the length of day in solution with sucrose (S) and without sucrose (O) (from LANDOLT 1957)

2.3.5.1.3. Light effects via the phytochrome system

There is extensive literature on the effect of red and far-red light of low intensity in Lemnaceae. KANDELER (1955, 1956) revealed that flowering of the long-day plant L. gibba is stimulated by far-red illumination (720-760 nm) and inhibited by red light (600-700 nm). Inversely, far-red light prevented flowering of the short-day plant L. aequinoctialis if applied at the end of the daily short-day (SCHUSTER 1968) (see also chapter 2.4.3.1.5.). Far-red light can be replaced by blue light, red light by UV irradiation (380 nm) (OHTANI and KUGAWA 1981) to reach appropriate effects on flowering of L. aequinoctialis.

Apart from the flowering effect, the phytochrome system regulates a series of further metabolic processes and the formation of different characteristics of Lemnaceae. ROMBACH (1976) lists the following characteristics of heterotrophically grown L. minor which are stimulated by light intensities far below the level needed for photosynthesis (light effect reversible by exposure to far-red irradiation): rate of cell division, elongation of fronds, and their multiplication. Far-red illumination given at the end of the daily main-light period on the other hand results in slower growth of roots, higher growth rate of fronds, and bigger fronds in the short-day plant L. aequinoctialis (SCHUSTER 1968). The production of resting fronds becomes inhibited by end of day far-red in L. perpusilla, another short-day plant (HUEGEL et al. 1979). The same treatment leads to a reduction of the growth rate, size of fronds and length of roots in the long-day plant L. gibba. A period of 5 minutes red light is able to neutralize the effect of far-red light (KANDELER 1963). The phytochrome control of root growth in L. gibba and L. minor is also described by CHAMURIS and NIELSEN (1983).

Furthermore, the phytochrome system regulates the nitrate reductase activity. According to LIEBERT and AUGSTEN (1986), there are low irradiance responses involved in the regulation of the enzyme in L. gibba and L. minor.

Exposure to far-red light (4-5 minutes applied about one hour after normal daylight) causes a hyperpolarization of the membrane potential of L. aequinoctialis as well as of L. gibba (LOEPPERT et al. 1978, KANDELER et al. 1980). The decomposition of starch is retarded by preceding application of far-red light (KANDELER et al. 1980). The vitamin B₁ synthesis in L. minor is stimulated by phytochrome (P_{fr}) (ROMBACH 1974a,b, 1976).

Phytochrome is involved in the photoregulation of anthocyanin production in S. polyrrhiza (MANCINELLI 1984, MANCINELLI and RABINO 1984). However, the authors suggest the involvement of a second photoreceptor (called cryptochrome) specific to light of the blue and UV region. Another phytochrome mediated regulation was detected by TOBIN (1981a,b). The phytochrome action rapidly affects the expression of the genes for the small subunit of RuDP carboxylase and chlorophyll a/b-protein in L. gibba. Red light induces rapid and specific increases in the amounts of RNA encoding these polypeptides. Therefore, phytochrome action is said to change either the transcription rate or the rate of degradation of these mRNAs (STIEKEMA et al. 1983a).

The phytochrome content (P_{fr}) diminishes in darkness after exposure to red light in L. gibba, L. minor, and L. aequinoctialis. However, after a longer dark period, it rises again (ROMBACH and SPRUIT 1968, ROMBACH 1978). PORATH and BEN-SHAUL (1973) obtained similar results with S. punctata.

2.3.5.1.4. Light effects not completely explainable by chlorophyll or phytochrome action

In this chapter, all light effects will be dealt with which are not understandable by action of the chlorophyll or phytochrome systems or which have not been investigated in this respect.

ZURZYCKA and ZURZYCKI (1950) and ZURZYCKI (1957a,b, 1962) studied the influence of light on the position of chloroplasts in L. trisulca. In darkness, the chloroplasts are distributed along the cell walls. At low light intensities (10-150 lux), the plastids are along the cell walls which are perpendicular to the light incidence. At high light, intensities (10000 lux and higher), they remain along the walls which are oriented parallel to the light incidence. The effect of low as well as high light intensities is caused mainly by blue light. Riboflavin is supposed to be responsible for the absorption of blue light. GABRYS et al. (1981) investigated chloroplast translocations in L. trisulca induced by a single strong pulse of blue light. With short pulses transient rearrangements of chloroplasts to a weak-light position occurred. With longer pulse duration, the initial movement to a partial strong-light position was followed by a wave of translocations to a weak-light arrangement. KAJFOSZ et al. (1983) report on chloroplast translocations in L. trisulca.

3.6. UTILIZATION AS TEST AND INDICATOR PLANT

The widespread use of Lemnaceae as test plants is due to the simple cultivation techniques, the need of little space, the fast growth rate and the genetical uniformity of its cultures which correspond to clones. Correspondingly, the literature is voluminous. The usefulness of Lemnaceae for investigation of physiological processes is dealt with in chapter 2. Lemnaceae are however not very suitable to show specific reactions towards different substances since only few distinguishing characteristics are present. In most studies, changes on growth rates are recorded. Also dry weight, size of frond, length of root, growth of root (LIEBERT 1986a), anthocyanin formation (reported as more sensitive than growth rate: EINHELLIG et al. 1985, LEATHER and EINHELLIG 1985, JUNG-NICKEL 1978, JUNG-NICKEL and AUGSTEN 1986), turion production and germination (JUNG-NICKEL 1978), gas exchange (MARTI et al. 1986) and CO₂ production (WUESTLING and BOEHM 1979) have been tested.

Tests have been developed and used for the following substances:

- a) heavy metals: AUGSTEN (1983), FISCHER (1981), LIEBERT (1986), NASU and KUGIMOTO (1981), SZABADOS et al. (1983), WANG (1986a);
- b) growth factors:
ABA: ENUKWESI and DUMBROFF (1978), GOLDBACH and MICHAEL (1976), JOHANSSON et al. (1982), LIEBERT (1980a), LIN and MATHES (1973), TAYLOR and DUMBROFF (1975), TILLBERG (1975), VAN STADEN and BORNWANN (1970a);
kinetin: LETHAM (1967);
different growth factors: JUNG-NICKEL and AUGSTEN (1986), RANE and TUCKEY (1972), WUESTLING and BOEHM (1979);
- c) herbicides: AUGSTEN (1983), BAHADIR and PFISTER (1985), BIRMINGHAM and COLMAN (1983), DAMANAKIS (1970, 1972), DECLEIRE and DE CAT (1977), FISCHER (1981), LIU and CEDENO-MALDONADO (1979), O'BRIEN and PRENDEVILLE (1978), PESTEMER (1979), RICHARDSON (1985), ZAWADZKI (1975);
- d) allelopathic substances: EINHELLIG et al. (1985), LEATHER and EINHELLIG (1985), MARTI et al. (1986), SAGGESE et al. (1985);
- e) drug substances: SHIMOMURA et al. (1981, 1982);
- f) further toxicants: DAVIS (1981), KING and COLEY (1985), SLOOF and CANTON (1983), WALBRIDGE (1977), WARD et al. (1981);

- g) phosphate (and nitrate) pollution: AEBLI (1986), FEKETE et al. (1976), JUNG-NICKEL (1978), POTT (1981)

The following Lemnaceae species have been used as test organisms:

- S. polyrrhiza: DAMANAKIS (1970, 1972), JUNG-NICKEL (1978), JUNG-NICKEL and AUGSTEN (1986), RICHARDSON (1985), WARD et al. (1981);
- S. punctata: KLAINE (1985);
- L. gibba: DAVIS (1981), KING and COLEY (1985), LIEBERT (1986a), TILLBERG (1975);
- L. minor: AEBLI (1986), BAHADIR and PFISTER (1985), BLACKMAN (1952), BISHOP and PERRY (1981), EINHELLIG et al. (1985), FEKETE et al. (1976), FISCHER (1981), FROMM (1946, 1951), FUNDERBURN and LAWRENCE (1963), KING and COLEY (1985), LEATHER and EINHELLIG (1985), LOCKHART and BLOUW (1979), MARTI et al. (1986), O'BRIEN and PRENDEVILLE (1978), OFFORD (1946), RANE and TUCKEY (1972), ROULET (1975), SAMPFORD (1952), SIMON and BLACKMAN (1953), SLOOFF and CANTON (1983), SZABADOS et al. (1983), WALBRIDGE (1977), WANG (1986a), WARD et al. (1981);
- L. obscura: EINHELLIG et al. (1985); LEATHER and EINHELLIG (1985);
- L. aquinoctialis: KING and COLEY (1985), LIU and CEDENO-MALDONADO (1979), NASU and KUGIMOTO (1981), ROWE et al. (1982), SHIMOMURA et al. (1981, 1982);
- W. arrhiza: AUGSTEN (1983), WUESTLING and BOEHM (1979).

Lemnaceae are instructive laboratory and experimental plants for school purposes (see RHODES 1968).

3.7. UTILIZATION FOR PRODUCTION OF ENERGY AND CHEMICAL COMPOUNDS

3.7.1. Energy production

WOLVERTON and McDONALD (1981) produced 0.14-0.22 m³ methane (CH₄) per kg dry weight of a mixture of water plants including Lemnaceae. SEKINE (1979) worked out a patent to obtain methane or ethanol from W. globosa (named as W. arrhiza) by fermentation with yeast. He got either 285 g ethanol per kg dry weight or 0.42 m³ methane per kg dry weight. He was able to cultivate the Wolffia on organic waste water treated first with photosynthetic bacteria, micro-algae and zooplankton before being inoculated with Wolffia. The author calculates that normal production of 1 liter ethanol in Japan amounts at present to 186 Yen. Production of ethanol via Wolffia is much cheaper (100 Yen per 1 liter). If all organic waste waters in Japan were used for Wolffia cultivation, a total production of 10⁶ t ethanol per year would be possible.

EL-HINNAWI (1983) proposes energy farms for fuel production in regions where duckweeds and water hyacinths are harvested. Also RAO (1984) suggests an aerobic digestion of the same water weeds for energy production. BAI (1985) investigated the biodegradability and methane production of cow-dung mixed with Lemna.

3.7.2. Production of pharmaceutical compounds

Lemnaceae have been known as medicinal plants for many centuries. DIOSCORIDES (first century) and GALEN (second century) used the name "phakos ho epi ton telmaton" or in latin "viparia" for L. minor. DIOSCORIDES and later M. ADANSON (18th century) and A. MORI (19th century) attributed the following pharmaceutical effects to Lemnaceae: soothing of freezing injuries, relief of aches of podagra and burns, healing of fractures (cited from BEAUVOIS 1816). Lemnaceae have also been used to cure hepatitis. Blood clots could be dispersed by applying L. trisulca dissolved in white wine. McCANN (1942) mentions that Lemnaceae are used in India to prevent blood loss during menstruation. A mixture of Lemnaceae with pepper is put on the eyes of unconscious typhus patients.

No positive effect of Lemnaceae against cough were observed by ANDRONOVA

(1972). In Canada, L. trisulca together with Stellaria media is applied by Iroquois Indians against swellings (ROUSSEAU 1945, ARNASON et al. 1981). According to WATT and BREYER-BRANDWIJK (1962), L. minor has been used in Africa as a remedy for dropsy and rheumatism. In China, it is used internally as a diuretic, antiscorbutic and antisyphilitic, and externally for eye diseases and carbuncles. The same authors report that Lemnaceae give negative antibacterial and antimalarial tests. On the other hand, STANGENBERG (1967) recorded a bacteriostatic effect of L. minor extracts against the gram-negative bacterium Sphaerotilus natans. No efficacy was observed against other bacteria, and not all clones of L. minor showed the same effect. Extracts of L. minor did not reveal any evident activity against Staphylococcus aureus, Mycobacterium smegmatis, Escherichia coli, Candida albicans and Fusarium roseum. However, Alter-naria sp. was inhibited moderately by methanol extracts of L. minor (SU et al. 1973b). HILLMAN (in lit. 1979) noted an algicidal effect (against a unicellular chlorophyte) of W. globosa (originating from Thailand) in Petri dishes on nutrient agar.

YONG and THO (1976) measured 0.095 mg per g dry weight of a cardiac glycoside-like substance present in the aqueous extract of the fronds of L. aequinoctialis (named as L. minor). The substance was able to produce the same effect on the chicken heart as standard digoxin. The authors isolated the reducing sugar digitoxose which is typical of cardiac glycosides.

The toxicity of L. minor for white mice was investigated by ANDRONOVA (1972). The smallest toxic level was 577 mg per kg live weight and the smallest lethal dosis 735 mg per kg. In conclusion, there is not much known on pharmaceutically important substances produced by Lemnaceae.

3.7.3. Enzyme production

AUGSTEN (1984a) gives a survey of the possibilities of using Lemnaceae in biotechnology. Lemnaceae can be successfully applied in enzyme reactors for continuous catalysis of metabolic processes. The enzyme is stored immobile within the plant (especially at the surface). The high activity of the enzymes is preserved for a long time. The harvested plants are stabilized and conserved by lyophilization, by drying with acetone or by other methods. Fig. 3.1 shows the activity of some of the




Fig. 3.1. Activity of some enzymes of *Wolffia arrhiza* in a reactor after having cultured the plants in a normal medium (white bars) or with limited phosphorus (black bars) (after AUGSTEN 1984a).

enzymes of *W. arrhiza* in a reactor used in Jena, GDR. Also turions of *S. polyrrhiza* have been investigated. TROMAC et al. (1984) cultivated *Lemnaea* to transform testosterone to androstenedione and 5α-androstane-3, 17-dione.

3.8. SPECIAL UTILIZATIONS

3.8.1. Cosmonautic plant

Since Lemnaceae are effective oxygen producers and CO₂ removers, some investigations have been performed to check the suitability as cosmonautic plants. EYSTER (1966) studied S. polyrrhiza in respect to a possible use in space flight. S. polyrrhiza was also investigated under simulated space-laboratory conditions. Early turion development was negatively affected by these conditions (KUTLJACHNEV et al. 1978). MILLER (1978) investigated the effect of vibrations (100 Gc, 0.6 mm) and ionizing irradiation (X-rays, 1-3 grad) on S. polyrrhiza. Vibrations affect the growth rate negatively. If applied together with irradiation, they increase the radiosensitivity of the plants. MEY (1960) and WILKS (1962) studied the conditions for the successful cultivation of water plants in spaceships. The efficiency of a duckweed system proved to be superior to one of algae. To produce enough oxygen for one person, about 2.5 m² of cultivation area are necessary. Photosynthetic and respiratory gas exchange of S. polyrrhiza was not affected by exposure to near weightlessness over a period of 230 hours (WARD et al. 1970).

3.8.2. Manure

In regions with mass development of Lemnaceae, the duckweed cover is distributed onto the fields and gardens as manure (e.g. ALIKUNHI et al. 1952). Already 1854 WELWITSCH reported (in herb.) that W. arhiza is used as manure in Angola. The cultivation of Lemnaceae for manure in China is described by TAI-HSING et al. (1975). In southern and eastern China, Lemnaceae are cultivated in ponds, channels, rice fields and other waters and harvested every 4 to 5 days to fertilize the fields (MERIAUX 1978, MCCALLA and PLUCKNETT 1981). In Mexico, the agricultural system of chinampa generates from aquatic vegetation containing mats of Lemnaceae mixed with other water plant. The detritus of these mats raises the soil level and further sedimentation accumulates by addition-

al duckweed cover and mud brought from the neighbouring channels. Eventually, a soil with excellent texture, and good water and cation exchange capacity develops and offers the possibility of harvesting four crops a year of vegetables or corn (LOT et al. 1979).

In the dry state, Lemnaceae contain 2-7% N, 0.5-3% P, and 2-5% K (CULLEY et al. 1978, cf. chapter 1.1). Mixed stands with Azolla are especially suited for manure production. Azolla has a symbiotic interrelation with the blue algae Anabaena azollae which is able to fix nitrogen from the air. PARK and YATAZAWA (1979) and ZUBERER (1981, 1982) demonstrated that Lemnaceae also live together with some nitrogen fixing heterotroph bacteria (e.g. Klebsiella) and Cyanophyta (e.g. Calothrix, Microchaete; Anabaena was only found in L. gibba). Up to 10⁵ individual organisms per g dry weight could be counted. They cannot supply more than 15-20% of the nitrogen needed by Lemnaceae (ZUBERER 1982). The nitrogen fixation in Azolla seems to be more efficient. SUTTON (1981b) states that up to 50% of the nitrogen need of rice in rice fields can be supplied by Azolla. According to EI-DIN (1982) a L. gibba association is able to fix 60 kg nitrogen per ha in 100 days which amounts to about 1/7 of the value of an Azolla cover. In Egypt the total N accumulation by L. gibba is 119-140 kg per ha and year.

3.8.3. Reduction of water loss in arid regions

Lemnaceae are possibly the only water plants that evapotranspire less than the same area of open water evaporates. The ratio evapotranspiration/evaporation amount to c. 0.9 for a L. obscura cover (named as L. minor) in southern USA (DEBUSK 1980, RYTHER et al. 1980, DEBUSK et al. 1983). BOYD (1975) noted a difference between different species. He measured a ratio of 0.9 for W. columbiana, and 0.85 for S. polyrrhiza. SEYBOLD (1930, cited from DEBUSK 1983) observed a lower ratio with greater wind velocities. Similarly, ORON et al. (1985) measured, in Israel, a distinctly lower ratio if the evaporation rate is higher than 4.5 mm per day: 0.75 for S. polyrrhiza and 0.7 for L. gibba. Most other water plants (e.g. Typha, Cyperus, Eichhornia) have a much greater evapotranspiration than the evaporation. Only in Pistia, Trapa, and Ipomoea it is not significantly higher than the evaporation (BREZNY et al. 1973). The water loss of ponds covered with Eichhornia turned out to be

3 to 5 times greater than in free surface ponds (ORON et al. 1985). The relatively low evapotranspiration of the Lemnaceae is due to the small frond surface in comparison with the water surface and with the surface of Eichhornia.

In conclusion, a Lemnaceae cover on the surface of water reservoirs and ponds in climatically dry regions, in order to prevent great water loss, may be advisable if it does not interfere with other purposes.

3.8.4. Reduction of mosquito breeding

A survey of possible use of Lemnaceae in fighting mosquito breeding is given by JENKINS (1964). In waters with a complete Lemnaceae cover JOHN-SON (1902) observed no mosquito breeding whereas in open spaces, he was able to detect larvae of Culex and Anopheles. According to ANCONA (1930), a closed cover of Lemnaceae mixed with Azolla prevented the development of some mosquito larvae. On the other hand, DYAS and KNAB (cited in MATHESON and HINMAN 1929) state that one of the most abundant breeding grounds of Culex salinarius was a large marsh completely covered by Lemna. BENTLEY (1910) reports from India that a cover of Lemna or Azolla is of no value in preventing the presence of mosquito larvae. However, a layer of Wolffia (probably W. globosa named as W. arrhiza) keeps the water free of larvae of Anopheles, Culex and Stegomyia. Larvae which were placed in water with a Wolffia cover died within several hours. Most authors observed a positive effect of Lemnaceae cover in preventing the breeding of mosquitoes (e.g. ADIE 1904, HILDEBRAND 1925, MATHESON and HINMAN 1929, HESS and HALL 1945, LAIRD 1956, JENKINS 1964, SMITH and ENNS 1967, SJOGREN 1968, KERBABAIEV et al. 1985). CULLEY and EPPS (1973) state that Spirodela hinders Anopheles mosquitoes from laying eggs. In addition, the larvae do not get enough oxygen below a duckweed cover. Moreover, Lemna covered ponds harbour numerous predatory insects which attack mosquito larvae. BRADLEY (1932) lists three main factors resulting in a decrease of Anopheles breeding in waters completely covered by Lemnaceae:

- 1) larval food supply is low due to a poor development of phytoplankton;
- 2) larvae cannot break the surface cover for respiration;
- 3) larvae are not very effectively hidden from their enemies in the water below the Lemnaceae cover.

FURLOW and HAYS (1972) report a decrease of Culicidae (Diptera) larvae below a S. punctata cover. If the cover was completely closed, no mosquito breeding occurred. In general, Anopheles species and Uranotaenia sapphirina reacted much more sensitively than Culex erraticus. Probably, the continuous surface mat represented a barrier to female oviposition. ANGERILLI (1980) investigated the influence of extracts of different water plants (included L. minor) on the development of the mosquito Aedes aegypti. An extract of L. minor had a toxic effect on the larvae. Into the vessel with the extract, only about 5% of the eggs were laid compared with the number in distilled water. JUDD and BORDEN (1980) demonstrated that aqueous and methanolic extracts of L. minor prevent the oviposition of Aedes aegypti but not of Culex pipiens at concentrations of 1000 and 10000 ppm. At concentrations of 1, 10, and 100 ppm, no effect showed up. In an outdoor experiment, in basins with L. minor, ANGERILLI and BEIRNE (1980) observed distinctly less eggs (c. 1/2) and larvae (c. 1/8) than in basins without Lemnaceae. This effect was studied in Culex inornata and Culex pipiens.

POSTSCRIPT - ACKNOWLEDGEMENTS

ACKNOWLEDGEMENTS

The invaluable contributions of friends, colleagues, and coworkers to this monographic treatment were already acknowledged in volume 1 (see LANDOLT 1986, p. 10-11). Sincere thanks go again to all these persons. The second volume was typed by Berti Brun, the lay-out was done by Anny Honegger, and the English kindly checked by Sonia Türler. Anita Hegi drew all the graphs and took care of the contact with the libraries. Many thanks for these laborious and careful contributions.

LIVING PLANT COLLECTIONS

Available clones of Lemnaceae

Both authors keep a collection of living clones at their university address. There are c. 1100 clones available in Zürich (Geobotanisches Institut ETH, Stiftung Rübel, Zürichbergstrasse 38, CH 8044 Zürich), and c. 120 clones in Vienna (Universität für Bodenkultur, Botanisches Institut, Gregor Mendel-Strasse 33, A-1180, Vienna). The clones are sent free of charge to all scientists. The collection in Zürich is to be liquidated after 1990.

CONCLUSIONS AND OUTLOOK

The aim of the present monographical treatment of the family of Lemnaceae was to bring together all the information known up to now on this family and to present a family of flowering plants from as many aspects as possible. We hope that we make available at least the main bulk of knowledge which is scattered around in many journals and in different scientific fields. We are aware of the fact that it is not possible for two persons to give a complete view of all the items. Much interesting information has been lost during the touching up and the transcriptions

of the present work. However, some of the facts and indications, though very often presented independently in different chapters, show up at places not expected by the user. In any case the reader is recommended to consult the original literature if he is interested in more details.

The family of Lemnaceae which consists only of four genera and 34 species represents one of the most thoroughly investigated families of flowering plants which is demonstrated by c. 3200 scientific papers dealing with Lemnaceae. The interest in this family is still unbroken: since 1947 the titles on Lemnaceae have been doubling about every twelve years.

Some of the advantages of Lemnaceae as a scientific test object are repeated herewith:

- smallness (length of 0.5 to several mm)
- easy handling in aseptic culture (controlled conditions)
- fast vegetative propagation (cloning)
- possible economic importance.

The Lemnaceae are an excellent example to show how important it is to integrate different scientific fields to solve biological problems. The investigations concern taxonomy, morphology, ecology, ontogeny, physiology, phytochemistry, molecular biology, application etc. After all the investigations we have a fairly consistent picture of the biology of Lemnaceae. We certainly can anticipate how complex the interplay of all factors and all processes is and how big the gaps of our knowledge to a full understanding still are. Some of the results can be summarized as follows:

Lemnaceae are a group of flowering plants which are extremely adapted to free floating life in the water. They are able to take advantage of their habitat by the following characteristics completely or partly realized within the family:

- rigid stomata
- absence of xylem and mechanical elements
- nutrient absorption with the whole surface immersed in the water
- independent individual floating of vegetatively propagating unities for fast spreading.

The following adaptations meet with the precariousness of floating life:

- excretion of phosphatases and RNases to improve availability of the minimum factor phosphorus
- storage of the phosphate in form of polyphosphates and phytin

- greater ability to accumulate micronutrients
 - protection against microorganisms by the formation of hardly decomposing cell walls (with apians) and by the excretion of phenols and flavonoids
 - formation of turions to survive shortage of nutrients and cold periods
 - formation of seeds to survive temporary drying.
- We hope that this work will give a stimulus to still more intense exploration of this fascinating family. The tiny little fronds of Lemnaceae which form the smallest of the flowering plants and are also contemptuously called weed are a marvellous example of well functioning and extremely adapted organisms. May the approach to this miraculous and many-sided plant fill all interested persons with enthusiasm.
- We close the work with the last sentence of H. BURGEFF in his studies on Marchantia (Genetische Studien an Marchantia, Fischer, Jena, 1954): "Wohl dem, der in der heutigen Periode des wissenschaftlichen Massenandrangs auf die Natur noch Gelegenheit hat, sich in Ruhe zu versenken, am Wachstum seines 'Materials' zu erfreuen und Beobachtungen zu machen, die, Fragen stellend, sie zugleich beantworten und den Frager zur Erleuchtung führen". ("Happy he who in the present period of scientific mass attack on nature still has the opportunity to peacefully enjoy the growth of his 'material' and make observations, asking questions and at the same time answering them, so leading the questioner to enlightenment").

BIBLIOGRAPHY

The bibliography comprises

- all the papers cited in the two volumes of the present work
- all papers dealing with Lemnaceae which came to the knowledge of the authors.

Since not all papers on Lemnaceae could be consulted in the original version, the references had to be taken from reviewing works or from literature services. Therefore, the indications may not always be complete. For the English, French, German, Italian, and Spanish papers, the titles are given in the original language. Titles of all other languages have been translated into English, and the original language is indicated in brackets. In very few cases it was not possible to translate the title. Most of the papers cited contain English summaries.

The papers are arranged alphabetically according to the first author. If two or more authors with the same surname occur, the alphabetical order of the initials is decisive. Surnames with prefixes as De, Van, Von, etc. are listed under the first letter of the prefix though this may sometimes be unfamiliar, especially if the prefix is a title of nobility.

Papers written by more than one author are listed as follows:

- papers with two authors are given in the alphabetical order
 - papers with more than two authors follow chronologically.
- This procedure is thought to facilitate the tracing of all citations in the text where the first author is followed by "et al."

- ABBA G., 1976: Apunti di Floristica Piemontese. Allonia 21, 97-103.
- ABDIEV M., 1971a: Duckweed communities in water bodies of Uzbekistane. (In Russian). Kultiv. Vodn. Vysch. Vodn. Rast. Uzbekistane 1971, 106-117.
- ABDIEV M., 1971b: Inflated duckweed (Lemna gibba L.) in culture. (In Russian). Kultiv. Vodn. Vysch. Vodn. Rast. Uzbekistane 131-136.
- ABDULLAEV D.A., 1969: The use of common duckweed as green feed for chickens. (In Russian). Uzbekskii Biol. Zh. (USSR) 13, 42-43.
- ABDULLAEV D.A., 1972: On the efficient utilization of aquatic swamp plants of the Fergana Valley in the economy. (In Russian). Kultiv. Vodn. Vysch. Vodn. Rast. Uzbekistane 119-132.
- ACCORSI C.A., MAZZANTI M.B., MARCHETTI D.B., FORLANI L. and SORBINI L., 1982: Primi cenni stratigrafici et e palinologici su sedimenti di pozzi nell'area di Verona. I. Sequenza di Avesa. G. Bot. Ital. 116 (suppl. 1), 143-144.
- ADABRA-MICHANOL Y., MONEGER R. and FRANCKE B., 1975: Effets de la kiné-tine à la lumière et à l'obscurité, sur les caroténoïdes et les chlorophylles des frondes de la Spirodela polyrrhiza (L.) Schleiden. Phytol. Veg. 13(3), 619-635.
- ADIE J.R., 1904: Lemna minor as a preventive against mosquitoes. Indian Med. Gaz. 39, 207-208.
- ABEILI D., 1986: Der Einfluss der Phosphat-Ernährung auf das Wachstum und die Entwicklung von Makrophyten. Institut für Pflanzenbau ETHZ. Diploma Thesis. 55 pp.
- AGAMI M., LIATAV M. and WAISEL Y., 1976: The effects of various components of water pollution on the behaviour of some aquatic macrophytes of the coastal rivers of Israel. Aquat. Bot. 2, 203-213.
- AGREN G.I., 1985: Limits to plant production. J. Theoret. Biol. 113(1), 89-92.
- AHMAD N. and YOUNUS M., 1979: Aquatic plants of Lahore. Pakistan Assoc. Advan. Sci. 41 pp.
- AHRENS J.F. and BLOCK P.M., 1972: Phytotoxicity and persistence of simazine in a shallow lake. Connecticut Agric. Exp. Stn. Valley Lab. Proc. NE Weed Soc. 26, 266.
- ALBANELLE E., PLAIZATS J. and ANDRES J., 1985: Interaction of abscisic acid and 6-benzylaminopurine on the metabolism of Lemna minor L. Plant Cell Physiol. 26(8), 1557-1564.
- ALBERGONI F.G. and BASSO B., 1985: An apparatus for the simultaneous measurement of photosynthetic ^{14}C fixation by a large number of leaf samples. Maydica 30(1), 75-84.
- ALBERSHEIM P., AUGUR C., DARVILL A.G., DOARES S.H., EBERHARD S., GOLLIN D.J., MARFA-RIERA V., MOHNER D., NURI W.A., O'NEILL R.A., TOUBART P. and YORK W.S., 1987: Oligosaccharines are active in defense responses, growth, and development. Lecture 2-096-2 at the 24th Int. Bot. Congress Berlin (West), Germany.
- ALGER G., 1956: Die Aufnahme von Harnstoff und Glycerin durch die Pflanzelle in Abhängigkeit von der Acidität der Aussenlösung und des Zellsaftes. Protoplasma 47, 78-102.
- ALIKHUNI S. ALIKHUNI
- ALIKHUNI K.H., RAMCHADRAN V. and CHAUDHURI H., 1952: On the role of duckweed (Lemna minor) in the preparation of carp nurseries. Sci. Cult. (Calcutta) 17, 436-437.
- ALLENBY K.G., 1967: The manganese and calcium contents of some aquatic plants and the water in which they grow. Hydrobiologia 29, 239-244.
- ALLENBY K.G., 1968: Some analyses of aquatic plants and waters. Hydrobiologia 32, 486-490.
- ALLENBY K.G., 1981: Some analyses of aquatic plants and their waters. Hydrobiologia 77(2), 177-189.

- ALLISON F.E., LOVE K.S., PINCK L.A. and GADDY V.L., 1948: Gaseous losses of nitrogen from green plants. I. Studies with Chlorella and Lemna. Plant Physiol. 23, 496-504.
- ALMADI L., 1961: Distribution of Wolffia arrhiza. (In Hung.). Bot. Közlem 49, 112-113.
- ALMESTRAND A. and LUNDH A., 1951: Studies on the vegetation and hydrochemistry of Scanian lakes I-II. Bot. Not. 2(3), 7-141.
- ALMQUIST E., 1929: The vegetation and flora of Uppland. (In Swedish). Acta Phytogeogr. Suecica 1, 622 pp.
- AL-RAIS A.H., MYERS A. and WATSON L., 1971: The isolation and properties of oxalate crystals from plants. Ann. Bot. 35(143), 1213-1218.
- AL-SHALAN I. and KANDELER R., 1978: Tageslängenabhängigkeit der Wirkung von Abscissinsäure und Benzyladenin auf das Wachstum von Lemna gibba Gl. Biochem. Physiol. Pflanzen 172, 521-529.
- AL-SHALAN I. and KANDELER R., 1979: Aufnahme und Verteilung von Gibberellin A₁ bei der Kurztagpflanze Lemna paucicostata 6746. Z. Pflanzenphys. 94, 257-262.
- AMADO R., MUELLER-HIEMEYER R. and MARTI U., 1980: Proteingehalt, Aminosäurezusammensetzung und Neutralzuckergehalt von Lemnaceen. Vorläufige Mitteilung. Veröff. Geobot. Inst. ETH, Stiftung Rühel, Zürich 70, 102-117.
- AMANOVA N.M., 1982: Boron in fodder plants of Tadzhikistan. (In Russian). Izv. Akad. Nauk Tadzh. SSR, Otd. Biol. Nauk 1, 64-70.
- AMBASHI R.S., 1970: Ecosystem study of a tropical pond in relation to primary production of different vegetational zones. Hydrobiologia 12, 57-61.
- AMBORSKI R.I. and LARKIN J.M., 1980: Human and animal health aspects. In: CULLEY D.D. and FRYE J.B. (eds.), US Dept. of Energy Final Rep. Baton Rouge. 15 pp. (Polycopy).
- AMBROSE K.B., 1978: The ecological significance and release of organic compounds by members of the Lemnaceae: A qualitative and quantitative investigation. M.S. Thesis, Univ. Pittsburgh. 83 pp.
- ANCONA H.L., 1930: Las Lemnaceas y las larvas de los mosquitos. Univ. Nac. Anal. Inst. Biol. (Mexico) 1, 33-37.
- ANDERSEN I.H., 1983: Effects of photosynthesis on Lemna gibba. (In Norwegian.) Hovedfagsoppgave, Bot. Inst., Fyottron. Univ. Oslo.
- ANDERSEN I.H., DONS C., NILSEN S. and HAUGSTAD M.K., 1985: Growth, photosynthesis and photorespiration of Lemna gibba: response to variations in CO₂ and O₂ concentrations and photon flux density. Photosynth. Res. 6(1), 87-96.
- ANDERSON A.C., ABDELGHANI A.A. and McDONELL D., 1980: Screening of four vascular aquatic plants for uptake of monosodium methanearsonate. Sci. Tot. Environ. 16, 95-98.
- ANDERSON J.L., THOMSON W.W. and SWADER J.A., 1973: Fine structure of Wolffia arrhiza. Can. J. Bot. 51, 1619-1622.
- ANDERSON J.W., 1980: Assimilation of inorganic sulfate into cysteine. In: STUMPF P.K. and CONN E.E. (eds.), The biochemistry of plants. Acad. Press, New York. 5, 203-223.
- ANDERSON J.W. and EMES M.J., 1985: Purification and properties of hydroxypyruvate reductase from Lemna minor. Int. J. Biochem. 17(10), 1085-1090.
- ANDERSON L.E., 1982: Isolation and purification of aldolase from Pisum sativum. In: EDELMAN M., HALLICK R.B. and CHUA N.H. (eds.), Methods in chloroplast molecular biology. Elsevier, North-Holland. Biomed. Press. 715-722.
- ANDRES J. and SMITH H., 1976: Evidence for a rapid effect of abscisic

- acid on amino acid metabolism in Lemna. *Plant.Sci.Lett.* 6, 315-318.
- ANDRONOVA L.M., 1972: Antitussive properties of certain medical plants. (In Russian). *Rastit.Resur.* 8(4), 588-591.
- ANGERILLI N.P.D., 1980a: Influences of extracts of freshwater vegetation on the survival and oviposition by *Aedes aegypti* (Diptera: Culicidae). *Can.Entomol.* 112, 1249-1252.
- ANGERILLI N.P.D., 1980b: A similarity dendrogram as an indicator of mosquito breeding sites. *Can.Mosq.News* 40(1), 117-118.
- ANGERILLI N.P.D. and BEIRNE B.P., 1974: Influences of some freshwater plants on the development and survival of mosquito larvae in British Columbia. *Can.J.Zool.* 52, 813-815.
- ANGERILLI N.P.D. and BEIRNE B.P., 1980: Influences of aquatic plants on colonization of artificial ponds by mosquitoes and their insect predators. *Can.Entomol.* 112(8), 793-796.
- ANGERILLI N.P.D. and BEIRNE B.P., 1982: Mortality of introduced mosquito larvae in natural and artificial ponds containing aquatic vegetation. *Prot.Ecol.* 4(4), 381-386.
- ANTANINENE A.S. and TRAINAUSKAITE I.Y., 1985: Production and biochemical characteristics of macrophytes from Lake Druksiai, Lithuanian-USSR, in 1982. (In Russian). *Liet.Tsr.Mokslu Akad.Darb.Ser.C.Biol.Mokslai* 1985(1), 68-75.
- ANTYPCHEUK A.P., 1974: Numbers of heterotrophic bacteria on some higher plants in carp ponds. *Hydrobiol.J.* 10(1), 48-49.
- ANTONIELLI M. and CAGIOTTI M.R., 1976: Presenza di clorofillide in idrofite ed alghe del lago Trasimeno e variabilità del pigmento in *Spirogyra*. *Riv.Idrobiol.* 14, 3-11.
- APPENROTH K.-J. and AUGSTEN H., 1987: An improvement of the protein determination in plant tissues with the dye-binding method according to Bradford. *Biochem.Physiol.Pflanz.* 182(1), 85-89.
- APPENROTH K.-J., AUGSTEN H., LIEBERMANN B. and FEIST H., 1982: Effects of light quality on amino acid composition of proteins in *Wolffia arrhiza* (L.) Wimm. using a specially modified Bradford method. *Biochem.Physiol.Pflanz.* 177(3), 251-258.
- APPENROTH K.-J., MOOTZ M. and AUGSTEN H., 1986: Effect of light quality on the alanine and aspartate aminotransferases in *Wolffia arrhiza* L. In: LAMBERS H., NEETESON J.J. and STULEN I. (eds.), *Fundamental, ecological and agricultural aspects of nitrogen metabolism in higher plants*. Nijhoff Publ., Dordrecht/Boston/Lancaster. 215-218.
- APPLETON C.C. and SHARP B.L., 1985: A preliminary study on the emergence of *Mansonia uniformis* from swamps at Richards Bay, Natal, South Africa. *J.Entomol.Soc.Southern Africa* 48(1), 179-184.
- ARAI S., 1986: Enzymatically modified proteins as new surfactants, their properties and functions. *J.An.Oil Chem.Spc.* 63(4), 420-421.
- ARBER A., 1919: On the vegetative morphology of *Pistia* and the *Lemna*-ceae. *Proc.R.Soc.* 91, 96-103.
- ARBER A., 1920: Water plants: a study of aquatic angiosperms. Cambridge Univ.Press. 435 pp.
- ARMITAGE K.B. and FASSETT N.C., 1971: Aquatic plants of El Salvador. *Arch.Hydrobiol.* 69(2), 234-255.
- ARMSTRONG W.P., 1981a: *Wolffia punctata* Griseb. (Lemnaceae). *Madroño* 28, 37.
- ARMSTRONG W.P., 1981b: *Wolffia columbiana* Karst. (Lemnaceae). *Madroño* 28, 187-188.
- ARMSTRONG W.P., 1982: Duckweeds, California's smallest wildflowers. *Fremontia* 10(3), 16-22.
- ARMSTRONG W.P., 1983: A marriage between a fern and an alga. *Environ. Southwest* 500, 20-24.

- ARMSTRONG W.P., 1984: *Wolffia globosa* (Roxb.) Hartog et Plas (Lemnaceae). *Madroño* 31(3), 191.
- ARMSTRONG W.P., 1985: A status report on the genus *Wolffia* in California. *Fremontia* 13(1), 11-14.
- ARMSTRONG W.P., 1986: Lemnaceae - Duckweed Family in California. 25 pp. (Polycopy).
- ARMSTRONG W.P. and THORNE R.F., 1984: The genus *Wolffia* (Lemnaceae) in California. *Madroño* 31(3), 171-179.
- ARNASON T., HEBDA R.J. and JOHNS T., 1981: Use of plants for food and medicine by native peoples of Eastern Canada. *Can.J.Bot.* 59, 2189-2325.
- ARNOLD H., WUESTLING W., HIEKEL H.G. and BOEHM H., 1979: Influence of substituents in various acetophenone derivatives on course of action on gas interchange inhibition of synchronous cultures *Chlorella vulgaris*. (In German). In: SCHUETTE H.R., (ed.), *Wirkungsmechanismen von Herbiziden und synthetischen Wachstumsregulatoren*. Fischer, Jena. 93-100.
- ARNER D.H., WESLEY D. and ANDING G., 1968: A quantitative study of invertebrates found in certain wetland plant communities in Mississippi. *pl. Mississippi State Univ. Water Resources Res.Inst.* 25 pp.
- ARNOTT H.J. and PAUTARD F.G.E., 1965: Development of raphide idioblasts in Lemna. *Amer.J.Bot.* 52, 618-619.
- ARNOTT H.J. and PAUTARD F.G.E., 1970: Calcification in plants. In: SCHRAER H. (ed.), *Biological calcification*. New York.
- ARO E.-H., 1982: A comparison of the chlorophyll-protein composition and chloroplast ultrastructure in two bryophytes and two higher plants. *Z.Pflanzenphysiol.* 108(2), 97-105.
- ASCHERSON P. and GRABNER P., 1902-1904: *Synopsis der mitteleuropäischen Flora*. Engelmann, Leipzig. 2/2, 390-397.
- ASHBY E., 1929a: The interaction of factors in the growth of Lemna. III. The interrelationship of duration and intensity of light. *Ann.Bot.* 43, 333-354.
- ASHBY E., 1929b: The interaction of factors in the growth of Lemna. IV. The influence of minute quantities of organic matter upon growth and reproduction. *Ann.Bot.* 43, 805-816.
- ASHBY E., 1935: An analysis of the influence of light and temperature on the assimilation rate and the rate of multiplication. *Ann.Bot.* 49, 309-336.
- ASHBY E., 1950: Leaf morphology and physiological age. *Sci.Prog.* 152, 678-685.
- ASHBY E. and OXLEY T.A., 1935: The interaction of factors in the growth of Lemna. VI. An analysis of the influence of light intensity and temperature on the assimilation rate and the rate of frond multiplication. *Ann.Bot.* 49, 309-336.
- ASHBY E. and WANGERMANN E., 1949: Senescence and rejuvenation in Lemna minor. *Nature* 164, 187.
- ASHBY E. and WANGERMANN E., 1950 s. 1949
- ASHBY E. and WANGERMANN E., 1951: Study in the morphogenesis of leaves. VII. 2. Correlative effects of fronds in Lemna minor. *New Phytol.* 50, 200-209.
- ASHBY E. and WANGERMANN E., 1954: The effects of meristem ageing on morphology and behaviour of fronds in Lemna minor. *Ann.N.Y.Acad.Sci.* 57, 476-483.
- ASHBY E., BOLAS B.D. and HENDERSON F.Y., 1928: The interaction of factors in the growth of Lemna. I. Method and technique. *Ann.Bot.* 42, 771-782.

- ASHBY E., WANGERMAN E. and WINTER E.J., 1949: Studies in the morphogenesis of leaves. III. Preliminary observations on vegetative growth in Lemna minor. New Phytol. 48, 374-381.
- ASTON H.I., 1973: Aquatic plants of Australia. Melbourne Univ. Press. 368 pp.
- ATHIAS-BINCHIE F. and FERNANDEZ N.A., 1986: Analyse démographique d'une population d'Hydrozetes lemnae Coggi, acarien oribate inféodé à la lentille d'eau Lemna gibba L. en Argentine. 2. Les relations prédateur/proie. Zool. Jb. Syst. 113(2), 229-249.
- ATHI F.R., 1983: Schwermetalle und Wasserpflanzen. Schriftenreihe Ver. Wasser-, Boden- u. Lufthygiene 55, 155 pp.
- AUGSTEN H., 1983: Zur Eignung der Lemnaceae als Testsystem für Herbizide. Wiss. Schriftenreihe Päd. Hochschule Potsdam, Reihe B, 35, 86-91.
- AUGSTEN H., 1984a: Lemnaceen: Aspekte ihrer Praxisrelevanz. Biol. Rundsch. 22(4), 225-235.
- AUGSTEN H., 1984b: Photosynthese und Biomassebildung bei Lemnaceen. Coll. Pfl. Physiol., Humboldt-Universität, Berlin, 7(2), 213-222.
- AUGSTEN H. and HUBALD M., 1980: Das rhythmische Verhalten der Nitratreduktase bei Lemnaceen und der Einfluss von Aminosäuren auf die Aktivität des Enzyms. Wiss.-Z. Ernst-Moritz-Arnt-Universität Greifswald, Math.-Natw. Reihe 29(1-2), 47-50.
- AUGSTEN H. and JUNGnickel F., 1979: Verfahren zur Ausleseprüfung und Bestimmung phytoaktiver Substanzen. German East Patent No. 133595. 21 pp.
- AUGSTEN H. and JUNGnickel F., 1983: Zur Steuerung der Turionenausbildung bei Spirodela polyrrhiza (L.) Schleiden durch Licht, Phosphat und Zink. Wiss.-Z. Ernst-Moritz-Arnt-Universität Greifswald, Math.-Natw. Reihe 32(3-4), 64-66.
- AUGSTEN H. and JUNGnickel F., 1984: Action of microelements on developmental processes in mass propagated axenic plants. Proc. Intern. Symp., Plant tissue and cell culture application to crop improvement. Prague. 497-498.
- AUGSTEN H., KUNZ E. and APPENROTH K.-J., 1987: Phytochrome-mediated germination of light and dark grown turions of Spirodela polyrrhiza. 14th Int. Bot. Congr. Berlin, Abstr., 124.
- AUSTIN C.F., 1967: Lemnaceae. In: GRAY A., Man. Bot., 478-480.
- AUSTIN C.F., 1970: Wolffia columbiana. Bull. Torr. Bot. Club 1, 36.
- AVENA G.C., BLASI C. and RUBICA L., 1975: Riccio fluitantis-Azolleum caroliniana, associazione nuova indicatrice ecologica per lo stato delle acque di un tratto del F. Tevere. Ann. Bot. (Roma) 34, 175-190.
- AVENA G.C., BLASI C. and SCOPPOLA A., 1982: Indagini ecologico-fittografiche sulle zone umide interne del Lazio. II. Sintassonomia delle comunità afferenti alla classe Lemneta minoris presenti nella Bonifica Pontina. Ann. Bot. (Roma) 40, 49-61.
- AVADI A., 1971: Effet du calcium sur l'absorption du potassium par la Lemna minor. Physiol. Veg. 9(1), 107-108.
- AVADI A. and THELLIER M., 1970: Formulation électrocinétique de l'effet de régulation par le calcium de l'absorption du rubidium chez la Lemna minor. C.R. Acad. Sci. Paris, D 271, 1280-1283.
- AVADI A., DEMUYTER P. and THELLIER M., 1971: Interprétation électrocinétique des interactions compétitives réciproques K/Rb lors de l'absorption de ces ions par la Lemna minor. C.R. Acad. Sci. Paris, D 273, 67-70.
- AVADI A., STELZ T., MONNIER A., LASSALLE L.P. and THELLIER M., 1974: Application of an electrokinetic formulation to the study of the effect of alkaline-earth cations on the absorption of K-ions by Lemna minor. Ann. Bot. 38, 677-696.

- BAJECZKO W., 1982: Wolffia arrhiza (L.) Wimm. in the West Pomerania area (Poland). (In Polish). Zesz. Nauk. Acad. Roln. Szcz. 29, 3-11.
- BAHADIR M., 1987: Safe formulations of agrochemicals. Chemosphere 16(2-3), 615-622.
- BAHADIR M. and KORTE F., 1987: New slow release application techniques with heterocyclic herbicides in aquatic weed control. Heterocycles (Tokyo) 25(1), 333-335.
- BAHADIR M. and PRISTER G., 1985: A comparative study on pesticide formulations for application in running waters. Ecotoxicology and Environ. Safety 10(2), 197-201.
- BAHL J., 1971: Sur la préparation de suspensions de chloroplastes isolés de Spirodela polyrrhiza (L.) Schleiden. C.R. Acad. Sci. Paris, D 272, 2185-2187.
- BAHL J., LECHEVALLIER D. and MONEGER R., 1971: Sur les lipides chloroplastiques des frondes de la Spirodela polyrrhiza (L.) Schleiden cultivées sur milieu minéral et sur milieu saccharosé. C.R. Acad. Sci. Paris, D 272(18), 2330-2323.
- BAI R., 1985: Biodegradability of substrate during anaerobic digestion of various biomass. Reg. J. Energy Heat Mass Transfer 7(1), 33-38.
- BAILEY D.R., 1971: Comparison of oats and Lemna oligorrhiza for diuron bioassay. Queensl. J. Agric. Anim. Sci. 27, 395-400.
- BAKAYA U. and KAUL V., 1973: Lemna-Salvinia, the noxious floating aquatic weed complex in Kashmir. Regional seminar on noxious aquatic vegetation in tropics and sub-tropics. Abstr., 21-22. UNESCO, New Delhi, India. 70 pp.
- BAKER J.H. and FARR I.S., 1987: Importance of dissolved organic matter produced by duckweed (Lemna minor) in a southern English river. Freshwat. Biol. 17(2), 325-330.
- BAKER J.H. and ORR D.R., 1986: Distribution of epiphytic bacteria on freshwater plants. J. Ecol. 74, 155-165.
- BAKHADIROVA Z., 1982: Flora and vegetation of fish ponds in the Tashkent fish nursery (Uzbek-SSR, USSR). (In Russian). Uzb. Biol. Zh. 1982(3), 33-34.
- BAKSHI I.S., 1983: Effects of asparagine, irradiation and NADH on the nitrate reductase in Wolffia microscopica. Biol. Plant 25(4), 259-265.
- BAKSHI I.S., FAROOQI A.H.A. and MAHESHWARI S.C., 1978: Circadian rhythm in nitrate reductase activity in the duckweed Wolffia microscopica Griff. Z. Pflanzenphysiol. 90(2), 165-169.
- BAKSHI I.S., FAROOQI A.H.A. and MAHESHWARI S.C., 1979: Control of circadian rhythm in nitrate reductase activity in Wolffia microscopica Griff. Plant Cell Physiol. 20(5), 957-963.
- BALDI B.G., SLOVIN J.P. and COHEN J.D., 1984: Synthesis and activity of halogen substituted indole-3-acetic acids: suitability for use in selection of genetic mutants. Plant Physiol. 75(suppl. 1), 107.
- BALDWIN J.D.C. and MYERS R., 1980: Duckweed harvesting and transport. Baton Rouge. 5 pp. (Polycopy).
- BALL G.A., BEAL E.O. and FLECKER E.A., 1967: Variation of chromatographic spot pattern of two species of clonal plants grown under controlled environmental conditions. Brittonia 19, 273-279.
- BANDERJEE M., 1971: Chromosome studies in Lemnaceae. Rev. Roum. Embryol. Ser. Cytol. 8(2), 21-27.
- BANGERTER E.B., 1978: New and interesting records of adventive plants from the Auckland Institute and Museum Herbarium. Part 4. Rec. Auckland Mus. 15, 27-36.
- BARBECK W., 1880: On the development of Lemna minor. Proc. Acad. Philadelphia 1880, 230-232.

- BARBER J.T. and CALDWELL C.R., 1976: Ascorbic acid and Lemna gibba: uptake, binding to macromolecules and effects upon RNA synthesis and flowering. *Plant Physiol.* 57(5), 21.
- BARBORA B.C., 1972: Chemical control of weeds in drains, drain sides and other non-tea pockets. *Two and a Bud* 19(1), 25-28.
- BARON D., WELLMANN E. and GRISEBACH H., 1972: Purification and properties of an enzyme from cell suspension cultures of parsley catalyzing the synthesis of UDP-apiose and UDP-D-xylose from UDP-D-glucuronic acid. *Biochim. Biophys. Acta* 258, 310-318.
- BASZYNSKI T., KROL M. and MOLINSKA D., 1981: Photosynthetic apparatus of Lemna minor L. as affected by chromatate treatment. In: AKOYUNOGLU G. (ed.), *Photosynthesis II. Electron transport and photophosphorylation*. Photosynth., Proc. Int. Congr., 5th, 2, 111-122.
- BATA J., 1973: The effect of pH, kinetin and sucrose on the branching pattern in Lemna trisulca L. *Bull. Inst. Jard. Bot. Univ. Beograd* 8, n.s., 45-52.
- BATA J. and NESCOVIC M., 1974: The effect of gibberellic acid and kinetin on chlorophyll retention in Lemna trisulca L. *Z. Pflanzenphys.* 73, 86-88.
- BATA J. and NESKOVIC M., 1982a: Effect of external factors on growth and morphogenesis in Lemna trisulca L. in axenic culture. I. Plant hormones. *Bull. Inst. Jard. Bot. Univ. Beograd* 15(1-3), 1-9.
- BATA J. and NESKOVIC M., 1982b: Effect of external factors on growth and morphogenesis in Lemna trisulca L. in axenic culture. II. Light conditions. *Bull. Inst. Jard. Bot. Univ. Beograd* 15(1-3), 10-13.
- BAUER A., SCHAEFER C. and ERTSMANN K.H., 1971: Zur Frage der Stickstoffinkorporation in Aminosäuren und Proteine bei Lemna minor unter Photosynthesebedingungen. *Verh. Schweiz. Naturf. Ges.* 151, 107-109.
- BAUER A., SCHLUNEGGER U. and ERTSMANN K.H., 1973: Untersuchungen zur Amoniumassimilation in Aminosäuren und Proteine bei Lemna minor L. unter Photosynthesebedingungen. *Verh. Schweiz. Naturf. Ges.* 153, 79-82.
- BAUER R., HUBER W. and SANKHLA N., 1976: Effect of abscisic acid on photosynthesis in Lemna minor L. *Z. Pflanzenphys.* 77, 237-246.
- BAUMEISTER W. and ERNST W., 1978: Mineralstoffe und Pflanzenwachstum. Fischer, Stuttgart/New York.
- BAUR R.J. and BUCK D.H., 1980: Active research on the use of duckweeds in the culture of grass carp, Tilapia, and fresh water prawns. *Ill. Nat. Hist. Survey, R.R.I., Kinmundy, Ill.* (Polycopy).
- BEAL E.O., 1977: A manual of marsh and aquatic vascular plants of North Carolina with habitat data. N.C. Agric. Exp. Stat. Tech. Bull., Raleigh, N.C., 247, 298 pp.
- BEAMS H.W., KESSEL R.G. and SHIH C.Y., 1979: Effects of ultracentrifugation on the mesophyll cells and chloroplasts of spinach leaf, and on the cells and chloroplasts of entire duckweed plants. *Biol. Cell.* 35, 87-96.
- BEARDSLEE H.C., 1877: *Wolffia*. Bot. Gaz. 2, 99.
- BEATSON M.E., 1955: Data for the study of postglacial history. XV. Subfossil pollen of Lemna in Quaternary deposit. *New Phytol.* 54, 208.
- BEAUMONT G., BASTIN R. and THERRIEN H.P., 1976: Effets physiologiques de l'atrazine à doses sublétales sur Lemna minor L. I. Influence sur la croissance, la teneur en chlorophylle, en protéines et en azote soluble et total. II. Influence sur la photosynthèse et sur la respiration. *Nat. Can.* 103, 527-533; 103, 535-541.
- BEAUMONT G., BASTIN R. and THERRIEN H.P., 1978: Effets physiologiques de l'atrazine à doses sublétales sur Lemna minor L. III. Influence sur les protéines solubles et les acides nucléiques. *Nat. Can.* 105, 103-113.

- BEAUMONT G., LORD A. and GRENIER G., 1980: Effets physiologiques de l'atrazine à doses sublétales sur Lemna minor. V. Influence sur l'ultrastructure des chloroplastes. *Can. J. Bot.* 58, 1571-1577.
- BEAUVOIS DE P., 1916: Mémoire sur les Lemna ou lentille d'eau, sur leur fructification et sur la germination de leur grain. *J. Phys. Chim. Hist. Nat.* 82, 101-115.
- BECHERER A., 1956: *Flora Vallesiacae Supplementum*. Denkschr. Schweiz. Naturf. Ges. 81, 556 pp.
- BECK E., 1964: Apian aus der Zellwand höherer Pflanzen. *Ber. Deutsch. Bot. Ges.* 77, 396-397.
- BECK E., 1967: Isolierung und Charakterisierung eines Apioalakturonans aus der Zellwand von Lemna minor. *Z. Pflanzenphysiol.* 57, 444-461.
- BECK E., 1969: Biosynthesis of branched chain sugars. *Xith Int. Bot. Congr.*, Seattle, Abstr., 10.
- BECK E., 1982: Branched-chain sugars. In: LOEWUS F.A. and TANNER W. (eds.), *Plant carbohydrates I*, Encycl. Plant. Physiol. Springer, Berlin/Heidelberg/New York. n.s. 13a, 124-157.
- BECK E. and KANDLER O., 1965: Apiose als Bestandteil der Zellwand höherer Pflanzen. *Z. Naturforsch.* 20b, 62-67.
- BECK E. and KANDLER O., 1966: Isotopenstudien zur Biosynthese von Apiose in Lemna. *Z. Pflanzenphysiol.* 55, 71-84.
- BEER S., 1985: Effects of CO₂ and O₂ on the photosynthetic O₂ evolution of Spirodela polyrrhiza turions. *Plant Physiol.* 79(1), 199-201.
- BEGUM S. and KHATUN R., 1983: Study on the occurrence of aquatic weeds and algae in Dhaka city and its suburbs. *Bangladesh J. Sci. Ind. Res.* 18(1/4), 136-145.
- BELL D.J., HARDWICK N.E., ISHERWOOD F.A. and CAHN R.S., 1954: D-(+)-Apiose from the monocotyledon Posidonia australis. *J. Chem. Soc. (London)* 1954, 3702.
- BELL W.A., 1949: Uppermost Cretaceous and Paleocene floras of Western Alberta. *Canada Dept. Mi. Res. Geol. Surv. Bull.* 13, 1-229.
- BELLANDO M., MONTACCHINI F. and AMEDEI G., 1978: L'azione fitotossica dell'Irpexil (acetidrossamioacetato di metile) sulle piante superiori. Nota prima: azione brachizzante ed inibizione mitotica. *G. Bot. Ital.* 112(1-2), 97-106.
- BENKOVA D., 1957: Blühende Wolffia (Wolffia arrhiza [L.] im nördlichen Kaukasus-Vorland. (In Russian). *Biologia* 12, 460-463.
- BENNINK G.J.H. and DE VRIES J.W.A., 1975: Flower development in Chrysanthemum and Lemna under long-day conditions with the cytokinin benzyladenine. *2nd Symp. Plant Growth Regulators, Sofia, 1975*. Abstr., 66-67.
- BENNINK G.J.H. and DE VRIES J.W.A., 1977: Flower development in Lemna perpusilla under long-day conditions with the cytokinin benzyladenine. In: KUDREV T., IVANOVA I. and KARANOV E. (eds.), *Plant Growth Regulators. Proc. 2nd Int. Symp., Sofia, 1975*. 544-547.
- BENNINK G.J.H., VAN DEN BERG R., KOOL H.J. and STEGWEE D., 1970: Flowering in Lemna minor. *Acta Bot. Neerl.* 19, 385-392.
- BENSON A.A., NISSEN P., COONEY R.V., HERRERA-LASSO J.M. and SUMMONS R. E., 1981: Arsenic metabolic paths in aquatic and terrestrial plants. *Proc. 13th Int. Bot. Congr. Abstr.*, 26.
- BEN-TAL Y. and CLELAND C.F., 1981: Investigation of the mode of action of salicylic acid in inducing flowering in Lemna obscura through the use of [¹⁴C] labeled salicylic acid. *Plant Physiol.* 67(4 suppl.), 103.
- BEN-TAL Y. and CLELAND C.F., 1982: Uptake and metabolism of [¹⁴C] salicylic acid in Lemna gibba strain G3. *Plant Physiol.* 70(1), 291-296.

- BENTLEY C.A., 1910: The natural history of Bombay malaria. J. Bombay Nat.Hist.Soc. 20, 392-422.
- BEPPU T., 1981: *Wolffia* in Japan. (In Japan.). Acta Phytotax.Geobot. 32, 197.
- BEPPU T. and TAKIMOTO A., 1981a: Geographical distribution and cytological variation of *Lemna paucicostata* Hegelm. in Japan. Bot.Mag.Tokyo 94, 11-20.
- BEPPU T. and TAKIMOTO A., 1981b: Further studies on flowering of *Lemna paucicostata* in Japan. Bot.Mag.Tokyo 94, 69-76.
- BEPPU T. and TAKIMOTO A., 1981c: Growth of various ecotypes of *Lemna paucicostata* in Japan under various temperature conditions, and their wintering forms. Bot.Mag.Tokyo 94, 107-114.
- BEPPU T. and TAKIMOTO A., 1983: Flowering behavior of the hybrids between strains 6746 and 371 of *Lemna paucicostata* Hegelm. Aquat.Bot. 17(3-4), 295-299.
- BEPPU T. and TAKIMOTO A., 1984, s. 1983
- BEPPU T., YANASE D., NOBUCHI T. and MURATA G., 1985: A revision of *Lemna paucicostata* group in Japan. (In Japanese). Acta Phytotax.Geobot. 36, 45-58.
- BERBERICH S., 1986: Duckweed may unlock sunflower's secret. Agric.Res., Washington DC, 34(7), 12.
- BERGAMINI P.G., PALMAS G., PIANTILLI F., SANI M., BANDITELLI P., PREVITERA M. and SODI F., 1979: Study of ¹³⁷Cs absorption by *Lemna minor*. Health Phys. 37, 315-321.
- BERNABEU C., TOBIN E.M., FOWLER A., ZABIN I. and LAKE J.A., 1983: Nascent polypeptide chains exit the ribosome in the same relative position in both eukaryotes and prokaryotes. J.Cell Biol. 96, 1471-1474.
- BERRIMAN D.J., 1973: The first record of *Spirodela punctata* (Lemnaceae) from Rhodesia. Kew Bull. 28, 325-326.
- BERRY C.J., Jr. and SCHRECK C.B., 1975: Toxicity of diquat and endothal separately and combined on several species of aquatic plants. Va.J.Sci. 26(4), 159.
- BERTOSSI M.F., 1950: L'idrazide maleica come fitormone. Atti.Ist.Bot. Univ.Pavia ser. 5, 8, 155-166.
- BERTSCH K., 1952: *Wolffia arrhiza* in Württemberg. Jb.Ver.Vaterl.Naturkde. Würtbg. 107, 133-136.
- BERUBE R., BEAUMONT G. and GRENIER G., 1982: Effets physiologiques de la benzyladénine sur la *Lemna minor*. I. Influence sur la composition lipidique. Can.J.Bot. 60(6), 758-764.
- BEST P.H., PIETERSE A.H., SOEKARJO R. and DE LANGE L., 1977: A preliminary study of the internal gas composition of *Lemna gibba* L. Acta Bot.Neerl. 26, 109-113.
- BEYER B.H., 1983: Ökologisch-systematische Differenzierung in der Familie der Lemnaceae in Bezug auf den Bedarf an Calcium und Magnesium. Diploma Thesis. Geobotanisches Institut ETH, Stiftung Rübel, Zürich. 137 pp. (Polycopy).
- BEZEMER-SYBRANDY S.M., 1969: Investigations on cytokinins. Studies with *Lemna minor* L. (In Dutch). Ph.D.Thesis. Univ.Leiden. 81 pp.
- BEZEMER-SYBRANDY S.M. and VELDSTRA H., 1971a: Investigations on cytokinins. III. Cytokinin activity in *Lemna minor* tRNA hydrolysates. Phytol.Plant. 24, 369-373.
- BEZEMER-SYBRANDY S.M. and VELDSTRA H., 1971b: Investigations on cytokinins. IV. The metabolism of 6-benzylaminopurine in *Lemna minor*. Phytol.Plant. 25, 1-7.
- BEZEMER-SYBRANDY S.M., TASSERON-DE JONG J.C. and VELDSTRA H., 1968: Effects of visible light on kinetin solutions. Biochem.Biophys.Acta 161, 568-571.

- BHALLA P.R., PIETERSE A.H. and SABHARWAL P.S., 1973: Some aspects of flowering, gibbosity and turion formation in Lemnaceae. Acta Bot. Neerl. 22, 433-445.
- BHALLA P.R. and SABHARWAL P.S., 1971: Investigations on the effect of tobacco smoke constituents on growth and morphogenesis of plant tissues. In Vitro 6(5), 402.
- BHALLA P.R. and SABHARWAL P.S., 1972a: Induction of flowering in *Lemna minor* by EDDHA. Acta Bot.Neerl. 21, 200-202.
- BHALLA P.R. and SABHARWAL P.S., 1972b: Induction of flowering in *Lemna gibba* G3 by aspirin. In Vitro 7(4), 281.
- BHALLA P.R. and SABHARWAL P.S., 1974: Effects of tobacco smoke and some of its constituents on growth and flowering of *Lemna gibba* G3. Environ.Pollut. 6, 59-66.
- BHALLA P.R. and SABHARWAL P.S., 1975: Induction of flowering in *Lemna gibba* G3 by aspirin. Experientia 31, 540-541.
- BHANTHUNNAVIN K. and MCGARRY M.G., 1971: *Wolffia arrhiza* as a possible source of inexpensive protein. Nature 232(5331), 495.
- BHAT P.K., DAGAR J.C., SINGH V.P. and MALL L.P., 1977: Studies on interspecific associations of aquatic plants. Biologia (Lahore) 23, 117-124.
- BHATNAGAR P. and TILLBERG E., 1982: Uptake of radioactivity by *Lemna gibba* from 1-¹⁴C-indole-3-acetic acid applied to the medium. Physiol. Plant. 54(2), 183-188.
- BHATTA H.L., SASTRY C.A. and NIGAM R.K., 1973: Control of aquatic weeds from water bodies using grass carp. Indian J.Environ.Health 15, 92-99.
- BIANCHETTI R., 1963: Azione della luce sull'assorbimento, l'utilizzazione ed il trasporto dei glucidi. G.Bot.Ital. 70, 329-337.
- BIEDLINGMAIER S., KOEST H.-P. and SCHMIDT A., 1986: Utilization of sulfonic acids as the only sulfur source for growth of photosynthetic organisms. Planta 169(4), 518-523.
- BIELECKI K. and SKRABKA H., 1976: Effect of some herbicides on photosynthesis of *Spirodela polyrrhiza*. (In Polish). Acta Agrobot. 29(1), 59-68.
- BIELECKI K., STACHURSKA A., and GRZYS E., 1981: Effect of the detergent Tween 20 on the phytotoxicity of linuron. (In Polish). Zesz.Nauk. Akad.Roln.Wroclawin 131, 99-108.
- BIELENSKI U., GARREC J.P., DEMARTY M., RIPOLL C. and THELLIER M., 1984a: Kinetic parameters in the compartmental analysis of lithium transport in *Lemna gibba* using the stable isotopes, ⁶Li and ⁷Li, as tracers. Physiol.Plant. 61(2), 236-242.
- BIELENSKI U., RIPOLL C., DEMARTY M., LUETTGE U. and THELLIER M., 1984b: Estimation of cellular parameters in the compartmental analysis of Li transport in *Lemna gibba* using the stable isotopes, ⁶Li and ⁷Li, as tracers. Physiol.Plant. 62(1), 32-38.
- BIELENSKI R.L., 1966a: Levels of phosphate in *Spirodela*. Plant Physiol. 43, 1297-1308.
- BIELENSKI R.L., 1966b: Effect of phosphorus deficiency on levels of phosphorus compounds in *Spirodela*. Plant Physiol. 43, 1309-1316.
- BIELENSKI R.L., 1971: Enzyme changes in plants following changes in their mineral nutrition. Proc.6th Int.Coll.Plant Analysis and Fertilizer Problems, Tel Aviv, 1970, 1, 143-153.
- BIELENSKI R.L., 1972: Turnover of phospholipids in normal and phosphorus-deficient *Spirodela*. Plant Physiol. 49(5), 740-745.
- BIELENSKI R.L., 1974: Development of an externally located alkaline phosphatase as a response to phosphorus deficiency. In: BIELENSKI R.L.,

- FERGUSON A.R. and CRESSWELL M.M. (eds.), Mechanisms of regulation of plant growth. Bull. 12, The Royal Soc. of New Zealand, Wellington, 165-170.
- BIELESKI R.L. and JOHNSON P.N., 1972: The external location of phosphate activity in phosphorus-deficient *Spirodela oligorrhiza*. Aust. J. Biol. Sci. 25(4), 707-720.
- BIERHUIZEN J.F., 1954: Observations on potassium deficiency in *Lemna minor* L. Meded. Landbouwhogeschool Wageningen 54, 311-319.
- BIERHACH W.N. and WINOGRADOV A.P., 1931: Elementary structure of *Lemna*; determination of species. (In Russian). DAN SSR 6.
- BIGGS R.H., 1983: Studies of ultraviolet-B radiation on growth and development of *Lemna minor*. Plant Physiol. 72(suppl.), 163.
- BIGGS R.H., 1984: Ultraviolet-B radiation alters proteins and asexual and sexual reproduction of *Lemna minor*. Plant Physiol. 75(suppl.), 135.
- BIGGS R.H. and KOSSUTH S.V., 1980: Developmental effects of Sandoz 6706 and ultraviolet-B radiation on *Spirodela punctata* and *Lemna minor*. Plant Physiol. 65(suppl.), 83.
- BIGGS R.H. and KOSSUTH S.V., 1985: Stress proteins in *Lemna minor*. Plant Physiol. 77(4 suppl.), 154.
- BINOT R.A., NAVEAU H.P. and NYNS E.-J., 1981: Methanogenic potential activity of mixed liquors (fluorometric monitoring). Biotechnol. Lett. 3(11), 623-628.
- BIRMINGHAM B.C. and COLMAN B., 1983: Potential phytotoxicity of diquat accumulated by aquatic plants and sediments. Water, Air, Soil Pollut. 19(2), 123-131.
- BISCOE T.D., 1973: The winter state of our duckweeds. Am. Nat. 7, 257-268.
- BISHOP W.E. and PERRY R.L., 1981: Development and evaluation of a flow-through growth inhibition test with duckweed (*Lemna minor*). ASTM Spec. Tech. Publ. 737, 421-435.
- BITCOVER E.H. and SIELING D.H., 1951: Effect of various factors on the utilization of nitrogen and iron by *Spirodela polyrrhiza* (L.) Schleid. Plant Physiol. 26, 290-303.
- BITTING L.E., Sr., 1974: Field uses of invert emulsions. Hyacinth Control J. 34-35.
- BJØRNDAL G., 1984: Growth performance, nutrient uptake and human utilization of duckweeds (Lemnaceae family). The Phytotron, Univ. Oslo and Agric. Res. Council, Norway, 102 pp.
- BJØRNDAL G. and NILSEN S., 1985: Growth potential of *Lemna gibba*: effect of CO₂ enrichment at high photon flux rate. Aquat. Bot. 22(1), 79-82.
- BLACK J.M., 1978: Lemnaceae. In: JESSOP J.P. (ed.), Flora of South Australia. (3rd ed.). Adelaide. 304-306.
- BLACKBURN K.B., 1933: Notes on chromosomes of the duckweeds (Lemnaceae) introducing the question of chromosome size. Proc. Univ. Durham Phil. Soc. 9, 84-90.
- BLACKBURN R.D., 1976: Effective control of an aquatic weed nuisance. Publ. Wks. 107(5), 58-60.
- BLACKBURN R.D. and SUTTON D.L., 1971: Growth of the white amur (Ctenopharyngodon idella Val.) on selected species of aquatic plants. Proc. Eur. Weed Res. Council 3, 87-93.
- BLACKBURN R.D. and WELDON L.W., 1965: The sensitivity of duckweeds (Lemnaceae) and Azolla to diquat and paraquat. Weeds 13, 147-149.
- BLACKMAN G.E., 1952: Studies in the principles of phytotoxicity. I. The assessment of relative toxicity. J. Exp. Bot. 3, 1-27.
- BLACKMAN G.E., 1956: Interrelationships between the uptake of 2:4-dichlorophenoxyacetic acid, growth, and ion absorption. In: WAIN R.L. and WIGHTMAN F. (eds.), The chemistry and mode action of plant growth substances. Butterworth, London. 253-259.
- BLACKMAN G.E., PARKE M.H. and GARTON G., 1954: The physiological activity of substituted phenols. Arch. Biochem. Biophys. 54, 45-71.
- BLACKMAN G.E. and PRASAD R., 1962: The influence of light intensity and temperature on the phytotoxicity of dalapon to *Lemna minor* and *Salvinia natans*. Res. Prog. Rept. Western Weed Control Conf., 91-92.
- BLACKMAN G.E. and ROBERTSON-CUNINGHAM R.C., 1953: The influence of pH on the phytotoxicity of 2:4 dichlorophenoxyacetic acid to *Lemna minor*. New Phytol. 52, 71-76.
- BLACKMAN G.E. and ROBERTSON-CUNINGHAM R.C., 1954: Interactions in the physiological effects of growth substances on plant development. J. Exp. Bot. 5, 184-203.
- BLACKMAN G.E. and ROBERTSON-CUNINGHAM R.C., 1955: Interrelationship between light intensity, temperature, and the physiological effects of 2:4-dichlorophenoxyacetic acid on the growth of *Lemna minor*. J. Exp. Bot. 6, 156-176.
- BLACKMAN G.E. and SARGENT J.A., 1959: The uptake of growth substances. II. The absorption and accumulation of 2:3:5-triiodobenzoic acid by the root and frond of *Lemna minor*. J. Exp. Bot. 10, 480-503.
- BLACKMAN G.E., SENN G., BIRCH W.R. and POWELL R.G., 1959: The uptake of growth substances. I. Factors controlling the uptake of phenoxyacetic acids by *Lemna minor*. J. Exp. Bot. 10, 33-54.
- BLACKWOOD G.C. and LEAVER C.J., 1977: The effect of light on protein synthesis in green leaves. Colloq. Int. C.N.R.S. No. 261: Acides nucléiques et synthèse des protéines chez les végétaux. 611-615.
- BLAKE C.H., 1938: *Wolffia floridana* in Massachusetts. Rhodora 40, 76.
- BLAKE G. and DUBOIS J.P., 1979: Rôle des macrophytes aquatiques dans l'épuration, synthèse bibliographique. Rapp. Ministère d'Environnement du Cadre de Vie. 91 pp.
- BLAKE G. and DUBOIS J.P., 1981: Water epuration: role of aquatic macrophytes in elimination mineral elements. Int. Coll. Aquat. Water Plants, Brussels, 1981, Abstr., 27.
- BLAKE G. and DUBOIS J.P., 1982: Epuration des eaux: rôles des macrophytes aquatiques dans l'élimination des éléments minéraux. In: SYMOENS J.J., HOOPER P. and COMPÈRE P. (eds.), Studies on aquatic plants. R. Bot. Soc. Belg. 315-323.
- BLAKE S.F., 1933: *Wolffia lingulata* in Louisiana. Rhodora 35, 226.
- BLAKE S.F., 1952: *Wolffia papulifera* in Texas. Rhodora 54, 306-307.
- BLAYTER E. and HALLBERG C., 1921: Species novae Indiae Orientalis. J. Ind. Bot. 2, 49-50.
- BLAZEY E.B. and MCCLURE J.W., 1968: The distribution and taxonomic significance of lignin in the Lemnaceae. Am. J. Bot. 55(10), 1240-1245.
- BLODGETT F.H., 1914: Development of the embryo and the germination in *Lemna perpusilla*. Science 39, 292.
- BLODGETT F.H., 1915: Morphology of the *Lemna* frond. Bot. Gaz. 60, 383-390.
- BLODGETT F.H., 1923: The embryo of *Lemna*. Am. J. Bot. 10, 336-342.
- BLUETHGEN J., 1964: Allgemeine Klimogeographie. De Gruyter, Berlin. 599 pp.
- BOCCHI A., 1981: Effetti del fitato di sodio su *Lemna minor*. I. Allungamento della radice in assenza di sali minerali nel mezzo di coltura. Ann. Fac. Med. Vet. Univ. Parma 1, 113-118.
- BOCCHI A., 1982: Effetti del fitato di sodio su *Lemna minor*. II. Influenza

- enza sul ciclo di sviluppo vegetativo e sulle caratteristiche delle colonie. Ann. Fac. Med. Vet. Univ. Parma 2, 71-80.
- BACION P.F., CATTANACH C.J., EGGENBERG P., GRESSEL J., HAGMANN M.-L., MALKIN S. and WENGER J., 1987: Synthesis and characterization of a group of dihydropyrimidinobenzimidazole photosystem II herbicides. *Pestic. Biochem. Physiol.* 28(1), 75-84.
- BOECHER M. and NOVACKY A., 1978: The active component of membrane potential of Lemna gibba effect of fusicoccin. *Plant Physiol.* 61(4 suppl.), 107.
- BOECHER M. and NOVACKY A., 1981: Effect of tentoxin on the membrane potential of Lemna gibba Gl. *Plant Sci. Lett.* 23(3), 269-276.
- BOECHER M., FISCHER E., ULLRICH-EBERUS C. and NOVACKY A., 1980: Effect of fusicoccin on the membrane potential, on the uptake of glucose and glycine, and on the ATP level in Lemna gibba Gl. *Plant Sci. Lett.* 18, 215-220.
- BOECKER E., 1936: Ueber eine Lemna mit regelmässig homodromer und antidromer Verzweigung. *Flora* 130, 438-440.
- BOESZÖRMENYI E., 1955: Contributions to the mineral nutrition of Lemna minor. (In Hung.). Ph.D. Thesis, Univ. Budapest.
- BOESZÖRMENYI E. and BOESZÖRMENYI Z., 1957: N and P nutrition and the physiological age of Lemna minor L. *Acta Bot. Acad. Sci. Hung.* 3, 1-7.
- BOGACKA T. and GROBA J., 1980: Toxicity and biodegradation of chlorofeninfos, carbaryl, and propoxur in an aquatic environment. (In Polish). *Bromatol. Chem. Toksykol.* 13(2), 151-158.
- BOGACKA T., WIKTOR J. and GROBA J., 1983: Toxicity and biodegradation of selected pesticides in water environment. (In Polish). *Bromatol. Chem. Toksykol.* 16(2), 145-159.
- BOGNER J., 1985: Wolffia welwitschii und Wolffia denticulata (Lemnaceae). *Aqua-Planta* 10(3), 3-5.
- BOLIGANO R.W., 1979: Effects of various changes in the nutrient medium on growth and potassium accumulation in Lemna. M.S. Thesis. James Madison Univ., Harrisonburg, Virginia.
- BOLLARD E.G., 1966: A comparative study of the ability of organic nitrogenous compounds to serve as sole sources of nitrogen for the growth of plants. *Plant and Soil* 25, 153-166.
- BOLLARD E.G. and COOK A.R., 1968: Regulation of urease in a higher plant. *Life Sci.* Part II, 7(20), 1091-1094.
- BOLLARD E.G., COOK A.R. and TURNER N.A., 1968: Urea as a sole source of nitrogen for plant growth. I. The development of urease activity in Spirodela oligorrhiza. *Planta* 83, 1-12.
- BOLLMANN O., STROTHER S. and HOFFMANN-OSTENHOF O., 1980: Studies on the biosynthesis of cyclitols. XL. The enzymes involved in the synthesis of phytic acid in Lemna gibba. *Mol. Cell Biochem.* 30, 171-175.
- BONEY S.E., 1977: Chemical and physical characterization of a two-stage animal waste lagoon supporting aquatic plants. M.S. Thesis. Louisiana State Univ., Baton Rouge, La.
- BONG C.L., COLE A.L.J., WALKER J.R.L. and PETERS J.A., 1979: Effect of sodium fluoroacetate ("Compound 1080") on the soil microflora. *Soil Biol. Biochem.* 11(1), 13-18.
- BONG C.L., WALKER J.R.L. and PETERS J.A., 1980: The effect of fluoroacetate ("Compound 1080") and fluoride upon duckweeds. *N.Z.J. Sci.* 23, 179-183.
- BONOMI A., QUARANTELLI A., SUPPERCHI P., SABBIONI A. and MALAVASI P., 1981: Contributo alla conoscenza della composizione chimico-bromatologica del giacinto d'acqua (Eichhornia crassipes) e della lenticchia d'acqua (Lemna minor) coltivati su reflui di porcellaia parzialmente trattati. *Ann. Fac. Med. Vet. Univ. Parma*, 1, 198-216.

- BORESCH K., 1935/36: Gehalt der Pflanzen an Mineralstoffen. *Tabulae Biologicae* 10, 315-353 (1935) and 11, 136-191 (1936).
- BORNKAMM R., 1964: Zur Oxalatsynthese von Lemna minor L. unter verschiedenen Anzuchtbedingungen. Vorläufige Mitteilung. *Ber. Deutsch. Bot. Ges.* 77, Sondernummer (187)-(193).
- BORNKAMM R., 1965: Die Rolle des Oxalats im Stoffwechsel höherer grüner Pflanzen. Untersuchungen an Lemna minor L. *Flora* A 156, 139-171.
- BORNKAMM R., 1966a: Die Oxalsäure im Stoffwechsel höherer Pflanzen. *Umschau* 1966, 262.
- BORNKAMM R., 1966b: Ein Jahresrhythmus des Wachstums bei Lemna minor L. *Planta* 69, 178-186.
- BORNKAMM R., 1969a: Typen des Oxalatstoffwechsels grüner Blätter bei einigen Familien höherer Pflanzen. *Flora* A 160, 317-336.
- BORNKAMM R., 1969b: Interactions and their influence on protein content of plants. 11th Int. Bot. Congr. and Int. Wood Chemistry Symp. 1969, Seattle. Abstr., 18.
- BORNKAMM R., 1970a: Dunkel-Assimilation von Nitrat bei Lemna minor L. *Planta* 92, 50-56.
- BORNKAMM R., 1970b: Ueber den Einfluss der Konkurrenz auf die Substanzproduktion und den N-Gehalt der Wettbewerbspartner. *Flora* A 159, 84-104.
- BORNKAMM R., 1970c: Ueber den Oxalat- und Stickstoffgehalt von Pflanzen aus verschiedenen Höhenstufen. *Oecol. Plant.* 5, 335-343.
- BORSTIAP A.C., 1970: Antagonistic effects of branched chain amino acids on the growth of Spirodela polyrrhiza (L.) Schleiden. *Acta Bot. Neerl.* 19(2), 211-215.
- BORSTIAP A.C., 1972: Changes in the free amino acids of Spirodela polyrrhiza (L.) Schleiden during growth inhibition by L-valine, L-isoleucine, or L-leucine. A gas chromatographic study. *Acta Bot. Neerl.* 21, 404-416.
- BORSTIAP A.C., 1974: Antagonisms between amino acids in the growth of Spirodela polyrrhiza due to competitive amino acid uptake. *Acta Bot. Neerl.* 23, 723-738.
- BORSTIAP A.C., 1975: Metabolic fate of exogenously supplied branched-chain amino acids in Spirodela polyrrhiza. *Acta Bot. Neerl.* 24(2), 203-206.
- BORSTIAP A.C., 1977a: Kinetics of the uptake of some neutral amino acids by Spirodela polyrrhiza. *Acta Bot. Neerl.* 26, 115-128.
- BORSTIAP A.C., 1977b: Effects of branched-chain amino acids on the growth of Spirodela polyrrhiza. Ph.D. Thesis. Univ. Utrecht. 116 pp.
- BORSTIAP A.C., 1978: Glycine and L-alanine as antagonists of growth inhibition by the branched-chain amino acids in Spirodela polyrrhiza. *J. Exp. Bot.* 29(110), 709-718.
- BORSTIAP A.C., 1981: Interactions between the branched-chain amino acids in the growth of Spirodela polyrrhiza. *Planta* 151, 314-319.
- BORSTIAP A.C. and VERNHOOG-GERRITSEN M., 1985: In-vivo control of valine and leucine synthesis in the duckweed Spirodela polyrrhiza. *Planta* 164(1), 129-134.
- BORSTIAP A.C., MEERNIS J.L.D., VAN ECK W.F. and BICKER J.T.E., 1986: Kinetics and specificity of amino acid uptake by the duckweed Spirodela polyrrhiza (L.) Schleiden. *J. Exp. Bot.* 37(180), 1020-1035.
- BOSS M.L., DIJKMAN M.J. and RUSSELL E., 1963b: Standardized culture of some Lemnaceae. *Quart. J. Florida Acad. Sci.* 26(4), 335-346.
- BOSS M.L., DIJKMAN M.J. and THEILING V.B., 1963a: Aging studies on Lemna. *Quart. J. Florida Acad. Sci.* 26, 175-183.
- BOTTOMLEY W.B., 1914: Some accessory factors in plant growth and nutrition. *Proc. R. Soc. B* 87, 237-247.

- BOTTOMLEY W.B., 1917: Some effects of organic growth promoting substances (auxinones) on the growth of Lemna minor in mineral culture solutions. *Proc.R.Soc. B* 89, 481-507.
- BOTTOMLEY W.B., 1919: The effect of nitrogen-fixing organisms and nucleic acid derivatives on plant growth. *Proc.R.Soc. B* 91, 83-95.
- BOTTOMLEY W.B., 1920a: The growth of Lemna plants in mineral solution and in their natural media. *Ann.Bot.* 34, 345-352.
- BOTTOMLEY W.B., 1920b: The effect of organic matter on the growth of various water plants in culture solution. *Ann.Bot.* 34, 353-367.
- BOUDET A., HUMPHREY T.J. and DAVIES D.B., 1975: The measurement of protein turnover by density labelling. *Biochem.J.* 152, 409-416.
- BOULGER G.S., 1881: The ancestry of duckweeds. (Reprint of an unknown journal: Gb. 17.9.1981). 1 p.
- BOUR W.S., 1932: Ecological and physiological studies on certain aquatic angiosperms. *Contrib.Boyce Thompson Inst.* 4(4), 425-496.
- BOUTIN C., 1983: Les macrophytes, leur rôle dans l'épuration des eaux usées et leur insertion dans les différents systèmes de traitement; étude sur site réel de l'exportation d'éléments nutritifs par les lentilles d'eau. *Rep. of ENSP to CEMAGREF*. (Polycopy).
- BOWEN H.J.M., 1958a: Variation in Lemna gibba. *Proc.Bot.Soc.Brit.* 13, 86-87.
- BOWEN H.J.M., 1958b: Two American species of Lemna. *Proc.Bot.Soc.Brit.* 13, 87.
- BOWKER D.W., 1978: Interactions of aquatic macrophytes and epiphytic algae in freshwater ecosystems. *Ph.D.Thesis*. Univ. of London.
- BOWKER D.W. and DENNY P., 1980: The seasonal succession and distribution of epiphytic algae in the phyllosphere of Lemna minor L. *Arch.Hydrobiol.* 90, 39-55.
- BOWKER D.W. and DUFFIELD A.N., 1981: Population growth of Lemna gibba L. in the Gwent levels drainage channels. *Proc.Conf.Aquat.Weeds and their Contr.* Nat.Veg.Res.Stat., Wellesbourne. 67-76.
- BOWKER D.W., DUFFIELD A.N. and DENNY P., 1980: Methods for the isolation, sterilization and cultivation of Lemnaceae. *Freshw.Biol.* 10, 385-388.
- BOWMAN C. and DYER T.A., 1979: 4.5S ribonucleic acid, a novel ribosome component in the chloroplasts of flowering plants. *Biochem.J.* 183, 605-614.
- BOYD C.E., 1968: Fresh-water plants: a potential source of protein. *Econ.Bot.* 22, 359-368.
- BOYD C.E., 1975: Competition for light by aquatic plants. *Circ.Agric. Exp.Stat.* Auburn Univ. 215, 19 pp.
- BOYD C.E. and SCARSBROOK E., 1975: Chemical composition of aquatic weeds. In: BREGONTIK P.L. and FOX J.L. (eds.), *Proc.Symp.Water quality management through biological control*, Gainesville. 144-150.
- BRADEN D.A. and CIALONE J.C., 1971: Toxicity of paraquat-treated plant tissue to Lemna minor L. *Proc.Northeastern Weed Sci.Soc. Abstr.* 25, 220.
- BRADLEY G.H., 1932: Some factors associated with the breeding of Anopheles mosquitoes. *J.Agric.Res.* 44, 381-399.
- BRAENDLE R., 1978: Downward transport of sulfur compounds from primary bean leaves (Phaseolus vulgaris) following gassing with sulfur-35 labelled hydrogen sulfide. *Translation 23884 of Al Ahram Center*. 9 pp.
- BRAENDLE R. and ERISMANN K.H., 1968: Photosynthese-abhängige Sulfidaufnahme grüner Pflanzen. *Naturwissenschaften* 55, 41.
- BRAENDLE R., STOECKLI B. and ERISMANN K.H., 1975: Beeinflussung der Thymidinphosphorylierung und Inkorporation durch Schwefeldioxid und Sul-

- fit bei der Wasserlinse (Lemna minor L.). *Experientia* 31, 511-513.
- BRAENDLE R., STRASSER R. and ERISMANN K.H., 1968: Intermediärprodukte der H₂S-Assimilation bei Lemna minor L. *Verh.Schweiz.Naturf.Ges.* 148, 122-124.
- BRAUN-BLANQUET J. and RUEBEL E., 1932-1935: Flora von Graubünden. *Veröff.Geobot.Inst.Rübel,Zürich* 7, 1695 pp.
- BRAUN S.A.M., 1986: Abscissinsäure und Indollessigsäure in Wasserpflanzen und deren Beziehung zum Wachstum unter natürlichen Umweltbedingungen. *Ph.D.Thesis*. TU München. 130 pp.
- BRAVO H., 1930: Las Lemnaceas del valle de Mexico. *An.Inst.Biol.(Mexico)* 1, 7-37.
- BRECHT E., 1986: The light-harvesting chlorophyll a/b-protein complex II of the higher plants. Results from a twenty-year research period. *Photobiochem.Photobiophys.* 12(1-2), 37-50.
- BREHM V., 1935: Die Organismenwelt der Wasserlinsenteppiche. *Mikrokosmos* 28, 109.
- BREZNY O. and SHARMA R.K., 1973: Studies on evapotranspiration of some aquatic weeds. *Weed Sci.* 21(3), 197-204.
- BRISTOW J.M., CARDENAS J., FULLERTON T.M. and SIERRA J., 1972: Aquatic weeds. *Columbian Agr.Inst.* and the Intern.Plant Protection Center, Oregon State Univ., Agency for Intern. Development.
- BROCK M.A. and LANE J.A.K., 1983: The aquatic macrophyte flora of saline wetlands in Western Australia in relation to salinity and permanence. *Hydrobiologia* 105, 63-76.
- BRONGNIART A., 1833: Notes sur la structure du fruit des Lemna. *Arch.Bot.* 2, 97-104.
- BROOKS J.S., 1940: The cytology and morphology of the Lemnaceae. *Ph.D. Thesis*. Cornell University.
- BROOKS R.E., 1981: Wolfia, new record for Colorado, USA. *Southwest Nat.* 26(3), 321.
- BROOKS R.E. and HAUSER L.A., 1978: Aquatic vascular plants in Kansas. I. Submersed and floating leaved plants. *Tech.Publ.State Biol.Surv.Kansas* 7, 70 pp.
- BROOKS R.E. and MCGREGOR R.L., 1979: New records and notes on the vascular flora of Kansas for 1978. *Tech.Publ.State Biol.Surv.Kansas* 8, 87-92.
- BROUARD F., BORIES A. and SAUZE F., 1983: Advance in anaerobic digestion of aquatic plants. (In French). *Comm.Eur.Communities EUR 8245*, 334-338.
- BROWN B.T. and RATTIGAN B.M., 1979: Toxicity of soluble copper and other metal ions to Elodea canadensis. *Environ.Pollut.* 20, 303-314.
- BRUCE B.D. and MALKIN R., 1987: Composition and function of photosystem I in Lemna. *Symp. on plant membranes. J.Cell Biochem. (suppl. 11 part B)*, 83.
- BRUMFIELD B.M., EVANS D.K. and BRANT A.E., 1982: Additions to the wetland flora of West Virginia. *Castanea* 47(2), 179-181.
- BRUNAUD A., 1974 s. 1974c
- BRUNAUD A., 1974a: Organisation de la pousse végétative d'une Lemnacee: Spirodela polyrrhiza (L.) Schleid. *C.R.Acad.Sci.Paris*, D 278, 1183-1186.
- BRUNAUD A., 1974b: Organisation végétative chez les Lemna minor L. et Wolffia arthiza (L.) Wimm. *C.R.Acad.Sci.Paris*, D 278(16), 2019-2022.
- BRUNAUD A., 1974c: Organisation de la pousse fertile chez le Lemna minor. *C.R.Acad.Sci.Paris*, D 278(23), 2913-2916.
- BRUNEL J.F., HIEPKO P. and SCHOLZ H., 1984: Flore analytique du Togo. *Phanérogames. Englera* 4, 751 pp.

- BRUNK D.G. and RHODES D., 1987: Effects of aminoxyacetate on amino acid metabolism of Lemna minor L. Plant Physiol. 83(4 suppl.), 24.
- BRUNOLD C., 1972: Regulation der Sulfataufnahme und der Sulfatassimilation durch Schwefelwasserstoff. Diss.Univ. Bern.
- BRUNOLD C. and ERISMANN K.H., 1968: Der Mechanismus der Photosynthesehemmung durch Acetat bei Lemna minor. Verh.Schweiz.Naturf.Ges. 148, 124-125.
- BRUNOLD C. and ERISMANN K.H., 1970: Die Hemmung der Sulfataufnahme durch H₂S bei Lemna minor. Verh.Schweiz.Naturf.Ges. 150, 238-239.
- BRUNOLD C. and ERISMANN K.H., 1972: H₂S als Schwefelquelle bei Lemna minor L. Verh.Schweiz.Naturf.Ges. 152, 145-147.
- BRUNOLD C. and ERISMANN K.H., 1974: H₂S als Schwefelquelle bei Lemna minor L.: Einfluss auf das Wachstum, den Schwefelgehalt und die Sulfataufnahme. Experientia 30, 465-467.
- BRUNOLD C. and ERISMANN K.H., 1975: H₂S as sulfur source in Lemna minor L. II. Direct incorporation into cysteine and inhibition of sulfate assimilation. Experientia 31, 508-509.
- BRUNOLD C. and ERISMANN K.H., 1976: Sulfur dioxide as a sulfur source in duckweeds (Lemna minor L.). Experientia 32, 296-297.
- BRUNOLD C. and SCHMIDT A., 1976: Regulation of adenosine 5'-phosphosulfate sulfotransferase activity by H₂S in Lemna minor L. Planta 133, 85-88.
- BRUNOLD C. and SCHMIDT A., 1978: Regulation of sulfate assimilation in plants. 7. Cysteine inactivation of adenosine 5'-phosphosulfate sulfotransferase in Lemna minor L. Plant Physiol. 61, 342-347.
- BRUNOLD C. and SUTER M., 1983: Aktivitätsmessung der Adenosin 5'-phosphosulfat-Sulfotransferase und ihre Anwendung bei der Untersuchung der de novo-Synthese des Enzyms. Bot.Helv. 93, 105-114.
- BRUNOLD C. and SUTER M., 1984a: Effect of sulfate concentration on nitrate and sulfate assimilating enzymes of Lemna minor. Plant Physiol. 75(suppl.1), 198.
- BRUNOLD C. and SUTER M., 1984b: Regulation of sulfate assimilation by nitrogen nutrition in the duckweed Lemna minor L. Plant Physiol. 76, 579-583.
- BRUNOLD C. and SUTER M., 1986: Regulation of adenosine 5'-phosphosulfate sulfotransferase activity of Lemna minor by sulfate. Experientia 42 (6), 712.
- BRUNOLD C., SUTER M. and LAVANCHY P., 1987: Effect of high and low sulfate concentrations on adenosine 5'-phosphosulfate sulfotransferase activity from Lemna minor. Physiol.Plant. 70(2), 168-174.
- BUCKINGHAM G.R., HAAG K.H. and HABECK D.H., 1986: Native insects enemies of aquatic macrophytes. Beetles. Aquatics 8, 28-34.
- BUCKLE D. and RALPH R.K., 1977: Control of growth of Spirodela oligorrhiza in darkness. Plant Sci.Lett. 9, 347-350.
- BUCEK J., 1984a: The occurrence of a nitrate reductase inactivating factor in extracts of Spirodela polyrrhiza. Acta Soc.Bot.Pol. 53(3), 411-417.
- BUCEK J., 1984b: The effect of ionic and non-ionic surfactants on the growth, nitrate reductase and nitrite reductase activities of Spirodela polyrrhiza (L.) Schleiden. Acta Soc.Bot.Pol. 53(4), 551-559.
- BUCEK J. and SULEJ J., 1986: Density gradient localization of vanadate- and NO₂-sensitive ATPase from sterile cultures of Spirodela polyrrhiza (L.) Schleiden. Acta Soc.Bot.Pol. 55, 253-262.
- BUDDHARI W., VIRABALIN R. and AIKAPHON K., 1984: Effects of external lead concentration on the uptake and distribution of lead in plants. In: THYAGARAJAN G. (ed.), Proc.Int.Conf. on Water Hyacinth. Nairobi, Kenya, UN Environ.Progr. 379-385.

- BUGNON F., 1974: Sur un type d'organisation morphologique des pousées fertiles ou de leurs dérivés, particulièrement répandu chez les Monocotylédones. C.R.Acad.Sci.Paris, D 278(12), 1553-1556.
- BUICKMA A.L., Jr., MCGINNIS M.J. and CAIRNS J., Jr., 1979: Phenolics in aquatic ecosystems - a selected review of recent literature. Mar. Environ.Res. 2(2), 87-182.
- BUMBY M.J., 1982: A survey of aquatic macrophytes and chemical qualities of nineteen locations in Costa Rica. Brenesia 19/20, 487-535.
- BURK C.J., LAUBERMANN S.D. and MESROBIAN A.L., 1976: The spread of sexual introduced or recently invading aquatics in Western Massachusetts. Rhodora 78(816), 767-772.
- BURKART A., 1957: The vegetation of the Parana River Delta. Darwiniana 11(3), 457-561.
- BURTON J.H., ISLINGER S. and MUZZAR A.J., 1975: The nutritional value of water plants. J.Anim.Sci. 41(1), 332.
- BURTON T.M., KING D.L. and ERVIN J.L., 1978: Aquatic plant harvesting as a lake restoration technique. Proc.Natl.Conf.on Lake Restoration, Minneapolis, 117-185.
- BUSTINE D.L. and KINDEL P.K., 1969: Biosynthesis of D-Apirose in a cell-free system from Lemna minor L. J.Biol.Chem. 244(5), 1382-1385.
- BUTLER G.W. and PETERSON P.J., 1967: Uptake and metabolism of inorganic forms of selenium-75 by Spirodela oligorrhiza. Austral.J.Biol.Sci. 20(1), 77-86.
- BYTNIWSKA K., 1977: Nitrogen and protein contents in some aquatic plant species. Acta Soc.Bot.Pol. 46(2), 165-172.
- BYTNIWSKA K. and MACIEJSKA-POTAPCZYK W., 1980: Amino acid composition and biological value of proteins in some aquatic plant species. Biochem.Physiol.Pflanz. 175(2), 172-175.
- BYTNIWSKA K. and POTAPCZYK W., 1981: Growth and contents of total and readily soluble protein in aseptic culture of Spirodela polyrrhiza (L.) Schleid. Acta Physiol.Plant. 3(3), 125-133.
- CADÉVAL J., 1933: Flora de Catalunya. Barcelona. 5, 396-398.
- CALDWELL O.W., 1899: On the life history of Lemna minor. Bot.Gaz. 27, 37-66.
- CALLAHAN F.E. and MATTOO A.K., 1987: Characterization of thylakoid protein acylation, comparison of in vivo and in vitro acylated proteins. Plant Physiol. 83(4 suppl.), 113.
- CALLAHAN F.E., MATTOO A.K., NORMAN H.A., ST JOHN J.B., WERGIN W.P., NELSON N. and EDELMANN M., 1986: Dynamics of chloroplast membrane metabolism. J.Cell.Biol. 103(5 part 2), 521A.
- CAMP W.H., 1933: Distribution and flowering in Wolffia papulifera. Ohio J.Sci. 33, 163.
- CANTON J.H., SLOOF W., KOOL H.J., STRUYS J., POOW T.J.M., WEGMAN R.C.C. and PITT G.J., 1985: Toxicity, biodegradability, and accumulation of a number of Cl/N-containing compounds for classification and establishing water quality criteria. Regul.Toxicol.Pharmacol. 5(2), 123-131.
- CAPOZZI A., 1952: Ricerche sul potere fitodinamico di alcuni insetticidi sistemici fosfororganici. Atti Ist.Bot.Univ.Lab.Crittogam.(Pavia), ser. 5, 10, 117-124.
- CARNEY K.J. and EBINGER J.E., 1982: Aquatic macrophytes in the Shelbyville moraine area of East-Central Illinois. Trans.Ill.Acad.Sci. 75 (1-2), 143-147.
- CASPERSON G., 1956: Wärmehaushaltstudien an Wasserpflanzen. Ber.Deutsch. Bot.Ges. 69, 479-486.

- CASSANI J.R., 1981: Feeding behaviour of underyearling hybrids of the grass carp, *Ctenopharyngodon idella*, female, and the bighead, *Hypophthalmichthys nobilis*, male, on selected species of aquatic plants. *J. Fish Biol.* 18(2), 127-133.
- CASSANI J.R. and CATON W.E., 1983: Feeding behaviour of yearling and older hybrid grass carp. *J. Fish Biol.* 22(1), 35-42.
- CASSANI J.R. and CATON W.E., 1986: Growth comparisons of diploid and triploid grass carp under varying conditions. *Progressive Fish-Culturist* 48(3), 184-187.
- CASSANI J.R., CATON W.E. and HANSEN T.H., Jr., 1982: Culture and diet of hybrid grass carp fingerlings. *J. Aquatic Plant Manage.* 20, 30-32.
- CAUSSE H., 1976: Etude du renouvellement des protéines végétales: mise au point et application d'une méthode de mesure de la vitesse de dégradation des protéines chez *Lemna minor* L. Ph.D. Thesis. Univ. Paul Sabatier, Toulouse. 63 pp.
- CAUX P.-Y., WEINBERGER P. and CARLISLE D.B., 1986: Dowanol, an environmentally safe adjuvant. *Environ. Toxicol. Chem.* 5(12), 1047-1054.
- CAYOETTE J., BERNARD J.-P., ROY C. and DUBE M., 1983: Plantes vasculaires nouvelles pour le Québec: additions, échappées de culture et éphémérophytes. *Nat. Can.* 110(3), 293-312.
- CEDENO-NALDONADO A. and LIU L.C., 1976: Effect of two substituted urea and two s-triazine type herbicides on the photosynthesis of *Lemna perpusilla* Torr. *J. Agric. Univ. Puerto Rico* 60, 369-374.
- CEDENO-NALDONADO A., LIU L.C. and DELGADO I., 1979: Effect of photosynthetic inhibitor herbicides on nitrate reductase activity of non-target species. *J. Agric. Univ. Puerto Rico* 63, 412-414.
- CELINSKI F., 1954: The smallest flowering plant, *Wolffia arrhiza* (L.) Wimm. (In Polish). *Wzschwat* 3(4), 75-79.
- CEMAGREF - Division qualité des eaux, pêche et pisciculture, 1982: Biomasse dans les lagunes d'épuration: biomasse végétale récoltable dans les lagunes d'épuration de Chaucenne (Doubs). Rep. COMES. 21 pp. (Polycopied; cited from HUBAC et al. 1984).
- CERVINKOVA H., 1976: Investigating of triazine residues by bioassays (abstract). *Sbornik Vedeckých Prací ze 6. Československé Konference o Ochrane Rostlin, Československá. Abstr.* 227-229.
- ČESKA A. and ČESKA O., 1979: Additions to the flora of British Columbia. *Can. Field-Nat.* 94, 69-74.
- CHABRECK R.H. and PALMISANO A.W., 1973: The effects of hurricane camille on the marshes of the Mississippi River Delta. *Ecology* 54, 1118-1123.
- CHAMP M.A., DAVIS F.S. and WAYLAND J.R., 1973: Ultrahigh-frequency electromagnetic radiation utilized for aquatic vegetation control. *Sport Fishery Abstr.* 19, 17584.
- CHAMURIS G.P. and NIELSEN P.T., 1980: Phytochrome control of root growth in *Lemna* species. *Va. J. Sci.* 31(4), 105.
- CHAMURIS G.P. and NIELSEN P.T., 1983: Phytochrome control of root growth rate in *Lemna* spp. *Am. J. Bot.* 70(6), 827-829.
- CHANG S.-M., YANG C.-C. and SUNG S.-C., 1977: The cultivation and the nutritional value of *Lemnaceae*. *Bull. Inst. Chem. Acad. Sin.* 24, 19-30.
- CHANG S.-M., YANG C.-C. and SUNG S.-C., 1978: The effects of plant hormones on the cultivation and chemical composition of *Lemnaceae*. *Bull. Inst. Chem. Acad. Sin.* 25, 19-34.
- CHANG W.-C. and CHIU P.-L., 1976: Induction of callus from fronds of duckweed (*Lemna gibba* L.). *Bot. Bull. Acad. Sin.* 17, 106-109.
- CHANG W.-C. and CHIU P.-L., 1978: Regeneration of *Lemna gibba* G3 through callus culture. *Z. Pflanzenphys.* 89, 91-94.
- CHANG W.-C. and HSING Y.-I., 1978: Callus formation and regeneration of

- frond-like structures in *Lemna perpusilla* 6746 on a defined medium. *Plant Sci. Lett.* 13, 133-136.
- CHAO T.-P., 1952: A study of phytotoxine action of 2,4-dichlorophenoxyacetic acid on certain aquatic plants. *Diss. Univ. of Michigan.*
- CHAPMAN K.S.R., TREWAVAS A. and VAN LOON L.C., 1975: Regulation of the phosphorylation of chromatin-associated proteins in *Lemna* and *Hordeum*. *Plant Physiol.* 55, 293-296.
- CHARPENTIER S. and GARNIER J., 1985: Etude de la multiplication et de la formation des colonies de *Spirodela polyrrhiza* L. C.R. Acad. Sci. Paris, III, 300(15), 587-590.
- CHARPENTIER S., GARNIER J. and FLAUGNATTI R., 1987: Toxicity and bioaccumulation of cadmium in experimental cultures of duckweed *Lemna polyrrhiza* L. *Bull. Environ. Contam. Toxicol.* 38(6), 1055-1061.
- CHARUDATTAN R. and LIN C.-Y., 1974: Isolates of *Penicillium*, *Aspergillus* and *Trichoderma* toxic to aquatic plants. *Hyacinth Control J.* 12, 70-73.
- CHASSANY-DE CASABIANCA M.-L., 1982a: Production de biomasse macrophytologique flottante et épuration des eaux résiduaires. *Tech. Eau Assainissement* 422, 17-26.
- CHASSANY-DE CASABIANCA M.-L., 1982b: De l'exploitation de la biomasse végétale aquatique aux nouveaux systèmes de production à macrophytes sur eaux résiduaires urbaines. *Eau Ind.* 64, 43-48.
- CHASSANY-DE CASABIANCA M.-L., et SAUZE F., 1981: Cultures de macrophytes sur eaux résiduaires; premières données écologiques; méthanisation des macrophytes aquatiques. *Comm. Eur. Communities*, EUR 7091, 665-677.
- CHATURVEDI R., HAUGSTAD M.K. and NIELSEN S., 1982: The relationship between photosynthetic electron transport and photorespiratory release after DCMU treatment in the duckweed *Lemna gibba*. *Physiol. Plant.* 56(1), 23-27.
- CHATURVEDI R., HAUGSTAD M.K., NIELSEN S. and SKOGEN D., 1985: Photoinhibition of photosynthesis: effect of light and selective excitation of photosystems on recovery. *Photosynthetica* 19(3), 382-387.
- CHESADLE V.I., 1942: The occurrence and types of vessels in the various organs of the plant in the Monocotyledoneae. *Am. J. Bot.* 29, 441-450.
- CHECHENKIN M.N., 1955: The distribution of high saturated fatty acids in the fats of fresh water plants. (In Russian). *Biokhimiya* 20, 249-250.
- CHEN K. and WILDMAN S.G., 1981: Differentiation of fraction I protein in relation to age and distribution of angiosperm groups. *Plant Syst. Evol.* 138(1-2), 89-113.
- CHEN P.K., LEATHER G.R. and KLAYMAN D.L., 1987: Allelopathic effect of artemisinin and its related compounds from *Artemisia annua*. *Plant Physiol.* 83(4 suppl.), 68.
- CHEN S.-J., HSU E.L. and CHEN Y.-L., 1982: Fate of the herbicide benthocarb (thiobencarb) in a rice paddy model ecosystem. *J. Pestic. Sci.* 7(3), 335-340.
- CHEN S.-S.C. and PARK W.-M., 1976: Dual effects of abscisic acid on the growth of a duckweed. *Taiwania* 21, 50-51.
- CHERRY D.S. and GUTHRIE R.K., 1975: Removal of ash basin effluent by the abiotic and biotic parameters of an aquatic drainage system. *ASB Bull.* 22(2), 46.
- CHERRY D.S. and GUTHRIE R.K., 1979: The uptake of chemical elements from coal ash and settling basin effluent by primary producers. II. Relation between concentrations in ash deposits and tissues of grasses growing on the ash. *Sci. Total Environ.* 13(1), 27-31.
- CHERRY D.S., GUTHRIE R.K. and RODGERS J.H., 1978: Cycling of elements in duckweed (*Lemna perpusilla*) in an ash settling basin and swamp drain-

- age system. Water Research 12(10), 765-770.
- CHESTER E.W., 1975: Range extensions and first report for some Tennessee vascular plants. Castanea 40(1), 56-63.
- CHITAPARNI L. and BISJACH M., 1968: On the chemical control of Lemna polytriza and other submerged weeds as a measure of prevention of such other plant infestations as Nelumbo speciosum and Nymphaea alba. Notiz.Mal.Piante 78-79, 141-157.
- CHITNIS P.R., HAREL E., KOHORN B., KARLIN-NEUMANN G., TOBIN E.M. and THORNER J.P., 1985: A model for the folding of the light-harvesting complex. II. Apoprotein of Lemna gibba. J.Cell.Biochem. Suppl.(9 part B), 118.
- CHITNIS P.R., HAREL E., KOHORN B., TOBIN E.M. and THORNER J.P., 1986: Assembly of the precursor and processed light-harvesting chlorophyll a/b protein of Lemna into the light-harvesting complex II of barley etioplasts. J.Cell.Biol. 102(3), 982-988.
- CHITNIS P.R., NECHUSHTAI R., HAREL E. and THORNER J.P., 1987: Some requirements for the insertion of the precursor of apoproteins of Lemna light-harvesting complex II into barley thylakoids. In: BIGGINS J. (ed.), Prog.Photosynth.Res., Proc.7th Int.Congr.Photosynth. Nijhoff, Dordrecht, Netherlands. 4, 573-576.
- CHIU M.M. and FALK R.H., 1975: Ultrastructural study on Lemna perpusilla. Cytologia 40, 313-322.
- CHOI H.O., ANN S.B. and KIM S.N., 1973: Survey of the distribution and amount of weeds emerged in paddy fields in the central part of Korea. Res.Rep.Off.Rural Dev. C 15, 69-75.
- CHOMCHALOW N., 1971: Miracle plant. Agric.Sci.(Thai Agric.Sci.Assoc., Bangkok) 4(4), 319-330.
- CHORNA H.A., 1978: Findings of rare aquatic plants in the Donets River (USSR). (In Russian). Ukr.Bot.Zh. 35(5), 476-478.
- CHORNA H.A., 1979: Flowering of Lemna in water bodies of the northern Donets River valley (USSR). (In Russian). Ukr.Bot.Zh. 36(4), 371-372.
- CHUA S.E. and DICKSON M.H., 1964: The effect of flashing light supplemented by continuous red and far-red light on the growth of Lemna minor L. in the presence of growth regulators. Can.J.Bot. 42, 57-64.
- CHUDYBA H., 1969: Water plants with high protein content. (In Russian). Wszechswiat 10, 253-255.
- CIPPERI R. and CIPPERI F., 1949: Effetto di alcune metilcumarine sul teste moltiplicazione fronde di Lemna. Atti Ist.Bot.Univ.Lab.Critico-gamico,Pavia Ser. 5(3), 322-325.
- CILLIE C.C., 1962: Operation and performance of sewage maturation ponds at Paarl, South Africa. Inst.Sewage Purif.J.Proc. 3, 234-240.
- CLAPHAM A.R., TUTIN T.G. and WARBURG E.F., 1962: Flora of the British Isles. Univ.Press, Cambridge. 1052-1054.
- CLARE P. and EDWARDS R.W., 1983: The macroinvertebrate fauna of the drainage channels of the Gwent Levels, South Wales, UK. Freshwater Biol. 13(3), 205-226.
- CLARK H.L., 1979: Lemnaceae. In: GODFREY R.K. and MOOTEN J.M. (eds.), Aquatic plants of Southeastern United States. Monocotyledons. 464-476.
- CLARK H.L. and THIERT J.W., 1968: The duckweeds of Minnesota. The Michigan Botanist 7, 67-76.
- CLARK J.R., VAN HASSEL J.H., NICHOLSON R.B., CHERRY D.S. and CAIRNS J., Jr., 1981: Accumulation and depuration of metals by duckweed (Lemna perpusilla). Ecotoxicol.Environ.Saf. 5, 87-96.
- CLARK N.A., 1925: The rate of reproduction of Lemna major as a function of intensity and duration of light. J.Phys.Chem. 29, 935-941.

- CLARK N.A., 1926: Plant growth promoting substances, hydrogenic concentration and the reproduction of Lemna. Plant Physiol. 1, 273-279.
- CLARK N.A., 1930: "Auximones" and the stimulation of Lemna by organic matter. Science 71, 268-269.
- CLARK N.A., 1932: Technique for the growth of Lemna under sterile conditions with controlled temperature and light. Iowa State Coll.J.Sci. 7, 13-16.
- CLARK N.A., 1933: Manganese and the growth of Lemna. Plant Physiol. 8, 157-163.
- CLARK N.A. and FLY C.L., 1930: The role of manganese in nutrition of Lemna. Plant Physiol. 5, 241-248.
- CLARK N.A. and FRAHM E.E., 1940a: Influence of auxins on reproduction of Lemna major. Plant Physiol. 15, 735-741.
- CLARK N.A. and FRAHM E.E., 1940b: The effect of various growth promoting substances on the reproduction of Lemna. Proc.Iowa Acad.Sci. 47, 239-247.
- CLARK N.A. and ROLLER E.M., 1931: The stimulation of Lemna major by organic matter under sterile and nonsterile conditions. Soil Sci. 31, 299-309.
- CLARK N.A., THOMAS B.A., 1934: Microorganisms and vitamin production in green plants. Science 79, 571-572.
- CLARK N.A., THOMAS B.A. and FRAHM E.E., 1938: The formation of vitamins A, B₁ and C in Lemna grown in the absence of organic matter. Iowa State Coll.J.Sci. 13, 9-16.
- CLATWORTHY J.N. and HARPER J.L., 1962: The comparative biology of closely related species living in the same area. 5. Inter- and intraspecific interference within cultures of Lemna ssp. and Salvinia natans. J.Exp.Bot. 13, 307-324.
- CLEAUS W.D., 1972: Lifespan and budding potential of Lemna as a function of age of the parent - a genealogic study. New Phytol. 71, 1081-1095.
- CLAVAUD A., 1878: Sur une particularité du Lemna trisulca L. Actes Soc. Linn.Bordeaux 31, 309-311.
- CLELAND C.F., 1967: The physiology of flowering of Lemna gibba L. strain G3. Ph.D.Thesis, Stanford Univ., USA.
- CLELAND C.F., 1970: The use of aphids in the search for the hormonal factors controlling flowering. Plant Physiol. 46(suppl.), 26.
- CLELAND C.F., 1971: Influence of cytokinins on flowering and growth in the long-day plant Lemna gibba G3. Plant Physiol. 47(suppl.), 13.
- CLELAND C.F., 1972a: Investigation into the hormonal control of flowering through the use of aphids. Plant Physiol. 49(suppl.), 62.
- CLELAND C.F., 1972b: The use of aphids in the search for the hormonal factors controlling flowering. In: CARR D.J. (ed.), Plant growth substances. Proc.Symp.,Canberra. 753-757.
- CLELAND C.F., 1974a: Isolation of salicylic acid from aphid honeydew and its effect on flowering. Plant Growth Subst., Proc.Int.Conf. 8th, 1973, Hirokawa, Tokyo. 119-125.
- CLELAND C.F., 1974b: Isolation of flower-inducing and flower-inhibitory factors from aphid honeydew. Plant Physiol. 54, 899-903.
- CLELAND C.F., 1974c: The influence of salicylic acid on flowering and growth in the long-day plant Lemna gibba G3. In: BIELESKI R.L., FERGUSON A.R. and CRESSWELL M.M. (eds.), Mechanisms of regulation of plant growth. The Royal Soc. of New Zealand, Wellington. Bull. 12, 553-557.
- CLELAND C.F., 1976: Synergistic interaction of red light and cytokinin in promotion of growth in Lemna gibba. Plant Physiol. 57(5 suppl.), 75.

- CLELAND C.F., 1977: Influence of photoperiod on induction of flowering by salicylic acid in Lemna. *Plant Physiol.* 59(suppl.), 47.
- CLELAND C.F., 1978: The flowering enigma. *BioScience* 28, 265-269.
- CLELAND C.F., 1979: Comparison of the flowering behavior of the long-day plant Lemna gibba G3 from different laboratories. *Plant Cell Physiol.* 20, 1263-1271.
- CLELAND C.F., 1982: The chemical control of flowering - a status report. In: WAREING P.F. (ed.), *Plant growth substances*. Acad. Press. 635-644.
- CLELAND C.F., 1985: Chemical control of flowering in the long-day plant Lemna gibba G3. *Biol. Plant* 27(4-5), 392-397.
- CLELAND C.F., 1986: Inhibition of flowering in the long-day plant Lemna gibba G3 by Hutner's medium and its reversal by medium modification. *Plant Cell Physiol.* 27(6), 1153-1158.
- CLELAND C.F. and AJAMI A., 1974: Identification of the flower-inducing factor isolated from aphid honeydew as being salicylic acid. *Plant Physiol.* 54, 904-906.
- CLELAND C.F. and BEN-TAL Y., 1982: Influence of giving salicylic acid for different time periods on flowering and growth in the long-day plant Lemna gibba G3. *Plant Physiol.* 70(1), 287-290.
- CLELAND C.F. and BEN-TAL Y., 1983: Hormonal regulation of flowering and sex expression. *Beltville Symp. Agric. Res.* 6, 157-180.
- CLELAND C.F. and BRIGGS W.R., 1967: Flowering responses of the long-day plant Lemna gibba G3. *Plant Physiol.* 42, 1553-1561.
- CLELAND C.F. and BRIGGS W.R., 1968: Effects of low-intensity red and far-red light and high-intensity white light on the flowering response of the long-day plant Lemna gibba G3. *Plant Physiol.* 43, 157-162.
- CLELAND C.F. and BRIGGS W.S., 1969: Gibberellin and CCC effects on flowering and growth in the long-day plant Lemna gibba G3. *Plant Physiol.* 44(4), 503-507.
- CLELAND C.F. and KANG B.G., 1985: Comparison of salicylic acid, benzoic acid, and p-hydroxybenzoic acid for their ability to induce flowering in Lemna gibba G3 and for their uptake and metabolism by Lemna gibba G3. 12th Int. Conf. Plant Growth Substances, Heidelberg, FRG. Abstr., 96.
- CLELAND C.F. and TANAKA O., 1979: Effect of day-length on the ability of salicylic acid to induce flowering in the long-day plant Lemna gibba G3 and the short-day plant Lemna paucicostata 6746. *Plant Physiol.* 64, 421-424.
- CLELAND C.F. and TANAKA O., 1980: Comparison of the ability of salicylic acid and ferricyanide to induce flowering in the long-day plant Lemna gibba G3. *Plant Physiol.* 65(suppl. 6), 95.
- CLELAND C.F. and TANAKA O., 1986: Inhibition of flowering in the long-day plant Lemna gibba G3 by Hutner's medium and its reversal by medium modification. *Plant Cell Physiol.* 27(6), 1153-1158.
- CLELAND C.F. and ZEEVAART J.A.D., 1970: The use of aphids in a search for the flowering stimulus. *Mich. State Univ. MSU/REC, Plant. Res. Lab. Annu. Rep.* 65-66.
- CLELAND C.F., TANAKA O. and FELDMAN L.J., 1982: Influence of plant growth substances and salicylic acid on flowering and growth in the Lemnaceae (duckweeds). *Aquat. Bot.* 13(1), 3-20.
- CLENDENNING K.A. and GORHAM P.R., 1950: Photochemical activity of isolated chloroplasts in relation to their source and previous history. *Can. J. Res. C* 28, 114-139.
- COATES J.B. and DAVIES D.D., 1983: The isolation and examination of protein samples of known age for the study of selectivity of protein de-

- gradation in Lemna minor. *J. Exp. Bot.* 34(146), 1155-1167.
- COCHRAN L.C. and NELSON R., 1933: Observations on the distribution and flowering of certain Lemnaceae in Michigan. *Pap. Michig. Acad. Sci.* 17, 67-68.
- COCUCCI A.E., 1966: Embriologia de Synandropadix vermitoxicus (Ara- ceae). *Kurtziana* 3, 157-181.
- CODY W.J., 1980: Wolffia columbiana (Lemnaceae), water-meal, new to Manitoba. *Can. Field-Nat.* 94(2), 193-194.
- COHEN J.D., SLOVIN J.P. and BIALEK K., 1985: Approaches to unraveling the mysteries of how plants regulate endogenous levels of indole-3-acetic acid. *Current Topics in Plant Biochemistry and Physiology* 4, 75-82.
- COHEN J.D., DALDI B.G. and SLOVIN J.P., 1986: ¹³C₆-(benzene ring)-indole-3-acetic acid. A new internal standard for quantitative mass spectral analysis of indole-3-acetic acid in plants. *Plant Physiol.* 80(1), 14-19.
- COHN F., 1972: Ueber parasitische Algen. *Beitr. Biol. Pflanzen* 1(2), 87-106.
- COLBAUGH P.F., 1981: Pathogenicity of Pythium aphanidermatum on duckweed and water meal. *Phytopathology* 71(8), 867-868.
- COLER R.A. and GUNNER H.B., 1969: The rhizosphere of an aquatic plant (Lemna minor). *Can. J. Mikrob.* 15, 964-969.
- COLER R.A. and GUNNER H.B., 1971: The response of a specialized aquatic ecosystem, the duckweed rhizosphere, to selected environmental influences. *Water Res.* 5, 329-333.
- COLLIER E., LEROITH D., ROTH J. and CLELAND C.F., 1986: Plants contain material that resembles vertebrate somatostatin. *Plant Physiol.* 80 (suppl. 4), 114.
- COLLIER E., WATKINSON A., CLELAND C.F. and ROTH J., 1987: Partial purification and characterization of an insulin-like material from spinach and Lemna gibba G3. *J. Biol. Chem.* 262(13), 6238-6247.
- COLMETRO D.M., 1989: Enumeración y Revisión de las Plantas de la Península Hispano-Lusitana 5, 7-9. Madrid.
- COLT L.C., Jr., HELLIQUIST C.B. and ZUHRIN W.J.L., 1971: An interesting association of rare aquatic plants from New Hampshire. *Rhodora* 73, 296-299.
- CONGDON R.A. and MCCOMB A.J., 1976: The nutrients and plants of Lake Joondalup, a mildly eutrophic lake experiencing large seasonal changes in volume. *J. R. Soc. Western Austr.* 59, 14-23.
- CONN W.M. and LANGWORTHY A.C., 1984: Practical operation of a small scale aquaculture. *Proc. Water Reuse Symp. III, AWWA Res. Found., Denver, Col.* 2, 703-712.
- CONRAD K., 1975: Evidence for the origin of zeatin riboside-like cytokinin in graminaceous tissue by several bioassays. *Biochem. Physiol. Pflanzen* 168(1-4), 341-347.
- COOK A.R., 1968: Urea as sole source of nitrogen for plant growth. II. Urease and the metabolism of urea in Spirodela oligorrhiza. *Planta* 83, 13-19.
- COOK A.R. and BIELESKI R.L., 1969: Fractionation of plant extracts by thin layer electrophoretic and chromatographic procedures. *Anal. Biochem.* 28(1-3), 428-435.
- COOKE R.J. and DAVIES D.D., 1980: General characteristics of normal and stress-enhanced protein degradation in Lemna minor (duckweed). *Biochem. J.* 192, 499-506.
- COOKE R.J., GREGO S., OLIVER J. and DAVIES D.D., 1979a: The effect of deuterium oxide on protein turnover in Lemna minor. *Planta* 146, 229-236.

- COOKE R.J., OLIVER J. and DAVIES D.D., 1979b: Stress and protein turnover in Lemna minor. *Plant Physiol.* 64, 1109-1113.
- COOKE R.J., ROBERTS K. and DAVIES D.D., 1980a: Model for stress-induced protein degradation in Lemna minor. *Plant Physiol.* 66, 1119-1122.
- COOKE R.J., GREGO S., ROBERTS K. and DAVIES D.D., 1980b: The mechanism of deuterium oxide-induced protein degradation in Lemna minor. *Planta* 148, 374-380.
- COOLEY W.E. and FOY C.L., 1986: Effects of SC-0224 and glyphosate on inflated duckweed (Lemna gibba) growth and EPS-synthase activity from Klebsiella pneumoniae. *Pestic. Biochem. Physiol.* 26(3), 365-374.
- COPE B.T., BOSE S., CRESPI H.L. and KATZ J.J., 1965: Growth of Lemna in H₂O-D₂ mixtures: Enhancement by kinetin. *Bot. Gaz.* 126, 214-221.
- COPELLI M., GHETTI P.F. and CORRADI M., 1982: Rimozione di azoto e fosforo de acque reflue di allevamenti suini mediante fitodepurazioni. *Quad. Ist. Ric. Acque* 60(4), 1-34.
- CORBET R.L., MUIR D.C.G. and WEBSTER G.R.B., 1983: Fate of 1,3,6,8-T4CDD in an outdoor aquatic system. *Chemosphere* 12(4-5), 523-527.
- CORDES H., 1980: Bericht über die Tagung der Floristisch-soziologischen Arbeitsgemeinschaft in Bremen vom 30.6.-3.7.1978. *Mitt. Florist.-Soz. Arb.* 22, 169-173.
- CORDO H.A., DELOACH C.J. and FERRER R., 1981: Biological studies on two weevils, Ochetina bruchi and Onychylis cretatus, collected from Pistia and other aquatic plants in Argentina. *Ann. Entomol. Soc. Am.* 74(4), 363-368.
- CORRADI M., COPELLI M. and GHETTI P.F., 1981: Colture di Lemna su scari-chi zootecnici. *Inquinamento* 23(10), 49-54.
- CORRADI M., GHETTI P.F. and COPELLI M., 1982: Colture di Lemnaee su reflui zootecnici parzialmente trattati. *Atti del convegno internazionale "Fitodepurazione e produzione di biomassa". Parma, 15-16 May, 1981.* Lorenzini, Bologna.
- CORRADI M., REALI M. and BASSI M., 1987: Morphological alterations induced by Cr in freshwater plants. *14th Int. Bot. Congr. Berlin, Abstr.*, 407.
- CORRELL D. and CORRELL H., 1975: Aquatic and wetland plants of South-western United States. Stanford Univ. Press, California, USA. 1, 563-578.
- COUCH R.W. and GANGSTAD E.O., 1974: The response of duckweed to CO₂-laser radiation. *Hyacinth Control J.* 12, 25-26.
- COUNTRYMAN W.D., 1968: Wolfia in New Hampshire. *Rhodora* 70(784), 491.
- COUTINHO A.X.P., 1939: Flora de Portugal. (2nd ed.). Bertrand, Lissabon. 134-135.
- COVEY S.N. and GRIERSON D., 1976: The measurement of plant polyadenylic acid by hybridisation with radioactive polyuridylic acid. *Planta* 131(1), 75-79.
- COWGILL U.M., 1970: The hydrochemistry of Linsley Pond, North Branford, Connecticut. I. Introduction, field work and chemistry by X-ray emission spectroscopy. *Arch. Hydrobiol.* 68, 1-95.
- COX P.A., 1983: Search theory, random motion, and the convergent evolution of pollen and spore morphology in aquatic plants. *Am. Nat.* 121(1), 9-31.
- CRAKER L.E., 1971: Effects of mineral nutrients on ozone susceptibility of Lemna minor. *Can. J. Bot.* 49, 1411-1414.
- CRAKER L.E., 1972: Influence of ozone on RNA and protein content of Lemna minor L. *Environ. Pollut.* 3, 319-323.
- CRESPI H.L., STEWART M.L., KLINE J.R. and KATZ J.J., 1972: Discrimination against tritium by fully deuterated algae. *Health Phys.* 23, 111-112.

- CROIZAT L., 1971: Gigantomachie botanique: La "théorie du Durian" contre la "théorie de la Lentille d'Eau". *Adansonia* 11(1), 47-76.
- CROQUIST A., HOLMGREN A.H., HOLMGREN N.L., REVEAL J.L. and HOLMGREN P.K., 1977: Intermountain Flora. Vascular plants of Intermountain West, USA. Columbia Univ. Press, New York. 6, 474-475.
- CROWDER A.A., BRISTOW J.M., KING M.R. and VANDERKLOET S., 1977a: Distribution, seasonality, and biomass of aquatic macrophytes in Lake Opinicon (Eastern Ontario). *Naturaliste Can.* 104, 441-456.
- CROWDER A.A., BRISTOW J.M., KING M.R. and VANDERKLOET S., 1977b: The aquatic macrophytes of some lakes in Southeastern Ontario. *Naturaliste Can.* 104, 457-464.
- CRUM G.H. and BACHMANN R.W., 1973: Submersed aquatic macrophytes of the Iowa Great Lakes region. *Iowa State J. Res.* 48, 147-173.
- CSEH E., BOEDDI B., BUJTAS C. and LAKATOS B., 1984: Uptake and utilization of iron by chlorotic and green plants. In: CRAM W.J. et al. (eds.), *Membrane transport in plants*. 389.
- CUFODONTIS G., 1971: List of plants of Ethiopia: Spermatophyta. *Bull. Jard. Bot. Nat. Belge* 41(3), 1483-1578.
- CULLEY D.D., Jr. and EPPS A.E., 1973: Use of duckweed for waste treatment and animal feed. *J. Water Pollut. Contr. Fed.* 45, 337-347.
- CULLEY D.D., Jr. and MYERS R.W., 1980: Effect of harvest rate on duckweed yield and nutrient extraction in dairy waste lagoon. *Baton Rouge. 6 pp.* (Polycopy).
- CULLEY D.D., Jr., GHOLSON J.H., CHISHOLM T.S., STANDIFER L.C. and EPPS E.A., 1978: Water quality renovation of animal waste lagoons utilizing aquatic plants. U.S. Environ. Prot. Agency, Ada, Oklahoma. 166 pp.
- CULLEY D.D., Jr., REJMANKOVA E., KVET J. and FRYE J.B., 1981: Production, chemical quality and use of duckweeds (Lemnaceae) in aquaculture, waste management, and animal feeds. *J. World Maric. Soc.* 12(2), 27-49.
- CURTIS H.L., 1971: A study of endogenously produced factors affecting growth and flowering in three species of Lemna. Ph.D. Thesis Ohio State Univ. 103 pp. Diss. Abstr. Int. B 32, 117.
- CURTIS H.L. and BENDIXEN L.E., 1974: Promotion and inhibition of flowering in Lemna. *ASB Bull.* 21(2), 50.
- CUSICK A.W., 1986: Significant additions to the vascular flora of western Maryland. *Castanea* 51(2), 129-136.
- CZERWINSKI W., GANCARZ R., PRZYBYLKA E. and WIECZOREK J.S., 1982: Preliminary estimation of physiological activity of 9-amino-9-fluorene-phosphonic acid derivatives. (In Polish). *Acta Agrobot.* 34(2), 253-260.
- CZERWINSKI W. and WEGRYN T., 1975: The effect of interaction of Venzar and bacteria on photosynthesis of plants. (In Polish). *Roczniki Gleboznawcze* 26(2), 73-77.
- CZOPEK M., 1959a: Cultivation of Polish Lemnaceae species in laboratory conditions. *Acta Biol. Cracov. Ser. Bot.* 2, 14-22.
- CZOPEK M., 1959b: Researches on the physiology of formation and germination of turions in Spirodela polyrrhiza (L.) Schleiden. *Acta Biol. Cracov. Ser. Bot.* 2, 75-90.
- CZOPEK M., 1962: The oligodynamic action of light on the germination of turions of Spirodela polyrrhiza (L.) Schleiden. *Acta Soc. Bot. Pol.* 31, 703-722.
- CZOPEK M., 1963a: Studies on the external factors inducing the formation of turions in Spirodela polyrrhiza (L.) Schleiden. *Acta Soc. Bot. Pol.* 32, 199-211.
- CZOPEK M., 1963b: Methods of Lemnaceae culture. (In Polish). *Wiadomosci Bot.* 7, 153-164.

- DAS R.R., 1965: Sprouting behaviour of turions of *Spirodela polyrrhiza* (L.) Schleid. P.L. (India), Ann. Rep. 480, 70-80.
- DAS R.R., 1966: Growth behaviour of *Spirodela polyrrhiza* (L.) Schleid. under three temperature conditions. P.L. (India), Ann. Rep. 480, 165-168.
- DAS R.R., 1967a: Distribution of *Spirodela polyrrhiza* (L.) Schleid. P.L. (India), Ann. Rep. 480, 197-202.
- DAS R.R., 1967b: Multiplication of *Spirodela polyrrhiza* (L.) Schleid. fronds. P.L. (India), Ann. Rep. 480, 203-206.
- DAS R.R., 1967c: The effect of photoperiod on dry matter production by *Spirodela polyrrhiza* (L.) Schleid. P.L. (India), Ann. Rep. 480, 207-212.
- DAS R.R., 1968: Growth and distribution of *Eichhornia crassipes* (Mart.) Solms. and *Spirodela polyrrhiza* (L.) Schleid. Ph.D. Thesis. Banaras Hindu Univ., Varanasi.
- DAS R.R., 1971: Productivity of *Spirodela polyrrhiza* (L.) Schleid. Int. Symp. Trop. Ecology Emphasizing Productivity. 177-184.
- DAS R.R. and GOPAL B., 1969: Vegetative propagation in *Spirodela polyrrhiza*. Trop. Ecol. 10, 270-277.
- DASSANAYAKE M.D. and FOSBERG F.R., 1981: A revised handbook to the flora of Ceylon. Balkema, Rotterdam. 2, 511 pp.
- DATKO A.H. and MUDD S.H., 1980: Methionine biosynthesis in Lemna: Inhibitor studies. Plant Physiol. 65(6 suppl.), 16.
- DATKO A.H. and MUDD S.H., 1982: Methionine biosynthesis in Lemna: Inhibitor studies. Plant Physiol. 69(5), 1070-1076.
- DATKO A.H. and MUDD S.H., 1983: Sulfate uptake and regulation in Lemna paucicostata Hegelm. 6746. Plant Physiol. 72(1 suppl.), 152.
- DATKO A.H. and MUDD S.H., 1984a: Sulfate uptake and its regulation in Lemna paucicostata Hegelm. 6746. Plant Physiol. 75(2), 466-473.
- DATKO A.H. and MUDD S.H., 1984b: Responses of sulfur-containing compounds in Lemna paucicostata Hegelm. 6746 to changes in availability of sulfur sources. Plant Physiol. 75(2), 474-479.
- DATKO A.H. and MUDD S.H., 1985: Uptake of amino acids and other organic compounds by Lemna paucicostata Hegelm. 6746. Plant Physiol. 77(3), 770-778.
- DATKO A.H. and MUDD S.H., 1986: Uptake of choline and ethanolamine by Lemna paucicostata. Plant Physiol. 81(1), 285-288.
- DATKO A.H., MUDD S.H. and GIOVANELLI J., 1977: Homocysteine biosynthesis in green plants. Studies of the homocysteine forming sulphydrylase. J. Biol. Chem. 252(10), 3436-3445.
- DATKO A.H., MUDD S.H., MACNICOL P.K. and GIOVANELLI J., 1978a: Phytostatin for the growth of Lemna in semicontinuous culture with low sulfate. Plant Physiol. 62, 622-628.
- DATKO A.H., MUDD S.H., GIOVANELLI J. and MACNICOL P.K., 1978b: Sulfur containing compounds in Lemna perpusilla 6746 grown at a range of sulfate concentrations. Plant Physiol. 62, 629-635.
- DATKO A.H., MUDD S.H. and GIOVANELLI J., 1980a: Lemna paucicostata Hegelm. 6746. Development of standardized growth conditions suitable for biochemical experimentation. Plant Physiol. 65, 906-912.
- DATKO A.H., MUDD S.H. and GIOVANELLI J., 1980b: Lemna paucicostata Hegelm. 6746. Life cycle and characterization of the colony types in a population. Plant Physiol. 65, 913-923.
- DAUBS E.H., 1962: The occurrence of *Spirodela oligorhiza* (Kurz) Hegelm. in the United States. Rhodora 64, 83-85.
- DAUBS E.H., 1965: A monograph of Lemnaceae. Univ. of Illinois Press, Urbana. Illinois Biol. Monogr. 34, 118 pp.
- DAVENPORT L.J. and HAYNES R.R., 1981: Aquatic and marsh plants of Alabama. II. Arceidae. Castanea 46(4), 291-299.

- CZOPEK M., 1964a: The action of kinetin, gibberellic acid and red light on the germination of turions of *Spirodela polyrrhiza*. Bull. Acad. Pol. Sci., II, 12, 177-182.
- CZOPEK M., 1964b: The course of photosynthesis and respiration in germinating turions of *Spirodela polyrrhiza*. Bull. Acad. Pol. Sci., II, 12, 463-469.
- CZOPEK M., 1964c: The oligodynamic action of light in the germination of turions of *Spirodela polyrrhiza*. Int. Symp. Physiol. Ecology and Biochem. of Germination, Greifswald, 1963. Abstract.
- CZOPEK M., 1967: Photosynthesis and respiration of turions and vegetative fronds of *Spirodela polyrrhiza*. Acta Soc. Bot. Pol. 36, 87-96.
- CZYGAN F.-C., 1962: Blütenbildung bei Lemna minor nach Zusatz von Oestrogenen. Naturwissenschaften 49(12), 285-286.
- DAFNI A. and AGAMI M., 1976: Extinct plants of Israel. Biol. Conserv. 10, 49-52.
- DAHLGREN R.M.T., CLIFFORD H.T. and YEO P.F., 1985: The families of the Monocotyledons. Springer, Berlin. 520 pp.
- DAIE J., SEELEY S.D. and CAMPBELL W.F., 1977: Effects of temperature on endogenous ABA levels in Lemna minor. 58th Ann. Meet. Pc. Div. AAAS.
- DALE H.M. and GILLESPIE T., 1976: The influence of floating vascular plants on the diurnal fluctuations of temperature near the water surface in early spring. Hydrobiologia 49, 245-256.
- DALE H.M. and MILLER G.E., 1978: Changes in the aquatic macrophyte flora of Whitewater Lake near Sudbury, Ontario from 1947 to 1977. Can. Field-Nat. 92, 264-270.
- DALEO G.R. and KINDEL P.K., 1979: A lipid derivative of galacturonic acid - a possible intermediate in pectic acid biosynthesis. Fed. Proc. 38(3/1), 790.
- DALGLIESH J.G., 1926a: Observations on the British Lemnaceae. J. Bot. (London) 64, 48-50.
- DALGLIESH J.G., 1926b: Notes on duckweeds. J. Bot. (London) 64, 272-274.
- DALRYMPLE R.L., 1971: Experiences with diuron for aquatic weed control. Proc. 24th Ann. Meet. South. Weed Sci. Soc. 333-337.
- DALY J.M. and BROWN A.H., 1954: The in vivo demonstration of cytochrome oxidase in leaves of higher plants. Arch. Biochem. Biophys. 52, 380-387.
- DAMANAKIS M., 1970: A bioassay for the determination of low concentrations of paraquat. Weed Res. 10(1), 77-80.
- DAMANAKIS M., 1972: A bioassay on Lemna polyrrhiza L. for determination of herbicide residues in soils and aqueous solutions. Ann. Inst. Phytopathol. Benaki N.S., 10(3), 248-255.
- DAMANAKIS M., 1976: Behaviour of glyphosphate in the soil (adsorption, leaching, degradation). Ann. Inst. Phytopathol. Benaki N.S. 11, 153-167.
- DAMANAKIS M., DRENNAN D.S.H., FRYER J.D. and HOLLY K., 1970: The adsorption and mobility of paraquat on different soils and soil constituents. Weed Res. 10, 264-277.
- DANN W., 1982: Vergleich von Klonen verschiedener Herkunft in ihrem Verhalten gegenüber Stickstoffkonzentrationen und Stickstoffformen am Beispiel von Lemna gibba. Diploma Thesis. Geobot. Inst. ETH, Stiftung Rübli, Zürich. 56 pp. (polycopy).
- DARVILL A., MCNEIL M. and ALBERSHEIM P., 1977: Structure of plant cell walls: fractionation and characterization of the pectic polymers. Plant Physiol. 59(6 suppl.), 17.
- DAS P., KUMAR D. and GUHA R.M.K., 1975: National demonstration on composite fish culture in West Bengal, India. J. Ind. Fish. Soc. India 7, 112-115.

to diets of different water content. *J. Arid. Environ.* 12(2), 169-174.

DECHI G.S. and EWEL K.C., 1984: Simulated effect of wastewater application on phosphorus distribution in cypress domes. In: EWEL K.C. and ODUM H.T. (eds.), *Cypress swamps*. Univ. Press, Florida. 102-111.

DE HEIJ H.T. and GROOT G.S.P., 1981: Spirodela oligorhiza chloroplast DNA codes for ATPase subunits α and β . Immunological evidence from a coupled transcription-translation system. *FEBS Lett.* 134(1), 6-10.

DE HEIJ H.T., LUSTIG H., MOESKOPF D.-J., BOVENBERG W.A., BISANZ C. and GROOT G.S.P., 1983: Chloroplast DNAs of *Spinacia*, *Petunia* and *Spirodela* have a similar gene organization. *Curr. Genet.* 7(1), 1-6.

DE HEIJ H.T., JOCHIMSEN A.G., WILLEMSSEN P.T.J. and GROOT G.S.P., 1984: Protein synthesis during chloroplast development in *Spirodela oligorhiza*. Coordinated synthesis of chloroplast-encoded and nuclear-encoded subunits of ATPase and ribulose-1,5-bisphosphate carboxylase. *Eur. J. Biochem.* 138(1), 161-168.

DE HEIJ H.T., LUSTIG H., VAN EE J.H., VOS Y.J. and GROOT G.S.P., 1985: Repeated sequences on mitochondrial DNA of *Spirodela oligorhiza*. *Plant Mol. Biol.* 4(4), 219-224.

DEKOCK P.C. and HALL A., 1981: Hormonal control of calcium uptake. *J. Sci. Food Agric.* 32, 989-996.

DEKOCK P.C. and INNES A.M., 1970: The effect of amitrole on duckweed. *Can. J. Bot.* 48, 1285-1288.

DEKOCK P.C., OHTA Y., INKSON R.H.E. and KNIGHT A.H., 1973: The effect of oxalate and ethylenediaminetetraacetic acid on the absorption of calcium into *Lemna*. *Physiol. Plant.* 28, 379-382.

DEKOCK P.C., GRABOWSKA F.B. and INNES A.M., 1974: The effect of salicylic acid on the growth of *Lemna gibba*. *Ann. Bot.* 38, 903-908.

DEKOCK P.C., VAUGHAN D. and HALL A., 1978: Effect of abscisic acid and benzyl adenine on the inorganic and organic composition of the duckweed, *Lemna gibba* L. *New Phytol.* 81, 505-511.

DEKOCK P.C., CHESHIRE M.V., MUNDIE C.M. and INKSON R.H.E., 1979: The effect of galactose on the growth of *Lemna*. *New Phytol.* 82, 679-685.

DE LA CRUZ A.A. and YARBROUGH J.D., 1982: Role of aquatic weeds in maintaining surface water quality. Rep. Mississippi State Univ. Gov. Rep. Announce. Index (US) 83(10), 2137. 64 pp.

DE LANGE L., 1972: An ecological study of ditch vegetation in the Netherlands. Ph.D. Thesis. Amsterdam. 112 pp.

DE LANGE L., 1974: Translocation experiments in the field with the *Lemna gibba* - *Lemna minor* complex. *Acta Bot. Neerl.* 23, 109-112.

DE LANGE L., 1975: Gibbosity in the complex *Lemna gibba*/*Lemna minor*: Literature survey and ecological aspects. *Aquat. Bot.* 1, 327-332.

DE LANGE L. and PIETERSE A.H., 1973: A comparative study of the morphology of *Lemna gibba* L. and *Lemna minor* L. *Acta Bot. Neerl.* 22(5), 510-517.

DE LANGE L. and REVIER J.M., 1982: An investigation into the status of *Spirodela polyrrhiza* (L.) Schleid. var. *masonii* Daubs. In: SYMOENS S., HOOPER S.S. and COMPERE P. (eds.), *Studies on aquatic vascular plants*. R. Soc. Bot. Belg. 82-87.

DE LANGE L. and SEGAL S., 1968: Ecological differences between *Lemna minor* and *Lemna gibba*. (In Dutch). *Gorteria* 4, 5-12.

DE LANGE L. and WESTINGA E., 1979: The distinction between *Lemna gibba* L. and *Lemna minor* L. on the basis of vegetative characters. *Acta Bot. Neerl.* 28(2/3), 169-176.

DE LANGE L., PIETERSE A.H. and VAN BAAREN-BECKERS I., 1981: The occurrence of mixed populations of different genotypes of the *Lemna gibba* - *Lemna minor* complex. *Acta Bot. Neerl.* 30(3), 191-197.

DAVIDSON D., 1979: Aspects morphologiques et enzymatiques de l'adaptation et de l'acclimatation au stress thermique des populations écologiques de *Spirodela polyrrhiza* (L.) Schleid. (Lemnaceae). M.S. Thesis. Univ. Montreal.

DAVIDSON D. and SIMON J.P., 1981a: Thermal adaptation and acclimation of ecotypic populations of *Spirodela polyrrhiza* (Lemnaceae): thermotolerance and apparent activation energy of NAD malate dehydrogenase. *Can. J. Bot.* 59(6), 1061-1068.

DAVIDSON D. and SIMON J.P., 1981b: Thermal adaptation and acclimation of ecotypic populations of *Spirodela polyrrhiza* (L.) Schleid. (Lemnaceae): Morphology and growth rates. *J. Therm. Biol.* 6(3), 121-128.

DAVIDSON D. and SIMON J.P., 1983: Thermal adaptation and acclimation of ecotypic populations of *Spirodela polyrrhiza* (L.) Schleid. (Lemnaceae): Temperature dependency of K_m of NAD malate dehydrogenase. *J. Therm. Biol.* 8(3), 289-296.

DAVIES D.D., 1978: The effect of stress on protein degradation in plants. In: SCHUETTE H.R. and GROSS D. (eds.), *Regul. Dev. Processes Plants*, Proc. Conf. 1977. Fischer, Jena. 13-35.

DAVIES D.D., 1979: Factors affecting protein turnover in plants. In: HEWITT E.J. and CUTTING C.V. (eds.), *Nitrogen assimilation of plants*. Acad. Press, New York.

DAVIES D.D. and HUMPHREY T.J., 1978: Amino acid recycling in relation to protein turnover. *Plant Physiol.* 61, 54-58.

DAVIS J.A., 1981: Comparison of static replacement and flow-through bioassays using duckweed, *Lemna gibba* G3. Final report. Gov. Rep. Announce. Index (US) 81(18), 3790. 106 pp.

DE BEAUVOIS P., 1816: Mémoire sur les Lemna ou lentille d'eau, sur leur fructification et sur la germination de leur grain. *J. Phys. Chim. Hist. Nat.* 82, 101-115.

DE BOLOS O. and MASCLANS F., 1955: La végétation de los arrozales en la region mediterranea. *Collectanea Botanica* 4(3,22), 415-434.

DEBUSK T.A., 1980: Evapotranspiration of some emergent freshwater plants. In: RYTER J.H. (ed.), *Cultivation of macroscopic marine algae and freshwater aquatic weeds*. Progress report for the period May 1, 1979 - December 15, 1979. Woodshole Oceanic Institution, Woods-hole, Mass., USA. 63-68.

DEBUSK T.A., RYTER J.H., HANISAK M.D. and WILLIAMS L.D., 1981: Effects of seasonality and plant density on the productivity of some freshwater water macrophytes. *Aquat. Bot.* 10(2), 133-142.

DEBUSK T.A., RYTER J.H. and WILLIAMS L.D., 1983: Evapotranspiration of *Eichhornia crassipes* (Mart.) Solms and *Lemna minor* L. in Central Florida: Relation to canopy structure and season. *Aquat. Bot.* 16(1), 31-39.

DECLERE M. and DE CAT W., 1977: Détection de traces d'herbicides dans l'eau par biotests. Etude de divers procédés d'extraction. *Rev. Agric.* 30(3), 589-594.

DECLERE M., DE CAT W. and BASTIN R., 1976: Détection rapide de divers herbicides dans l'eau par la mesure in vivo de l'activité de la nitratre réductase de *Lemna minor* L. *Z. Pflanzenphysiol.* 77, 315-322.

DEGANI N., 1981: DNA repair synthesis in carrot root tissue irradiated with UV light. *Proc. Int. Bot. Congr.* 13, 83.

DEGANI N., BEN-HUR E. and RIKLIS E., 1980: DNA damage and repair: induction and removal of thymine dimers in ultraviolet light irradiated intact water plants. *Photochem. Photobiol.* 31(1), 31-36.

DEGEN A.A., 1987: Responses of two Negev Desert Israel phasianids: the chukar, *Alectoris chukar*, and the sand partridge, *Ammodramus hesperis*.

DEVI S.L. and MAHESHWARI S.C., 1979: Diurnal fluctuations in the activity of the enzyme nitrate reductase in *Lemna paucicostata*. *Physiol. Plant.* 45, 467-469.

DEVIDE Z., 1956: A new locality of *Wolffia arrhiza* (L.) Wimm. in Croatia. *Acta Bot. Zagreb* 14/15, 184-186.

DEYL M., 1955: The evolution of the plants and the taxonomy of the monocotyledons. *Acta Mus. Nat. Prag.* 118(6), 143 pp.

DEYSSON G., 1959: Action de la kinétine et de la thiokinétine sur la croissance de la lentille d'eau (*Lemna minor* L.). *C.R.Acad.Sci.Paris*, 248, 841-843.

D'HARLINGUE A., 1976: Stérols, acides gras, lipochromes et nucléotides pyridiniques et adényliques des frondes de *Spirodela* cultivées en présence d'acide gibbérellique. *Physiol.Vég.* 14(4), 713-723.

D'HARLINGUE A., LECHÉVALIER D. and MONGER R., 1976: Nucléotides pyridiniques, chlorophylles et stérols des frondes de *Spirodela* cultivées en présence de saccharose. *Physiol.Vég.* 14(2), 367-376.

DICHT N., KOPP A., FELLER U. and ERISMANN K.H., 1976: Einfluss von Ammonium und Nitrat auf den Proteingehalt von *Lemna minor* L. unter Photosynthesebedingungen. *Biochem.Physiol.Pflanz.* 170, 531-534.

DICKSON H., 1938a: The occurrence of long and short cycles in growth measurements of *Lemna minor*. *Ann.Bot.N.S.* 2, 97-106.

DICKSON H., 1938b: Sampling as the cause of the apparent growth cycles of *Lemna minor*. *Ann.Bot.N.S.* 2, 793-806.

DIEKJOEST H., 1983a: Ein vorübergehendes Vorkommen von *Wolffia arrhiza* in Westfalen. *Natur und Heimat* 43(3), 65-73.

DIEKJOEST H., 1983b: Zur gegenwärtigen Verbreitung von *Lemna minor* in der Unteren Erft. *Gött.Flor.Rundbr.* 17, 168-173.

DIEKJOEST H., 1984: *Pistia stratiotes* L. und *Lemna aquinoctialis* Welwitsch vorübergehend im Gebiet der unteren Erft. *Gött.Flor.Rundbr.* 18, 90-95.

DJIKMAN M.J., BOSS M.L. and RUSSELL E.R., 1964: Extension of life span of *Lemna perpusilla* clone 6746 by ultraviolet irradiation. *Radiat. Res.* 22, 662-667.

DINGES R., 1973: Ecology of *Daphnia* in stabilization ponds. *Texas Dept. Health,Austin,Texas* 1, 143 pp.

DINGES R., 1976a: A proposed integrated biological wastewater treatment system. In: TOURBIER J. and PIERSON R.W. (eds.), *Biological control of water pollution*. Univ.Pennsylvania Press,Philadelphia,PA. 225-230.

DINGES R., 1976b: Who says sewage treatment plants have to be ugly? *Water and Wastes Engineering* 13(4), 20-23.

DINGES R., 1976c: Water hyacinth culture for wastewater treatment. *Texas Dept.Health,Austin,Texas*. 143 pp.

DINGES R., 1981: Natural systems for water pollution control. *Van Nostrand,Wokingham,UK*. 102-110.

DINGES R., 1982: Aquatic plant systems - an unconventional approach to removal of toxic materials. 10th Water Resources Symp., "Toxic Materials - Methods of Control", Austin,Texas. 23 pp. (Polycopy).

DOCAUER D.M., 1983: A nutrient basis for the distribution of the *Lemna* ceae. *Ph.D.Thesis*. Univ.Michigan. 223 pp.

DOCAUER D.M., 1984: Problems in differentiating *Lemna turionifera* (Lam.) from *Lemna minor* (Hegel.). *Am.J.Bot.* 71(5/2), 164.

DOHERRY H.M., SELVEDRAN R.R. and BOWLES D.J., 1984: The role of oligosaccharides in cell-signalling events in plants. Poster 2-109-1 at the 14th Int.Bot.Congr.,Berlin.

DOLBERG F., SAADULLAH M. and HAQUE M., 1981: A short review of the feeding value of water plants. *Trop.Animal Prod.* 6(4), 322-326.

DONE W.G., 1957: *Wolffia* in Canada. *Can.Field-Nat.* 71, 10-16.

DE LANGE L., PIETERSE A.H. and WETSTEYN L.P.M.J., 1984: On the occurrence of the flat form of *Lemna gibba* L. in nature. *Acta Bot.Neerl.* 33(4), 469-474.

DELAY C., 1947: Recherches sur la structure des nouveaux quiescents chez les phanérogames. *Rev.Cytol.Cytophysiol.Vég.* 9, 129-223; 10, 103-229.

DELILE A., 1813: Flore d'Egypte. Paris. 75.

DELPINO F., 1982: Rivista di Botanica dell'anno 1881. Milano. 33 pp.

DE MARTONNE E., 1927: Traité de Géographie physique. I. Notions générales. Hydrographie. Colin, Paris. 495 pp.

DEMARTY M., AYADI A., MONNIER A., MORVAN C. and THELLIER M., 1977: Electrochemical properties of isolated cell walls of *Lemna minor* L. In: THELLIER M. et al. (eds.), *Transmembrane ionic exchanges in plants*. CNRS et Univ. Rouen, Paris et Mont Saint Aignan. 61-73.

DEMARTY M., MORVAN C. and THELLIER M., 1978: Exchange properties of isolated cell walls of *Lemna minor* L. *Plant Physiol.* 62, 477-481.

DEMARTY M., RIPPOLL C. and THELLIER M., 1980: Ion exchange in plant cell walls (*Lemna minor*). *Developments in plant biology* 4. Elsevier, North-Holland Biomed.Press. 33-47.

DEN HARTOG C., 1968: The flat form of *Lemna gibba*, still a problem. (In Dutch). *Gorteria* 4, 90-92.

DEN HARTOG C., 1969: Proposal to conserve the generic name 976 *Wolffia* Horkel ex Schleiden (1844) (Lemnaceae) versus *Wolffia* Schreiber (Flacourtiaceae), and by retypification against *Wolffia* Horkel ex Schleiden (1839) (Lemnaceae). *Taxon* 18, 591-592.

DEN HARTOG C., 1970: Proposal of the conservation of the generic name *Spirodela* Schleiden versus *Lenticularia* Seguiet (Lemnaceae). *Taxon* 19(4), 647-648.

DEN HARTOG C., 1975: Thoughts about the taxonomical relationships within the Lemnaceae. *Aquat.Bot.* 1, 407-416.

DEN HARTOG C., 1978: Subjective phytosociology of aquatic plants. *Aquat.Bot.* 4, 96-98.

DEN HARTOG C. and SEGAL S., 1964: A new classification of the water-plant communities. *Acta Bot.Neerl.* 13, 367-393.

DEN HARTOG C. and VAN DER PLAS F., 1970: A synopsis of the Lemnaceae. *Blumea* 18, 355-368.

DEN HARTOG C. and VAN DER PLAS F., 1972: The Australian species of *Wolffia* (Lemnaceae). *Blumea* 20, 151-153.

DENNY P., HARMAN J., ABRAHAMSON J. and BRYCESON I., 1978: Limnochemical and phytoplankton studies on Nyumba ya Munga reservoir. Tanzania. *Biol.J.linn.Soc.* 10, 24-48.

DENTON J.B., 1966: Relationship between the chemical composition of aquatic plants and water quality. M.S.Thesis. Auburn Univ., Auburn. 14 pp.

DE SLOOVER J.-L., 1961: Note sur le pollen de *Lemna minor* L. Pollen et Spores 3, 5-10.

DE SLOOVER J.-L., 1964: Sur la floraison des Lemnaceae. *Nat.Mosana* 17, 73-81.

DE SLOOVER J.-L., 1966: La fronde, la graine et la germination d'une *Lemna*. *Nat.Belg.* 47, 443-456.

DE SLOOVER J.-L., 1963: Lemnaceae du Rwanda. *Bull.Jard.Bot.Nat.Belg.* 43, 361-368.

DEUTCH B. and RASMUSSEN O., 1974: Growth chamber illumination and photomorphogenetic efficacy. I. Physiological action of infrared radiation beyond 750 nm. *Physiol.Plant.* 30, 64-71.

DEVARAJ K.V., RAO D.S.K. and KESHAVAPPA G.V., 1981: Utilization of duckweed *Lemna minor* and waste cabbage leaves in the formulation of fish feed. *Mysore J.Agric.Sci.* 15(1), 132-135.

- DORMAN-PRZYBYL D., STRITTMATTER G. and KOSSEL H., 1986: The region distal to the rRNA operon from chloroplasts of maize contains genes coding for TRNA^{Arg}(AGC) and TRNA^{Asn}(GUU). *Plant Mol. Biol.* 7(6), 419-431.
- DOROFEY P.I., 1963: The tertiary floras of Western Siberia. (In Russian). *Bot. Inst. Komarov, Akad. Nauk SSSR*. 129-130.
- DOSS R.P., 1973: Inhibition of flowering in Lemna by non-flowering fronds. *Plant Physiol.* 51(suppl.), 29.
- DOSS R.P., 1974: Factors which influence the flowering of Lemna perpusilla Torr., strain 6746. Ph.D. Thesis. Univ. Calif., Davis. 112 pp. *Int. Diss. Abstr. Int. B* 35, 1525.
- DOSS R.P., 1975a: Influence of temperature on the flowering of Lemna perpusilla 6746 grown under skeleton photoperiods. *Plant Physiol.* 55, 108-109.
- DOSS R.P., 1975b: Influence of timing and number of consecutive inductive photoperiodic cycles on the flowering of Lemna. *Plant Physiol.* 55, 110-111.
- DOSS R.P., 1975c: Reversal of the effects of a night interruption in Lemna by inhibitors of ribonucleic acid synthesis. *Plant Physiol.* 55, 112-113.
- DOSS R.P., 1975d: Influence of short term inhibitor treatment on the flowering of Lemna perpusilla 6746. *Plant Physiol.* 56, 360-363.
- DOSS R.P., 1978: Handedness in duckweed: Double flowering fronds produce right- and left-handed lineages. *Science* 199, 1465-1466.
- DOUGALL D.X., 1977: Current problems in the regulation of nitrogen metabolism in plant cell cultures. In: BARZ W., REINHARD E. and ZENK M.H. (eds.), *Proceedings in Life Science. Plant tissue culture and its biotechnological application*. Springer, New York/Berlin. 76-84.
- DRAGENDORFF G., 1898: *Die Heilpflanzen der verschiedenen Völker und Zeiten*. Enke, Stuttgart. 108.
- DRIVER E.A. and PEDEN D.G., 1977: The chemistry of surface water in prairie ponds. *Hydrobiologia* 53(1), 33-48.
- DROTAR A., PHELPS P. and FALL R., 1985: Evidence for glutathione peroxidase activities in cultured plant cells. *Plant Sci.* 42(1), 35-40.
- DUBINA D.V. and PROTOPOVA V.V., 1983: Lemna minuscula new record for the USSR flora. (In Ukrainian). *Ukr. Bot. Zh.* 40(5), 28-31.
- DUDLEY J.L., 1983: Responses of some Lemna strains to different environmental conditions. M.S. Thesis. Univ. Pittsburgh, 78 pp.
- DUDLEY J.L., 1987: Turion formation in strains of Lemna minor (6591) and Lemna turionifera (6573, A). *Aquat. Bot.* 27(2), 207-215.
- DUFF R.B., 1965: The occurrence of apiose in Lemna (duckweed) and other angiosperms. *Biochem. J.* 94, 768-772.
- DUFF R.B. and KNIGHT A.H., 1963: The occurrence of apiose in Lemna (duckweed) and other angiosperms. *Biochem. J.* 88, 33P-34P.
- DUFFIELD A.N., 1981: The impact of Lemna on the oxygen resources of channels of potential value as fisheries. *Natl. Veget. Res. Sta., Wellesbourne, Warwick, UK, Assoc. Applied Biologists* 257-264.
- DUFFIELD A.N. and EDWARDS R.W., 1981: Predicting the distribution of Lemna spp. in a complex system of drainage channels. *Natl. Veget. Res. Sta., Wellesbourne, Warwick, UK, Assoc. Applied Biologists* 59-65.
- DUROVA E., 1970: Contribution to the production ecology of duckweeds. (In Czech.). Diploma Thesis. Prace Karlovy Univ.
- DUKE S.H., 1975: Glutamate dehydrogenase: studies of the enzyme in Pisum and Pastinaca roots, Lemna clones and Glycine root nodules in relation to distribution, red and far-red illuminations, oscillations in activity, urea utilization and carbohydrate metabolism. Ph.D. Thesis. Univ. Minnesota. 144 pp. *Int. Diss. Abstr. B* 36, 3196.
- DUKE S.H. and KOUKKARI W.L., 1977: Glutamate dehydrogenase activity in

- Lemna perpusilla 6746: The effects of sucrose, glucose and fructose addition to growth media. *Physiol. Plant.* 39, 67-72.
- DUKE S.H., WENZLER H.C. and KOUKKARI W.L., 1975: The effects of sucrose on glutamate dehydrogenase activity in Lemna perpusilla 6746. *Plant Physiol.* 56(2 suppl.), 66.
- DUONG T.P. and TIEDJE J.M., 1985: Nitrogen fixation by naturally occurring duckweed-cyanobacterial associations. *Can. J. Microbiol.* 31(4), 327-330.
- DUTAILLY G., 1978: Sur la nature réelle de la fronde et du cotylédon des Lemna. *Bull. Mens. Soc. Linn. Paris* 19, 147.
- DUTHU G.S. and KILGEN R.H., 1975: Aquarium studies in the selectivity of 16 aquatic plants as food by fingerling hybrids of the cross between Ctenopharyngodon idella ♂ (grass carp) and female Cyprinus carpio ♀. *J. Fish Biology* 7, 203-208.
- DUVAL Y., THELLIER M., HEURTEAUX C. and WISSOCQ J.C., 1980: Detection of stable isotopes with a (n,α) nuclear reaction: Application to the measurement of unidirectional fluxes of borate in a plant. *J. Radioanal. Chem.* 55, 297-306.
- DVORAKOVA-HLADKA J., 1964: Respiration of intact and damaged plants of Spirodela oligorhiza Schl. *Biol. Plant. (Praha)* 6, 198-201.
- DYAR M.T., 1953: Studies on the reduction of a tetrazolium salt by green plant tissue. *Am. J. Bot.* 40, 20-25.
- DYER T.A., 1982: Isolation of low molecular weight RNAs from chloroplasts. In: EDELMAN M., HALLICK R.B. and CHUA N.-H. (eds.), *Methods Chloroplast Mol. Biol.* Elsevier, North-Holland Biomed. Press. 359-368.
- DYER T.A. and BOWMAN C.M., 1976: A sequence analysis of low-molecular-weight RNA from chloroplasts of flowering plants. In: BUECHER Th. et al. (eds.), *Genetics and biogenesis of chloroplasts and mitochondria*. Elsevier, North-Holland Biomed. Press. 645-651.
- DYER T.A. and BOWMAN C.M., 1979: Nucleotide sequences of chloroplast 5S ribosomal ribonucleic acid in flowering plants. *Biochem. J.* 183, 595-604.
- DYKJOVA D., 1979: Selective uptake of mineral ions and their concentration factors in aquatic higher plants. *Folia Geobot. Phytotax.* 14, 267-325.
- DYKJOVA D. and KVET J., 1978: Growth, production and nutrient uptake of duckweeds in fishponds and in experimental cultures. In: *Pond littoral ecosystems*. Ecol. Stud. No. 28. Springer, Berlin. 278-285.
- DYKJOVA D. and KVET J., 1982: Mineral nutrient economy in wetlands of the Trebon Basin Biosphere Reserve, Czechoslovakia. In: GOPAL B. et al. (eds.), *Wetlands ecology and management*. Nat. Inst. Ecol. Int. Sci. Publ., Jaipur (India) 335-355.
- DYLIK J., LEZNICKA B., MAJEWSKA J., PIECUCH I., PRZADKA W. and WIDLICKA B., 1979: Feeding of Aythya ferina L. and platyrhynchos L. in seminaural conditions and their mineral metabolism. *Acta Physiol. Pol.* 30(1), 63.
- EMMES C. and POSNER H.B., 1974: Abnormal flowering responses and a lack of photosynthesis in a mutant of Lemna perpusilla. *Plant Physiol.* 54(suppl.), 3.
- EATON R.J., 1939: Wolffia columbiana in Concord, Massachusetts. *Rhodora* 41, 42-43.
- EATON R.J., 1947: Lemna minor as an aggressive weed in the Sudbury River. *Rhodora* 49, 165-171.
- EBERLE G., 1933: Wasserlinse. *Natur und Museum* 63, 209-211.
- ECHLIN P., 1981: The analysis of organic surfaces. *Scanning Electron Microsc.* 1981(1), 1-20.

- ECHLIN P., 1982: Low-temperature X-ray microanalysis of solid biological samples: prospects and problems. *Proc. Meet. Electron Microsc. Soc. America*, 40, 380-383.
- ECHLIN P. and BURGESS A., 1977: Cryofracturing and low-temperature scanning electron microscopy of plant material. In: JOHARI O.M. (ed.), *Scanning Electron Microsc.* 1977(1), 491-500.
- ECHLIN P. and HAYES T.L., 1986: The distribution of elements in developing roots of Lemna minor L. *Microbeam Anal.* 21, 245-247.
- ECHLIN P., PAWLEY J.B. and HAYES T.L., 1979a: Freeze-fracture scanning electron microscopy of Lemna minor L. (duckweed). *Scann. Electron Microsc.* 1979(5), 69-76.
- ECHLIN P., SAUBERMANN A., SKAER H. and FRANKS F., 1979b: The use of polymeric cryoprotectants in low-temperature X-ray microanalysis of biological specimens. *Cryobiology* 16(6), 591-592.
- ECHLIN P., FRANKS F., SAUBERMANN A., LAI C.E. and SKAER H., 1980a: The use of high molecular weight polymers in the cryofixation of cells and tissues for ultrastructural and analytical studies. *Electron Microsc.*, *Proc. Eur. Congr.* 7th, 2, 714-715.
- ECHLIN P., HAYES T., LAI C. and HOOK G., 1980b: Elemental analysis of phloem tissue in Lemna minor root tips. *Proc. Ann. Meet. Electron Microsc. Soc. America* 38, 798-799.
- ECHLIN P., LAI C.E., HAYES T.L. and HOOK G., 1980c: Elemental analysis of frozen-hydrated differentiating phloem parenchyma in roots of Lemna minor L. *Scanning Electron Microsc.* 1980(2), 383-394.
- ECHLIN P., LAI C.E., HAYES T.L. and SAUBERMANN A., 1980d: Cryofixation of Lemna minor roots for morphological and analytical studies. *Cryo Lett.* 1, 289-298.
- ECHLIN P., LAI C.E., SAUBERMANN A., HAYES T.L., SKAER H. and FRANKS F., 1980e: Cryofixation of plant material for ultrastructural and analytical studies (Lemna minor). *Proc. Meet. Electron Microsc. Soc. America* 38, 796-797.
- ECHLIN P., SAUBERMANN A., HAYES T.L. and LAI C.E., 1980f: Elemental analysis of differentiating phloem tissue in the root-tips of Lemna minor. *Electron Microsc.*, *Proc. Eur. Congr.* 7th, 3, 124-125.
- ECHLIN P., LAI C.E. and HAYES T.L., 1981: The distribution and relative concentration of potassium in the root-tips of Lemna minor L. analyzed by low temperature X-ray microanalysis. *Scanning Electron Microsc.* 1981(2), 489-498.
- ECHLIN P., LAI C.E. and HAYES T.L., 1982a: Low-temperature X-ray microanalysis of the differentiating vascular tissue in root tips of Lemna minor L. *J. Microsc.* 126(3), 285-306.
- ECHLIN P., LAI C.E. and HAYES T.L., 1982b: The uptake and distribution of potassium in roots of duckweed (Lemna minor L.) as measured using low temperature X-ray microanalysis. *Proc. Meet. Electron Microsc. Soc. America* 40, 408-409.
- ECHLIN P., LAI C.E. and HAYES T.L., 1982c: Uptake and distribution of rubidium into root tips of Lemna minor measured using low temperature X-ray microanalysis. *Proc. 10th Int. Congr. Electron Microsc.*, Hamburg, 401-402.
- ECHLIN P., HAYES T.L. and MCKOON M., 1983: Analytical procedures for bulk frozen-hydrated biological tissues. *Microbeam Anal.* 18, 243-246.
- ECHLIN P., GEE W. and CHAPMAN B., 1985: Very low voltage sputter coating. *J. Microsc.* 137(2), 155-170.
- EDELMANN M., MARDER J.B. and MATTOO A.K., 1983: A compendium of characteristics for the rapidly metabolized 32 kD protein of the chloro-

- plast membrane (Spirodela oligorhiza). *NATO Adv. Study Inst. Ser. A* 63, 187-192.
- EDELMANN M., MATTOO A.K. and MARDER J.B., 1984: Three hats of the rapidly metabolized 32-kD protein of thylakoids. In: ELLIS R.J. (ed.), *Chloroplast biogenesis. Soc. Exp. Biol. Sem. Ser.* 21, 283-302.
- EDELMANN M. and REISFELD A., 1978: Characterization, translation and control of the 32,000 dalton chloroplast membrane protein in Spirodela. In: AKOYUNOGLU G. and ARGYROUDI-AKOYUNOGLU J.H. (eds.), *Chloroplast development. Elsevier North-Holland, Amsterdam.* 641-652.
- EDELMANN M. and REISFELD A., 1980: Synthesis, processing and functional probing of P-32000, the major membrane protein translated within the chloroplast. In: LEAVER C.J. (ed.), *Genome organization and expression in plants. Plenum, New York.* 353-362.
- EDMONDSON Y.H. and THIMANN K.V., 1950: The biogenesis of the anthocyanins. II. Evidence for the mediation of copper in anthocyanin synthesis. *Arch. Biochem.* 25, 79-90.
- EDWARDS D.J., 1974: Weed preference and growth of young grass carp in New Zealand. *New Zealand J. Marine and Freshwater Res.* 8(2), 341-350.
- EDWARDS D.J., 1975: Taking a bite at the waterweed problem. *New Zealand J. Agric.* 130(1), 33-36.
- EDWARDS J. and FLETCHER J., 1978: Influence of light on amino acid and protein synthesis in Lemna minor. *Plant Physiol.* 61(4 suppl.), 20.
- EDWARDS J.L., 1932: *Wolffia florida* in northern New Jersey. *Torreyana* 32, 100.
- EDWARDS P., 1980: Food potential of aquatic macrophytes. ICLARM, Manila, Philippines. 51 pp.
- EFRAI Y., ARZEE T. and PORATH D., 1975: Duckweeds (Lemnaceae) in culture, a morphogenetic system for studying the leaf-root relationship. *Isr. J. Bot.* 24(1), 50.
- EFRAI Y., PETERSE A.H. and BOUMAN F., 1977: A comparative study of developing air chambers in flat and gibbous fronds of Lemna gibba L. *Acta Bot. Neerl.* 26, 343-347.
- EGGERS G., 1985: Zum natürlichen Standort von *Wolffia* welwitschii in Senegambia. *Aqua Planta* 10(3), 6.
- EHMKE A. and HARTMANN T., 1976: Properties of glutamate dehydrogenase from Lemna minor. *Phytochem.* 15, 1611-1617.
- EHMKE A. and HARTMANN T., 1978: Control of glutamate dehydrogenase from Lemna minor by divalent metal ions. *Phytochem.* 17, 637-641.
- EHMKE A., FLOSSDORF J., HABICHT W., SCHIEBEL H.M. and SCHULTEN H.R., 1980: A new technique for determining the hydride transfer stereospecificity and NAD linked oxido reductases electron impact and field desorption mass spectrometry. *Anal. Biochem.* 101(2), 413-420.
- EHRENBURG L., 1945: Chromosome numbers of some vascular plants. (*In Swed.*). *Bot. Not.* 1954(4), 430-437.
- EHRHART F., 1979: Wiedergefundene Blüte der dicken Wasserlinse (Lemna gibba L.). *Hannoversch. Magazin* 17, 1057-1068; reprinted in EHRHART F., *Beitr. Naturk.* 1, 43-51 (1978).
- EHRlich S., 1966: Two experiments in the biological clarification of stabilization-pond effluents. *Hydrobiologia* 27, 70-80.
- EICHENBERGER W., 1975: Ueber den Steringehalt pflanzlicher Mikrosomen. *Chimia* 29(3), 132-133.
- EICHORN M., 1983: Zur Aktivität der Glucose-6-phosphat-Dehydrogenase im Ernst-Moritz-Arndt Univ. Greifsw. Math.-Natw. Reihe 32(3-4), 34-36.
- EICHORN M., 1984: Untersuchungen zur Populationsdynamik und zur Regulation einiger Stoffwechselparameter bei Chemostat-Kulturen von *Wolffia*

- arrhiza unter besonderer Berücksichtigung des Lichtfaktors. Diss.B. Friedrich-Schiller-Univ. Jena.
- EICHORN M., 1986: Untersuchungen zur Populationsdynamik und zur Regulation einiger Stoffwechselparameter bei Chemostat-Kulturen von *Wolffia arrhiza* unter besonderer Berücksichtigung des Lichtfaktors. Biol. Rundsch. 24, 71-73.
- EICHORN M., 1987: Zur Populationsdynamik der Wasserlinse *Wolffia arrhiza*. Biol. Rundsch. 25, 1-4.
- EICHORN M. and AUGSTEN H., 1969: Continuous cultivation of *Wolffia arrhiza* in a chemostat. Flora A 160(6), 576-580.
- EICHORN M. and AUGSTEN H., 1974: Der Einfluss des Bors auf verschiedene altrige Populationen von *Wolffia arrhiza* (L.) Wimm. in Chemostaten-Kultur. Biochem. Physiol. Pflanzen 165(4), 371-385.
- EICHORN M. and AUGSTEN H., 1977a: Der Einfluss löslicher Kohlenhydrate auf die Aktivität der Glucose-6-phosphat-Dehydrogenase verschiedener altriger *Wolffia*-Populationen unter Berücksichtigung von Energiecharge, O₂-Austausch und Pyruvat-Gehalt. Z. Pflanzenphysiol. 84(1), 37-48.
- EICHORN M. and AUGSTEN H., 1977b: Die Wirkung von Blau- und Rotlicht auf die Aktivität der Glucose-6-phosphat-Dehydrogenase und das Adenylsystem bei *Wolffia arrhiza* unter steady state Bedingungen. Z. Pflanzenphysiol. 85, 147-152.
- EICHORN M. and AUGSTEN H., 1980: Einfluss verschiedener Lichtqualitäten auf das endogene Zytokinin-Spektrum bei Lemnaceen-Populationen in Abhängigkeit von der Wachstumsrate. Wiss. Z. Ernst-Moritz-Arndt Univ. Greifsw., Math.-Natw. Reihe 29(1-2), 51-53.
- EICHORN M. and AUGSTEN H., 1983a: Hill reaction activity in isolated chloroplasts from *Wolffia arrhiza* cultivated under white, blue or red radiation under different chemostat steady states. Photosynthetica 17(1), 28-33.
- EICHORN M. and AUGSTEN H., 1983b: Changes in the polyadenylated RNA from *Wolffia arrhiza* continuously illuminated with white, blue or red light. Biochem. Physiol. Pflanzen 178(2-3), 183-192.
- EICHORN M. and AUGSTEN H., 1984: Untersuchungen an isolierten Lemnaceen-Chloroplasten. Coll. Pflanzenphysiol. Humboldt-Univ., Berlin, 7, 149-152.
- EICHORN M. and AUGSTEN H., 1986: Photoregulated RNA metabolism aspects of the mRNA processing in *Wolffia arrhiza*. (In German). Wiss. Z. Ernst-Moritz-Arndt-Univ. Greifsw., Math.-Natw. Reihe 35(5), 577-590.
- EICHLER A.W., 1875: Blutendiagramme I. Leipzig. 73-80.
- EINHELLIG F.A., LEATHER G.R. and HOBBS L.L., 1985: Use of *Lemna minor* L. as a bioassay in allelopathy. J. Chem. Ecol. 11(1), 65-72.
- EINOR L.O., 1984: Characteristics of the formation of oxygen status and pH of natural water in presence of submerged vegetation under conditions in the Ivankovo Reservoir. (In Russian). Vodn. Resur. 2, 122-131.
- EISENMEIER S. and SPILLER H., 1973: Verhinderung der Nitrat-Reductase-Inaktivierung bei *Lemna minor* im Verlaufe der Anreicherung. Hoppe Seyler's Z. Physiol. Chem. 354(10-11), 1184.
- ELAMZON R., 1977: Clarosan IG for control of submerged vegetation. Swed. Weed Conf. 18(1), K1-K3.
- EL-DIN M.N.A., 1982: Biofertilizers - requirements and application. FAO Soils Bull. 45, 164-174.
- EL-DIN M.N.A., FAWAZ K., ABDEL-GHAFFAR A.S. and HASHIM F., 1978: Nitrogenase activity under some water plants in Egypt. 1. *Lemna gibba* and *Eichhornia crassipes* (cited in EL-DIN 1982 as in press).
- EL-FAWARIS A.H., 1985: Bioaccumulation of stable chromium by duckweed

- plants in stream microcosm. In: LEKKAS T.D. (ed.), Heavy Met. Environ., Proc. 5th Int. Conf., Tripoli. CEP Consult., Edinburgh. 2, 268-270.
- EL-GHAZAL R.A.K. and RIEMER D.N., 1986: Germination suppression by extracts of aquatic plants. J. Aquat. Plant Manage. 24, 76-79.
- EL-HINNAWI E., 1983: Energy farms: future prospects. Mazingira 7(3), 4-13.
- ELLIOT D.C., 1977: Induction by EDTA of anthocyanin synthesis in *Spirodela oligorrhiza*. Austral. J. Plant Physiol. 4, 39-49.
- ELLIS L., 1955: Preliminary notes on the correlation between alkalinity and the distribution of some free-floating and submerged aquatic plants. Ecology 36(4), 763-764.
- ELLIS P.C. and DAVIS G.J., 1980: Analysis of phytoplankton and algal species diversity of municipal wastewater following treatment with alligatorweed and duckweed communities. J. Elisha Mitchell Sci. Soc. 96(1), 250.
- ELLIS R.J., 1969: Sulfate activation in higher plants. Planta 88(1), 34-42.
- EL-SHINAWY R.M.K. and ABDEL-MALIK W.E.Y., 1980: Retention of radionuclides by some aquatic freshwater plants. Hydrobiologia 69(1-2), 125-129.
- ELZENGA J.T.M., DE LANGE L. and PIETERSE A.H., 1980: Further indications that ethylene is the gibbosity regulator of the *Lemna gibba*/Lemna minor complex in natural waters. Acta Bot. Neerl. 29, 225-229.
- EMES M.J. and ERISMANN K.H., 1981: Effect of nitrogen nutrition on the structure and activity of glycolate oxidase from *Lemna minor*. Plant Physiol. 67(4 suppl.), 85.
- EMES M.J. and ERISMANN K.H., 1982: The influence of nitrogen supply on the structure and activity of glycolate oxidase in *Lemna minor*. Plant Sci. Lett. 27(1), 103-109.
- EMES M.J. and ERISMANN K.H., 1984: Purification and properties of glycolate oxidase from *Lemna minor*. Int. J. Biochem. 16(12), 1373-1378.
- ENGELMANN G., 1870: *Spirodela*. Bull. Torr. Club 1, 42-43.
- ENGELMANN G., 1871a: Notes of *Lemna*. Bull. Torr. Club 2, 10-11.
- ENGELMANN G., 1871b: *Arctostaphylos*. Bull. Torr. Club 2, 34-35.
- ENGELMANN G., 1871c: *Spirodela polyrrhiza*. Bull. Torr. Club 2, 46-47.
- ENGLANDER A.J. and KAIGATE B., 1981: Removal of persistent heavy metals by vascular aquatic plant systems. Ann. Meet. Am. Inst. Chem. Energy., New Orleans.
- ENGLER A., 1877: Vergleichende Untersuchungen über die morphologischen Verhältnisse der Araceae. Nova Acta K. Leop.-Carol. Deutsch. Akad. Natf. 39, 135-224.
- ENGLER A., 1889: Lemnaceen. In: ENGLER A. and PRANTL K. (eds.), Natürliche Pflanzenfamilien II. 3, 154-164.
- ENGLER A., 1920: Araceae-Pistioideae. In: ENGLER A., Das Pflanzenreich IV. Engelmann, Leipzig. 23 F, 250-262.
- ENTZERTH M., MEAD D.J., PATTERSON G.M.L. and MOORE R.E., 1985: A herbicidal fatty acid produced by *Lyngbya aestuarii*. Phytochem. 24(12), 2875-2876.
- ENU-KWESI L. and DUMBROFF E.B., 1978: Changes in abscisic acid in the embryo and covering structures of *Acer saccharum* during stratification. Z. Pflanzenphysiol. 86, 371-377.
- EPSTEIN E. and BENTAL Y., 1985: Involvement of indoleacetic acid in the promotion of flowering of *Lemna gibba* G3. Abstr. 12th Int. Conf. Plant Growth Substances, Heidelberg, 7.
- EPDTWAN G., 1952: Pollen morphology and plant taxonomy: I. Angiosperms. Almquist and Wiksells. 539 pp.

- ERICKSON L.C. and WEDDING R.T., 1956: Effects of ozonated hexene on photosynthesis and respiration of Lemna minor. *Am.J. Bot.* 43, 32-36.
- ERICSSON T., LARSSON C.M. and TILLBERG E., 1982: Growth responses of Lemna to different levels of nitrogen limitation. *Z. Pflanzenphysiol.* 105(4), 331-340.
- ERISMANN K.H., 1972: Untersuchungen zur Analyse des CO₂-Outburst bei der Photorespiration von Lemna minor. *Verh. Schweiz. Naturf. Ges.* 152, 151-153.
- ERISMANN K.H. and BRUNOLD C., 1967: Die Probeentnahme in kinetischen Stoffwechseluntersuchungen mit Wasserlinse Lemna minor L. (*Lemnaceae*). *Experientia* 23, 235.
- ERISMANN K.H. and BRUNOLD C., 1973: Die Verwendung einer neuen Lemna-Kulturanlage in Wachstums- und Stoffwechseluntersuchungen mit gelösten und gasförmigen Schwefelverbindungen. *Ber. Schweiz. Bot. Ges.* 83, 213-222.
- ERISMANN K.H. and FANKHAUSER M., 1967: Die Veränderung des Stärke-, Protein- und RNS-Gehaltes von Lemna minor unter dem Einfluss von Kinetin (6-Furfurylaminopurin). *Experientia* 23, 621-622.
- ERISMANN K.H. and FINGER A., 1968: Lemnaceen in kontinuierlicher Kultur. *Ber. Schweiz. Bot. Ges.* 78, 5-15.
- ERISMANN K.H. and KIRK M.R., 1967: Kinetische Untersuchungen zur Säure- und Proteinsynthese durch Lemna minor L. im Licht unter steady-state-Bedingungen. *Verh. Schweiz. Naturf. Ges.* 147, 141-143.
- ERISMANN K.H. and KIRK M.R., 1969: The influence of nitrogen source on metabolic intermediates in steady-state photosynthesis by Lemna minor L. In: METZNER H. (ed.), *Progress Photosynth. Res.*, Tübingen 3, 1538-1545.
- ERISMANN K.H. and MARTI J., 1986: Untersuchung zur Unterdrückung der Photorespiration bei der Wasserlinse Lemna minor L. *Schweiz. Bot. Ges.*, *Beil. Bull.* 4, 8-9.
- ERISMANN K.H., MARTI J. and RUTTER J.C., 1986: Nitrogen nutrition and photorespiration in the C3-plant Lemna minor. *Schweiz. Bot. Ges.*, *Beil. Bull.* 4, 7-8.
- ERISMANN K.H., STRASSER R. and BRAENDLE R., 1967: Untersuchungen zur Photosyntheseabhängigen Sulfidoxydation durch Algen und höhere Pflanzen, insbesondere durch Lemna minor L. *Verh. Schweiz. Naturf. Ges.* 147, 139-141.
- ERISMANN K.H. and WEGNER F., 1967: Der Einfluss einer wachstumshemmenden Kinetinkonzentration auf Chlorophyllgehalt, Photosyntheserate und Stärkeproduktion von Lemna minor L. *Flora* 158, 433-442.
- ERNST W.H.O. and MARQUENIE-VAN DER WERFF M., 1978: Aquatic angiosperms as indicators of copper contamination. *Arch. Hydrobiol.* 83(3), 356-366.
- ESASHI Y. and ODA Y., 1964: Effect of light intensity and sucrose on the flowering of Lemna perpusilla. *Plant Cell. Physiol.* 5, 513-516.
- ESASHI Y. and ODA Y., 1966: Two light reactions in the photoperiodic control of flowering of Lemna perpusilla and L. gibba. *Plant Cell Physiol.* 7, 59-74.
- ESASHI Y., SHIBASAKI T. and SAITO K., 1972: Flowering responses of Lemna perpusilla and L. gibba in relation to nitrate concentration in the culture medium. *Plant Cell Physiol.* 13, 623-631.
- ESHEL A. and BEER S., 1986: Inorganic carbon assimilation by Spirodela polyrrhiza. *Hydrobiologia* 131(2), 149-153.
- ESKUCHE U., 1986: Bericht über die 17. Internationale pflanzengeographische Exkursion durch Nordargentinien (1983). *Veröff. Geobot. Inst. ETH, Stiftung Rübel, Zürich* 91, 12-117.

- ESKUCHE U. and ROMERO FONSECA L., 1982: Contribucion a la biologia de Wolffia linguata (Lemnaceae). *Bol. Soc. Argent. Bot.* 21, 259-268.
- EVANS D.M., 1978: Aquatic weed control with the isopropylamine salt of N-phosphonomethyl glycine. *Proc. European Weed Res. Soc. Symp. Aquatic Weeds* 5, 171-178.
- EVANS G.C., 1972: The quantitative analysis of plant growth. Blackwell, Oxford. 186-254.
- EVANS I.T., 1969: The induction of flowering, some case histories. McMillan Co., South Melbourne. 488 pp.
- EVANS O.D., 1970: Some observations on the Lemnaceae or "Duckweeds" of New South Wales. *Contrib. N.S. Wales Nat. Herb.* 4, 87-94.
- EVANS S. and WALKER J.R.L., 1978: Effect of quaternary ammonium compounds on some aquatic plants. *Marine Pollut. Bull.* 9(5), 136-137.
- EVERARD M. and DENNY P., 1985: Flux of lead in submerged plants and its relevance to a freshwater system. *Aquat. Bot.* 21, 181-193.
- EVERSULL H., ACHACOSO A.S. and SORBERT R.H., Jr., 1980: Ensiling duckweed with high dry matter corn crop. *Dairy Sci.* 65(suppl.1), 140.
- EWEL K.C., 1984: Effects of fire and wastewater on understory vegetation in cypress domes. In: EWEL K.C. and ODUM H.T. (eds.), *Cypress swamps*. Univ. Press Florida. 119-126.
- EWEL K.C. and ODUM H.T., 1978: Cypress swamps for nutrient removal and wastewater recycling. In: WANIELISTA M.P. and ECKENFELDER W.W., Jr. (eds.), *Advances in water and wastewater treatment: Biological nutrient removal*. Ann Arbor Sci. Publishers. 181-198.
- EYSTER C., 1966: Optimization of the physiological activity (growth rate) of the giant duckweed, Spirodela polyrrhiza. *Final Report*. Monsanto Res. Corp. Dayton Lab., Dayton, Ohio. 200 pp. (Polycopy).
- EYSTER C., 1978: Nutrient concentration requirements for Chlorella sorokiniana. *Ohio J. Sci.* 78, 79-81.
- EYSTER C., 1981: Mineral nutrient requirements for the growth of the giant duckweed, Spirodela polyrrhiza. *Proc. Int. Bot. Congr.* 13, 168.
- EYSTER C., BROWN T.E., TANNER H.A. and HOOD S.L., 1958: Manganese requirement with respect to growth, Hill reaction and photosynthesis. *Plant Physiol.* 33, 235-241.
- FAERBER E., 1984: Die Dichte der Pflanzendecke als Faktor für den Aethylenhaushalt bei Lemna. *Mitt. band Botaniker-Tagung Wien* 42 (0405).
- FAERBER E., KOENIGSHOFER H. and KANDELER R., 1986: Ethylene production and overcrowding in Lemnaceae. *J. Plant Physiol.* 124(3-4), 379-384.
- FAGERLIND F. and MASSALSKI A., 1974: The development of cell walls and intercellulars in the root of Lemna minor L. *Sven. Bot. Tidskr.* 68, 64-93.
- FALTER C.M. et al., 1974: Aquatic macrophytes of the Columbia and Snake River drainages (US). U.S. Army Corps of Engrs., Walla Walla Distr.
- FANKHAUSER H., 1975: Untersuchung von Auswirkungen subtoxischer Schwefeldioxid-Konzentrationen auf Wachstum und Stoffproduktion bei Lemna minor L. unter Standardbedingungen. Lic., Univ. Berne. (Polycopy).
- FANKHAUSER H., BRUNOLD C. and ERISMANN K.H., 1974: Der Einfluss von SO₂ im Bereich der Grenzkonzentration auf die Netto-Stoffproduktion bei Lemna minor L. *Verh. Schweiz. Naturf. Ges.* 154, 198-199.
- FANKHAUSER H., BRUNOLD C. and ERISMANN K.H., 1976: The influence of sublethal concentrations of sulfur dioxide on morphology, growth and production yield of the duckweed Lemna minor L. *Oecologia* 23, 201-209.
- FANKHAUSER M. and ERISMANN K.H., 1967: Durch Kinetin induzierte Veränderungen im Protein- und Nukleinsäure-Stoffwechsel von Lemna minor L. *Verh. Schweiz. Naturf. Ges.* 147, 141.

- FEOLI E. and GERDOL R., 1982: Evaluation of syntaxonomic schemes of aquatic plant communities by cluster analysis. *Vegetatio* 49, 21-27.
- FERGUSON A.R., 1966: Responses of *Spirodela oligorhiza* to changes in its nutrient environment. M.S. Thesis. Univ. Ruckland, N.Z. (Polycopy).
- FERGUSON A.R., 1969a: The nitrogen metabolism of *Spirodela oligorhiza*. II. Control of the enzymes of nitrate assimilation. *Planta* 88, 353-363.
- FERGUSON A.R., 1969b: Control of nitrate utilization in a higher plant. Abstr. XI Int. Bot. Congr. and Int. Wood Chemistry Symposium, Seattle. 59.
- FERGUSON A.R., 1970: Nitrogen metabolism of *Spirodela oligorhiza*. III. Amino acids and the utilization of nitrate. *Planta* 90, 365-369.
- FERGUSON A.R. and BOLLARD E.G., 1969: Nitrogen metabolism of *Spirodela oligorhiza*. I. Utilization of ammonium, nitrate and nitrite. *Planta* 88, 344-352.
- FERGUSON A.R. and KNYPL J.S., 1974: Specificity of induction of nitrate reductase in plants. In: WEHRMANN J. (ed.), Plant analysis and fertilizer problems. Proc. 7th Int. Coll. Plant Analysis and Plant Fertilizer Problems, Hannover. 1, 101-109.
- FERGUSON I., TURNER N. and BOLLARD E.G., 1980: Problems in fractionating calcium in plant tissue. *J. Sci. Food Agric.* 31, 7-14.
- FERNALD M.L., 1941: Another century of additions to the Flora of Virginia. *Rhodora* 43, 547.
- FERNANDEZ D.B. and BALDOS S.C., 1981: On water pollution: I. Absorption of nitrates by four aquatic plants. *WSSP Newsletter* 9(2), 5.
- FERNANDEZ D.B., BALDOS S.C. and BELTRAN M.M., 1983: Absorption of inorganic pollutants by four aquatic plants. *NRCP Res. Bull.* 38(1), 1-37.
- FERNANDEZ N.A. and ATHIAS-BINCHE F., 1986: Analyse démographique d'une population d'*Hydrozotes lemnae* Coggi, Acarien Oribate inféodé à la lentille d'eau *Lemna gibba* en Argentine. I. Méthodes et techniques, démographie d'*H. lemnae* et comparaison avec d'autres Oribates. *Zool. Jb. Syst.* 113(2), 213-228.
- FERNANDEZ O.A. and MUJICA B., 1973: Effects of some environmental factors on the differentiation of stomata in *Spirodela intermedia* W. Koch. *Bot. Gaz.* 134, 117-121.
- FERNANDEZ O.A., ORDONEZ M.E., CURVETTO N., FERNANDEZ L.M. and RIGATO M., 1972: Absorción del ácido 2,4-diclorofenoxiacético por *Spirodela intermedia* W. Koch. *Phyton (Argentina)* 29, 105-117.
- FERRARA L., FORGIONE P., SCETTINO O. and RULLO V., 1985: Utilizzazione di piante acquatiche nella correzione dell'inquinamento idrico da agenti chimici. Assorbimento di zirconio e cadmio. *Boll. Soc. Ital. Biol. Sper.* 61(9), 1343-1348.
- FERRERA R.B. and DAVIES D.D., 1986: Is protein degradation correlated with either the charge or size of Lemna proteins? *Planta* 169(2), 278-288.
- FERRERA R.B. and DAVIES D.D., 1987a: Protein degradation in Lemna with particular reference to ribulose biphosphate carboxylase I. The effect of light and dark. *Plant Physiol.* 83(4), 869-877.
- FERRERA R.B. and DAVIES D.D., 1987b: Protein degradation in Lemna with particular reference to ribulose biphosphate carboxylase II. The effect of nutrient starvation. *Plant Physiol.* 83(4), 878-883.
- FILBIN G.J. and HOUGH R.A., 1979: The effects of excess copper sulfate on the metabolism of the duckweed *Lemna minor*. *Aquat. Bot.* 7, 79-86.
- FILBIN G.J. and HOUGH R.A., 1984: Extraction of C-labeled photosynthesate from aquatic plants with dimethyl sulfoxide (DMSO). *Limnol. Oceanogr.* 29(2), 426-428.

- FANKHAUSER M. and ERISMANN K.H., 1969a: Ueber den Einfluss des Kinetins (6-Purifyl-Amino-Purin) auf den Zuckergehalt von *Lemna minor* L. *Flo-ra* A 160, 342-349.
- FANKHAUSER M. and ERISMANN K.H., 1969b: Ueber die Wirkung kurzfristiger Kinetinaben auf Protein, Aminosäuren und RNS bei *Lemna minor*. *Planta* 88(4), 332-343.
- FARAGO M.E. and PARSONS P.J., 1985: The recovery of platinum metals by the water hyacinth. *Environ. Techn. Lett.* 6, 165-174.
- FASSETT C.N., 1972: A manual of aquatic plants. 6th ed. Univ. Wisc. Press Madison. 405 pp.
- FEDER W.A. and SULLIVAN F., 1969: Ozone: depression of frond multiplication and floral production in duckweed. *Science* 165(3900), 1373-1374.
- FEDER W.A., MIKA J.S., VARDARO P. and PERKINS I., 1980: The biological activity of the O₃ protectant, DPX. *Phytopathology* 70(5), 462.
- FEKETE A., 1973: The release of phosphorus from pond sediments and its availability to *Lemna minor* L. New Brunswick, N.J., Rutgers Univ., Springfield, Va. 94 pp.
- FEKETE A. and RIEMER D.N., 1973: Effects of varying phosphorus concentration on *Lemna minor* L. *Coll. Agric. Environ. Sci., Rutgers Univ.*, New Brunswick, N.J., 27, 109-114.
- FEKETE A., RIEMER D.N. and MOTTO H.L., 1976: A bioassay using common duckweed to evaluate the release of available phosphorus from pond sediments. *J. Aquat. Plant Manage.* 14, 19-25.
- FELDMANN A., 1968: Beiträge zur Strahlenstimulation. I. Versuche zu einer Reproduktion von Stimulationserscheinungen bei *Lemna minor* L. *Radiat. Bot.* 8, 425-437.
- FELDMANN A., 1969: Beiträge zur Strahlenstimulation. II. Ueber den Einfluss der Tageslänge auf die strahleninduzierte Wachstumsförderung bei *Lemna minor* L. *Radiat. Bot.* 9, 459-471.
- FELDMANN A., 1971: Beiträge zur Strahlenstimulation. III. Einfluss der Temperatur auf die strahleninduzierte Wachstumsförderung bei *Lemna minor* L. *Radiat. Bot.* 11, 59-65.
- FELDMANN A., 1975: Beiträge zur Strahlenstimulation. V. Der Einfluss von Licht- und Temperaturänderungen sowie des Bestrahlungszeitpunktes auf die strahleninduzierte Wachstumsförderung bei *Lemna minor* L. *Radiat. Bot.* 15, 49-58.
- FELLE H. and BERTL A., 1986: The fabrication of proton-selective liquid-membrane micro-electrodes for use in plant cells. *J. Exp. Bot.* 37(182), 1416-1428.
- FELLER U., 1975: Untersuchungen zur Ermittlung von Angriffspunkten der Stoffwechselregulation durch Ammonium und Nitrat bei *Lemna minor* L. *Diss. Univ. Bern.* (Polycopy).
- FELLER U. and ERISMANN K.H., 1971: Einfluss der Beleuchtungsstärke auf die Ammonium- und Nitrataufnahme bei *Lemna minor*. *Verh. Schweiz. Natf. Ges.* 151, 96-99.
- FELLER U. and ERISMANN K.H., 1973: Wechselwirkungen zwischen Stickstoffquelle und Ionenhaushalt bei *Lemna minor* L. unter Photosynthesebedingungen. *Verh. Schweiz. Natf. Ges.* 153, 75-79.
- FELLER U. and ERISMANN K.H., 1976: Einfluss der Aminosäuren Ornithin, Citrullin und Arginin auf das Wachstum von *Lemna minor* bei gleichzeitigem Angebot von Ammonium oder Nitrat. *Ber. Schweiz. Bot. Ges.* 86, 129-135.
- FELZINES J.C., 1977: Analyse des relations entre la minéralisation des eaux douces stagnantes et la distribution des végétaux qui les peuplent. *Ann. Sci. Nat., Bot. Biol. Vég., Paris, 12e sér.* 18, 221-249.

- FILBIN G.J. and HOUGH R.A., 1985: Photosynthesis, photorespiration, and productivity in Lemna minor L. *Limnol. Oceanogr.* 30(2), 322-334.
- FILNER P., 1966: Regulation of nitrate reductase in cultured tobacco cells. *Biochim. Biophys. Acta* 118, 299-310.
- FINKLE L.R. and SBELEY H.W., 1978: Nitrogen fixation (acetylene reduction) by epiphytes of freshwater macrophytes. *Appl. Environ. Microbiol.* 36, 129-138.
- FINLAYSON M., CHICK A., VAN OERTZEN I. and MITCHELL D., 1987: Treatment of piggery effluent by an aquatic plant filter. *Biol. Wastes* 19(3), 179-196.
- FINTHA I., 1979: Revision of the home distribution of *Wolffia arrhiza* (L.). *Tiscia* 14, 71-79.
- FISCHER E. and LUETTGE U., 1980: Membrane potential changes related to active transport of glycine in *Lemna gibba* Gl. *Plant Physiol.* 65, 1004-1008.
- FISCHER E. and NOVACKY A., 1980: Effect of PCMBs and DCCD on membrane potential of *Lemna gibba* Gl. *Dev. Plant Biol.* 4, 553-554.
- FISCHER G.W., 1953: *Manual of North American smut fungi*. Ronald Press, New York. 343 pp.
- FISCHER H., 1949: Plasmolyseform und Mineralsalzgehalt in alternierenden Blättern. II. Untersuchungen an Land- und Schwimmpflanzen. *Planta* 37, 244-292.
- FISCHER S., 1981: Rapid toxicological test using the duckweed *Lemna minor*. (In Swedish). *Inst. Vatten-Luftvaardsforsk. IVL B-641*, 27 pp.
- FISCHER W., 1957: *Unsere einheimischen Wasserlinsen*. Märkische Heimat 2. Potsdam.
- FISCHER Z., 1968: Food selection in grass carp *Ctenopharyngodon idella* under experimental conditions. *Pol. Arch. Hydrobiol.* 15(1), 1-8.
- FITZGERALD G.P., 1968: Detection of limiting or surplus nitrogen in algae and aquatic weeds. *J. Phycol.* 4(2), 121-126.
- FITZGERALD G.P., 1969: Some factors in the competition of antagonism among bacteria, algae, and aquatic weeds. *J. Phycol.* 5(4), 351-359.
- FITZGERALD G.P., 1971: Comparative rates of phosphorus sorption and utilization by algae and aquatic weeds. *J. Phycol.* 7(suppl.), 11.
- FITZGERALD V.M., 1918: The botany of the Kimberleys, North-west Australia. *J. Proc. R. Soc. West-Australia* 3, 102-224.
- FIUSSELLO N., 1973: Lead pollution effects on chlorophyll. *Inf. Bot. Ital.* 5(1), 107-108.
- FLETCHER F. and ARNOTT H.J., 1963: The venation of *Lemna minor* in sterile culture. *Am. J. Bot.* 50, 621.
- FLETCHER F. and ARNOTT H.J., 1964: Statistical analysis of *Lemna minor* venation. *Am. J. Bot.* 51, 675.
- FLETCHER N.F., 1965: An experimental study of the venation of *Lemna minor* L. *Diss. Abstr.* 25(11), 6178.
- FLORES A.Q., 1981 s. QUIROZ F.A., 1981
- FLORES H.E. and GALSTON A.W., 1981: Polyamines in plants. Determination of microquantities by thin-layer chromatography and high performance liquid chromatography. *Plant Physiol.* 67(4 suppl.), 111.
- FLORES H.E. and GALSTON A.W., 1982: Analysis of polyamines in higher plants by high performance liquid chromatography. *Plant Physiol.* 69(3), 701-706.
- FLORES S. and TOBIN E.M., 1985: Cytokinin-modulated regulation of mRNA levels for the chlorophyll a/b protein in *Lemna*. *J. Cell Biochem. (suppl. 9 part C)*, 253.
- FLORES S. and TOBIN E.M., 1986a: Post-transcriptional regulation of LHCP gene expression by benzyladenine. *J. Cell Biochem. (suppl. 10 part B)*, 21.

- FLORES S. and TOBIN E.M., 1986b: Benzyladenine modulation of the expression of two genes for nuclear-encoded chloroplast proteins in *Lemna gibba*: Apparent post-transcriptional regulation. *Planta* 168(3), 340-349.
- FLUHR R., KUHLEMEIER C., NAGY P. and CHUA N.-H., 1986: Organ-specific and light-induced expression of plant genes. *Science* 232(4754), 1106-1112.
- FLY C.L., 1935: Organic iron and hydrogen-ion concentration as associated factors affecting the rate of reproduction of *Lemna major*. *Oklahoma Acad. Sci. Proc.* 15, 77-80.
- FOCHT J., 1972: Additions to the flora of Rockland County, New York. *Bull. Torrey Bot. Club* 99(5), 249-250.
- FONTAINE S., ALIBERT G. and BOUDET A.M., 1978: Antagonisme métabolisme protéique - métabolisme phénolique chez les végétaux supérieurs. *Bull. Liaison, Groupe Polyphenols* 8, 91-105.
- FOOTE R.H. and COOK D.R., 1959: Mosquitoes of medical importance. *US Dept. Agric., Washington DC, Agric. Handbook* 152, 158 pp.
- FORSLUND M. and RUNDLOEF S., 1986: *Lemna minor* new record and *Schoenus ferrugineus* new record in Asele, Lappland, Northern Sweden. (In Swedish). *Sven. Bot. Tidskr.* 80(3), 185.
- FOSSBERG F.R. and FALANRUM M.V.C., 1980: Noteworthy micronesian plants. 4. *Micronesica* 16(2), 201-210.
- FOURST G.G., 1968: Anatomy and structure of some aquatic plants. (In Russian). *Bull. Bot. Zh. (USSR)* 71, 67-74.
- FOWLER M.C., 1985: The results of introducing grass carp, *Ctenopharyngodon idella* Val., into small lakes. *Aquacult. Fish. Manage.* 16, 189-202.
- FOX R.L. and ALBRECHT W.A., 1958: Calcium-boron interactions - demonstrated by *Lemna minor* on clay suspensions. *Univ. Missouri Agr. Expt. Sta. Res. Bull.* 663, 15 pp.
- FOY C.L. and COOLEY W.E., 1987: Studies on comparative modes(s) of herbicidal action of SC-0224 and glyphosate. 14th Int. Bot. Congr. Berlin, Abstr., 71.
- FRANK E.E., 1938: Plant promoting substances on vitamin content and reproduction of *Lemna*. *Iowa State Coll.* 13, 63-66.
- FRANCESCHI V.R., 1985: Pathways for formation of oxalate in *Lemna minor*. *Plant Physiol.* 77(4 suppl.), 25.
- FRANCESCHI V.R., 1987: Calcium oxalate formation in *Lemna minor* is a rapid and reversible process. *Plant Physiol.* 83(4 suppl.), 42.
- FRANCESCHI V.R. and SCHUREN A.M., 1986: Incorporation of strontium into plant calcium oxalate crystals. *Protoplasma* 130(2-3), 199-205.
- FRANCHET A., 1864: Note sur le mode de reproduction de la *Bruniera vivipara* (*Lemna arrhiza* L.). *Biblioth. 1*, 25-31.
- FRANK P.A., 1975: Competitive interactions among aquatic plants. In: BREZONIK P.L. and FOX J.L. (eds.), *Proc. Symp. Water Quality Management through biological control*. Rept. No. ENV-07-75-1, 24-27.
- FRASER G.G., 1974: Sulfur nutrition in duckweed, *Spirodela oligorrhiza*. *Mauri Ora* 2, 147-156.
- FREDERICK S.E., GRUBER P.J. and TOLBERT N.E., 1973: The occurrence of glycolate dehydrogenase and glycolate oxidase in green plants. An evolutionary survey. *Plant Physiol.* 52, 318-323.
- FREEMAN T.E., 1975: Rhizoctoniosis of aquatic plants. In: MCGRAW (ed.), *Hill Encyclopedia. Sci. Technol. Yearbook*, 7-28.
- FREY M., RALL S., ROTH A. and HEMLEBEN V., 1980: Evidence for uptake of plasmid DNA into intact plants (*Lemna perpusilla*) proved by an E. coli transformation assay. *Z. Naturforsch. C* 35, 1104-1106.

- FRICK H., 1972: A study of inhibitor-induced bleaching in *Lemna minor*. *Diss. Abstr. Int.* B 33, 4670.
- FRICK H., 1975: Phytochrome control of the lag phase of chlorophyll accumulation in *Lemna minor*. *Can. J. Bot.* 53, 2405-2410.
- FRICK H., 1978: Pyrimidine metabolism in *Lemna minor*. 1. Functional compartmentation of chloroplast pyrimidine metabolism in a higher plant. *Plant Physiol.* 61, 989-992.
- FRICK H., 1985a: Boron tolerance and accumulation in the duckweed, *Lemna minor*. *J. Plant Nutrition* 8(12), 1123-1129.
- FRICK H., 1985b: Micronutrient tolerance and accumulation in the duckweed, *Lemna*. *J. Plant Nutrition* 8(12), 1131-1145.
- FRICK H. and JONES R.F., 1975: Inhibition of chlorophyll synthesis in *Lemna minor* by nalidixic acid. *Can. J. Bot.* 53, 2319-2324.
- FRICK H. and JONES R.F., 1976: Physiology and plastid fine structure of deetiolating *Lemna minor*. *Can. J. Bot.* 54, 1819-1826.
- FRICK H. and MOHR H., 1973: Phytochrome-mediated growth responses in green and etiolated *Lemna minor*. *Planta* 109, 281-292.
- FRITS E.M., 1985: Angiosperm fruits and seeds from the middle miocene of Jutland (Denmark). *Biol. Skr. Dan. Vid. Selsk.* 24, 165 pp.
- FROEMING E., 1952: Ueber das Verhalten unserer Wassertschnecken gegenüber Lemnaceen. *Arch. Molluskenkunde* 81, 45-48.
- FROMM F., 1943: Growth stimulation by ammonium sulfamate in low concentration. *Science* 98, 391-392.
- FROMM F., 1946: El empleo de *Lemna minor* L. en ensayos rápidos de toxicidad. *Ciencia (Mex.)* 7, 214-218.
- FROMM F., 1948(1949): La acción del ácido furilacrilico sobre *Spirodela polyrrhiza*. *Ciencia (Mex.)* 9, 40-42.
- FROMM F., 1951: A quantitative evaluation of the *Lemna* test for herbicides. *Bot. Gaz.* 113, 86-90.
- FROMM F., 1955: La acción de ácidos acrílicos substituidos sobre *Lemna minor*. *Acta Cienc. Venezolana* 6, 121-124.
- FROMM F., 1958: The toxicity of some amines for duckweed, *Lemna minor*. *Ohio J. Sci.* 58, 354-356.
- FROMM F., 1960: A modification of the *Lemna* test for phytotoxicity. *J. Agric. Univ. Puerto Rico* 44, 93-102.
- FROMM F. and O'DONNELL M.L., 1951: The action of $-SO_3NH_2$ derivatives on duckweed. *Proc. Pennsylv. Acad. Sci.* 25, 85-88.
- FROMM F. and O'DONNELL M.L., 1952: The influence of p-aminobenzoic acid on the growth of duckweed (*Lemna minor*). *Proc. Pennsylv. Acad. Sci.* 26, 50-53.
- FROMM F. and O'DONNELL M.L., 1953: Acción simultánea del ácido p-aminobenzoico y derivados del grupo SO_3NH_2 sobre lentejas de agua (*Lemna minor*). *Acta Cienc. Venezolana* 4, 66-67.
- FROMM F. and O'DONNELL M.L., 1955: The action of tauramide, 2-acetyl-amino-1,3,4-thiadiazole-5-sulfonamide, and N-substituted sulfanilamides on duckweed, *Lemna minor*. *Proc. Pennsylv. Acad. Sci.* 29, 135-140.
- FROMM F. and PACE A., 1957: The phytotoxicity of 2-amino-1,3,4-thiadiazole-5-sulfonamide. *Bol. Col. Quím. Puerto Rico* 14(1).
- FROMM F., VIDAL I.M. and COLEMAN J., 1949: The action of herbicides on Lemnaceae. *Proc. Pennsylv. Acad. Sci.* 23, 85-90.
- FROMM H., DEVIC M., FLUHR R. and EDELMAN M., 1985: Control of PSB-A gene expression in mature *Spirodela oligorrhiza* chloroplasts. Light regulation of 32-kilodalton protein synthesis is independent of transcript level. *EMBO J.* 4(2), 291-296.
- FRYE J.B. and CULLEY D.D., 1980: A feasibility study of the conversion of animal feedlot wastes to useful energy in Louisiana. Final Report 1978-1980. Baton Rouge. 11 pp. (Polycopy).

- FUHRER J., 1983: Light-inhibition of dark respiration in *Lemna minor*. *Bot. Helv.* 93(1), 67-75.
- FUHRER J., 1984: The effect of CO_2 on the conversion of l-aminocyclopropane-l-carboxylic acid (ACC) in *Lemna minor* L. *Plant Physiol.* 75 (1 suppl.), 127.
- FUHRER J., 1985: Production and release of ethylene from l-aminocyclopropane-l-carboxylic acid in *Lemna minor* L. in the dark and at different carbon dioxide compensation concentrations. *J. Plant Physiol.* 117(4), 307-317.
- FUHRER J. and ERISMANN K.H., 1983: Net CO_2 assimilation and dry weight accumulation in *Lemna minor* L. *Plant Physiol.* 72(1 suppl.), 10.
- FUHRER J. and ERISMANN K.H., 1984: Steady-state carbon flow in photosynthesis and photorespiration in *Lemna minor* L.: The effect of temperature and ammonium nitrogen. *Photosynthetica* 18(1), 74-83.
- FUJIKAWA R. and KYO S., 1973: Efficacy of chlomethoxymil in rice. *Proc. 4th Asian Pacific Weed Science Soc. Conf., Rotorua.* 130-133.
- FUJIOKA S., YAMAGUCHI I., MUROFUSHI N., TAKAHASHI N., KAIHARA S. and TAKIMOTO A., 1983a: Flowering and endogenous levels of benzoic acid in *Lemna* species. *Plant Cell Physiol.* 24(2), 235-239.
- FUJIOKA S., YAMAGUCHI I., MUROFUSHI N., TAKAHASHI N., KAIHARA S. and TAKIMOTO A., 1983b: The role of plant hormones and benzoic acid in flowering of *Lemna paucicostata* 151 and 381. *Plant Cell Physiol.* 24(2), 241-246.
- FUJIOKA S., YAMAGUCHI I., MUROFUSHI N., TAKAHASHI N., KAIHARA S., TAKIMOTO A. and CLELAND C.F., 1985: The role of benzoic acid and plant hormones in flowering of *Lemna gibba* G3. *Plant Cell Physiol.* 26(4), 655-659.
- FUJIOKA S., YAMAGUCHI I., MUROFUSHI N., TAKAHASHI N., KAIHARA S., TAKIMOTO A. and CLELAND C.F., 1986a: Isolation and identification of nicotinic acid as a flower-inducing factor in *Lemna*. *Plant Cell Physiol.* 27(1), 103-108.
- FUJIOKA S., YAMAGUCHI I., MUROFUSHI N., TAKAHASHI N., KAIHARA S., TAKIMOTO A. and CLELAND C.F., 1986b: The influence of nicotinic acid and plant hormones on flowering in *Lemna*. *Plant Cell Physiol.* 27(1), 109-116.
- FUJIOKA S., SAKURAI A., SUGIMOTO N., YAMAGUCHI I., MUROFUSHI N., TAKAHASHI N., KAIHARA S. and TAKIMOTO A., 1986c: Isolation and identification of lutein from *Lemna paucicostata* 381 as an inhibitor of benzoic acid-induced flowering in *Lemna paucicostata* 151. *Agric. Biol. Chem.* 50(8), 2053-2059.
- FUJIOKA S., SAKURAI A., YAMAHUCHI I., MUROFUSHI N., TAKAHASHI N., KAIHARA S., TAKIMOTO A. and CLELAND C.F., 1986d: Flowering and endogenous levels of plant hormones in *Lemna* species. *Plant Cell Physiol.* 27(7), 1297-1307.
- FULLER G. and STUMPF P.K., 1984: Lipid biosynthesis in *Lemna minor*. *Plant Physiol.* 75(1 suppl.), 101.
- FULLER G. and STUMPF P.K., 1987: Fatty acids in plants: a model system. *ACS Symp. Ser.* 325, 44-52.
- FULLER R.C., ANDERSON I.C. and NATHAN H.A., 1958a: Pteridines in photosynthesis. *Proc. Natl. Acad. Sci. US* 44, 239-244.
- FULLER R.C., ANDERSON I.C. and NATHAN H.A., 1958b: Pteridines in photosynthesis. An artifact of paper chromatography. *Proc. Natl. Acad. Sci. US* 44, 518-519.
- FUNDERBURN H.H. and LAWRENCE J.M., 1963: A sensitive method for the determination of low concentrations of diquat and paraquat. *Nature* 199, 1011-1012.

- FURLOW B.M. and HAYS K.L., 1972: Some influences of aquatic vegetation on the species and number of Culicidae (Diptera) in small pools of water. *Mosquito News* 32(4), 595-599.
- FURST G.G., 1968: The anatomical structure of some aquatic plants. (In Russian). *Byull. Gl. Bot. Sada* 71, 67-74.
- FURUYA M. and THIMANN K.V., 1963: Effects of gibberellic acid on anthocyanin synthesis and growth in *Spirodela*. *Plant Physiol.* 38 (suppl.), 44.
- FURUYA M. and THIMANN K.V., 1964: The biogenesis of anthocyanins. XI. Effects of gibberellic acid in two species of *Spirodela*. *Arch. Biochem. Biophys.* 108, 109-116.
- GABA V., MARDER J.B., GREENBERG B.M., MATTOO A.K. and EDELMAN M., 1987: Degradation of the 32-kDa herbicide binding protein in far red light. *Plant Physiol.* 84(2), 348-352.
- GABALDON A., ULLOA G. and PULIDO J., 1981: Distribucion geografica, ecologia y etologia de *Aedeomyia squamipennis*, importante vector natural de malaria aviaria en Venezuela. *Bol. Direc. Malariologia y Saneamiento Ambiental* 21(2), 103-113.
- GABALDON et al. 1983 see 1981
- GABRYS H., 1978: The application of the interference microscopy for the refractive index determination of the cell wall and cytoplasm in plant cells. *Microsc. Acta* 80(3), 215-218.
- GABRYS H., WALCZAK T. and ZURZYCKI J., 1981: Chloroplast translocations induced by light pulses. Effects of single light pulses. *Planta* 152(6), 553-556.
- GABRYS-MIZERA H., 1976a: Refractive index determination of the cell wall and cytoplasm by interference microscopy. *Bull. Acad. Pol. Sci. Biol.* 24, 181-184.
- GABRYS-MIZERA H., 1976b: Model considerations of the light conditions in non-cylindrical plant cells. *Photochem. Photobiol.* 24, 453-461.
- GRAEVSKAYA N.S., 1969: The role of higher aquatic plants in the nutrition of the animals of freshwater basins. (In Russian, translated by MUELLER G.M.). *Natl. Lending Library Sci. Technol.* 3 vols. 629 pp.
- GAIGHIER I.G., FORATH D. and GRANTH G., 1984: Evaluation of duckweed (*Lemna gibba*) as feed for Tilapia (*Oreochromis niloticus* x *O. aureus*) in a recirculating unit. *Aquaculture* 41(3), 235-244.
- GALKINA N.V., ABDULAEV D.A. and ZACHAROVA V.L., 1965: Biological and feed features of duckweeds. (In Russian). *Uzb. Biol. Zh.* 3, 44-47.
- GANCARZ R., WIELKOPOLSKI W., JASKULSKA E., KAFARSKI P., LEJCZAK B., MASZALCZAK P. and WIECZOREK J.S., 1985: Phosphonic analogs of morphactins 4. 9-aminofluoren-9-ylphosphine oxides. *Pestic. Sci.* 16(3), 234-238.
- GANGSTAD E.O., SEAMAN D.E. and NELSON M.L., 1972: Potential growth of aquatic plants of the Lower Mekong River Basin, Laos, Thailand. *Hyacinth Control* J. 10, 4-13.
- GANNING B. and WULF F., 1970: Measurements of community metabolism in some Baltic brackish water rockpools by means of dial oxygen curves. *Oikos* 21, 292-298.
- GAPONENKO V.I. and SPAZHETSKII V., 1969: Change in the rate of photosynthesis and chlorophyll content in duckweed in connection with age and conditions of illumination. *Sov. Plant. Physiol.* 16(6), 825-832.
- GARANOV S.A., 1965: Lemnaceae; Vermehrung in Massenkulturen. (In Russian). *VNIRO* 10.
- GASPARRINI G., 1856: Osservazioni morfologiche sopra taluni organi della *Lemna minor*. Napoli.
- GAUDINET A., RIPOLL C., THELLIER M. and KRAMER D., 1984: Morphometric

- study of *Lemna gibba* in relation to the use of compartmental analysis and the flux-ratio equation in higher plant cells. *Physiol. Plant.* 60(4), 493-501.
- GAULT J. and VAULOT O., 1977: Récupération de déchets par l'aquaculture. Centre Technique du Génie Rural des Eaux et des Forêts, Gaziinet (France). 49 pp.
- GAUSMAN H.W., 1977: Reflectance of leaf components. *Remote Sens. Environ.* 6, 1-9.
- GAUTIER B., 1980: Les limites phytogéographiques du Saint-Laurent. *Prochancheria* 11, 103 pp.
- GEARHEART R.A., FINNEY B.A., WILBUR S. and HULL D., 1984: The use of wetland treatment processes in water reuse. *Proc. Water Reuse Symp. III. ANWA Res. Found.*, Denver, Col. 2, 617-638.
- GEGER G., 1984: Karyosystematische Untersuchung der Gattungen *Spirodela* und *Lemna* (Lemnaceae). *Mitt. Bd. Bot.-Tag. Wien*, 9 (1016).
- GEISSERT F., SIMON M. and WOLFF P., 1985: Investigations floristiques et faunistiques dans le nord de l'Alsace et quelques secteurs limitrophes. *Bull. Assoc. Philom. Alsace Lorr.* 21, 111-127.
- GEISSMAN T.A. and JURD L., 1955: The anthocyanin of *Spirodela oligorhiza*. *Arch. Biochem. Biophys.* 56, 259-263.
- GELLINI R. and PICCARDI E.B., 1981: Sulla possibilità di assorbimento di metalli pesanti da parte di alcune piante acquatiche. *Inquinamento* 23(2), 45-46.
- GERGEL J., HARTWICH P. and HOREJSI J., 1985: The use of duckweed in the nutrition of ducks. (In Czech.). *SB Prov. Econ. Fak. Cesk. Bude. Zootech. Rada* 2(1), 79-88.
- GERVAIS, 1982: Adaptation et productivité de la lentille d'eau (*L. minor*). Application à la valorisation du lisier de porc. *Mém. fin d'études Esap.* 162 pp. (cited from HUBAC et al. 1984).
- GESSENER F., 1955: *Hydrobotanik*. 1. Energiehaushalt. *Deutsch. Verl. d. Wiss., Berlin*. 517 pp.
- GHATNEKAR S.D., PANDEY R.P., KULKARNI U.B., UDAY B. and IYER S.P., 1980: Biomangement of malathion effluent in Excel Industries Limited (India). *Indian Chem. Manuf. Annu.* 12, 17 pp.
- GHETTI P.P., CORRADI M. and COPELLI M., 1982: Culture de Lemnaceae pendant huit mois sur des déchets zootecniques partiellement traités. In: SYMOENS J.J., HOOPER S.S. and COMPERE P. (eds.), *Studies on aquatic vascular plants*. 370-374.
- GIANFAGNA T. and FOY C.L., 1975: The effect of glyphosate on the growth of *Lemna perpusilla*. *Va. J. Sci.* 26(2), 63.
- GIARDELLI M.L., 1935: Las flores de *Wolffia oblonga*. *Rev. Argentina Agron.* 2, 17-20.
- GIARDELLI M.L., 1937: Una nueva especie de Lemnaceae de la Flora Argentina. *Notas del Museo de la Plata* 2, *Botanica* 12, 97-100.
- GIARDELLI M.L., 1939 see 1939a
- GIARDELLI M.L., 1939a: El florecimiento de *Spirodela intermedia* W. Koch. *Notas del Museo de la Plata* 4, *Botanica* 26, 317-322.
- GIARDELLI M.L., 1939b: Nuevas especies de Lemnaceae para la Flora Argentina. *Physis* 15, 323-329.
- GIARDELLI M.L., 1941: *Lemna disperma*, especie nueva para la Flora Argentina. *Darwiniana* 5, 190-193.
- GIARDELLI M.L., 1947: Nota sobre una Lemnaceae tropical poco conocida: *Wolffia Welwitschii* Hegelm. *Darwiniana* 7, 552-556.
- GIARDELLI M.L., 1959: *Lemna aquinoctialis* Welwitsch nueva para la flora de America y de las Islas Filipinas. *Darwiniana* 11, 584-590.
- GIARDELLI M.L., 1970: Lemnaceae. In "Flora Patagonica" INTA - Coleccion Cientifica 8/2, 93-101.

- GODFREY R.K. and WOOTEEN J.W., 1979: Aquatic and wetland plants of South-eastern United States. The Univ. of Georgia Press, Athens.
- GODFRIAUX B.L., VALKENBURG H.J., VAN RIPER A., GUERRA C.R., EBLE A.F. and CAMPBELL P., 1975: Power plant heated water use in aquaculture. In: LANGWORTHY V.W. (ed.), Proc. 3rd Ann. Poll. Contr. Conf. of the Water and Wastewater Equipment Manuf. Assoc., Ann Arbor Sci. Publ. 233-250.
- GODWIN H., 1923: Dispersal of pond floras. J. Ecol. 11, 160-164.
- GODZIEMBA-CZYŻ J., 1969: Characteristic of vegetative and resting forms of *Wolffia arrhiza* (L.) Wimm. 1. Growth and dynamics of their mutual transformations. Acta Soc. Bot. Pol. 38, 437-452.
- GODZIEMBA-CZYŻ J., 1970: Vegetative and resting forms of *Wolffia arrhiza*. 2. Anatomy, physical and physiological properties. Acta Soc. Bot. Pol. 39, 421-443.
- GODZIEMBA-CZYŻ J., 1973: Certain aspects of the chemotaxis reaction of chloroplasts in *Funaria hygrometrica*. Acta Soc. Bot. Pol. 42, 453-459.
- GODZIK S., 1964: The action of crotonic aldehyde, 'sulfol' and 'nekal' on the respiration intensity of Lemna minor, *Riccia fluitans* and *Cladophora glomerata*. Acta Biol. Crac. ser. Bot. 7, 37-54.
- GOEBEL K., 1921: Zur Organographie der Lemnaceen. Flora 114, 278-305.
- GOEBEL K., 1928-1933: Organographie der Pflanzen. (3rd ed.). Jena.
- GOLDBACH H. and MICHAEL G., 1976: Abscissic acid content of barley grains during ripening as affected by temperature and variety. Crop Sci. 16 (6), 797-799.
- GOLDSBOROUGH L.G. and ROBINSON G.G.C., 1985: Seasonal succession of diatom epiphyton and dense mats of Lemna minor. Can. J. Bot. 63(12), 2332-2339.
- GOLLE B. and LUETTGE U., 1983: Inhibition of the glucose and amino acid carriers of Lemna gibba by pretreatment with HgCl₂. Physiol. Plant 57(1), 62-66.
- GOLLIN D.J., DARVILL A.G. and ALBERSHEIM P., 1984: Plant cell wall fragments inhibit flowering and promote vegetative growth in Lemna gibba. Biol. Cell 51, 275-280.
- GOLUBINSKY I.N., GAPON S.V. and SIKACH N.A., 1987: The case of the mass flowering of Lemna minor (Lemnaceae) in the Poltava region. (In Russian). Bot. Zh. 72(6), 780-781.
- GOLUJEK C.G., 1979: Aquaculture in resource recovery. Compost Sci. 20 (3), 16-23.
- GOODMAN B.A. and DEKOCK P.C., 1982: Moessbauer studies of plant materials. I. Duckweed, stocks, soybean and pea. J. Plant Nutr. 5(4-7), 345-353.
- GOODMAN B.A., DEKOCK P.C. and RUSH J.D., 1982: Moessbauer studies of plant materials. II. Spectra of Fe-enriched duckweed at low temperatures. J. Plant Nutr. 5(4-7), 355-362.
- GOODWIN T.W. and MERCER E.I., 1972: Introduction to plant biochemistry. Pergamon Press, Oxford, UK. 359 pp.
- GORDON W.R., 1977: An investigation of phenylalanine ammonia-lyase during growth and development in members of the Lemnaceae under varying conditions of axenic culture. Univ. Microfilms Int., Ann Arbor, Order No. 77-18,991, 174 pp. Diss. Abstr. Int. B, 38, 989.
- GORDON W.R. and KOUKKARI W.L., 1977: A circadian rhythm in the activity of L-phenylalanine ammonia-lyase. Chronobiologia 4(2), 112.
- GORDON W.R. and KOUKKARI W.L., 1978: Circadian rhythmicity in the activities of phenylalanine ammonia-lyase from Lemna perpusilla and Spirodela polyrrhiza. Plant Physiol. 62, 612-615.
- GORDON W.R., SCHWEMMER S.S. and HILLMANN W.S., 1978: Nickel and the metabolism of urea by Lemna paucicostata Hegelm. 6746. Planta 140, 265-268.

- GIARDELLI M.L., 1972: *Wolffia brasiliensis*, especie de Lemnaceas nueva para la flora argentina. Darwiniana 17, 596-597.
- GICKLHORN J., 1958: Wasserlinsen. Universum, Natur und Technik 13, 24-27.
- GILBERT H.C., 1937: Lemnaceae in flower. Science 86, 308.
- GILLMAN H., 1871a: L. trisulca in flower. Am. Naturalist 5, 651-652.
- GILLMAN H., 1871b: L. polyrrhiza in flower. Am. Naturalist 5, 652-653.
- GILLMAN H., 1881: L. polyrrhiza again discovered in flower on Detroit River. Am. Natur. 15, 896-897.
- GIOVANELLI J., 1981: Homocysteine and methionine metabolism. Proc. Int. Bot. Congr. 13, 18.
- GIOVANELLI J., DATKO A.H., MUDD S.H. and THOMPSON G.A., 1983: In vivo metabolism of 5'-methylthioadenosine in Lemna. Plant Physiol. 71(2), 319-326.
- GIOVANELLI J. and MUDD S.H., 1985: Radioactive methionine determination and distribution of radioactivity in the sulfur methyl and 4-carbon moieties. J. Biochem. Biophys. Methods 11(1), 1-12.
- GIOVANELLI J., MUDD S.H. and DATKO A.H., 1980a: Homocysteine biosynthesis in plants. In: CAVALLINI D.G., GAULL G.E. and ZAPPALÀ V. (eds.), Natural sulfur compounds: Novel biochemical and structural aspects. Plenum Press, New York/London. 81-92.
- GIOVANELLI J., MUDD S.H. and DATKO A.H., 1980b: Sulfur amino acids in plants. In: STUMPF P.K. and CONN E.E. (eds.), The biochemistry of plants. Acad. Press, New York. 5, 453-505.
- GIOVANELLI J., MUDD S.H. and DATKO A.H., 1981: Recycling of methionine sulfur in a higher plant by two pathways characterized by either loss or retention of the 4-carbon moiety. Biochem. Biophys. Res. Comm. 100 (2), 831-839.
- GIOVANELLI J., VELUTHAMBI K., THOMPSON G.A., MUDD S.H. and DATKO A.H., 1984: Threonine synthase of Lemna paucicostata Hegelm. 6746. Plant Physiol. 76(2), 285-292.
- GIOVANELLI J., MUDD S.H. and DATKO A.H., 1985a: In vivo regulation of de novo methionine biosynthesis in a higher plant (Lemna). Plant Physiol. 77(2), 450-455.
- GIOVANELLI J., MUDD S.H. and DATKO A.H., 1985b: Quantitative analysis of pathways of methionine metabolism and their regulation in Lemna. Plant Physiol. 78(3), 555-560.
- GIOVANELLI J., MUDD S.H., DATKO A.H. and THOMPSON G.A., 1986: Effects of orthophosphate and adenosine 5'-phosphate on threonine synthase and cystathionine γ -synthase of Lemna paucicostata Hegelm. 6746. Plant Physiol. 81(2), 577-583.
- GIUGA G., 1973: Vita segreta di Lemnaceae. Lemna symmeter G. Giuga - species nova. Blarion, Napoli. 19 pp.
- GLEASON F.K. and CASE D.E., 1986: Activity of the natural alginate cyanobacterin on angiosperms. Plant Physiol. 80(4), 834-837.
- GLANDON R.P. and McNABB C.D., 1978: The uptake of boron by Lemna minor. Aquat. Bot. 4, 53-64.
- GLICKMAN L.T., 1966: Some factors affecting the dark processes in the germination of seeds of Lemna perpusilla 6746. M.S. Thesis. State Univ. of New York at Binghamton, N.Y.
- GLICKMAN L.T. and POSNER H.B., 1966: Some factors affecting the dark processes in the germination of seeds of Lemna perpusilla 6746. Plant Physiol. 41(suppl.), 31.
- GODEANU S., GODEANU M., OLTEAN M., DIACONU M.O.I. and GITTA V., 1978: Bioceneses installation in the pools of biological epuration naturally insemminated or planted with aquatic macrophytes. Trav. Mus. Hist. Nat. "Grigore Antipa" 19, 125-129.

- GORHAM P.R., 1941: Measurements of the response of Lemna to growth promoting substances. *Am.J.Bot.* 28, 98-101.
- GORHAM P.R., 1945: Growth factor studies with *Spirodela polyrrhiza* (L.) Schleid. *Am.J.Bot.* 32, 496-505.
- GORHAM P.R., 1950: Heterotrophic nutrition of seed plants with particular reference to Lemna minor L. *Can.J.Research C* 28, 356-381.
- GOTO K., 1978: Mutually inverse rhythmic and sigmoidal changes in activity of cytoplasmic and chloroplast glycerolaldehyde 3-phosphate dehydrogenase in Lemna gibba G3. *Plant Cell Physiol.* 19, 749-758.
- GOTO K., 1979a: Modes of control by the circadian oscillator and the hourglass mechanism of the activities of cytoplasmic and chloroplast glycerolaldehyde 3-phosphate dehydrogenases in Lemna gibba G3. *Plant Cell Physiol.* 20, 513-521.
- GOTO K., 1979b: Mechanism of control by a circadian oscillator of chloroplast NADP-linked glycerolaldehyde 3-phosphate dehydrogenase in Lemna gibba G3. *Plant Cell Physiol.* 20, 523-532.
- GOTO K., 1983: A model for the circadian oscillator in Lemna gibba G3. *Plant Physiol.* 72(1 suppl.), 86.
- GOTO K., 1984: Causal relationships among metabolic circadian rhythms in Lemna gibba. *Z.Naturforsch.* 39C(1-2), 73-84.
- GOWER R.A. and POSNER H.B., 1979: Effects of light and 3-(3,4-dichlorophenyl)-1,1-dimethylurea on levels of ATP in Lemna paucicostata 6746 and a photosynthetic mutant with abnormal flowering responses. *Plant Physiol.* 63, 548-551.
- GRAF B., 1987: Nährwert der Wasserlinse als Komponente von Geflügelfutter. Semesterarbeit. Institut f.Nutzwissenschaften.ETHZ. 17 pp. (Poly-copy).
- GRAVIS A., 1935: Observations anatomiques et éthologiques sur les Cactées et les Lennacées. *Acad.R.Belg.Cl.Sci.* 8(14), fasc. 6. 70 pp.
- GRAY A., 1968: Manual of the Botany of the Northern United States. (5th ed.). Lennaceae. Ivison, Phinney, Blakeman Co., New York. 478-480.
- GRBOVIC M., RADENOVIC C. and DAMJANOVIC Z., 1975: Bioelectric potential of the root of Lemna minor. *Acta Bot.Croat.* 34, 182-183.
- GREENBERG B.M., MATTOO A.K., GABA V. and EDELMAN M., 1986: In vivo breakdown products of the 32-kDa thylakoid herbicide binding protein. *Plant Physiol.* 80(4 suppl.), 47.
- GREENBERG G.M., GABA V., MATTOO A.K. and EDELMAN M., 1987: Effect of spectral light quality on turnover of the 32 kDa PSII reaction center protein. *Plant Physiol.* 83(4 suppl.), 27.
- GREENHAM C.G., 1986: Influence of pH on toxicity of the non-electrolytes coumarin and ethyl-N-phenylcarbamate to *Spirodela polyrrhiza*. *Plant Prot.Q.* 1(3), 111-114.
- GREENWOOD D.J. and NELDER J.A., 1964: The effect of drugs on the growth of Lemna minor C. *Ann.Bot.* 28(112), 711-715.
- GRENIER G. and BEAUMONT G., 1983: Effets physiologiques de l'atrazine à doses sublétales sur Lemna minor. VII. Incorporation d'acétate-1,2-¹⁴C dans les groupes de lipides et leurs acides gras. *Physiol. Plant.* 57, 477-484.
- GRENIER G., MARIER J.P. and BEAUMONT G., 1979: Effets physiologiques de l'atrazine à doses sublétales sur Lemna minor L. IV. Influence sur la composition lipidique. *Can.J.Bot.* 57, 1015-1020.
- GRENIER G., MARIER J.P. and BEAUMONT G., 1982: Effets physiologiques de l'atrazine à doses sublétales sur Lemna minor L. VI. Influence sur les classes importantes de phospholipides. *Physiol.Végétale* 20(2), 179-185.

- GRESSEL J., 1978: Light requirements for the enhanced synthesis of a plastid mRNA during *Spirodela* greening. *Photochem.Photobiol.* 27, 167-169.
- GRESSEL J., 1982a: Triazine herbicide interaction with 32000 MW thylakoid protein-alternative possibilities. *Plant Sci.Lett.* 25(1), 99-106.
- GRESSEL J., 1982b: Is the site of triazine herbicide action really kilodalton thylakoid protein? *Photoparasitica* 10(4), 250.
- GRESSEL J. and COHEN N., 1977: Effects of dikegulac, a new growth regulator, on RNA synthesis in *Spirodela*. *Plant.Cell.Physiol.* 18(1), 255-259.
- GRESSEL J., ROSNER A. and COHEN N., 1975: Temperature of acrylamide polymerization and electrophoretic mobilities of nucleic acids. *Anal. Biochem.* 69(1), 84-91.
- GRIFFITH W., 1851: Notulae ad plantas asiaticas 3, 216, 220, 221, 229. *Icones plantarum asiaticarum* 3, plate 264, fig. 13, 16. Bishop College Press, Calcutta.
- GRISEBACH A.H.R., 1864: Flora of the British West Indian Islands. London. P. 512.
- GRISEBACH H., 1967: Biosynthetic pattern in microorganisms and higher plants. Wiley, New York.
- GRISEBACH H., 1980: Branched-chain sugars: Occurrence and biosynthesis. In: STUMPF P.K. and CONN E.E. (eds.), *The biochemistry of plants*. Acad.Press, New York. 3, 171-197.
- GRISEBACH H., BARON D., KELLEHER W.J. and WELLMANN E., 1972a: Biosynthesis of D-apiose and D-xylose in parsley and Lemna minor and their regulation by light. *Hoppe-Seyler's Z.Physiol.Chem.* 353, 1519-1520.
- GRISEBACH H., BARON D., SANDERMANN H. and WELLMANN E., 1972b: Formation of UDP-apiose from UDP-glucuronic acid. In: GINSBURG V. (ed.), *Methods in enzymology*. Complex carbohydrates, Part B. Acad.Press, New York/London. 28, 439-446.
- GRITSENKO N.V. and DYMINA G.D., 1976: Species of flora new for the Amur Oblast. (In Russian). *Bot.Zh.* (Leningr.) 61(2), 239-242.
- GROB E.C. and EICHENBERGER W., 1969: Ueber die Galactolipide und Fettsäuren in normalen und glucosestimulierten Kulturen von *Spirodela oligorhiza*. *FEBS Lett.* 5(5), 335-337.
- GROB E.C. and EICHENBERGER W., 1973: Beiträge zum Stickstoffmetabolismus in grünen Pflanzen. I. Die reversible Plasteridenumwandlung in glucosestimulierten Kulturen von *Spirodela oligorhiza*, eine Folge von Stickstoffmangel. *Experientia* 29(4), 398-400.
- GROB E.C., HODLER M. and EICHENBERGER W., 1973: Beiträge zum Stickstoffmetabolismus grüner Pflanzen. II. Der Einfluss exogener Glucose auf die Aufnahme von Nitrat und Ammonium aus der Nährlösung und auf den Proteingehalt von *Spirodela oligorhiza*. *Experientia* 29(4), 401-402.
- GROB E.C. and RUFENER J., 1969: Influence of sugar containing nutrients on ultrastructure and photosynthesis activity of *Spirodela oligorhiza* chloroplasts. In: METZNER H. (ed.), *Progress in photosynthesis research*. Tübingen. 1, 55-62.
- GROOT G.S.P. and VAN HARTEN-LOOSBROEK N., 1981: Physical mapping of 4S RNA genes on chloroplast DNA of *Spirodela oligorhiza*. *Curr.Genet.* 4(3), 187-190.
- GRUNTZEL S., 1982: Die Wirkung unterschiedlicher Stickstoffernährung auf den Cytokinin-Gehalt von Lemna aequinoctialis 6746. Diploma thesis. Institute of Pharmacognosy, Univ. Vienna, Austria. 62 pp.
- GUDKOVA E.V., MITROFANOV Y.A. and SHAVEL ZON R.A., 1970: Cytogenetic effect of 2,6-diaminopurine. (In Russian). *Izv.Akad.Nauk SSSR.Ser.Biol.* 4, 508-514.

- GUERN J., 1960: Mise en évidence de phénomènes de corrélation au cours de la croissance des frondes de *Lemna trisulca* L. C.R.Acad.Sci.Paris, 251, 894-896.
- GUERN J., 1962: Observations sur la production de frondes axillaires par les frondes de *Lemna trisulca* L. C.R.Acad.Sci.Paris, 254, 343-345.
- GUERN J., 1963a: Caractéristiques de la croissance des frondes de *Lemna trisulca* L. C.R.Acad.Sci.Paris, 256, 2220-2222.
- GUERN J., 1963b: Modifications expérimentales de l'intensité de la dominance entre frondes de *Lemna trisulca*. C.R.Acad.Sci.Paris, 256, 2686-2688.
- GUERN J., 1963c: Remarques à propos de l'action de la kinétine sur le mode de germination des turions de *Spirodela polyrrhiza* Schleid. C.R. Acad.Sci.Paris, 257, 3464-3467.
- GUERN J., 1965: Corrélations de croissance entre frondes chez *Lemnacées*. Ann.Sc.Nat.Bot.Paris, 12e Sér. 6, 1-156.
- GUERN J., GUERN N. and TRAPY F., 1964: Action de quelques purines substituées sur la levée de dominance entre frondes filles formées au cours de la germination des turions de *Spirodela polyrrhiza* Schleid. C.R.Acad.Sci.Paris, 258, 4329-4332.
- GULATI D.K., CHAMBERS C.L., ROSENTHAL G.A., and SABHARWAL P.S., 1981: Comparative toxicity of some naturally occurring and synthetic non-protein amino acids. Environ.Exp.Bot. 21(2), 225-230.
- GULATI D.K., ROSENTHAL G.A. and SABHARWAL P.S., 1977: Effect of cana-line-urea cycle amino acids on macromolecular synthesis by *Lemna minor* L. Plant Physiol. 59(suppl.), 70.
- GULATI D.K., ROSENTHAL G.A. and SABHARWAL P.S., 1978: Rapid analysis of macromolecular metabolism in *Lemna minor*. Plant Physiol. 61(4 suppl.), 20.
- GULATI D.K., ROSENTHAL G.A. and SABHARWAL P.S., 1979: An improved method for estimating macromolecular synthesis. J.Exp.Bot. 30(118), 919-924.
- GULLIVER, 1967: *Wolffia* has no raphides in contrast to *Lemna*. (In Dutch). J.Bot. 5, 22.
- GUMINSKI S., MIAZGA T. and NOWAK J., 1978: The effects of sodium humate and detergent DBSS and their joint effect on cultures of *Spirodela polyrrhiza* (L.) Schleiden. Ekol.Pol. 26, 231-240.
- GUNTA W.J., 1987: -amylase in *L. minor*: characterization, intracellular distribution and its role in glucosyl-metabolism. 14th Int.Bot.Congr. Berlin, Abstr., 57.
- GUNKEL G., 1983: Untersuchungen zur Ökotoxikologischen Wirkung eines Herbizids in einem aquatischen Modellökosystem. I. Subletale und letale Effekte. Arch.Hydrobiol., suppl. 65(2-3), 235-267.
- GUNKEL G., 1984: Untersuchungen zur Ökotoxikologischen Wirkung eines Herbizids in einem aquatischen Modellökosystem. II. Nahrungskettenprobleme und Schadstoffbilanzierung. Arch.Hydrobiol., suppl. 69(1), 130-168.
- GUNNER H.B. and COLE R.A., 1971: A microbiotic ecoassay for environmental pollutants. In: WESTLEY B. (ed.), Identification and measurement of environmental pollutants. Nat.Res.Council, Canada. 314-319.
- GUPPY H.B., 1984: On the habits of *Lemna minor*, L. gibba and *L. polyrrhiza*. J.Linn.Soc. (London) 30, 323-330.
- GUPPY H.B., 1966: Observations of a naturalist in the Pacific. McMillan, London. 2, 407-408.
- GUPTA B.L., 1935: Studies on the development of the pollen grain and embryo sac of *Wolffia arrhiza*. Curr.Sci. 4, 104-105.
- GUPTA S. and MAHESHWARI S.C., 1969: Induction of flowering by cytokinins in a short-day plant, *Lemna paucicostata*. Plant Cell Physiol. 10, 231-233.

- GUPTA S. and MAHESHWARI S.C., 1970a: Growth and flowering of *Lemna paucicostata*. I. General aspects and role of chelating agents in flowering. Plant Cell Physiol. 11, 83-95.
- GUPTA S. and MAHESHWARI S.C., 1970b: Growth and flowering of *Lemna paucicostata*. II. Role of growth regulators. Plant Cell Physiol. 11, 97-106.
- GUSKOVA V.N., BRAGINA A.N., ZASEDATELEV A.A., ILYIN B.N., KUPRIYANOVA V.M., MASHREVA N.I., RODIONOVA L.F., SUKALSKAYA S.Y. and TIKHONOVA A.I., 1970: Effects of uranium mixture fusion products on sanitary conditions and hydrobionts of weakly mineralized freshwater basins. (In Russian). Gidrobiol.Zh. 6(4), 5-11.
- GUSTINE D.L., 1969: Biosynthesis of D-apiose by an enzyme system isolated from *Lemna minor* L. Ph.D.Thesis. Michigan State Univ. 106 pp.
- GUSTINE D.L. and KINDEL P.K., 1969: Biosynthesis of D-apiose in a cell-free system from *Lemna minor* L. J.Biol.Chem. 244(5), 1382-1385.
- GUSTINE D.L., YUAN D.H.F. and KINDEL P.K., 1975: Uridine diphosphate D-glucuronic acid cyclase and uridine diphosphate D-glucuronic acid carboxy-lyase I from *Lemna minor*; purification, characterization, and separation from uridine diphosphate D-glucuronic acid carboxy-lyase II. Arch.Biochem.Biophys. 170, 82-91.
- GUTHRIE R.K. and CHERRY D.S., 1976: Pollutant removal from coal ash basin effluent. Water Resour.Bull. 12(5), 889-902.
- GUTHRIE R.K. and CHERRY D.S., 1979a: The uptake of chemical elements from coal ash and settling basin effluent by primary producers. I. Relative concentrations in predominant plants. Sci.Total Environ. 12, 217-222.
- GUTHRIE R.K. and CHERRY D.S., 1979b: Trophic level accumulation of heavy metals in a coal ash basin drainage system. Water Resour.Bull. 15(1), 244-248.
- HAAG R.W., 1983: Emergence of seedlings of aquatic macrophytes from lake sediments. Can.J.Bot. 61(1), 148-156.
- HABECK D.H., 1983: The potential of *Parapoxyn stratiotata* L. as a biological control agent for Eurasian watermilfoil. J.Aquat.Plant Manage. 21, 26-29.
- HABERLANDT G., 1987: Zur Kenntnis des Spaltöffnungsapparates. Flora 70, 97-110.
- HAERTEL O., 1972: Die Zwergrinse, *Wolffia arrhiza* (L.) Wimm. auch noch 1971 wieder im Curauer und Malkendorfer Moor. Kieler Notizen zur Pfl.kunde in Schleswig-Holstein 4, 10-11.
- HAJRA A. and TRIPATHI S.D., 1985: Nutritive value of aquatic weed, *Spirodela polyrrhiza* (Linn.) in grass carp. Indian J.Anim.Sci. 55(8), 702-705.
- HAKONSON T.E. and WHICKER F.W., 1975: Cesium kinetics in a montane lake ecosystem. Health Phys. 28, 699-706.
- HALABAN R., 1972: Mitotic index and cell cycle of *Lemna perpusilla* under different photoperiods. Plant Physiol. 50, 308-310.
- HALABAN R. and HILLMAN W.S., 1970a: Response of *Lemna perpusilla* to periodic transfer to distilled water. Plant Physiol. 46, 641-644.
- HALABAN R. and HILLMAN W.S., 1970b: Rhythmic response of *Lemna perpusilla* to periodic transfer to distilled water. Plant Physiol. 46 (suppl.), 26.
- HALABAN R. and HILLMAN W.S., 1971: Factors affecting the water-sensitive phase of flowering in the short-day plant *Lemna perpusilla*. Plant Physiol. 48, 760-764.
- HALABAN R. and HILLMAN W.S., 1974: Photoperiodic time measurement of

- flowering in a short-day plant: effects of the external medium. In: SCHEVING L.E. et al. (eds.), Chronobiology. 666-670.
- HALLAKSY E. see VON HALACSY E.
- HALLER W.T., SUTTON D.L. and BARLOWE W.C., 1974: Effects of salinity on growth of several aquatic macrophytes. Ecology 55, 891-894.
- HAMASHIMA S., 1971: Lemna minor found in Aichi prefecture, Japan. (In Japan.). J. Geobot. 19(1-2), 28.
- HAMASHIMA S., 1974: A new naturalized duckweed, Lemna gibba L. (In Japan.). J. Jap. Bot. 49, 359.
- HAMASHIMA S., 1978a: Seed germination of three Lemna species. (In Japan.). J. Jap. Bot. 53, 28-31.
- HAMASHIMA S., 1978b: On the flower of Wolffia arrhiza. (In Japan.). J. Jap. Bot. 53, 62.
- HAMASHIMA S., 1983: Relationship between some environmental conditions and the distribution of aquatic weeds in the irrigation canals of the Nobi Plain. (In Japan.) Weed Res., Japan 28(4), 266-270.
- HAMBRECH R.N., 1968: Control of duckweed and Azolla with diuron. Weed Sci. Soc. Am. Meet. Abstr., 65-66.
- HAMNER K.C. and HODSON H.K., 1969: The relationship of floral inducing extracts to flowering gibberellins and inhibitors. XI Intern. Bot. Congr. Abstr., 83.
- HANES G., 1971: Wolffia II. Trail and Landscape 5, 100-106.
- HANNA R.B. and MENDICINO J., 1969: Synthesis of isomers and nucleoside diphosphate derivatives of D-apio-furanosyl-1 phosphate. Fed. Proc. 28 (2), 856.
- HANNA R.B., PICKEN M. and MENDICINO J., 1973: Purification of a specific D-apitol dehydrogenase from a Micrococcus isolated from the surface of germinating parsley seeds. Biochim. Biophys. Acta 315(2), 259-271.
- HANSTEEN B., 1896: Beiträge zur Kenntnis der Eiweißbildung und der Bedingungen der Realisierung dieses Prozesses im Phanerogamen Pflanzenkörper. Ber. Deutsch. Bot. Ges. 14, 362-371.
- HANSTEEN B., 1899: Ueber Eiweißsynthese in grünen Phanerogamen. Jb. Wiss. Bot. 33, 417-486.
- HANTKE R., 1954: Die fossilen Floren der obermiozänen Fundstelle Schrotzberg (Schlienerberg, Südbaden). Denkschr. Schweiz. Naturf. Ges. 80, 30-118.
- HARA H., 1976: Comments on the East Asiatic plants. (3). (In Japan.). J. Jap. Bot. 52(8), 225-229.
- HARBORNE J.B., 1986: The natural distribution in angiosperms of anthocyanins acylated with aliphatic dicarboxylic acids. Phytochemistry 25(8), 1887-1894.
- HARBORNE J.B. and TURNER B.L., 1984: Plant chemosystematics. Acad. Press, London. 562 pp.
- HARGROVE C.L., 1976: Operation duckweed. Weeds, Tress and Turf 15(6), 30-34.
- HARRIS S.K. and BEAN R.C., 1951: Wolffia columbiana in Methuen, Massachusetts. Rhodora 53, 272.
- HARRISON D.E., 1964: The taxonomic significance of the effect of nutrient media, photoperiod, and light intensity on the morphological features of the genus Spirodela Schleid. M.S. Thesis. North Carolina State Univ., Raleigh.
- HARRISON D.E. and BEAL E.O., 1963 see 1964
- HARRISON D.E. and BEAL E.O., 1964: The Lemnaceae (duckweeds) of North Carolina. J. Elisha Mitchell Sci. Soc. 80, 12-18.
- HART D.A., 1969: Apiogalacturonans from the cell wall of Lemna minor. Univ. Microfilms Int., Ann Arbor, Order No. 70-15,048; 136 pp. Diss. Abstr. Int. B, 31, 507.

- HART D.A. and KINDEL P.K., 1970a: A novel reaction involved in the degradation of apiogalacturonans from Lemna minor and the isolation of apiose as a product. Biochemistry 9, 2190-2196.
- HART D.A. and KINDEL P.K., 1970b: Isolation and partial characterization of apiogalacturonans from the cell wall of Lemna minor. Biochem. J. 116, 569-579.
- HARTMAN P., KRAL V. and TISCHLER O., 1982: Device and methods for biological treatment of wastewater especially from agricultural large-scale production. (In Czech.). Patent CS 196593 B. 6 pp.
- HARTMAN R.T., 1960: Algae and metabolites of natural waters. In: TRIVON C.A. and HARTMAN R.T. (eds.), The ecology of algae. The Pygmatunng Symposia in Ecology, Univ. Pittsburgh. 2, 28-55.
- HARTMAN R.T. and ENGLISH S.M., 1959: Wolffia floriadana in Western Pennsylvania. Castanea 24, 45-47.
- HARTMAN W.A. and MARTIN D.B., 1984: Effect of suspended bentonite clay on the acute toxicity of glyphosate to Daphnia pulex and Lemna minor. Bull. Environ. Contam. Toxicol. 33(3), 355-361.
- HARTMAN W.A. and MARTIN D.B., 1985: Effect of four agricultural pesticides on Daphnia pulex, Lemna minor, and Potamogeton pectinatus. Bull. Environ. Contam. Toxicol. 35(5), 646-651.
- HARTMANN A., BIELENSKI U., LUETTGE U., GARREC J.P., THOIRON A. and THELTER M., 1983: Measurements of unidirectional fluxes of lithium: application to the study of lithium ion/proton interactions with the transmembrane exchanges of Lemna gibba Gl. Stud. Phys. Theor. Chem. 24, 591-597.
- HARTMANN T. and EHMKE A., 1976 see EHMKE A. and HARWANN T., 1976
- HARTUNG W. and KANDELER R., 1976: Die Wirkung abendlicher Dunkelrotbestrahlung auf die Aufnahme und Verteilung markierter Phytohormone in kurztagkultivierten Lemna gibba Gl.-Pflanzen. Z. Pflanzenphysiol. 79, 360-367.
- HARTUNG W. and WITT J., 1968: Ueber den Einfluss der Bodenfeuchtigkeit auf den Wuchsstoffgehalt von Anastatica hierochuntica und Helianthus annuus. Flora B 157, 603-614.
- HARTUNG W., ULIRICH-EBERUS C.I., LUETTGE U., BOECHER M. and NOVACKY A., 1980: Effect of abscisic acid on membrane potential and transport of glucose and glycine in Lemna gibba Gl. Planta 148, 256-261.
- HARTZ P., ROCHLING Y. and MARIQUW-SMIT F., 1972: 2-dichloroacetamido-3-chlor-1,4-naphthaquinon - ein neues Algizid für den Einsatz in Reis und anderen Kulturen. Medel. Fak. Landbouwetenschappen Gent 37(2), 699-704.
- HARVEY R.M. and FOX J.L., 1973: Nutrient removal using Lemna minor. J. Water Poll. Control Fed. 45, 1928-1938.
- HASIMI R.Y. and OMER S., 1986: Lemnaceae. In: NASIR E. and ALI S.I. (eds.), Flora of Pakistan. 173. 11 pp.
- HASUNUMA K., 1986: Cyclic nucleotides in Lemna paucicostata. (In Japan.). IDEN 40(8), 35-40.
- HASUNUMA K. and FUNADERA K., 1987: GTP-binding proteins in green plant, Lemna paucicostata. Biochem. Biophys. Res. Commun. 143(3), 908-912.
- HASUNUMA K., SHINOHARA Y., FUNADERA K. and FURUKAWA K., 1987: Circadian rhythmicity in the oscillations of cAMP and cGMP concentrations in Neurospora and Lemna. 14th Int. Bot. Congr. Berlin, Abstr., 142.
- HATANAKA A., SEKIYA J. and KAJIWARA T., 1978: Distribution of an enzyme system producing cis-3-hexenal and n-hexanal from linolenic and linoleic acids in some plants. Phytochemistry 17(5), 869-872.
- HAUPT W., 1982: Light mediated movement of chloroplasts. Ann. Rev. Plant Physiol. 33, 205-233.

- HAUPT W. and WEISENSEE M., 1966: Die Temperaturabhängigkeit der Chloroplastenbewegungen bei *Lemna trisulca*. Naturwiss. 53, 411-412.
- HAUPT W. and WEISENSEE M., 1967: Chloroplastenbewegung bei *Lemna trisulca* in polarisiertem Licht. Naturwiss. 54, 48-49.
- HAUSER L.A., 1977: The Lemnaceae collection of C.H. Thompson in the University of Kansas Herbarium. Rep.State Biol.Surv. Kansas 10, 12 pp.
- HAYEK A. see VON HAYEK A.
- HAYES T.L., 1980: Biophysical aspects of scanning electron microscopy. Scann.Electron.Microsc. 1980(11), 1-10.
- HEALY W.B. and MCCOLL R.H.S., 1974a: Clay particles as sources of phosphorus for *Lemna* and their role in eutrophication. (In Hung.). Agrokem.Talajtan 23, 407-417.
- HEALY W.B. and MCCOLL R.H.S., 1974b: Clay particles as sources of phosphorus for *Lemna* and their role in eutrophication. New Zealand J. Sci. 17, 409-420.
- HECKMANN C.W., 1984: Erstfund von *Lemna turionifera* Landolt 1975, in Europa: Haseldorfer Marsch. Kieler Notiz.Pflk.Schl.-Holst. 16(1/2), 1-3.
- HECKMANN R.A., WINGET R.N., INFANGER R.C., MICKELSEN R.W. and HENDERSEN J.M., 1984: Warm water aquaculture using waste heat and water from zero discharge power plants in the Great Basin. Great Basin Nat. 44(1), 75-82.
- HEDTKE S.F., WEST C.W., ALLEN K.N., NOSBERG-KING T.J. and MOUNT D.I., 1986: Toxicity of pentachlorophenol to aquatic organisms under naturally varying and controlled environmental conditions. Environ.Toxicol.Chem. 5(6), 531-542.
- HECKMANN 1984 see HECKMANN 1984
- HEGELMAIER F., 1865: Welwitschii iter angolense IV. Lemnacearum a cl. Fr. Welwitsch in Africae aequinoctialis territorio angolensi collectum descriptio. J.Bot. 3, 110-115.
- HEGELMAIER F., 1868: Die Lemnaceen. Eine monographische Untersuchung. Engelmann, Leipzig. 169 pp.
- HEGELMAIER F., 1871: Ueber die Fruktifikationstheile von *Spirodela*. Bot.Z. 29(38), 621-630; 645-666.
- HEGELMAIER F., 1878: Lemnaceae. In: Martius Flora Brasiliensis 3/2, 1-23.
- HEGELMAIER F., 1885: Wolfia microscopica. Bot.Z. 43(16), 241-249.
- HEGELMAIER F., 1895: Systematische Uebersicht der Lemnaceen. Bot.Jb. 21, 268-305.
- HEGNAUER R., 1963: Chemotaxonomie der Pflanzen. Monocotyledonae. Birkhäuser, Basel/Stuttgart. 2, 73-79, 267-269, 483.
- HEGNAUER R., 1986: Chemotaxonomie der Pflanzen. (Nachtr. Bd.1 und 2). Birkhäuser, Basel/Boston/Stuttgart. 7, 681-685.
- HEJNY S., 1960: Ökologische Charakteristik der Wasser- und Sumpfpflanzen in den slowakischen Tiefebene (Donau- und Theissgebiet). Slow. Akad.Wiss.Bratislava. 487 pp.
- HEJNY S., 1967: Wintering of plants in water tanks. (In Czech.). Ziva 15(5), 162-164.
- HEJNY S., 1968: Bemerkungen zur Klassifikation einiger Makrophytengesellschaften der stehenden Gewässer. In: TUEXEN R. (ed.), Pflanzensoziologische Systematik. Junk, The Hague. 230-238.
- HEJNY S. and HROUDOVA Z., 1984: Plant adaptations to shallow water habitats. 14 pp. (Polycopy).
- HEJNY S. and HUSAK S., 1978: Higher plant communities. In: DYKJOVA D. and KVET J. (eds.), Pond littoral ecosystems. Berlin. 23-64.
- HEJNY S., KVET J. and DYKJOVA D., 1981: Survey of biomass and net pro-

- duction of higher plant communities in fishponds. Folia Geobot.Phytotax. 16, 73-94.
- HELDWEIN R. and KANDELER R., 1981: Adaptation to end-of-day far red and malate accumulation in *Lemna*. Z.Pflanzenphysiol. 102(2), 141-146.
- HELLMUND D., METZLAFF M. and SERFLING E., 1984: A transfer RNAARG gene of *Pelargonium chloroplasts*, not a 5S RNA gene, is efficiently transcribed after injection into *Xenopus oocyte* nuclei. Nucleic Acids Res. 12(21), 8253-8268.
- HELLQUIST C.B., 1972: Range extensions of vascular aquatic plants in New England. Rhodora 74(797), 131-141.
- HELLQUIST C.B. and CROW G.E., 1982: Aquatic vascular plants of New England. V. Araceae, Lemnaceae, Xyridaceae, Eriocaulaceae and Pontederiaceae. Station Bull. 523, 46 pp.
- HENLEBEN V., 1972: Untersuchungen zur Biosynthese und Funktion von Nucleinsäuren in höheren Pflanzen. Thesis for Habilitation. Univ. Tübingen. 99 pp.
- HENMERLING J. and MEUSEL W., 1968: Wolfia arthiza, die kleinste Blütenpflanze. Monatschr.Ornithol.Vivarienk. B 15(4), 124.
- HENDERSON J.M., HECKMANN R.A. and WINGET R.N., 1984: Multiple use systems for aquaculture. Great Basin Nat. 44(3), 471-481.
- HENEBRY M.S., CAIRNS J., SCHWINTZER C.R. and YOUNGUE W.H., 1981: A comparison of vascular vegetation and protozoan communities in some freshwater wetlands of northern Lower Michigan. Hydrobiologia 83, 353-375.
- HENNINGS P., 1991: *Lemna trisulca* var. *pygmaea*. Verh.Bot.Ver.Prov.Brandenburg 33, 8-9.
- HENSLOW G., 1911: The origin of Monocotyledons from Dicotyledons through self-adaptation to a moist or aquatic habitat. Ann.Bot. 25, 717-744.
- HENSSON A., 1954: Die Dauerorgane von *Spirodela polyrrhiza* (L.) Schleid. in physiologischer Betrachtung. Flora 141, 523-566.
- HEPHER B. and PRUGNIN Y., 1979: Guide to fish culture in Israel. 4. Fertilization, manuring and feeding. Foreign Training Dept., Israel. 61 pp.
- HEPPER F.N., 1968: Lemnaceae. In: HUTCHINSON J. and DALZIEL J.M. (eds.), Flora of West Tropical Africa. (2nd ed.). Crown Agents, London. 127-129.
- HEPPER F.N., 1973: Lemnaceae. In: POLLHILL R.M. (ed.), Flora of Tropical East Africa. East African Community, Arusa, Tanzania. 9 pp.
- HEPPER F.N., 1981: Lemnaceae. In: DASSANAYAKE M.D. and FOSBERG F.R. (eds.), A revised handbook to the flora of Ceylon. Balkema, Rotterdam. 2, 397-402.
- HEPPER F.N., LAUNERT E., VERDCOURT B., BALLY F.R.O., RADCLIFFE-SMITH A. and NEWTON L.E., 1973: Tropical African plants. Part 33. Kew Bull. 28(2), 319-326.
- HERBST R.P. and HARTMAN R.T., 1981: Phytoplankton distribution of a duckweed covered pond. J.Freshwater Ecol. 1(1), 97-112.
- HERTEL W., 1986: Morphologische und chemische Charakterisierung der Licht- und Dunkelkulturen von *Spirodela polyrrhiza* (L.) Schleiden. Wiss.Z.Friedrich-Schiller-Univ.Jena, Naturw.R. 35(5), 591-598.
- HERTER W.G., 1954: Flora del Uruguay. 1. Lemnaceae. Rev.Sudam.Bot. 9, 184-186.
- HESS A.D. and HALL T.F., 1945: The relation of plants to malaria control on impounded waters with a suggested classification. J.Nat.Malar.Soc. 4, 20-46.
- HESS W.J., 1986: Wolfia papulifera Thompson (Lemnaceae), new to Michigan. Sida 11(4), 407-411.

- HESENLAND M. and FROMM F., 1932: Die Wirkung von Natriumchlorat auf Wasserpflanzen. Chem.-Z. 56, 326.
- HESENLAND M., FROMM F. and SAALMAN L., 1933: Die Wirkung von Chlorat, Bromat und Jodat auf pflanzenwuchs. Ang.Chemie 46, 577-579.
- HEVLY R.H., 1961: Notes on aquatic flowering plants with four additions to the Arizona flora. Plateau 33, 115-119.
- HICKS L.E., 1930: Physiological experiments with the Lemnaceae. Proc. Ohio Acad.Sci. 8, 393-394.
- HICKS L.E., 1932a: Flower production in the Lemnaceae. Ohio J.Sci. 32, 115-127.
- HICKS L.E., 1932b: Ranges of pH tolerance of the Lemnaceae. Ohio J.Sci. 32, 237-244.
- HICKS L.E., 1937: The Lemnaceae of Indiana. Am.Midl.Nat. 18, 774-789.
- HICKS P., 1934: Interaction of factors in the growth of Lemna. V. Some preliminary observations upon the interaction of temperature and light on the growth of Lemna. Ann.Bot. 48, 515-523.
- HIDAKA S. and SHIBA H., 1983: The chemical characters of the aqueous endosperm in the rice fields with polluted water. Effect of Lemna and viroment in the rice fields with polluted water. (In Japan.) Jap.J.Soil Sci.Plant Nutr. 54(5), 429-433.
- HIGHAM B.M. and SMITH H., 1969: The induction of flowering by abscisic acid in Lemna perpusilla 6746. Life Sci. 8(20/2), 1061-1065.
- HILBIG W., 1971: Übersicht über die Pflanzengesellschaften des südlischen Teiles der DDR. I. Die Wasserpflanzengesellschaften. Hercynia N.F. 8, 4-33.
- HILD J. and REHNELT K., 1965: Oeko-soziologische Untersuchungen an einjährigen niederrheinischen Kolken. Ber.Deutsch.Bot.Ges. 78, 289-304.
- HILDEBRAND S.F., 1925: A study of the top minnow Gambusia holbrooki in its relation to mosquito control. Publ.Health Bull. 156, 98-103.
- HILLMAN W.S., 1954: On the mechanism of action of benzimidazole on Lemna minor. Ph.D.Thesis. Yale Univ.
- HILLMAN W.S., 1955: The action of benzimidazole on Lemna minor. Plant Physiol. 30, 535-542.
- HILLMAN W.S., 1957: Nonphotosynthetic light requirement in Lemna minor and its partial satisfaction by kinetin. Science 126, 165-166.
- HILLMAN W.S., 1959: Photoperiodic control of flowering in Lemna perpusilla. Nature 181, 1275.
- HILLMAN W.S., 1959a: Experimental control of flowering in Lemna. I. General methods. Photoperiodism in L. perpusilla 6746. Am.J.Bot. 46, 466-473.
- HILLMAN W.S., 1959b: Experimental control of flowering in Lemna. II. Some effects of medium composition, chelating agents and high temperatures on flowering in L. perpusilla 6746. Am.J.Bot. 46, 489-495.
- HILLMAN W.S., 1960a: Growth promotion by kinetin of Wolffia columbiana grown in excessively concentrated medium. Phytion(Argentina) 14, 43-46.
- HILLMAN W.S., 1960b: Effects of gibberellic acid on flowering, frond size and multiplication rate of Lemna perpusilla. Phytion(Argentina) 14, 49-54.
- HILLMAN W.S., 1961 see 1961a
- HILLMAN W.S., 1961a: The Lemnaceae, or duckweeds. A review of the descriptive and experimental literature. Bot.Rev. 27, 221-287.
- HILLMAN W.S., 1961b: Photoperiodism, chelating agents, and flowering of Lemna perpusilla and L. gibba in aseptic culture. In: MELROY W.D. and GLASS B. (eds.), A symposium on light and life. John Hopkin's Press, Baltimore. 673-686.

- HILLMAN W.S., 1961c: Experimental control of flowering in Lemna. 3. A relationship between medium composition and the opposite photoperiodic responses of L. perpusilla 6746 and L. gibba G3. Am.J.Bot. 48, 413-419.
- HILLMAN W.S., 1961d: Test-tube studies on flowering: experiments with the Lemnaceae, or duckweeds. Bull.Forr.Bot.Club 88, 327-336.
- HILLMAN W.S., 1961e: Heavy metals and the photoperiodic control of flowering in Lemna. Plant Physiol. 36(suppl.), 53.
- HILLMAN W.S., 1962: Experimental control of flowering in Lemna. 4. Inhibition of photoperiodic sensitivity by copper. Am.J.Bot. 49, 892-897.
- HILLMAN W.S., 1963: Photoperiodism: an effect of darkness during the light period on critical night length. Science 140, 1397-1398.
- HILLMAN W.S., 1964: Endogenous circadian rhythms and the response of Lemna perpusilla to skeleton photoperiods. Am.Natur. 98, 323-328.
- HILLMAN W.S., 1965: Red light, blue light, and copper ion in the photoperiodic control of flowering in Lemna perpusilla 6746. Plant Cell Physiol. 6, 499-506.
- HILLMAN W.S., 1966: Photoperiodism in Lemna: reversal of night-interruption depends on color of the main photoperiod. Science 154, 1360-1362.
- HILLMAN W.S., 1967: Blue light, phytochrome and the flowering of Lemna perpusilla 6746. Plant Cell Physiol. 8, 467-473.
- HILLMAN W.S., 1969 see 1969c
- HILLMAN W.S., 1969a: Analysis of photoperiodic timing by the use of a rhythmic component in the carbon dioxide evolution of Lemna perpusilla. Plant Physiol. 44(suppl.), 14.
- HILLMAN W.S., 1969b: Analysis of photoperiodic timing by the use of a rhythmic component in the carbon dioxide evolution of Lemna. 11ch Int.Bot.Congr. Abstr., 91.
- HILLMAN W.S., 1969c: Lemna perpusilla Torr., strain 6746. In: EVANS L.T. (ed.), The induction of flowering. Mac Millan of Australia. 186-204.
- HILLMAN W.S., 1969d: Experimental control of flowering in Lemna II. Some effects of medium composition, chelating agents and high temperatures on flowering in Lemna perpusilla. Am.J.Bot. 46, 489-495.
- HILLMAN W.S., 1970a: Carbon dioxide output as an index of circadian timing in Lemna photoperiodism. Plant Physiol. 45, 273-279.
- HILLMAN W.S., 1970b: Effects of the medium on the entrainment of Lemna perpusilla carbon dioxide output by 24 hr light dark cycles. Plant Physiol. 46(suppl.), 26.
- HILLMAN W.S., 1971a: Nitrate and the course of Lemna perpusilla carbon dioxide output under daily photoperiodic cycles. Plant Physiol. 47, 431-434.
- HILLMAN W.S., 1971b: Entrainment of Lemna CO₂ output through phytochrome. Plant Physiol. 47(suppl.), 13.
- HILLMAN W.S., 1971c: Entrainment of Lemna CO₂ output through phytochrome. Plant Physiol. 48, 770-774.
- HILLMAN W.S., 1972a: Photoperiodic entrainment patterns in the CO₂ output of Lemna perpusilla 6746 and of several other Lemnaceae. Plant Physiol. 49, 907-911.
- HILLMAN W.S., 1972b: Interaction of light and rhythms. Book of Abstracts, 6th Int.Photobiol.Congr., Bochum, FRG. 027.
- HILLMAN W.S., 1974: Towards a real-time analysis of photoperiodism: effects of nitrogen source on timing and phytochrome response in Lemna carbon dioxide output. Am.J.Bot. 61(suppl.), 28.
- HILLMAN W.S., 1975a: Photoperiodism in seedling strains of Lemna perpusilla.

- silla: juvenility without obvious morphological correlates? *Am.J.Bot.* 62, 537-540.
- HILLMAN W.S., 1975b: Effects of inorganic nitrogen on the response of Lemna carbon dioxide output to light quality and timing. *Photochem. Photobiol.* 21, 39-47.
- HILLMAN W.S., 1975c: Light timer interactions in photoperiodism and carbon dioxide output. *Plant Physiol.* 56(2 suppl.), 74.
- HILLMAN W.S., 1976a: Calibrating duckweeds: light, clocks, metabolism, flowering. Special characteristics of Lemnaceae may offer unique insights into plant development. *Science* 193, 453-458.
- HILLMAN W.S., 1976b: A metabolic indicator of photoperiodic timing. *Proc.Nat.Acad.Sci. USA* 73, 501-504.
- HILLMAN W.S., 1976c: Light/timer interactions in photoperiodism and carbon dioxide output patterns: towards a real-time analysis of photoperiodism. In: Smith H. (ed.), *Light and plant development*. Butterworth, London/Boston. 383-397.
- HILLMAN W.S., 1977a: The effects of light and nitrogen sources on the carbon dioxide output patterns of heterotrophic Lemna are paralleled in oxygen uptake as measured by intermittent flow. *Plant Physiol.* 59 (6 suppl.), 92.
- HILLMAN W.S., 1977b: The control of chronobiological patterns in plant respiration. *Chronobiologia* 4(2), 119.
- HILLMAN W.S., 1977c: Control of plant respiration through non-photosynthetic light action. *Nature* 266, 833-835.
- HILLMAN W.S., 1979a: Temporal compartmentation in Lemna paucicostata: Photoperiodism, respiration, nitrogen nutrition and heterotrophic growth of different strains. *Am.J.Bot.* 66, 1021-1028.
- HILLMAN W.S., 1979b: Temperature sensitivity of daily respiratory patterns entrained through phytochrome action in Lemna paucicostata strain 6746. *Physiol.Plant.* 47, 56-60.
- HILLMAN W.S. and CULLEY D.D., 1978a: The uses of duckweed. *Am.Sci.* 66, 442-451.
- HILLMAN W.S. and CULLEY D.D., 1978b: Cultivating duckweeds. 32 pp. (Photocopy).
- HILLMAN W.S. and LAMM S.S., 1981: Occurrence and reversal in Lemna paucicostata 6612 of a condition that prevents nitrate utilization. *Plant Sci.Lett.* 20, 325-331.
- HILLMAN W.S. and POSNER H.B., 1971: Ammonium ion and the flowering of Lemna perpusilla. *Plant Physiol.* 47, 586-587.
- HILTIERAN R.C., 1973: Duckweed control. III. *Nat.Hist.Surv., Aquatic Biol., Urbana.* 2 pp.
- HILTIERAN R.C., 1978: The chemical control of some aquatic plants. III. *Nat.Hist.Surv., Aquatic Biol., Urbana* 12(suppl.), 11 pp.
- HILTON-BIGGS R. and KOSSUTH S.V., 1980: Developmental effects of Sandoz 6706 and UV-B radiation on *Spirodela punctata* and Lemna minor. *Plant Physiol.* 65(6 suppl.), 83.
- HIMES C.L., HERBERT M., BROWN W. and FEESE W., 1967: A limnological study of a shallow alkaline pond. *Proc.PA.Acad.Sci.* 41, 49-53.
- HINDS H.R., 1983: The rare vascular plants of New Brunswick. *Syllogues* 50, 38 pp.
- HORGAND R.E., 1978: The effects of N-(phosphonomethyl)glycine on selected enzymes in Lemna gibba L. *Proc. 31st Ann.Meet.Southern Weed Sci.Soc., Abstr.* 285.
- HORGAND R.E. and DUKE S.O., 1982: Biochemical effects of glyphosate (N-phosphonomethylglycine). *ACS Symp.Ser.* 181, 175-205.

- HORGAND R.E. and PAUL R., 1978: Ultrastructural effects of glyphosate on Lemna gibba L. *Meet.Weed Sci.Soc.America, Abstr.* 78.
- HODGSON G.L., 1970: Effects of temperature on the growth and development of Lemna minor, under conditions of natural daylight. *Ann.Bot.* 34, 365-381.
- HODSON H.K. and HAMNER K.C., 1970: Floral inducing extract from Xanthium. *Science* 167(3917), 384-385.
- HODSON H.K. and HAMNER K.C., 1971: A comparison of the effects of autoclaved and non-autoclaved gibberellic acid on Lemna perpusilla 6746. *Plant Physiol.* 47, 726-728.
- HOESEL W., FREY G., TEUFEL E. and BARZ W., 1972: Ueber Vorkommen und Lokalisation eines flavonolumwandelnden Enzyms in Pflanzen. *Planta* 103, 74-86.
- HOFFMAN-PALK H., MATTOO A.K., MARDER J.B., EDELMAN M. and ELLIS R.J., 1982: General occurrence and structural similarity of the rapidly synthesized, 32000-dalton protein of the chloroplast membrane. *J.Biol.Chem.* 257(8), 4583-4587.
- HOFFMANN J.F., 1840a: Beiträge zur näheren Kenntnis von Lemna arthiza nebst einigen Bemerkungen über L. polyrrhiza, L. gibba, L. minor und L. trisulca. *Wiegmanns Arch.Naturgeschichte(Berlin)* 6, 138-163.
- HOFFMANN J.F., 1840b: Matériaux pour servir à la connaissance de Lemna arthiza, avec quelques observations sur les autres espèces de ce genre. *Ann.Sci.Nat.*, 2e sér., 14, 223-242.
- HOFFMANN-OSTENHOF O., PITTMER F. and KOLLER F., 1978: Some enzymes of inositol metabolism, their purification and their mechanism of action. In: WELLS W.W. and EISENBERG F., Jr. (eds.), *Cyclitols and phosphoinositides*. *Proc.Symp.East Lansing, Mich., USA. Acad.Press, New York.* 233-247.
- HOFMEISTER W., 1861: Neue Beiträge zur Kenntnis der Embryobildung der Phanerogamen. 2. Monokotyledonen. *Abh.Math.-Phys.Cl.K.Sächs.Ges.Wiss.* 5, 716 pp.
- HOGWEG P. and BRENKERT A.L., 1969: Structure of aquatic vegetation: A comparison of aquatic vegetation in India, the Netherlands and Czechoslovakia. *Trop.Ecol.* 10, 139-162.
- HOLLAND K., 1983: Extracts of Cortinarius speciosissimus affecting the photosynthetic apparatus of Lemna minor. *Trans.Br.Mycol.Soc.* 81(3), 633-635.
- HOLMQUIST C., 1971: Northerly localities for three aquatic plants, Lemna trisulca L., Ceratophyllum demersum L., and Myriophyllum spicatum L. *Bot.Not.* 124, 335-342.
- HOLST R.W. and YOPP J.H., 1979: Comparative utilization of inorganic and organic compounds as sole nitrogen sources by the submerged duckweed, Lemna trisulca L. *Biol.Plant.* 21, 245-252.
- HOMES M.V., 1963: The method of systematic variations. *Soil Sci.* 96, 380-386.
- HONDA S.I., 1983a: Cyclic AMP and cyclic GMP effects on chloroplasts. *Plant Physiol.* 72(1 suppl.), 55.
- HONDA S.I., 1983b: Some effects of cyclic nucleotide treatment on chloroplast growth and replication in Wolffia arrhiza 7014. *J.Cell Biol.* 97(5/2), 363a.
- HONDA Y., KOGA T., MORISHIMA H., SATO Y., KITAWAKI S. and NISHIWAKI Y., 1971: Uptake and loss of radioactive ruthenium by freshwater organisms. (In Japan.). *Radioisotopes* 20(8), 376-382.
- HOOGERS B.J. and VAN DER WEIJ H.G., 1971: Time tables as a method to determine the dynamic nature of ditch vegetation. Wageningen. (In Dutch). 12 pp. From *Excerpta Bot.* B 14, 41. (Photocopy).

- HOPKINS D.W., GOLDBERG G. and KLEIN W.H., 1978: Simulation of UV-B expected at various atmospheric ozone levels and its effect on growth of plants. *Plant Physiol.* 61(4 suppl.), 74.
- HOPKINS E.F., 1931: Manganese and the growth of *Lemna minor*. *Science* 74, 551-552.
- HOPKINS E.F., 1934: Manganese an essential element for green plants. *Cornell Univ. Agr. Exp. Stat. Mem.* 151, 3-40.
- HORN AF RANTZEN H., 1951: Certain aquatic plants collected by Dr. J.T. Baldwin Jr. in Liberia and the Gold Coast. *Bot. Not.* 1919, 368-398.
- HORST K., 1978: Die Zwerg-Wasserlinse (*Wolffia arrhiza*) - kleinste Blü-
tenpflanze der Welt im Maujahn bei Dannenberg. *Hannov. Wendland* 7, 9-18.
- HORWITZ B.A. and SAMISH Y.B., 1975: Light-stimulated bioelectric re-
sponse in *Spirodela oligorrhiza* and its relation to photosynthesis. *Z. Pflanzenphysiol.* 76(2), 182-189.
- HOSSETTI B.B. and PATIL H.S., 1986: Impact of *Lemna minor* on the effluent quality of sewage stabilization pond. *Geobios (Jodhpur)* 13(6), 244-247.
- HOSHINO T., 1979: Simulation of acetylcholine action by -indole acetic acid in inducing diurnal change of floral response to chilling under continuous light in *Lemna gibba* G3. *Plant Cell Physiol.* 20(1), 43-50.
- HOSHINO T. and OOTA Y., 1978: The occurrence of acetylcholine in *Lemna gibba* G3. *Plant Cell Physiol.* 19(5), 769-776.
- HOSSELL J.C. and BAKER J.H., 1976: The distribution and characterization of bacteria on the surfaces of some river macrophytes. *J. Appl. Bacteriol.* 41(3), 14-15.
- HOSSELL J.C. and BAKER J.H., 1979a: Estimation of the growth rates of epiphytic bacteria and *Lemna minor* in a river. *Freshw. Biol.* 9, 319-327.
- HOSSELL J.C. and BAKER J.H., 1979b: A note on the enumeration of epiphytic bacteria by microscopic methods with particular reference to two freshwater plants. *J. Appl. Bacteriol.* 46(1), 87-92.
- HOUGH R.A. and WETZEL R.G., 1977: Photosynthetic pathways of some aquatic plants. *Aquat. Bot.* 3, 297-313.
- HOWARD R.A., 1979a: Nomenclatural notes on some lesser Antillean monocotyledones. *J. Arnold Arb.* 60, 290-301.
- HOWARD R.A., 1979b: Flora of the lesser Antilles 3. Monocotyledonae. *Arnold Arb.* Harvard Univ., Jamaica Plain. 401-403.
- HOWARD-WILLIAMS C., 1977: A check-list of the vascular plants of Lake Chilwa, Malawi, with special reference to the influence of environmental factors in the distribution of taxa. *Kirkia* 10, 563-580.
- HOWARD-WILLIAMS C., 1979: The distribution of aquatic macrophytes in Lake Chilwa: Annual and long-term environmental fluctuations. In: KALLIK M., MCLACHLAN A.J., HOWARD-WILLIAMS C. (eds.), *Lake Chilwa, studies of change in a tropical ecosystem*. Monogr. Biol. 35, 105-122.
- HUBAC J.-M., BEUFFE H., BLAKE G., CORRADI M., DUTARTRE A., VAUCOULOUX M. and VUILLOT M., 1984: Les plantes aquatiques utiles: Les Lentilles d'Eau (*Lemnaceae*), utilisation en phyto-épuration et valorisation. *Assoc. Fr. Etude des Eaux*, Paris. 115 pp.
- HUBALD M., 1975: Das Wachstum und die Entwicklung der Wasserlinse (*Lemna gibba* L. G1) unter besonderer Berücksichtigung des Einflusses verschiedener Zucker und Stickstoffverbindungen. Ph.D. Thesis. Jena.
- HUBALD M. and AUGSTEN H., 1977a: Einfluss verschiedener Zucker und Stickstoffverbindungen auf Wachstum und Entwicklung von *Lemna gibba* L. *Beitr. Biol. Pflanz.* 53, 91-102.

- HUBALD M. and AUGSTEN H., 1977b: The ultrastructure of duckweed chloroplasts (*Lemna gibba* L. G1) influenced by glycine and deficiency conditions. *Acta Biol. Med. Exp.* 2, 61-64.
- HUBALD M. and AUGSTEN H., 1979: Die Beziehungen zwischen Licht-Dunkelwechsel und Nitratreduktaseaktivität bei *Wolffia arrhiza* (L.) Wimm. *Biochem. Physiol. Pflanzen* 174(7), 555-561.
- HUBALD M., AUGSTEN H. and FEIST H., 1979: Einfluss von Aminosäuren auf den Aminosäure- und Proteingehalt bei *Lemna gibba* L. *Biol. Rundschau* 17, 196-199.
- HUBER W., 1982: Ueber die Wirkungen eines Salzstresses auf die Stickstoffassimilation von *Lemna minor* L. *Biochem. Physiol. Pflanzen* 177(3), 259-265.
- HUBER W., 1985: Xanthoxin - ein neues "Stresshormon" in Pflanzen? *Verh. Ges. Oekol.* 13, 499-503.
- HUBER W. and SANKHLA N., 1979: Effect of sodium chloride on photosynthesis of *Lemna minor* L. *Z. Pflanzenphysiol.* 91, 147-156.
- HUBER W. and SANKHLA N., 1981 see 1979
- HUBER W., SCHUBERT V. and SAUTTER C., 1982: Effects of pentachlorophenol on the metabolism of the aquatic macrophyte *Lemna minor* L. *Environ. Pollut. A* 29(3), 215-224.
- HUEGEL B., 1974: In vitro-Kultur der Blütenstandsanlagen von *Lemna gibba* und *Lemna paucicostata*. Vergleichende Untersuchungen des Hormonbedarfs für Entwicklung und Geschlechtsausprägung beider Arten. *Diss. Würzburg*. 80 pp.
- HUEGEL B., 1976a: Gegensätzliche Geschlechtsausprägung von Blütenstandsanlagen der Langtagpflanze *Lemna gibba* und der Kurztagpflanze *Lemna paucicostata* in vitro. *Z. Pflanzenphysiol.* 77, 395-405.
- HUEGEL B., 1976b: Wirkung von Gibberellin-A₃, CCC, Ethrel und Indol-essigsäure auf die Geschlechtsausprägung isolierter Blütenstandsanlagen von *Lemnaceen*. *Z. Pflanzenphysiol.* 80, 283-297.
- HUEGEL B., 1976c: Wirkung von Kinetin und Abscisinsäure auf die Entwicklung von *Lemnaceen*-Blütenstandsanlagen in vitro. *Z. Pflanzenphysiol.* 80, 298-305.
- HUEGEL B. and KANDELER R., 1974: Hormonbedarf in vitro bei Blütenanlagen einer Kurztagpflanze (*Lemna paucicostata* 6746) und einer Langtagpflanze (*Lemna gibba* G1). *Zusammenfass. Vorträge Deutsch. Bot. Ges. u. Verein Angew. Bot.*, Tagung Würzburg. 17-18.
- HUEGEL B., ROTTENBURG T. and KANDELER R., 1979: Phytochromsteuerung der Turionbildung und anderer Entwicklungsprozesse bei *Lemna perpusilla* P 146. *Biochem. Physiol. Pflanzen* 174, 761-771.
- HUFFMAN D.C., 1980: Economic feasibility of methane generation and production of duckweed for feed on dairy farms in Southeast United States. *Baton Rouge*. 37 pp. (Polycopy).
- HUFFMAN E.W.D., Jr. and ALLAWAY W.H., 1973: Growth of plants in solution culture containing low levels of chromium. *Plant Physiol.* 52(1), 72-75.
- HULTEN E., 1927: Flora of Kamtschatka and the adjacent islands I. *Almqvist and Wiksell*, Stockholm. 331.
- HULTEN E., 1968: Flora of Alaska and neighboring territories. *Stanford Univ. Press.*, Stanford. 282.
- HULTEN E., 1971: Atlas of the distribution of vascular plants in North-western Europe. (2nd ed.). (in Swed.). *Stockholm*. 531 pp.
- HUMBOLDT see VON HUMBOLDT
- HUMPHREY T.J. and DAVIES D.D., 1974: A new method for the measurement of protein turnover. *Biochem. J.* 148, 119-128.
- HUMPHREY T.J. and DAVIES D.D., 1976: A sensitive method for measuring

protein turnover based on the measurement of two tritium labeled amino acids in protein. Biochem.J. 156(3), 561-568.

HUMPHREY T.J., SARAWAK S. and DAVIES D.D., 1977: The effect of nitrogen deficiency on the growth and metabolism of Lemna minor L. Planta 137, 259-264.

HUNT R., 1978: Plant growth analysis. Studies in biology 96. Edward Arnold. ISBN 0-7171-2696-5

HUNTER R.D., 1976: Changes in carbon and nitrogen content during decomposition of three macrophytes in freshwater and marine environments. Hydrobiol. 51, 119-128.

HUNYADI K., 1973: The biotest method for determining bipyridyl herbicides. Keszthelyi Mezogazdasagutmonayi Kar Kozlemenyei 15(4), 12 pp.

HUPPATZ J.L. and PHILLIPS J.N., 1981: Cyanoacrylate herbicides. CSIRO Div.Pl.Ind. 1, 85.

HUSAK S. and OTAHELOVA H., 1982: Wolffia arrhiza (L.) Horkel ex Wimmer in Slovakia. (In Slovak.). Biologia (Bratislava) 37(9), 933-935.

HUTCHINSON J., 1934: The families of flowering plants. (2nd ed.). Vol. 2. Monocotyledons. Clarendon Press, Oxford.

HUTCHINSON T.C., 1974: Heavy metal toxicity and synergism to floating aquatics. In: MARSHALL K.E. (ed.), 19th Congr.Int.Assoc.Limnology, Winnipeg, Canada. 91.

HUTCHINSON T.C. and CZYRSKA H., 1975: Heavy metal toxicity and synergism to floating aquatic weeds. Verh.Int.Ver.Limnol. 19, 2102-2111.

HUTNER S.H., 1953: Comparative physiology of heterotrophic growth. In: LOOMIS W.E. (ed.), Growth and differentiation in plants. Iowa State Coll.Press, 417-446.

HUTNER S.H., PROVANSOLI L., SCHATZ A. and HASKINS C.P., 1950: Some approaches to the study of the role of metals in the metabolism of microorganisms. Proc.Am.Phil.Soc. 94, 150-170.

IAKOVLEVA N.V., 1980: Sensitivity of plants to sulfur dioxide. (In Russian). Gaz.Rast. 18, 153-154.

IBRAHIM A.R. and JUNGNIKKEL F., 1980: Utilization of nitrate in Lemna minor under autotrophic and mixotrophic conditions. Acta Biol.Med. Exp. 5, 11-17.

ICE J. and COUCH R., 1987: Nutrient absorption by duckweed. J.Aquat. Plant Manage. 25, 30-31.

IKUSIMA I., 1955: Growth of duckweed populations as related to frond density. (In Japan.). Physiol.Ecol.Jap. 6, 69-81.

IKUSIMA I., 1962: Biology of duckweeds with special reference to their growth. I. (In Japan.). Physiol.Ecol.Jap. 10, 130-164.

IKUSIMA I., 1963a,b: Biology of duckweeds with special reference to their growth. II and III. (In Japan.). Physiol.Ecol.Jap. 11, 84-102; 11, 120-137.

IKUSIMA I. and KIRA T., 1958: Effect of light intensity and concentration of culture solution on the frond multiplication of Lemna minor L. (In Japan.). Physiol.Ecol.Jap. 8, 50-60.

IKUSIMA I., SHINOZAKI K. and KIRA T., 1955: Intraspecific competition among higher plants. III. Growth of duckweed, with a theoretical consideration on the C-D effect. J.Inst.Polytech. Osaka City Univ. D, 6, 107-119.

IMANAKA M., HINO S., KENMOCHI K., MATSUNAGA K. and ISHIDA T., 1986: Studies on toxicants in food. 10. Bioaccumulation of oxadiazon in the aquatic environment. (In Japan.). Okayama-ken Kankyo Hoken Senta Nenpo 10, 122-124.

INGEMARSSON B., 1986: Patterns of N uptake, accumulation and assimilation in Lemna gibba when grown with exponentially increasing daily doses of N. In: LAMBERS H., NETESON J.J. and STULEN I. (eds.), Fundamental, ecological and agricultural aspects of nitrogen metabolism in higher plants. Nijhoff Publishers, Dordrecht, Netherlands. 65-69.

INGEMARSSON B., JOHANSSON L. and LARSSON C.-M., 1983: N utilization, photosynthesis and growth in N-limited cultures of Lemna. In: CRAM W.J. et al. (eds.), Membrane transport in plants. Wiley, New York. 402-403.

INGEMARSSON B., JOHANSSON L. and LARSSON C.-M., 1984: Photosynthesis and nitrogen utilization in exponentially growing nitrogen-limited cultures of Lemna gibba. Physiol.Plant. 62(3), 363-369.

INGEMARSSON B., OSCARSON P. and LARSSON C.-M., 1985: Intracellular compartmentation of NO₃⁻ in Lemna; effects on NO₃⁻ uptake and assimilation. Physiol.Plant. 64(2), 17A.

INHUEISEN D. and NIEMEYER R., 1975: Kondensierte Phosphate in Lemna minor L. und ihre Beziehungen zu den Nucleinsäuren. Planta 124, 159-167.

INHUEISEN D. and NIEMEYER R., 1978: Inosit-Phosphate aus Lemna minor L. Z.Pflanzenphys. 88, 103-116.

INOUE Y. and SHIBATA K., 1973: Light induced chloroplast rearrangements and their action spectra as measured by absorption spectrophotometry. Planta 114(4), 341-358.

IRBE I.K. and VORONINA O.Y., 1983: Action of 2,4-D herbicide on the photosynthetic apparatus of higher aquatic plants. (In Russian). Mezhevuz.SB.Nauchn.Tr. 204, 48-52.

IRBE I.K. and VORONINA O.Y., 1986: Effect of 2,4-D on pigment system of duckweed in relation to pH of aquatic environment. (In Russian). Ref. Zh., Biologiya, 14-22.

ISENSEE A.R. and JONES G.E., 1975a: Distribution of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in aquatic model ecosystem. Environ.Sci.Techn. 9(7), 668-672.

ISENSEE A.R. and KEARNEY P.C., 1975b: Distribution of carbaryl and 3,5-xylylmethylcarbamate in an aquatic model ecosystem. J.Agric.Food Chem. 23(4), 760-763.

ISHIGURI Y. and ODA Y., 1972: The relationship between red and far-red light on flowering of the long-day plant, Lemna gibba. Plant Cell Physiol. 13, 131-138.

ISHIGURI Y. and ODA Y., 1974: Flowering of the long-day plant, Lemna gibba, under short-day schedules composed of red and far-red light. Plant Cell Physiol. 15, 287-293.

ISHIGURI Y. and ODA Y., 1976: Photoreversibility of flower initiation in Lemna perpusilla as affected by the light quality of the main light period. Plant Cell Physiol. 17(2), 255-264.

ISHIGURI Y. and ODA Y., 1979: Diurnal changes of red and far-red reversible reaction inducing flowering of a long-day plant, Lemna gibba. Rep.Inst.Agric.Res., Tohoku Daigaku, 30, 13-19.

ISHIGURI Y., ODA Y. and INADA K., 1975: Spectral dependences of flowering in Lemna perpusilla and L. gibba. Plant Cell Physiol. 16, 521-523.

IUNUSOV I. and KELDIBEKOV S., 1972: On the mass culture of Wolffia arrhiza Wimmer in open pools of water. (In Russian). Kuit.Vod.Vyss. Vod.Rast.Uzb. 116-119.

IVANOVA I.E., 1970: Certain characteristic features of flowering and pollination in duckweeds (Lemnaceae: S. Gray). (In Russian). Bot.Zh. (USSR) 55, 649-659.

- IVANOVA I.E., 1973: On the taxonomy of Lemnaceae S. Gray. (In Russian). Bot. Zh. (USSR) 58, 1413-1423.
- IVERSEN J., 1929: Studien über die pH-Verhältnisse dänischer Gewässer und ihren Einfluss auf die Hydrophytenvegetation. Bot. Tidsskr. 40, 277-333.
- IVERSEN J. and OLSON S., 1946: Die Verbreitung der Wasserpflanzen in Relation zur Chemie des Wassers. Bot. Tidsskr. 46, 136-145.
- IVES J.M. and POSNER H.B., 1982: Epinephrine, propranolol, and the sucrose ammonium inhibition of flowering in *Lemna paucicostata* 6746. Plant Physiol. 70(1), 311-312.
- JACOBS D.L., 1947: An ecological life history of *Spirodela polyrrhiza* (greater duckweed) with emphasis on the turion phase. Ecol. Monogr. 17, 437-469.
- JACOBS D.L., 1949: *Wolffia papulifera* in Florida. Am. Midl. Nat. 42, 110-111.
- JACOBS D.L., 1950: Ecological and taxonomical status of Georgia duckweeds. J. Tenn. Acad. Sci. 25, 228.
- JACQUET J. and BOUTIBONNES P., 1970: Effets des flavocoumarines sur quelques animaux et végétaux. C.R. Soc. Biol. 164(11), 2239-2244.
- JACQUET J., BOUTIBONNES P. and SAINT S., 1971: Effets biologiques des flavocoumarines d'*Aspergillus parasiticus* ATCC 15517. II. Végétaux. Rev. Immunol. 35(4-5), 159-186.
- JAEGER E., 1964: Zur Deutung des Arealbildes von *Wolffia arrhiza* (L.) Wimmer und einiger anderer ornithochorer Wasserpflanzen. Ber. Deutsch. Bot. Ges. 77, 101-111.
- JAEGER E., 1965: *Wolffia arrhiza* (L.) Wimmer, gefunden im Herbst. 1963 bei Hartmannsdorf, Kreis Lübben/Spreewald. Niederlaus. Florist. Mitt. 1, 40.
- JANAS K.M., 1980: The growth retarding effect of daminozide in *Spirodela oligorrhiza* and its reversal by gibberellin. Acta Soc. Bot. Pol. 49 (4), 471-476.
- JANAUER G.A., 1977: Amino acids in aquatic macrophytes. Z. Pflanzenphysiol. 82, 45-50.
- JANAUER G.A., 1982: The organic and mineral composition of some aquatic plants. In: SYMOENS S., HOOPER S.S. and COMPERE P. (eds.), Studies on aquatic vascular plants. R. Soc. Bot. Belg., Bruxelles. 136-142.
- JANES G.A., 1975: Controlled release copper herbicides. Creative Biol. Lab. Inc., Barberton, Ohio, USA, 326-333.
- JARCZYK L., REUTLINGER G. and ULLRICH W.R., 1984: Nitrataufnahme und Aktivität der Nitratreduktase bei *Lemna gibba* unter verschiedener Begasung. Mitt. bd. Bot.-Tag. Wien, 55(0514).
- JAWORSKI E.G., 1971: Gamma methylthiobutyric acid, a growth stimulant. Life Sci., II (Biochem. Gen. Mol. Biol.) 10(11), 613-616.
- JAWORSKI E.G., 1972: Mode of action of N-phosphonomethylglycine: inhibition of aromatic amino acid biosynthesis. J. Agric. Food Chem. 20(6), 1195-1198.
- JEFFERIES R.L., LAYCOCK D., STEWART G.R. and SIMS A.P., 1969a: The properties of mechanisms involved in the uptake and utilization of calcium and potassium by plants in relation to an understanding of plant distribution. In: RORISON I.H. (ed.), Ecological aspects of the mineral nutrition of plants. Blackwell, Oxford. 281-307.
- JEFFERIES R.L., SIMS A.P. and YOUNG M., 1969b: Regulation of potassium transport in *Lemna minor*. 11th Int. Bot. Congr., Abstr., 102.
- JENKINS D.W., 1964: Pathogens, parasites and predators of medically important arthropods. Bull. WHO 30(suppl.), 150 pp.

- JENSEN S., 1979: Classification of lakes in southern Sweden on the basis of their macrophyte composition by means of multivariate methods. Vegetatio 39(3), 129-146.
- JENTSCH H., 1979: Vorkommen und Vergesellschaftung von *Wolffia arrhiza* (L.) Horkel ex Wimmer im Spreewald. Gleditschia 7, 251-254.
- JEREMIE J. and RAYNAL-ROQUES A., 1981: Dynamique de la végétation des mares de dolines aux Petites Antilles. Adansonia 3, 259-280.
- JERVIS R.A., 1969: Primary production in the freshwater marsh ecosystem of Troy Meadows, New Jersey. Bull. Torr. Bot. Club 96, 209-231.
- JOHANSSON B., 1980: On developing embryo sacs of the Angiosperms. (In Swed.). Lunds. Univ. Arskr. 16(5), 86 pp.
- JOHANSEN D.A., 1950: Plant embryology. Chronica Botanica. Waltham, Mass.
- JOHANSSON L. and LARSSON C.-M., 1985: Carbon and nitrogen metabolism in *Lemna* as affected by methionine sulfoximine (MSO). Plant Physiol. 64 (suppl. 2), 17a.
- JOHANSSON L. and LARSSON C.-M., 1986a: Relationship between inhibition of CO₂ fixation and glutamine synthetase inactivation in *Lemna gibba* L. treated with L-methionine-D,L-sulfoximine (MSO). J. Exp. Bot. 37(175), 221-229.
- JOHANSSON L. and LARSSON C.-M., 1986b: Effects of the glutamine synthetase inhibitor methionine sulfoximine on CO₂ fixation in *Lemna gibba*. Plant Soil 91(3), 373-376.
- JOHANSSON L., ANDERSSON B. and ERIKSSON T., 1982: Improvement of anther culture technique: Activated charcoal bound in agar medium in combination with liquid medium and elevated CO₂ concentration. Physiol. Plant. 54(1), 24-30.
- JOHNSON E.L., 1941: Effect of x-radiation upon the growth of *Lemna minor* L. Colorado Univ. Stud., D, Phys. Biol. 1, 165-175.
- JOHNSON H.P., 1902: A study of certain mosquitoes in New Jersey and a statement of the mosquito-malaria-theory. Ann. Rept. N.J. Agr. Exp. Sta.
- JOHRI T.S. and SHARMA P.N., 1980: Studies on utilization of dried duckweed (*Lemna minor*) in chicks. Indian Poul. Gaz. 64(11), 24-27.
- JONES C.N., 1963: Flora of Illinois. (2nd ed.). Notre Dame Univ. Press. 91-92.
- JORDE D.G., KRAPU G.L. and CRAWFORD R.D., 1983: Feeding ecology of mallards wintering in Nebraska USA. J. Wildl. Manage. 47(4), 1044-1053.
- JOUKOVSKY A.V., 1935: La floraison de *Lemna*. Beih. Bot. Zentralblatt Abt. A 53, 620-626.
- JOVET P. and JOVET-AST S., 1966: *Lemna valdiviana* Philippi, espèce signalée pour la première fois en Europe. Bull. Centre Etud. Rech. Sci., Biarritz 6, 57-64.
- JOVET P. and JOVET-AST S., 1967: Floraison, fructification, germination du *Lemna valdiviana* au Lac Marion (B.P.). Bull. Cent. Etud. Rech. Sci. Biarritz 6, 729-734.
- JOVET-AST S., 1968: Contribution à l'étude des eaux douces de l'Ennedi. 2. Lemnaceae. Bull. Inst. Fondam. Afr. Noire 30(3), 830-847.
- JOY K.W., 1968: Enzymes of nitrate assimilation in *Lemna*. Plant Physiol. 43, 8.
- JOY K.W., 1969a: Nitrogen metabolism of *Lemna minor*. I. Growth, nitrogen sources, and amino acid inhibition. II. Enzymes of nitrate assimilation and some aspects of their regulation. Plant Physiol. 44, 845-848, 849-853.
- JOY K.W., 1969b: Glutamate dehydrogenases in higher plants. 11th Int. Bot. Congr., Abstr., 105.
- JOY K.W., 1971: Glutamate dehydrogenase changes in *Lemna* not due to enzyme induction. Plant Physiol. 47, 445-446.

- JOYNER B.G. and FREEMAN T.E., 1973: Pathogenicity of *Rhizoctonia solani* to aquatic plants. *Phytopathology* 63, 681-685.
- JUDD G.J.R. and BORDEN J.H., 1980: Oviposition deterrents for *Aedes aegypti* in extracts of Lemna minor. *J. Entomol. Soc. B.C.* 77, 30-33.
- JUNSON M., 1976: Studies on inhibition factors and the role of phytochrome in the floral induction in short-day plants. *Korean J. Bot.* 19 (1), 14-18.
- JUNG K.D. and LUETTGE U., 1980: Amino acid uptake by Lemna gibba by a mechanism with affinity to neutral L- and D-amino acids. *Planta* 150, 230-235.
- JUNG K.D. and LUETTGE U., 1982a: Mechanisms of amino-acid transport in Lemna gibba L. *Dev. Plant Biol.* 7, 21-28.
- JUNG K.D. and LUETTGE U., 1982b: Inhibition of the amino-acid co-transport carrier of Lemna gibba by incorporation of p-fluorophenylalanine. *Plant Physiol.* 55(2), 231-235.
- JUNG K.D., LUETTGE U. and FISCHER E., 1982: Uptake of neutral and acidic amino acids by Lemna gibba correlated with the H⁺-electrochemical gradient at the plasmalemma. *Plant Physiol.* 55(3), 351-355.
- JUNGnickel F., 1978: Phosphatbedarf und Mangelsymptome bei einigen axenisch kultivierten Lemnaceen. *Limnologica* 11, 469-478.
- JUNGnickel F., 1981: Regulatory properties of surface located hydrolases in *Candida utilis* and in some Lemnaceae. *Abh. Akad. Wiss. DDR, Abt. Math., Natw., Techn.* N3, 421-426.
- JUNGnickel F., 1984: Zur Manganwirkung bei einigen Lemnaceen. *Biol. Rundsch.* 22(4), 255-257.
- JUNGnickel F., 1986a: Turion formation and behaviour in *Spirodela polyrrhiza* at two levels of phosphate supply. *Biol. Plant.* 28(3), 168-173.
- JUNGnickel F., 1986b: Zur Beziehung zwischen Modell- und Praxisobjekten bei der axenischen Mikrovermehrung von Blütenpflanzen. *Wiss. Z. Friedrich-Schiller-Universität Jena, Natw.* 35(5), 599-611.
- JUNGnickel F. and ANDREAS W., 1987: Clonal differences of biomass and turion production in *Spirodela polyrrhiza* (L.) Schleiden as influenced by phosphate. *Acta Biol. Med. Exp. (Priatina)*, in press.
- JUNGnickel F. and AUGSTEN H., 1984: Biotechnical use of immobile cormo-phyte enzymes and multienzyme/cofactor complexes in fixed bed enzyme reactors. 16th Meeting Fed. European Biochem. Soc. Abstr., 30.
- JUNGnickel F. and AUGSTEN H., 1986: *Spirodela polyrrhiza* - ein Biotest-System zur Erfassung phytoaktiver Substanzen. *Wiss. Z. Friedrich-Schiller-Universität Jena, Natw.* 35(5), 613-629.
- JUNGnickel F. and GEBHARD A., 1986: Zur Manganwirkung bei Lemna minuscula unter autotrophen und mixotrophen Ernährungsbedingungen. *Beitr. Biol. Pflanzen* 61, 179-189.
- JUNGnickel F. and IBRAHIM A.R., 1983: Limitation der Biomasse-Bildung bei autotrophen und mixotrophen Kulturen von Lemna minor L. und L. minor F. *Leptophylla* Domin. *Wiss. Z. Ernst-Moritz-Arndt Univ. Greifswald, Math.-Natw.* 32(3-4), 42-44.
- JUNGnickel F. and VOELKSCH B., 1979: Multiple forms of orthophosphoric monoester phosphohydrolase in phosphate starved *Wolffia arrhiza*. *Biochem. Physiol. Pflanz.* 174(1), 44-50.
- JUNGnickel F. and AUGSTEN H. and MICHEL D., 1982: Verfahren zur Stoffwandlung durch immobile Biokatalysatoren. Patent DD 158040, Int. Cl. C 12N 11/16 and C 12N 11/18.
- JUNK W.J., 1982: Zur Entwicklung aquatischer Makrophyten in Curua-Una, dem ersten Stausee in Zentralamazonien. *Arch. Hydrobiol.* 95, 169-180.
- JURD L., GEYSSMAN T.A. and SEIKEL M.K., 1957: The flavonoid constituents of *Spirodela oligorrhiza*. II. The flavone constituents. *Arch. Biochem. Biophys.* 67, 284-297.

- KADONO Y., 1982: Occurrence of aquatic macrophytes in relation to pH, alkalinity, Ca⁺⁺, Cl⁻ and conductivity. *Jap. J. Ecol.* 32(1), 39-44.
- KAFARSKI P., LEJCZAK B., GANCARZ R., JAKULSKA E., MASTALERZ P., WIECZO-REK J.S. and ZBYRYT I., 1985: Phosphonic analogs of morphactins 5. Peptides containing 9-amino-fluoren-9-ylphosphine oxides. *Pestic. Sci.* 16(3), 239-243.
- KAIHARA S. and TAKIMOTO A., 1985a: Flower-inducing activity of vitamin K in Lemna paucicostata. *Plant Cell Physiol.* 26(1), 89-98.
- KAIHARA S. and TAKIMOTO A., 1985b: Induction of flowering in Lemna paucicostata by dicoumarol and 4-hydroxycoumarin. *Plant Cell Physiol.* 26(8), 1465-1472.
- KAIHARA S., WATANABE K. and TAKIMOTO A., 1981: Flower-inducing effect of benzoic and salicylic acids in various strains of Lemna paucicostata and L. minor. *Plant Cell Physiol.* 22(5), 819-825.
- KAJFOSZ J., WALCZAK T. and ZURZYCKI J., 1983: Chloroplast translocations in Lemna trisulca induced by two successive blue light pulses. *Physiol. Veg.* 21(3), 519-526.
- KAK A.M., 1986: Family Lemnaceae in the Kashmir Himalayas. *J. Bombay Nat. Hist. Soc.* 83(1), 279-281.
- KAK A.M., BAKAVA U. and JAVED G.N., 1978: *Wolffia papulifera* Thomps. and *Wolffia columbiana* Karsten - two new plant records for India. *Indian For.* 104(4), 282-285.
- KALBERLAH A., 1985: Das Blüten der Wasserlinsen. *Z. Natw.* 68, 136-138.
- KAMEOKA H., YANAMOTO H. and MIYAZAWA M., 1985: Studies on removal of the pollutants from wastewater by aquatic plants from rivers (IV). Removal of metal elements. (In Japan.). *Kankyo Kagaku Kenkyusho Kenkyu Hokoku* 13, 25-30.
- KAMSHILOV M.M., 1977: Some results of studies on the cycle of substances and biological self-purification of bodies of water. (In Russian). *Gidrobiol. Zh.* 13, 5-13.
- KANAZAWA J., ISENSEE A.R. and KEARNEY P.C., 1975: Distribution of carbaryl and 3,5-xylol methylcarbamate in an aquatic model ecosystem. *J. Agric. Food Chem.* 23, 760-763.
- KANDELER R., 1955: Ueber die Blütenbildung bei Lemna gibba L. I. Kulturbedingungen und Tageslängenabhängigkeit. *Z. Bot.* 43, 61-71.
- KANDELER R., 1956: Ueber die Blütenbildung bei Lemna gibba L. II. Das Wirkungsspektrum von blühenförderndem Schwachlicht. *Z. Bot.* 44, 153-174.
- KANDELER R., 1961: Lemnaceen als Forschungsobjekt. *Ber. Phys. Med. Ges. Würzburg* 70, 81-86.
- KANDELER R., 1962: Die Aufhebung der photoperiodischen Steuerung bei Lemna gibba. *Ber. Deutsch. Bot. Ges.* 75, 431-442.
- KANDELER R., 1963: Phytochrom-Wirkung auf die vegetative Entwicklung von Lemna gibba. *Naturwiss.* 50, 551-552.
- KANDELER R., 1964a: Wirkungen des Kohlendioxids auf die Blütenbildung von Lemna gibba. *Naturwiss.* 51, 561-562.
- KANDELER R., 1964b: Zweifache Wirkung von Bikarbonat auf die Lichtsteuerung der Blütenbildung von Lemna gibba. *Ber. Deutsch. Bot. Ges.* 77, 140-142.
- KANDELER R., 1966: Trennung zweier Dunkelrotwirkungen bei der Lichtsteuerung der Sprossvermehrung von Lemna gibba. *Z. Pflanzenphysiol.* 54, 161-173.
- KANDELER R., 1967: The role of photophosphorylation in flower initiation of the long-day plant Lemna gibba. *European Photobiology Symp.* Hvar, YU. Book of Abstracts, 45.
- KANDELER R., 1968: Blühinduktion bei Lemnaceen. *Biol. Rundschau* 6, 49-57.
- KANDELER R., 1969a: Hemmung der Blütenbildung von Lemna gibba durch Anmonium. *Planta* 84, 279-291.

- KANDELER R., 1969b: Förderung der Blütenbildung von *Lemna gibba* durch DCMU und ADP. Z. Pflanzenphysiol. 61, 20-28.
- KANDELER R., 1970: Die Wirkung von Lithium und ADP auf die Phytochromsteuerung der Blütenbildung. Planta 90, 203-207.
- KANDELER R., 1971a: Die Wirkung von Ascorbinsäure, NADH und NADPH auf die Blütenbildung von *Lemna perpusilla* 6746 im Dauerlicht. Z. Pflanzenphysiol. 64, 278-280.
- KANDELER R., 1971b: Die Bedeutung des Photosynthese für die Blütenbildung. Umsch. Wiss. Tech. 71(3), 88.
- KANDELER R., 1972: Die Wirkung von Acetylcholin auf die photoperiodische Steuerung der Blütenbildung bei *Lemnaceen*. Z. Pflanzenphysiol. 67, 86-92.
- KANDELER R., 1973: *Lemnaceae*. Vervielfältigtes Manuskript für G. HEGI: Illustrierte Flora von Mitteleuropa. (3rd ed.). 2/1, 22 pp.
- KANDELER R., 1974 see KANDELER et al., 1974
- KANDELER R., 1975: Species delimitation in the genus *Lemna*. Aquat. Bot. 1, 365-376.
- KANDELER R., 1979: Familie: *Lemnaceae*. S.F. Gray 1821, Natur-Arrangm. Br. Fl. 2: 729 ("Lemnaceae"). Wasserlinsengewächse. In: HEGI G. (ed.), Illustrierte Flora von Mitteleuropa. Parey, Berlin/Hamburg. 2/1, (Lief. 5), 335-346.
- KANDELER R., 1983 see 1985
- KANDELER R., 1984: Flowering in the *Lemna* system. Phytion (Austria) 24 (1), 113-124.
- KANDELER R., 1985: *Lemnaceae*. In: HALEVY A.H. (ed.), Handbook of flowering. CRC Press, Boca Raton, FL, USA. 3, 251-279.
- KANDELER R. and FAERBER E., 1987: Short plant contact as stimulus for ethylene evolution in *Lemnaceae*. 14th Int. Bot. Congr. Berlin, Abstr., 126.
- KANDELER R. and HELDWEIN R., 1979: Significance of photosynthesis, N-deficiency, ABA and pH for synthesis of malate in *Lemna*. In: MARCELLE R., CLUJSTERS R. and VAN POUCKE M. (eds.), Photosynthesis and plant development. Junk, The Hague. 103-110.
- KANDELER R. and HUEGEL B., 1973: Blütenbildung bei *Lemna paucicostata* 6746 durch kombinierte Anwendung von Abscisinsäure und CCC. Plant Cell Physiol. 14, 515-520.
- KANDELER R. and HUEGEL B., 1974 see 1974a
- KANDELER R. and HUEGEL B., 1974a: Wiederentdeckung der echten *Lemna* perpusilla Torr. und Vergleich mit *L. paucicostata* Hegelm. Plant Syst. Evol. 123, 83-96.
- KANDELER R. and HUEGEL B., 1974b: Development in vitro of flower primordia of *Lemnaceae*. 3rd Int. Congr. Plant Tissues and Cell Culture, Leicester. Abstr., 160.
- KANDELER R., HUEGEL B. and ROTTENBURG T., 1974: Gegensätzliche Wirkung der Sprossalterung auf die Blütenbildung bei *Lemna paucicostata* und *Lemna gibba*. Biochem. Physiol. Pflanzen 165, 331-336.
- KANDELER R., HUEGEL B. and ROTTENBURG T., 1975: Relations between photosynthesis and flowering in *Lemnaceae*. In: MARCELLE R. (ed.), Environ. Biol. Contr. Photosynth. Junk, The Hague. 161-169.
- KANDELER R., LOEPFERT H., ROTTENBURG T. and SCHARFETER E., 1980: Early effects of phytochrome in *Lemna*. In: DE GREEF J. (ed.), Photoreceptors and Plant Development, Proc. Ann. Europ. Symp. Plant Photomorphogenesis, Antwerpen. Antwerpen Univ. Press. 485-492.
- KANG B.C. and CLELAND C.F., 1985: Investigation of senescence in *Lemna gibba* G3. A determinate growth system. Plant Physiol. 77(4 suppl.), 126.

- KANNAIYAN S., THANGARAJU M. and OBLISAMI G., 1984a: Azolla - a potential biofertilizer for rice production. Practical application of Azolla for rice production. Junk, Den Hague. 119-122.
- KANNAIYAN S., THANGARAJU M. and OBLISAMI G., 1984b: Effect of Azolla application on grain yield of rice. Agric. Res. J. Kerala 22(2), 183-185.
- KAR B.K., 1947: Methaxone as eradicator of waterhyacinth and other aquatics. Sci. Cult. 12, 545-550.
- KARELS L.A. and LEMBI C.A., 1973: Studies on the life history and control of watermeal (*Wolffia* sp.). Proc. North Central Weed Contr. Conf. 28, 107.
- KARLIN-NEUMANN G.A., KOHORN B.D., THORNBER J.P. and TOBIN E.M., 1985: A chlorophyll a/b-protein encoded by a gene containing an intron with characteristics of transposable element. J. Mol. Appl. Genet. 3(1), 45-61.
- KARLIN-NEUMANN G.A. and TOBIN E.M., 1985: Transit peptides of nuclear-encoded chloroplast proteins share a common amino-acid framework. EMBO 5(1), 9-14.
- KARPATI I., KARPATI V. and VOLF F., 1985: Role of water and wetland macrophytes in protection of water quality. Sborn. Vys. Skoly Zem. Praze, Fak. Agronomicka A 42, 25-45.
- KARPATI V. and POMOGYI P., 1979: Accumulation and release of nutrients by aquatic macrophytes. Symp. Biol. Hung. 19, 33-42.
- KARSTEN H., 1965: Ueber die Geschlechtsfähigkeit der Pflanzen. Bot. Unters. Phys. Lab. Berlin 1, 84-112.
- KARZEL R., 1926: Ueber die Nachwirkungen der Plasmolyse. Jb. Wiss. Bot. 65, 551-591.
- KASINOV V.B., 1966: Corymb inversion in irradiated colonies of the duckweed *Lemna* minor. (In Russian). Dokl. Akad. Nauk (USSR) 167, 201-204.
- KASINOV V.B., 1968a: Study of individual configurations in hereditarily persistent left-handed and right-handed clones of *Spirodela polyrrhiza*. (In Russian). Biofizika 13(6), 1092-1096.
- KASINOV V.B., 1968b: Interaction of mother and daughter fronds during the growth process of *Lemna* minor colonies. (In Russian). Bot. Zh. 53 (11), 1558-1568.
- KASINOV V.B., 1968c: On the inheritance of the left- and right-handedness in *Lemnaceae*. (In Russian). Genetika (USSR) 4, N11, 11-21.
- KASINOV V.B., 1969: On the inheritance of the left- and right-handedness in *Lemnaceae* and other organisms. (In Russian). Genetika (USSR) 5, N2, 22-29.
- KASINOV V.B., 1972: Concept of generation and the problem of variability within one generation in organisms reproducing by budding. (In Russian). Ontogenez. (USSR) 3, 360-370.
- KASINOV V.B., 1973: Handedness in *Lemnaceae*. On the determination of left and right types of development in *Lemna* clones and on its alteration by means of external influences. Beitr. Biol. Pflanz. 49, 321-337.
- KASINOV V.B., 1978: Genetical and morphogenetical effect of repeated 2,4-dichlorophenoxyacetic acid treatment in *Lemna* minor. (In Russian). Bot. Zh. (USSR) 63, 986-990.
- KASINOV V.B., 1980: The action of arginine, asparagine and atabrine stereomers upon the left and right *Lemna* minor plants. Biol. Plant 22, 321-326.
- KASINOV V.B., 1981: Influence of premature removal of daughter individuals upon the lifetime of maternal individual of *Lemna* minor. (In Russian). Bot. Zh. (USSR) 66(11), 1606-1613.

- KASINOV V.B. and KASINOVA G.V., 1971a: Retention of radiation-induced deformations in vegetative offspring of *Lemna gibba*. Dokl. Biol. Sci. 199(1-6), 476-478.
- KASINOV V.B. and KASINOVA G.V., 1971b: Dynamics of radiation damage to the plant organism based on the example of common duckweed (*Lemna minor*). (In Russian). Radiobiol. 11, 580-586.
- KASINOV V.B. and KASINOVA G.V., 1971c: Permanent polymorphic strains of duckweed as a model for studying some problems of developmental biology. (In Russian). Ontogenez. 2, 555-564.
- KASINOV V.B. and KASINOVA G.V., 1974: The reproduction rhythm in *Lemna*-ceae: A possible link with right and left handedness. Int. J. Chronobiol. 2, 47-52.
- KASINOV V.B. and PAVLOVA L.E., 1970: The interaction of sister fronds in a duckweed (*Lemna minor* L.) colony with a special reference to the right- and left-handedness. (In Russian). Bot. Zh. (USSR) 55, 1748-1763.
- KASINOV V.B. and PAVLOVA L.E., 1975: Hereditary reversion of chirality in duckweed under the effect of the herbicide 2,4-D: Condition of competent embryos. (In Russian). Tezisy Dokl. Vses. Soveshch. Embriol. 5, 78-79.
- KASINOV V.B. and PAVLOVA L.E., 1977: Dependence of morphogenetic reactions of *Lemna* frond primordia at the action of 2,4-dichlorophenoxyacetic acid on the age of primordium and its position in the plant. To the problem of ontogenetic control of heredity. (In Russian). Bot. Zh. (USSR) 62, 625-634.
- KATANSKAJA W.M., 1979: Flora of cooling water reservoirs of the USSR. (In Russian). Nauka, Leningrad. 278 pp.
- KATO A., 1979a: Effect of interruption of the nyctiperiod with a red to far-red-mixture of various ratios of red and far-red light on flowering in *Lemna gibba* G3. Plant Cell Physiol. 20, 1273-1283.
- KATO A., 1979b: Maintenance of high pfr-level in the dark period in relation to flowering in *Lemna gibba* G3. Plant Cell Physiol. 20, 1285-1293.
- KATO A., 1982: Kinetic studies of growth and flowering of *Lemna gibba* G3 under continuous light: effects of night interruptions with red and far-red light. Plant Sci. Lett. 27(2), 203-212.
- KATO A. and NAKASHIMA H., 1979: The effects on RNA synthesis in a long-day duckweed, *Lemna gibba* G3, of irradiation with different ratios of red and far red light during the prolonged dark period. Z. Pflanzenphysiol. 91, 109-117.
- KATO A. and NAKASHIMA H., 1980: The effect of temperature and dinitrophenol on the change of proportion of far-red absorbing phytochrome controlling RNA synthesis in *Lemna gibba* G3. Z. Pflanzenphysiol. 98, 177-182.
- KATO R. and FUJII T., 1985: Some characteristics of membrane-associated protein kinase in *Lemna paucicostata*. Plant Cell Physiol. 26(7), 1379-1386.
- KATO R., UNO I., ISHIKAWA T. and FUJII T., 1983: Effects of cyclic AMP on the activity of soluble protein kinases in *Lemna paucicostata*. Plant Cell Physiol. 24(5), 841-848.
- KATO R., UNO I., ISHIKAWA T. and FUJII T., 1984: Some characteristics of protein kinases in *Lemna paucicostata*. Plant Cell Physiol. 25(5), 691-696.
- KAUFMANN N., 1867: Ueber die aufsteigenden Axen einiger Lemnaceae. (In Russian). Papers of a Meeting of Russian Botanists, St. Petersburg.
- KAUFMANN N., 1868: Entwicklungsgeschichtliche Untersuchungen über die Lemnaceen. Bot. Z. 26, 382-384.

- KAUL R.B., 1976: Anatomical observations on floating leaves. Aquat. Bot. 2(3), 215-234.
- KAUL V. and BAKAYA U., 1973: The noxious floating, Lemna-Salvinia aquatic weed complex in Kashmir (India). In: VARSHNEY C.K. and RZOSKA J. (eds.), Aquatic weeds in South East Asia. Proc. Region. Seminar Noxious Aquatic Vegetation, Dew Delhi. Junk, The Hague. 183-192.
- KAUL V. and ZUTSHI D.P., 1963: *Lemna trisulca* L. A new record for Kashmir. Trop. Ecol. 4, 95-96.
- KAVANAGH F., 1941: New photoelectric fluorimeter and some applications. Industr. Engin. Chem. Ann. 13, 108-111.
- KAWABATA Z., TATSUKAWA R. and SATO K., 1986: Growth of duckweed and nutrient removal in a paddy field irrigated with sewage effluent. Int. J. Environm. Studies 27(3/4), 227-285.
- KAWAMATSU S., 1967: Electron microscope observation of plastids in the root cells of some hydrophytes. Cytologia 32(2), 157-164.
- KEARNY T.H. and PEEBLES R.H., 1973: Arizona Flora. (4th. ed.). Univ. California Press, Berkeley. 165-166.
- KEATES R.A.B. and TREWAVAS A.J., 1973: Protein kinase activity associated with ribosomes. Plant Physiol. 51(suppl.), 65.
- KEATES R.A.B. and TREWAVAS A.J., 1974: Protein kinase activity associated with isolated ribosomes from peas and Lemna. Plant Physiol. 54, 95-99.
- KEDDY P.A., 1976: Lakes as islands: The distributional ecology of two aquatic plants, *Lemna minor* L. and *L. trisulca* L. Ecology 57(2), 353-359.
- KELLEHER W.J. and GRISEBACH H., 1971: Hydride transfer in the biosynthesis of uridine diphospho- α -glucose from uridine diphospho-D-glucuronic acid with an enzyme preparation of *Lemna minor*. Eur. J. Biochem. 23, 136-142.
- KELLY D.G., 1980: Toxic algae in dairy lagoons. In: FRYE J.B. and CULLEY D.D. (eds.), U.S. Dept. Energy Final Report. Louisiana State Univ., Baton Rouge, LA. 4 pp. (Polycopy).
- KELLY D.G. and MYERS R.W., 1980: Decomposition of dairy lagoon sludge. In: FRYE J.B. and CULLEY D.D. (eds.), U.S. Dept. Energy Final Report. Louisiana State Univ., Baton Rouge, LA. 3 pp. (Polycopy).
- KELLY D.G., GOUGH R.H., GHOLSON J.H., HONEY S.E. and MYERS R.W., 1978: Effect of duckweed growth on nutrient utilization from dairy farm waste lagoons. J. Dairy Sci. 61(1), 206-207.
- KEMPERS-VEENSTRA A.E., VAN HEERIKUIZEN H., MUSTERS W., KLOOTWIJK J. and PLANTA R.J., 1984: Transcription of an artificial ribosomal RNA gene in yeast. EMBO J. 3(6), 1377-1382.
- KENNEY-WALLACE G. and BLACKMAN G.E., 1972: The uptake of growth substances. XIV. Patterns of uptake by *Lemna minor* of phenoxycetic and benzoic acids following progressive chlorination. J. Exp. Bot. 23, 114-127.
- KENNARD C.P., 1973: Control of weeds in direct seeded rice with some of the newer herbicides. Internat. Rice Comm. Newsletter 22(1), 15-21.
- KEPCZYNSKI K., 1960: New localities of *Wolffia arrhiza* (L.) Wimm. in the district Lipno. (In Polish). Zesz. Nauk. Univ. Torun. Biol. 5, 115-124.
- KEPCZYNSKI K., 1968: New localities of *Wolffia arrhiza* (L.) Wimm. in the district Wabro. (In Polish). Zesz. Nauk. Univ. Torun. Biol. 11, 269-272.
- KEPCZYNSKI K., 1972: Further localities of *Wolffia arrhiza* (L.) Wimm. on the Dobryzn diluvial plateau (Poland) in different plant communities. (In Polish). Zesz. Nauk. Univ. Torun. Biol. 15, 11-18.
- KEPCZYNSKI K. and FABISZAK S., 1972: *Salvinia natans* (L.) All. and *Spirodela-Salvinietum Slavnic* 1956 in the district Bydgoszcz. (In Polish). Zesz. Nauk. Univ. Torun. Biol. 15, 33-40.

KEPCZYNSKI K. and FABISZAK S., 1974: New localities for the rootless *Wolffia arrhiza*; new record in the Chelmo District. (in Polish). Acta Univ. Nicolai Copernici Biol. 16, 95-98.

KERBAEV E.B., SHCHERBAN Z.P., NAZAMOV N.D. and NAGIEV G., 1985: A non-mechanical method of reservoir treatment to control the larvae of blood-sucking mosquitoes. (In Russian). Uzb.Biol.Zh. 5, 65-66.

KERN H. and NAEF-ROTH S., 1975: Zur Bildung von Auxinen und Cytokinen durch *Taphrina*-Arten. Phytopath.z. 83, 193-222.

KESER M., 1955: Papierchromatographische Untersuchungen über das Auftreten freier und gebundener Aminosäuren in höheren Pflanzen. Planta 45, 273-288.

KESSLER B. and STEINBERG N., 1973: Cyclic mononucleotide-gibberellin interactions in the flowering and proliferation of the long-day plant *Lemna gibba* G3. Physiol.Plant. 28, 548-553.

KEUS R.J.A., DEKKER A.F., VAN ROON M.A. and GROOT G.S.P., 1983a: The nucleotide sequences of the regions flanking the genes coding for 23S, 16S and 4.5S ribosomal RNA on chloroplast DNA from *Spirodela oligorhiza*. Nucleic Acids Res. 11(18), 6465-6474.

KEUS R.J.A., ROOVERS D.J., DEKKER A.F. and GROOT G.S.P., 1983b: The nucleotide sequence of the 4.5S and 5S rRNA genes and flanking regions from *Spirodela oligorhiza* chloroplasts. Nucleic Acids Res. 11(10), 3405-3410.

KEUS R.J.A., ROOVERS D.J., VAN HEERIKHUIZEN H. and GROOT G.S.P., 1983c: Molecular cloning and characterization of the chloroplast ribosomal RNA genes from *Spirodela oligorhiza*. Curr.Genet. 7(1), 7-12.

KEUS R.J.A., DEKKER A.F., KREUK K.C.J. and GROOT G.S.P., 1984a: Transcription of ribosomal DNA in chloroplasts of *Spirodela oligorhiza*. Curr.Genet. 9(1), 91-97.

KEUS R.J.A., SYAM N.J., ZWIERS T., DE HEIJ H.T. and GROOT G.S.P., 1984b: The nucleotide sequences of the genes coding for rRNA and UCU, rRNA and AGC and rRNA on *Spirodela oligorhiza* chloroplast DNA. Nucleic Acids Res. 12(14), 5639-5646.

KHAKIMOVA V.K. and GALKINA N.V., 1973: Extraction of copper from solutions by the common duckweed. (In Russian). Akad.Nauk Usbek.Viniti 73 (5390), 5 pp.

KHAKIMOVA V.K. and GALKINA N.V. and TAUBAEV T.T., 1971: Lemna minor absorption of cobalt from ammonios. (In Russian). Khim Sel.Khoz 9(10), 67-68.

KHARE L.J., 1977: Effect of four herbicides on some aquatic weeds. Geobios 4(6), 271-272.

KHARE L.K., 1979: Effect of four commercial herbicides on some aquatic weeds. Geobios 6(2), 75-77.

KHUDAIRI A.K. and HEMBERG T., 1974: Serine involvement in the flowering of Lemna during photoperiodic induction. J.Exp.Bot. 25, 740-744.

KHUDAIRI A.K. and MAENG J., 1973: Studies on the flowering mechanism in Lemna. II. The dark reaction of the short-day plant *Lemna perpusilla*. Physiol.Plant. 28, 271-277.

KHURANA J.P., 1982: In vitro control of flowering in duckweeds. Ph.D. Thesis. Univ. of Delhi, Delhi.

KHURANA J.P. and MAHESHWARI S.C., 1978: Induction of flowering in Lemna paucicostata by salicylic acid. Plant.Sci.Lett. 12, 127-131.

KHURANA J.P. and MAHESHWARI S.C., 1980: Some effects of salicylic acid on growth and flowering in *Spirodela polyrrhiza* SP. Plant Cell Physiol. 21, 923-927.

KHURANA J.P. and MAHESHWARI S.C., 1983a: Floral induction in *Wolffia microscopica* by salicylic acid and related compounds under non-inductive long days. Plant Cell Physiol. 24(5), 907-912.

KHURANA J.P. and MAHESHWARI S.C., 1983b: Promotion of flowering in Lemna paucicostata 6746 (a short-day duckweed) by cytokinins. Plant Cell Physiol. 24(5), 913-918.

KHURANA J.P. and MAHESHWARI S.C., 1983c: Effect of 8-hydroxyquinoline on flowering and endogenous levels of iron and copper in Lemna paucicostata strain LP-6. Plant Cell Physiol. 24(7), 1251-1254.

KHURANA J.P. and MAHESHWARI S.C., 1984: Floral induction in short-day Lemna paucicostata 6746 by 8-hydroxyquinoline, under long days. Plant Cell Physiol. 25(1), 77-83.

KHURANA J.P. and MAHESHWARI S.C., 1986a: Induction of flowering in the duckweed Lemna paucicostata under non-inductive photoperiods by tannic acid. Plant Physiol. 66(3), 447-450.

KHURANA J.P. and MAHESHWARI S.C., 1986b: A comparison of the effects of chelates, salicylic acid and benzoic acid on growth and flowering of *Spirodela polyrrhiza*. Plant Cell Physiol. 27(5), 919-924.

KHURANA J.P. and MAHESHWARI S.C., 1986c: Floral induction in a photoperiodically neutral duckweed, Lemna paucicostata strain LP₆: role of chelating agents and iron. Plant Cell Physiol. 27(7), 1217-1224.

KHURANA J.P. and MAHESHWARI S.C., 1986d: Floral induction in a photoperiodically neutral duckweed, Lemna paucicostata LP₆: interaction of iron, EDTA and cytokinins. Biochem.Physiol.Pflanzen 161(8), 559-564.

KHURANA J.P., TANOT B.K. and MAHESHWARI S.C., 1986: Induction of flowering in a duckweed, *Wolffia microscopica*, under non-inductive long days, by 8-hydroxyquinoline. Plant Cell Physiol. 27(2), 373-376.

KICKUTH R. and TITZTNER T., 1974: Makrophysten limnischer Standorte und ihr Verhalten gegenüber p-Toluolsulfonsäure im Substrat. I. Ueber die Toleranz von Sumpf- und Wasserpflanzen gegenüber p-Toluolsulfonsäure. Angew.Bot. 48(3-4), 185-194.

KIENER W.B., 1944: Duckweed in relation to wildfowl in Nebraska. Proc. Nebraska Acad.Sci. 54th Ann.Meet., 11.

KIKUCHI E. and KURIHARA Y., 1981: Effects of tubificids on the biological and chemical characteristics of submerged rice field ecosystems. Verh.Int.Ver.Limnol. 21(2), 1194.

KIKUCHI E. and KURIHARA Y., 1982: The effects of the oligochaete *Branchiura sowerbyi* Beddard (Tubificidae) on the biological and chemical characteristics of overlying water and soil in a submerged ricefield soil system. Hydrobiologia 97(3), 203-208.

KIM I.B. and KHANG S., 1982: Growth of *Tilapia* in a closed water recirculating system without filter bed. (In Korean). Bull.Korean Fish Soc. 15(1), 47-51.

KINDEL P.K., 1973: Occurrence and metabolism of D-apiose in Lemna minor. Biogenesis Plant Cell Wall Polysaccharides, Proc.Symp. 1972, 85-94.

KINDEL P.K. and WATSON R.R., 1973: Synthesis, characterization and properties of uridine 5'-(α -D-apio-D-furanosyl pyrophosphate). Biochem. J. 133, 227-241.

KINDEL P.K., GUSTINE D.L. and WATSON R.R., 1970: Enzymatic synthesis of UDP-D-Apiose 1 α C. Plant Physiol. 46(suppl.), 27.

KING J.M. and COLEY K.S., 1985: Toxicity of aqueous extracts of natural and synthetic oils to three species of Lemna. In: BARNER R.C. and HANSEN D.J. (eds.), Aquatic toxicology and hazard assessment. ASTM 891, 302-309.

KINSER P.D. and NEUNZIG H.H., 1981: Description of the immature stages and biology of *Syncytia tinealis*. J.Lepid.Soc. 35(2), 137-146.

KINZEL H., 1982: Pflanzenökologie und Mineralstoffwechsel. Ulmer, Stuttgart. 534 pp.

KIRKLAND L.L., 1974: A study of photoperiodic inhibition of flower de-

- velopment in Lemna perpusilla 6746. Ph.D. Thesis. State Univ. Binghamton, N.Y. 170 pp. Diss. Abstr. B 35, 1525-1526.
- KIRKLAND L. and POSNER H.B., 1974: The role of light in the photoperiodic inhibition of flower development in Lemna perpusilla 6746. Plant Physiol. 53(suppl.), 3.
- KIRPENKO N.I., 1986: Phytopathic properties of toxin in blue-green algae. (In Russian). Gidrobiol. Zh. 22(1), 48-50.
- KLAINE S.J., 1985: Toxicity of coal gasifier solid waste to the aquatic plants Selenastrum capricornutum and Spirodela oligorhiza. Bull. Environ. Contam. Toxicol. 35(4), 551-555.
- KLASS D.L. and GOSH S., 1984: Methane production from and beneficiation of anaerobic digestion of aquatic plant material. Patent US 4424064. 5 pp.
- KLEBS G., 1993: In: ARTARIA: Chlorosarcina consociata (Klebs) Smith (= Chlorosphaera consociata) endophytic in thalli of Lemna collected at Pacific Grove. Bull. Soc. Imp. Nat. Moscou NS6, 222-262.
- KLEINSCHMIDT H.E., 1969: Effect of granular 2,4-D on some water-weeds and its persistence. Queensl. J. Agr. Anim. Sci. 26, 587-592.
- KLICH M.G., 1986: Un estudio sobre la lemnacea Spirodela intermedia W. Koch y la accion de algunos reguladores vegetales sobre su crecimiento y diferenciacion estomatica. M.S. Thesis. Univ. Nac. del Sur, Bahia Blanca. 150 pp.
- KLICH M.G. and MUJICA M.B., 1985: Cleistanthery in Spirodela intermedia W. Koch. Plant Cell Incompatibility Newsletter. 17, 20-23.
- KLICH M.G., MUJICA M.B. and FERNANDEZ O.A., 1985: The effect of gibberellic acid on the buoyancy of Spirodela intermedia W. Koch. Aquat. Bot. 21, 63-69.
- KLICH M.G., MUJICA M.B. and FERNANDEZ O.A., 1986: Stomatal morphology and ontogeny in Spirodela intermedia W. Koch. Aquat. Bot. 21(1-2), 155-164.
- KLICH M.G., FERNANDEZ O.A. and MUJICA M.B., 1987a: Influencia de algunas sustancias reguladores sobre el crecimiento de Spirodela intermedia W. Koch. I. Efectos del acido abscisico. Phytol. (Argentina) 47(1/2), 1-7.
- KLICH M.G., FERNANDEZ O.A. and MUJICA M.B., 1987b: Influencia de algunas sustancias reguladores sobre el crecimiento de Spirodela intermedia W. Koch. II. Efectos de dos auxinas y una antiauxina. Phytol. (Argentina) 47(1/2), 9-16.
- KLICH M.G., FERNANDEZ O.A. and MUJICA M.B., 1987c: Influencia de algunas sustancias reguladores sobre el crecimiento de Spirodela intermedia W. Koch. III. Efectos de dos citocininas. Phytol. (Argentina) 47(1/2), 17-24.
- KLICH M.G., FERNANDEZ O.A. and MUJICA M.B., 1987d: The effect of exogenous gibberellic acid on stomatal differentiation in Spirodela intermedia W. Koch. Aquat. Bot. 28(3-4), 393-396.
- KLINE L. and MCCUNE B., 1987: Factors influencing the distribution of Wolffia columbiana and Wolffia punctata (Lemnaceae). Northwest Sci. 61(1), 41-43.
- KLOSE H., 1963: Zur Limnologie von Lemna-Gewässern. Wiss. Z. Univ. Leipzig, Math.-Nat. Reihe 12, 233-259.
- KLOSE H., 1964 see 1963
- KNAPP R. and STOFFERS A.L., 1962: Ueber die Vegetation von Gewässern und Ufern im mittleren Hessen und Untersuchungen über den Einfluss von Pflanzen auf Sauerstoffgehalt, Wasserstoff-Ionen-Konzentration und die Lebensmöglichkeit anderer Gewächse. Ber. Oberhess. Ges. Nat.-Heilkunde, Giessen, 32, 90-141.

- KNIGHT R.L., WINCHESTER B.H. and HIGMAN J.C., 1985: Ecology, hydrology, and advanced wastewater treatment potential of an artificial wetland in north-central Florida. Wetlands 5, 167-180.
- KNOBLOCH E., 1981: New paleobotanical studies in the pannonian and pannonian of the Moravian Czechoslovakia part of the Vienna Basin. (In German). SB. Nar. Muz. Praha. Rada. B. Prir. Vedy 37(3-4) 7, 205-228.
- KNOBLOCH K., 1966: Photosynthetische Sulfid-Oxydation grüner Pflanzen. I. Mitteilung. Planta 70, 73-86.
- KNUDSON V., 1968: Comparative effects of aquatic herbicides on duckweed. Abstr. Meet. Weed Sci. Soc. Am., 66.
- KNUTH P., 1909: Handbook of flower pollination 3, order 121: Lemnaceae (translat. by J.R.A. Davis), Clarendon Press, Oxford.
- KNYPL J.S., 1976: Culture of Spirodela oligorhiza in ammonium-media buffered with calcium carbonate or calcium phosphate. Biochem. Physiol. Pflanzen 170, 243-252.
- KNYPL J.S., 1977a: The growth-retarding effect of alden in Spirodela oligorhiza. Experimentia 33(6), 725-726.
- KNYPL J.S., 1977b: Induction of ribonuclease activity by N,N-dimethylmorpholinium chloride and CCC in Spirodela. Biochem. Physiol. Pflanzen. 171(4), 289-298.
- KNYPL J.S., 1978: Reversal of the symptoms of phosphate deficiency in Spirodela by RNA and adenosine monophosphates. Z. Pflanzenphysiol. 90(3), 265-277.
- KNYPL J.S., 1979a: Molecular forms of phosphatase and ribonuclease in phosphate deficient and N,N-dimethylmorpholinium chloride treated Spirodela oligorhiza (Lemnaceae). Acta Soc. Bot. Pol. 48(1), 65-85.
- KNYPL J.S., 1979b: Time-course of phosphatase activity in phosphate deficient Spirodela. Bull. Acad. Pol. Sci., Biol. 27(2), 149-153.
- KNYPL J.S., 1980: Mutual exchange of phosphate amongst phosphate-deficient Spirodela oligorhiza plantlets. Z. Pflanzenphysiol. 96, 49-57.
- KNYPL J.S., 1982: Enzymatic adaptation of Spirodela oligorhiza to phosphate deficiency. 9th Int. Plant Nutrition Colloq., Warwick Univ., England. Abstr., 294.
- KNYPL J.S. and JANAS K.M., 1980: Stimulatory effect of fusococcin on growth and ribonuclease activity in Spirodela oligorhiza. Plant Sci. Lett. 19, 43-46.
- KNYPL J.S. and JANAS K.M., 1986: Physiological activity of l-amino-2-phenylethylphosphonic acid, a substrate analogue of phenylalanine. Biol. Plant. 28(2), 91-94.
- KNYPL J.S. and KABZINSKA E., 1977: Growth, phosphatase and ribonuclease activity in phosphate-deficient Spirodela oligorhiza cultures. Biochem. Physiol. Pflanzen 171(4), 279-287.
- KNYPL J.S. and OSWIECIMSKA M., 1986: Growth regulation activity of N-dodecylmethylmorpholinium and N-dodecylmethylmorpholine-N-methylmorpholinium chlorides in Lactuca sativa and Spirodela oligorhiza. J. Plant Physiol. 123(2), 127-133.
- KNYPL J.S. and SOBOLEWSKA J., 1978: Age dependent changes of phosphatase activity in phosphate supplied and phosphate deficient Spirodela. Physiol. Veg. 16(4), 773-784.
- KNYPL J.S., WITEK S. and OSWIECIMSKA M., 1976: Growth-retarding effect of N,N-dimethylmorpholinium chloride and CCC chlorocholine chloride in Spirodela oligorhiza. Z. Pflanzenphysiol. 79(1), 53-61.
- KNYPL J.S., WITEK S. and OSWIECIMSKA M., 1977: Growth-retarding effect and induction of ribonuclease activity by N,N-dimethylmorpholinium chloride in Spirodela oligorhiza. Proc. Int. Symp. Plant Growth Regulation, 411-416.

- KNYPL J.S., JANAS K.M. and WOLSKA M., 1986: Rhythmicity of L-phenylalanine ammonia-lyase activity in *Spirodela oligorrhiza*. Effects of darkening, abscisic acid and 1-amino-2-phenylethylphosphonic acid. *Physiol. Plant.* 66(3), 543-549.
- KOBAYASHI M., FURUTA T., ITAGAKI H. and KOMORI K., 1977: Studies on the hydraulic characteristics of the orifice type screen for weeding Lemna in irrigation canal. (In Japan.). *Res. Bull. Fac. Agric., Gifu Univ.* 40, 117-123.
- KOBUSZEWSKA D.M., 1973: Experimentally increased fish stock in the pond type Lake Wąrnak. 13. Distribution and biomass of the Lemnaceae and the fauna associated with them. *Ekol. Pol.* 21(39), 611-629.
- KOCH W., 1932: Beitrag zur Lemnaceen-Flora Mittel- und Südamerikas. *Ber. Schweiz. Bot. Ges.* 41, 113-118.
- KOCH W., 1933: *Spirodela biperforata*, eine neue Teichlinse aus Surinam. *Ber. Schweiz. Bot. Ges.* 42, 186-189.
- KOCH W., 1934 see 1932
- KOCH W., 1952: Zur Flora der oberitalienischen Reisfelder. *Ber. Schweiz. Bot. Ges.* 62, 628-663.
- KOCH W., 1954: Pflanzensozologische Skizzen aus den Reisfeldgebieten des Piemont (Po-Ebene). *Vegetatio* 5-6, 487-493.
- KOHLER A. and ZELTNER G.-H., 1974: Verbreitung und Ökologie von Makrophyten in Weichwasserflüssen des Oberrheinischen Waldes. *Hoppea* 33, 171-232.
- KOHOEN B.D. and TOBIN E.M., 1986a: Mutational analysis of light harvesting chlorophyll-protein. The major chlorophyll-protein component of the antenna of photosystem. *Photochem. Photobiol.* 43(suppl.), 113 pp.
- KOHOEN B.D. and TOBIN E.M., 1986b: Chloroplast import of light-harvesting chlorophyll a/b proteins with different amino termini and transit peptides. *Plant Physiol.* 82(4), 1172-1174.
- KOHOEN B.D., HAREL E., CHITNIS P.R., THORNER J.P. and TOBIN E.M., 1986: Functional and mutational analysis of the light-harvesting chlorophyll a/b protein of thylakoid membranes. *J. Cell Biol.* 102(3), 972-981.
- KOIE A. and KOIE M., 1939: Distribution of Geraniaceae, Araceae, Lemnaceae and Droseraceae in Denmark. (In Danish). *Bot. Tidsskr.* 45, 73-96.
- KOLES S.M., 1986: Modeling of duckweed (family Lemnaceae). Biomass production and water quality improvement. M.E. Thesis. Univ. Florida, Gainesville. 135 pp.
- KOLES S.M., PETRELLI R.J. and BAGNALL L.O., 1986: Duckweed culture for removal of NH₄ and for reduction of TSS from algal rich waters. *Proc. 26th Ann. Meet. Aquat. Plant Manage. Soc., Sarasota, Florida.* July 13-16. Abstr.
- KOLKOWITZ R., 1933: Zur Ökologie der Pflanzenwelt Brasiliens. *Ber. Deutsch. Bot. Ges.* 51, 396-406.
- KONDO T., 1978: Diurnal change in leakage of electrolytes from a long-day duckweed Lemna gibba G3, under osmotic stress induced by water treatment. *Plant Cell Physiol.* 19, 985-995.
- KONDO T., 1982a: Persistence of the potassium uptake rhythm in the presence of exogenous sucrose in Lemna gibba G3. *Plant Cell Physiol.* 23(3), 467-472.
- KONDO T., 1982b: Uptake and leakage of some ions in relation to potassium uptake rhythm in Lemna gibba G3. *Plant Cell Physiol.* 23(5), 901-908.
- KONDO T., 1982c: Correlation between potassium uptake rhythm and nitrate uptake rhythm in Lemna gibba G3. *Plant Cell Physiol.* 23(5), 909-915.
- KONDO T., 1982d: Phase control of the potassium uptake rhythm by the

- light signals in Lemna gibba G3. *Z. Pflanzenphysiol.* 107(5), 395-407.
- KONDO T., 1983a: Phase shift in the potassium uptake rhythm of the duckweed Lemna gibba G3 caused by an azide pulse. *Plant Physiol.* 73(3), 605-608.
- KONDO T., 1983b: Phase shifts of potassium uptake rhythm in Lemna gibba G3 due to light, dark or temperature pulses. *Plant Cell Physiol.* 24(4), 659-665.
- KONDO T., 1984a: Removal by a trace of sodium of the period lengthening of the potassium uptake rhythm due to lithium in Lemna gibba G3. *Plant Physiol.* 75(4), 1071-1074.
- KONDO T., 1984b: The period of circadian rhythm in Lemna gibba G3 is influenced by the substitution of rubidium for potassium. *Plant Cell Physiol.* 25(7), 1313-1317.
- KONDO T. and NAKASHIMA H., 1979: Content of adenosine phosphate compounds in a long-day duckweed, Lemna gibba G3, under different light and nutritional conditions. *Physiol. Plant.* 45, 357-362.
- KONDO T. and TSUDZUKI T., 1978: Rhythm in potassium uptake by a duckweed, Lemna gibba G3. *Plant Cell Physiol.* 19, 1465-1473.
- KONDO T. and TSUDZUKI T., 1980a: Phase progress under low temperature treatment of the potassium uptake rhythm in a duckweed, Lemna gibba G3. *Plant Cell Physiol.* 21, 95-103.
- KONDO T. and TSUDZUKI T., 1980b: Energy supply for potassium uptake rhythm in a duckweed, Lemna gibba G3. *Plant Cell Physiol.* 21, 433-443.
- KONDO T. and TSUDZUKI T., 1980c: Participation of a membrane system in the potassium uptake rhythm in a duckweed Lemna gibba G3. *Plant Cell Physiol.* 21, 627-635.
- KOPP A., FELLER U. and ERISMANN K.H., 1974a: Hat bei Lemna minor L. die Stickstoffquelle (Ammonium/Nitrat) einen Einfluss auf den Gehalt der Stärke und der freien Zucker? *Verh. Schweiz. Naturf. Ges.* 154, 196-198.
- KOPP A., FELLER U. and ERISMANN K.H., 1974b: Untersuchungen zur Regulation der Stickstoffassimilation von Lemna minor im Uebergang von Ammonium- auf Nitrat- bzw. Nitrat- auf Ammoniumernährung unter Photosynthesebedingungen. *Z. Pflanzenphysiol.* 73, 456-460.
- KORDAKOW J., 1970: New localities of Limnathemum nymphoides (L.) Link, Salvinia natans (L.) All., Lemna gibba L. and Wolffia arrhiza (L.) Wimm. in the valley of Wisla. (In Polish). *Bad. Fizjogr. Pol. Zach. Ser. B* 23, 243-250.
- KORDUS-WALANKIEWICZ B., 1978: The new stands of Wolffia arrhiza (L.) Wimm. on Wysoczyzna Siedlecka. *Fragm. Flor. Geobot.* 24, 273-275.
- KORSAN N.B. and MYAKUSHKO V.K., 1980: Vegetation of Tuzdinsk Reservoir (Kazakh-SSR, USSR) and its production. (In Russian). *Gidrobiol. Zh.* 16(1), 31-37.
- KORSAN N.B. and MYAKUSHKO V.K., 1981: Formation of water quality in shallows of the Yuzhny Reservoir (Dnjepr Krivoi Rog Canal, Ukrainian-SSR, USSR). under the effect of higher aquatic plant overgrowth. (In Russian). *Gidrobiol. Zh.* 17(1), 48-54.
- KOSTYTSCHEV S. and BERG V., 1929: Die Form der Calciumverbindung in lebenden Pflanzengeweiben. *Planta* 8, 55-67.
- KOUKAKI W.L. and DUKE S.H., 1973: Regulating growth of an aquatic plant: Lemna perpusilla. *J. Minn. Acad. Sci.* 39, 12-14.
- KOVACS M., NYARY I. and TOTTH L., 1984: The microelement content of some submerged and floating aquatic plants. *Acta Bot. Hung.* 30(1-2), 173-185.
- KRAJNCIC B., 1972: Photoperiodic relations in Lemnaceae of north-eastern Slovenia. (In Slov.). *M.S. Thesis. Univ. Zagreb.* 106 pp.

- KRAJNCIC B., 1974a: Study of photoperiodic reactions in Lemnaceae of Slovenia. (In Slov.). 4th Biol. Congr. Yugoslavia, Sarajevo, Abstr., 84.
- KRAJNCIC B., 1974b: Photoperiodic responses of Lemnaceae from Northeastern Slovenia. Acta Bot. Croat. 33, 81-88.
- KRAJNCIC B., 1974c: Contribution to the knowledge of Lemnaceae from northeastern Slovenia. (In Slov.). Biol. Vestn. 22, 21-28.
- KRAJNCIC B., 1975: The mechanisms of floral induction in Lemnaceae from Slovenia. Acta Bot. Croat. 34, 189.
- KRAJNCIC B., 1976: The mechanism of floral induction in Lemnaceae from Slovenia. (In Serbo-Croat.). Ph.D. Thesis. Univ. Zagreb. 192 pp.
- KRAJNCIC B., 1976b: Lemnaceae in the region of Slovenia. (In Slov.). Biol. Vestn. 24, 133-144.
- KRAJNCIC B., 1982: Effects of kinetin on floral induction and floral development in the species *Lemna minor* and *Spirodela polyrrhiza*. Biol. Vestn. 30(1), 85-104.
- KRAJNCIC B., 1983: The effects of cytokinins on flowering in the long-short-day plant *Wolffia arrhiza* (L.) Wimm. Z. Pflanzenphysiol. 112(3), 281-286.
- KRAJNCIC B., 1985: Regulation of the floral induction with ABA and EDDHA. (In Slov.). Biol. Vestn. 33(2), 39-52.
- KRAJNCIC B. and BAVEC F., 1986: Research of abscisic acid effects on flower morphogenesis in the species *Lemna minor*. (In Slov.). Biol. Vestn. 34(1), 61-68.
- KRAJNCIC B. and DEVIDE Z., 1977: The morphogenesis of flowers in *Lemna* from the territory of Slovenia, Yugoslavia. Biol. Vestn. 25(2), 180.
- KRAJNCIC B. and DEVIDE Z., 1979: Flower development in *Spirodela polyrrhiza* (Lemnaceae). Plant. Syst. Evol. 132, 305-312.
- KRAJNCIC B. and DEVIDE Z., 1980: Report on photoperiodic responses in Lemnaceae from Slovenia. Ber. Geobot. Inst. ETH, Stiftung Rübel, Zürich 47, 75-86.
- KRAJNCIC B. and DEVIDE Z., 1982: see 1982c
- KRAJNCIC B. and DEVIDE Z., 1982a: Photoperiodic responses in Lemnaceae from North Croatia. Acta Bot. Croat. 41, 57-63.
- KRAJNCIC B. and DEVIDE Z., 1982b: The action of amino acids on floral induction in species *Lemna minor* and *Spirodela polyrrhiza*. (In Serbo-Croat.). Bio. Vestn. 30(1), 105-120.
- KRAJNCIC B. and DEVIDE Z., 1982c: Distribution of Lemnaceae in North Croatia. (In Serbo-Croat.). Acta Bot. Croat. 41, 175-180.
- KRAUSE A., 1979: Zur Kenntnis des Wasserpflanzenbestandes der westdeutschen Mittelgebirgsflüsse Fulda, Ahr, Sieg und Saar. Dechenia 132, 15-28.
- KRAUSE J., 1978: Die Zimtsäurederivate von *Spirodela polyrrhiza* (L.) Schleiden. Z. Pflanzenphysiol. 88, 465-470.
- KRAUSE J. and STRACK D., 1979a: Malonyl cyanidin 3-monoglucoside in *Spirodela polyrrhiza* (L.) Schleiden. Z. Pflanzenphysiol. 95, 183-187.
- KRAUSE J. and STRACK D., 1979b: High performance liquid chromatographic separation of glucose esters and quinic-acid esters of hydroxy cinamic acids. J. Chromatogr. 176(3), 465-469.
- KRAUSS F., 1945: Pflanzen des Cap- und Natallandes. Flora 28, 344.
- KREWER B.P., 1983: Die kleinste Blütenpflanze der Welt. Mikrok. 72(1), 5-8.
- KRIVOKAPIC K. and ERIC Z., 1984: The influence of dinitro-ortho-cresol (DNOC) on some cytological and physiological changes in plants. (In Macedon.). Godisnjak 37, 51-59.

- KRONBERGER W., 1983: pH-dependency of fluorine uptake by different plants. Aquilo Ser. Bot. 19(2), 304.
- KRNIK-RASOL M. and RENDIC L., 1975: Effects of some herbicides on the growth of *Lemna gibba* and *Spirodela polyrrhiza*. (In Croat.). Acta Bot. Croat. 34, 190.
- KRNIK-RASOL M. and RENDIC L., 1977: The effect of some triazine derivatives on the growth and development of duckweeds. (In Croat.). Acta Bot. Croat. 36, 75-82.
- KRUGMANN-RANDOLF I., 1978: Forschung für die Landwirtschaft der Entwicklungsländer. Wasserpflanzen als Futter nutzen, statt chemisch vernichten. Entwicklung und Zusammenarbeit 11, 13-14.
- KRULL J.N., 1969: Factors affecting plant die-offs in shallow water areas. Am. Midland Natur. 69(82), 293-295.
- KRZECZOWSKA M. and ZIMNA J., 1972: Effect of kinetin and β -indolylacetic acid on the content of nitrogen compounds in *Lemna minor* cultures. (In Polish). Zesz. Nauk. Univ. Lodz., Ser. 2, 47, 95-100.
- KRZECZOWSKA M., ZIMNA J., BYTNIWSKA K. and MACIEJEWSKA-POTAPCZYK W., 1975: Attempts at establishing the culture conditions for *Lemna minor* L. (In Polish). Acta Soc. Bot. Pol. 44(2), 243-254.
- KRZEMINSKI S.F., BRACKETT C.K. and FISHER J.D., 1975: Fate of microbicides dal 3-isothiazolone compounds in the environment: modes and rates of dissipation. J. Agric. Food Chem. 23(6), 1060-1068.
- KUCHAR K.W., 1954: Bakteriologische und limnologische Untersuchungen an einem Lemnagesässer. Arch. Hydrobiol. 49, 329-334.
- KUECHLER L., 1986: Analyse der Wasserpflanzenentwicklung in Grabensystemen zur Grundwasserregulierung. Acta Hydrochim. Hydrobiol. 14, 47-57.
- KUESEL H., 1955: Die Entenlinse (*Wolffia arrhiza* Wimm.), eine neue Pflanze der nordwestdeutschen Flora. Küsel, Lahausen, Mitt. 1, 10-11.
- KUHN D.J., 1969: The duckweed. Am. Biol. Teacher 31(5), 328-329.
- KULAKOVA I.A. and ANDREEV V.Y., 1981: Effect of phytohormones on ion transport into plant cells. (In Russian). Rost. Rast. Puti. Ego. Regul. 35-43.
- KULIKOV N.V., CHEBOTINA M.Y. and LYUBIMOVA S.A., 1980: Mobility of strontium-90 and cesium-137 in water-freshwater plant and water-soil systems. (In Russian). Radiobiologiya 20(1), 146-148.
- KULSHRESHTHA M. and GOPAL B., 1983: Allelopathic influence of *Hydrilla verticillata* (L.F.) Royle on the distribution of *Ceratophyllum species*. Aquat. Bot. 16, 207-209.
- KUMAR T.S., VASUDEVAN P. and PATWARDHAN S.V., 1983: Growth studies of *Azolla pinnata* in a permanent pond. Int. J. Environ. Stud. 21(3-4), 297-300.
- KUNII H., 1984: Seasonal changes in water quality and surface cover of aquatic plants in pond Ojaga-like, Chiba, 1978-1980. Mem. Fac. Sci., Shimane Univ., Matsue, Japan 18, 59-68.
- KUNII H., TSUCHIYA T., MATSUI K. and IKUSIMA I., 1985: Present state of aquatic plants in Lake Biwa, Japan, and its surrounding water bodies. Jap. J. Limnol. 46(3), 215-218.
- KUNTII K.C.S., 1841: Enumeratio Plantarum omnium hucusque cognitarum, 3, 2-7. Stuttgart and Tübingen.
- KUPIYANOVA L.A. and TARASEVICH V.F., 1984: The ultrastructure of the surface of the pollen grain wall in some genera of the family Lemnaceae and the related genera of the family Araceae. (In Russian). Bot. Zh. 69(12), 1656-1662.
- KURIMO U., 1970: Effect of pollution on the aquatic macroflora of the Varkans area, Finnish Lake District. Ann. Bot. Fenn. 7, 213-254.
- KURZ H. and CROWSON D., 1949: The flowers of *Wolffia floridana* (J.D. Sm.) Thompson. Quart. J. Florida Acad. Sci. 11, 87-98.

- KURZ S., 1966: Enumeration of Indian Lemnaceae. J. Linn. Soc. Bot. 9, 264-268.
- KURZ S., 1967: Enumeration of Australian Lemnaceae. J. Bot. 5, 115.
- KURZ S., 1971: On some new or imperfectly known Indian plants. 109. Lemnaceae. J. As. Soc. Bengal 40(2), 78.
- KUTLACHMEDEV Y. A., SOKURKA G. S. and GROBZINSKY D. M., 1978: Investigations of the influence of comonautic factors on the anabiotic stage of turions of Spirodela polyrrhiza. (In Russian). Cosmic Invest. in Ukraine, Kiev. 49-54.
- KVET J., 1978: Growth analysis of fish pond littoral communities. In: DYKJOVA D. and KVET J. (eds.), Ecological studies, analysis and synthesis. Springer, New York/Berlin. 28, 198-206.
- KVET J., REJMANKOVA E. and REJMANEK M., 1979: Higher aquatic plants and biological wastewater treatment. The outline of possibilities. Proc. Aktiv Jihocesky Vodooh. Conf. 9 pp.
- LACHAVANNE J.-B., CROZET B., JUZE R., NOETZLIN A. and PERFEYTA J., 1984: Etude des macrophytes du Vierwaldstaettersee. Unité de Biol. Aquat., Genève. 230 pp.
- LACOR M. A. M., 1968: Flowering of Spirodela polyrrhiza (L.) Schleiden. Acta Bot. Neerl. 17, 357-359.
- LACOR M. A. M., 1969: On the influence of gibberellic acid and kinetin on the germination of turions of Spirodela polyrrhiza (L.) Schleiden. Acta Bot. Neerl. 18, 550-557.
- LACOR M. A. M., 1970: Some physiological and morphogenetic aspects of flowering of Spirodela polyrrhiza (L.) Schleiden. Acta Bot. Neerl. 19, 53-60.
- LAGERBERG T., 1947: Wild plants in the North. 1. Lemnaceae. (In Swed.). Natur och Kultur, Stockholm. 302-309.
- LAGMAN L. H., NELSON F. P. and RICHARDSON G. E., 1980: Water quality and aquatic vegetation in the Back River Reservoir, Berkeley Co., South Carolina. Columbia, SC, Water Res. Comm. Rept. No. 130, 62 pp.
- LAI C. E. and HAYES T. L., 1980: Calibration of SEM frozen-hydrated analyses using standard salt solutions and peak to background measurements. Proc. Ann. Meet. Electr. Microsc. Soc. America 38, 800-801.
- LAIRD M., 1956: Studies of mosquitoes and freshwater ecology in the South Pacific. Bull. R. Soc. New Zealand 6, 213 pp.
- LAKATOS E., 1978: Wolffia arrhiza in northern Hungary. (In Hung.). Botanikai Közlemények 65(3), 177-179.
- LAM E. and MALKIN R., 1985: Characterization of a photosynthetic mutant of Lemna lacking the cytochrome b_6 -f complex. Acta Biochim. Biophys. 810(1), 106-109.
- LANCIANI C. A., 1987: Rearing immature Mesovelia mulsanti (Hemiptera mesoveliidae) on a substratum of duckweed. Fla. Entomol. 70(2), 286-288.
- LANCASTER T. L., 1930: On the occurrence of Lemna oligorrhiza Kurz in New Zealand. Trans. Proc. New Zealand Inst. 60, 563-564.
- LANDNER L. and JERNELOV A., 1969: Cadmium in aquatic systems. Metals and ecology, Symposium. Ecol. Res. Comm. Bull. 5, 47-54.
- LANDOLT E., 1955: Ueber das Wachstum in der Dunkelheit bei einigen Lemnaceen. Verh. Schweiz. Natf. Ges. 135, 135-136.
- LANDOLT E., 1957: Physiologische und ökologische Untersuchungen an Lemnaceen. Ber. Schweiz. Bot. Ges. 67, 271-410.
- LANDOLT E., 1975: Morphological differentiation and geographical distribution of the Lemna gibba - Lemna minor group. Aquat. Bot. 1, 345-363.
- LANDOLT E., 1977 see 1957

- LANDOLT E., 1979: Lemna minuscula Herter (= L. minima Phil.), eine in Europa neu eingebürgerte amerikanische Wasserpflanze. Ber. Geobot. Inst. ETH, Stiftung Rübel, Zürich 46, 86-89.
- LANDOLT E. (ed.), 1980a: Biosystematische Untersuchungen in der Familie der Wasserlinsen (Lemnaceae). Veröff. Geobot. Inst. ETH, Stiftung Rübel, Zürich 70, 247 pp.
- LANDOLT E., 1980b: Key to the determination of taxa within the family of Lemnaceae. Veröff. Geobot. Inst. ETH, Stiftung Rübel, Zürich 70, 13-21.
- LANDOLT E., 1980c: Description of six new species of Lemnaceae. Veröff. Geobot. Inst. ETH, Stiftung Rübel, Zürich 70, 22-29.
- LANDOLT E., 1980d: Bibliographie der Familie der Lemnaceae. Veröff. Geobot. Inst. ETH, Stiftung Rübel, Zürich 70, 142-204.
- LANDOLT E., 1981: Distribution pattern of the family Lemnaceae in North Carolina. Veröff. Geobot. Inst. ETH, Stiftung Rübel, Zürich 77, 112-148.
- LANDOLT E., 1982a: Distribution pattern and ecophysiological characteristics of the European species of the Lemnaceae. Ber. Geobot. Inst. ETH, Stiftung Rübel, Zürich 49, 127-145.
- LANDOLT E., 1982b: Distribution patterns within the family Lemnaceae. In: SYMOENS J. J., HOOPER S. S. and COMPERE P. (eds.), Studies on aquatic plants. Abstr., 313.
- LANDOLT E., 1984 see 1984a
- LANDOLT E., 1984a: Flowers and fruits in the genus Wolffia (Lemnaceae). Ber. Geobot. Inst. ETH, Stiftung Rübel, Zürich 51, 164-172.
- LANDOLT E., 1984b: Verbreitungsmuster in der Familie der Lemnaceae und ihre ökologische Deutung. Verh. Ges. Ökol. (Bern 1982) 12, 241-253.
- LANDOLT E., 1986: The family of Lemnaceae - a monographic study (vol. 1). Veröff. Geobot. Inst. ETH, Stiftung Rübel, Zürich 71, 566 pp.
- LANDOLT E., 1987: Eco-geographical differentiation in some aquatic plants: the Lemnaceae. In: URBANSKA K. (ed.), Differentiation patterns in higher plants. Acad. Press, London. 201-215.
- LANDOLT E. and DANN W., 1983: Vergleich von 10 Klonen von Lemna gibba bei verschiedenen Stickstoffkonzentrationen. Ber. Geobot. Inst. ETH, Stiftung Rübel, Zürich 50, 86-97.
- LANDOLT E. and KANDELER R., 1987: The family of Lemnaceae - a monographic study (vol. 2). Veröff. Geobot. Inst. ETH, Stiftung Rübel, Zürich 95, 638 pp.
- LANDOLT E. and URBANSKA-WORYTKIEWICZ K., 1980: List of the studied Lemnaceae samples: origin and chromosome numbers. Veröff. Geobot. Inst. ETH, Stiftung Rübel, Zürich 70, 205-247.
- LANDOLT E. and WILDI O., 1977: Ökologische Felduntersuchungen bei Wasserlinsen (Lemnaceae) in den südwestlichen Staaten der USA. Ber. Geobot. Inst. ETH, Stiftung Rübel, Zürich 44, 104-146.
- LANGE see DE LANGE
- LAPHAM V. T., 1967: A new technique for duckweed control. Proc. S. Weed Conf. 20, 339-341.
- LASS B. and ULLRICH-EBERIUS C. I., 1984a: Evidence for proton/sulfate cotransport and its kinetics in Lemna gibba Gl. Planta 161(1), 53-60.
- LASS B. and ULLRICH-EBERIUS C. I., 1984b: Mechanism of sulfate uptake in Lemna gibba Gl.: 3 H⁺/sulfate cotransport and its biphasic kinetics. In: CRAM W. J. et al. (eds.), Membrane transport in plants. Symp. Abstr., 412.
- LASS B., THIEL G. and ULLRICH-EBERIUS C. I., 1986: Electron transport across the plasmalemma of Lemna gibba Gl. Planta 169(2), 251-259.
- LASSALLE J. P., AYADI A., MONNIER A., STELZ T. and THELLIER M., 1973: Application des relations et de la loi d'Onsager à la formulation d'interactions compétitives lors de l'absorption d'ions alcalins par la Lemna minor. C. R. Acad. Sci. Paris, D 276, 1053-1056.

- LASSOCSINSKI W., 1982: Substrate specificity of ribonucleases of phosphate-supplied and phosphate-deficient *Spirodela*. *Biochem. Physiol. Pflanzen* 177(2), 125-136.
- LAUBE H.R. and WOHLER J.R., 1973: Studies on the decomposition of a duckweed (*Lemnaceae*) community. *Bull. Torr. Bot. Club* 100, 238-240.
- LAUTNER V. and MUELLER Z., 1954: Feeding value of some water plants I. (In Czech.). *Spornik Cesk. Akad. Zemedl. Ved* 27A, 333-354.
- LAWALREE A., 1943: La multiplication végétative des *Lemnaceae* en particulier chez *Wolffia arrhiza*. *Cellule* 49, 337-382.
- LAWALREE A., 1945: La position systématique des *Lemnaceae* et leur classification. *Bull. Soc. Roy. Bot. Belg.* 77, 27-38.
- LAWALREE A., 1952 see 1952a
- LAWALREE A., 1952a: L'embryologie des *Lemnaceae*. Observations sur *Lemna minor* L. *Cellule* 54, 305-326.
- LAWALREE A., 1952b: Compositae, *Leaceae*, *Lemnaceae* et *Vitaceae*. Exploration hydrobiologique du Lac Tanganyika. *Inst. Roy. Sci. Nat. Belg., Résultat Sci.* 4, 53-82.
- LAWALREE A., 1961: La pollinisation de *Lemna minor* L. *Natural. Belg.* 42, 164-165.
- LAWALREE A., 1980: *Lemnaceae*. In: TUTIN T.G., HEYWOOD V.H., BURGESS N.A., MOORE D.M., VALENTINE D.H., WALTERS S.M. and WEBB D.A. (eds.), *Flora Europaea*. Cambridge Univ. Press. 5, 273.
- LAWALREE A., 1984: *Lemna minuscula* Herter, nouveau pour la flore belge. *Dumortiera* 28, 5-7.
- LAWSON T.B., 1973: Methods of drying duckweed. M.S. Thesis. Louisiana State Univ., Shreveport, LA.
- LAWSON T.B., BRAUD H.J. and WRATTEN F.T., 1974: Methods of drying duckweed, *Lemnaceae*. *Am. Soc. Agr. Eng.*, Winter Meeting, Chicago. 12 pp.
- LEATHER G.R. and EINHELLIG F.A., 1985: Mechanisms of allelopathic action in bioassay. The chemistry of allelopathy. *Biochemical interactions among plants*. *Amer. Chem. Soc.* 197-205.
- LEATHER G.R. and EINHELLIG F.A., 1986: Bioassays in the study of allelopathy. In: PUTMAN A.R. and TANG C.S. (eds.), *The science of allelopathy*. Wiley, New York. 133-145.
- LEBEDEVA G.D. and KUZNETSOVA G.A., 1968: Accumulation and excretion of manganese-54 by freshwater fish and other hydrobiants. (In Russian). *Biol. Nauk* 11(10), 42-44.
- LECHEVALLIER D., 1966: Les lipides des *Lemnaceae*: Analyse des acides gras des lipides des frondes de *Spirodela polyrrhiza*. *C.R. Acad. Sci. Paris*, D 263, 1849-1852.
- LECHEVALLIER D., 1967: Analyse des lipides polaires des frondes de *Spirodela polyrrhiza*. *C.R. Acad. Sci. Paris*, D 264, 2110-2113.
- LECHEVALLIER D., 1969: Sur les paraffines des frondes de *Spirodela polyrrhiza* Schleid. et de *Lemna trisulca* L. *C.R. Acad. Sci. Paris*, D 268 (2), 314-317.
- LECHEVALLIER D., 1970: Sur les stérols des frondes de *Spirodela polyrrhiza* Schleid. et de *Lemna trisulca* L. *C.R. Acad. Sci. Paris*, D 271, 591-594.
- LECHEVALLIER D., 1977a: Lipides, nucléotides pyridiniques et nucléotides adényliques de tissus et de plastides isolés de *Spirodèles* cultivées sur milieu enrichi en calcium. *Physiol. Vég.* 15(1), 95-119.
- LECHEVALLIER D., 1977b: Effets du polyéthylène glycol sur les lipides et les lipochromes des colonies de *Spirodèle*. *Physiol. Vég.* 15(3), 387-402.
- LECHEVALLIER D., BAHL J. and MONEGER R., 1971: Lipid components of chloroplasts isolated from *Spirodela polyrrhiza* light-cultivated on su-

- crose-containing media. In: FORTI F., AVRON M. and MELANDRI A. (eds.), *Photosynthesis. Proc. 2nd Int. Congr. Photosynth. Res.*, 1647-1652.
- LECHEVALLIER D., BAHL J. and MONEGER R., 1976: Qualité spectrale de la lumière et lipides plastidiaux du blé et de la *Spirodèle*. In: JACQUES R. (ed.), *Etudes de biologie végétales*, hommage au Professeur Pierre Chouard. Impr. Louis-Jean, Paris. 521-535.
- LECHEVALLIER D., VERMEERSCH J. and MONEGER R., 1977: Micro-analyse du NADP et du NAD réduits et oxydés dans les tissus foliaires et dans les plastides isolés de *Spirodèle* et de blé. 2. Méthode d'analyse des nucléotides pyridiniques de tissus végétaux. Application à l'étude des effets du saccharose, de la lumière rouge et de l'obscurité. *Physiol. Vég.* 15(1), 63-93.
- LEDBETTER M. and PORTER K., 1970: Introduction to the fine structure of plant cells. Springer, Berlin/Heidelberg/New York. 148-150.
- LEDL G., JANAUER G.A. and HORAK O., 1981: Die Anreicherung von Schwermetallen in Wasserpflanzen aus einigen Oesterreichischen Fließgewässern. *Acta Hydrochim. Hydrobiol.* 9(6), 651-663.
- LEGGETT W.H., 1870a: *Lemna*. *Bull. Torr. Bot. Club* 1, 29.
- LEGGETT W.H., 1870b: *Spirodela*. *Bull. Torr. Bot. Club* 1, 37-38.
- LEHMAN P.W., SILK W.K. and KNIGHT A.W., 1981: Protein and nitrate content of *Lemna* sp. as a function of developmental stage and incubation temperature. *Plant Physiol.* 68(1), 127-132.
- LEINBACH E.D., 1975: Characterization of a particulate UDP-galacturonate: acceptor D-galacturonosyltransferase from *Lemna minor* and studies on the physical properties of the product. Ph.D. thesis, Michigan State Univ. 109 pp. *Diss. Abstr. Int.* B 36, 6134.
- LEINERTE M.P., 1968: Experiments involving the exposure of *Lemna minor* to gamma-rays and fast neutrons. (In Russian). *Radioobiologiya* 8(1), 156-157.
- LEINERTE M.P., 1969: Influence of external γ -irradiation of higher aquatic plants on their accumulation of ^{86}Sr , ^{137}Cs , and ^{144}Ce . *Radioobiologiya* 9(3), 427-432. *Translation UDC* 577.391, 131-138.
- LEMBI C.A., 1976: Aquatic weed control results in Indiana in 1975. *Proc. North Central Weed Contr. Conf.* 30, 169.
- LEMBI C.A., 1978: Results of 1978 aquatic herbicide trials in Indiana. *Proc. North Central Weed Contr. Conf.* 33, 102-103.
- LEONARD S.W., 1972: New records and notes on the flora of the Carolinas. *J. Elisha Mitchell Sci. Soc.* 88, 265-266.
- LE PABIC C., 1972: Ultrastructure et différenciation des chloroplastes de *Spirodela polyrrhiza*. *C.R. Acad. Sci. Paris*, D 275(21), 2339-2342.
- LE PABIC C., 1976a: Modifications de la croissance et de la pigmentation de *Spirodela polyrrhiza* (Schleid.) induites par une cytokinine, la 6-benzylaminopurine. *Physiol. Vég.* 14(1), 77-88.
- LE PABIC C., 1976b: Modifications de la pigmentation et de l'ultrastructure des chloroplastes de *Spirodela polyrrhiza*. *J. Microsc. Biol. Cell* 25, 181-186.
- LE PABIC C., 1980: Composition en lipides polaires et analyse des acides gras des frondes de *Spirodèle* traitées par la 6-benzylaminopurine. *Plant Sci. Lett.* 17, 303-310.
- LE PABIC C., 1982b see 1972
- LEPAGE E., 1976: Additions and extension of distribution in the genera *Eriophorum*, *Spirodela*, *Atriplex* and *Hieracium* in Quebec, Canada. *Natur. Can.* 103(3), 235-238.
- LEPAGE E., 1978 see 1976
- LEPPARD G.G. and LEAN D.R.S., 1976: Ion exchange fibrils in inland wa-

- ters, their production and environmental significance. Proc. Can. Fed. Biol. Soc. 19, 86.
- LEWIS D., PICKENS W., WILSON G.L., MILLER B., BERELWITZ M., VINIK A.I., COLLIER E. and CLELAND C.F., 1985: Somatostatin-like material is present in flowering plants. Endocrinology 117(5), 2093-2097.
- LESCH S. and ULLRICH W., 1984: Ammonium-Aufnahme + oder NH₃? Mitt. Bd. Carriertransport und/oder Diffusion von NH₄ Bot.-Tag. Wien, 52 (0508).
- LESLIE A.C. and WALTERS S.M., 1983: The occurrence of *Lemma minuscula* Herter in the British Isles. Watsonia 14(3), 243-248.
- LETHAM D.S., 1967: Regulators of cell division in plant tissues. V. A comparison of the activities of zeatin and other cytokinins in five bioassays. Planta 74, 228-242.
- LEUCHTMANN A., 1979: Physiologische Differenzierungen in der Familie der Lemnaceen. Diploma Thesis. Geobot. Inst. ETH, Stiftung Rüböl, Zürich. 86 pp. (Polycopy).
- LEUTE G.H. and MUELLER I.E., 1979: Potamogeton acutifolius Link und Lemna trisulca L., zwei verschollen geglaubte Vertreter der Hydrophytenflora in Kärnten wiederentdeckt. Carinthia 2(169/89), 137-142.
- LEUTWILER L.S. and TOBIN E.M., 1985: Characterization of the multigene family encoding the light-harvesting chlorophyll a/b-protein in *Arabidopsis thaliana*. J. Cell. Biochem. (suppl. 9 part C), 236.
- LEWKOWICZ see LEWKOWICZ
- LEWIN E., 1977: Effects of phenytoin on the release of ¹⁴C adenine derivatives. Epilepsia 18(3), 349-356.
- LEWIN R.A., 1984: Culture and taxonomic status of *Chlorochytrium lemnae*, green algae endophyte. Br. Phycol. J. 19(2), 107-116.
- LEWINSOHN E. and GRESSEL J., 1983: The determination of chlorophylls a and b together with CO₂ dioxide fixation in the same plant tissue samples. Anal. Biochem. 135(2), 438-442.
- LEWINSOHN E. and GRESSEL J., 1984a: Benzyl viologen suppression of diquat and paraquat phytotoxicities. Plant Physiol. 75(1 suppl.), 51.
- LEWINSOHN E. and GRESSEL J., 1984b: Benzyl viologen-mediated counteraction of diquat and paraquat phytotoxicities. Plant Physiol. 76(1), 125-130.
- LEWINSOHN E. and GRESSEL J., 1985: Protection against paraquat and diquat damage by benzyl viologen in a plant model system. Phytoparasitica 13(3/4), 233.
- LEWIS D.H., 1984: Storage carbohydrates in vascular plants. Distribution, physiology and metabolism. Cambridge Univ. Press. 284 pp.
- LEWIS W.M. and BENDER M., 1961: Effect of a cover of duckweeds and the alga *Phytolacca* upon the dissolved oxygen and free carbon dioxide of small ponds. Ecology 42, 602-603.
- LEWKOWICZ M. and LEWKOWICZ S., 1977: Restoration of a pond after a five year period of fertilization with beet sugar factory wastes. Chemical factors and zooplankton. Acta Hydrobiol. 19(4), 315-333.
- LI H.-L., LIU T., HUANG T., KOYAMA T. and DE VOL C.E., 1978: Flora of Taiwan. Epoch Publ. Co., Taipei, Taiwan. 5, 816-818.
- LI Z., HUANG G. and NI J., 1985: The stabilizing factor of the nitrate reductase activity from the leaves of higher plants. (In Chinese). Zhiwu Shengli Xuebao 11(1), 1-8.
- LID J., 1952: Flora of Norway. (In Norw.). Norske Samlaget, Oslo. 180.
- LIEBERT H.-P., 1977: Einfluss von Abscisinsäure auf Wachstum und Mineralstoffgehalt bei *Lemma gibba* L. Biol. Rundsch. 15, 180-182.
- LIEBERT H.-P., 1980a: Zur Kontrolle des lichtinduzierten Wurzelwachstums bei Heterotrophkulturen von *Lemma gibba* L. durch Phytohormone. Biol. Rundsch. 18(3), 169-171.

- LIEBERT H.-P., 1980b: Der Kalium-, Calcium- und Magnesiumgehalt verschiedener alter Sprosse von *Lemma gibba* L. Biol. Rundsch. 18, 229-232.
- LIEBERT H.-P., 1986a: Der Einfluss von Schwermetallen auf das Wurzelwachstum heterotroph vorkultivierter *Lemma gibba* L. Biol. Rundsch. 24, 59-60.
- LIEBERT H.-P., 1986b: Einfluss von Kalium und Licht auf den Mineralstoffgehalt im Dunkeln vorkultivierter Pflanzen von *Lemma gibba* L. Wiss. Z. Friedrich-Schiller-Univ. Jena, Natw. R. 35(5), 639-645.
- LIEBERT H.-P. and AUGSTEN H., 1986: Der Einfluss von Licht auf die Nitratreduktase-Aktivität bei *Lemma gibba* L. und *Wolffia arrhiza* (L.) Wimm. in Abhängigkeit von Kulturbedingungen. Wiss. Z. Friedrich-Schiller-Univ. Jena, Natw. R. 35(5), 647-652.
- LIEDTKE M.P. and OHMANN E., 1969: Synthese und Inaktivierung von alkalischer Phosphatase in *Lemma minor*. Flora A 160(4), 378-390.
- LIEN B.C., COLE A.L.J., WALKER J.R.L. and PETERS J.A., 1979: Effect of sodium fluoroacetate ("Compound 1080") on the soil microflora. Soil Biology and Biochemistry 11(1), 13-18.
- LIEN B.C., COLE A.L.J., WALKER J.R.L. and PETERS J.A., 1981: The effect of fluoroacetate ("Compound 1080") and fluoroide on duckweeds. New Zealand Sci. 23, 179-184.
- LIN A. and MATHES M.C., 1973: The in-vitro secretion of growth regulators by isolated callus tissue. Am. J. Bot. 60(1), 34-41.
- LINDEMAN W., 1951: The influence of phosphate on the photosynthesis of *Lemma minor* L. Proc. Kon. Ned. Acad. Wetensch. C 54, 287-295.
- LINDEMAN W., 1952: The influence of phosphate on the photosynthesis of *Lemma minor* L. (In Dutch). Ph.D. Thesis. Univ. Amsterdam.
- LINDEMAN W., 1972: Reactivation of photosynthesis in dependence on wavelength in phosphate deficient *Lemma minor*. Acta Bot. Neerl. 21, 86-94.
- LINDEMAN W., 1973: Emerson enhancement effect and the reactivation of photosynthesis in phosphate deficient *Lemma minor*. Acta Bot. Neerl. 22, 553-568.
- LINDEMAN W., 1979: Inhibition of photosynthesis in *Lemma minor* by illumination during chilling in the presence of oxygen. Photosynthetica 13, 175-185.
- LINN J.G., GOODRICH R.D., MEISKE J.C. and STABA E.J., 1973: Aquatic plants from Minnesota. Part IV. Nutrient comparison. Univ. Minn. Water Resources Res. Center Bull. 56, 22 pp.
- LINNE C.S. see VON LINNE
- LIPKIN Y., 1973: *Wolffia arrhiza* (L.) Hork. ex Wimmer on the Golan Plateau. Israel J. Bot. 22, 175-177.
- LIPSER H. and DOBERAUER G., 1958: Ein zweiter reicher Bestand der Zwergwasserlinse (*Wolffia arrhiza* [L.] Wimm.) im Offenbacher Gebiet. Hess. Flor. Br. 7, 3-4.
- LITTLE C.S., 1979: Handbook of utilization of aquatic plants. FAO Fisheries Techn. Paper 187, 176 pp.
- LIU L.C. and CEDENO-MALDONADO A., 1974: Effects of fluometron, prometryne, ametryne, and diuron on growth of two *Lemma* species. J. Agric. Univ. Puerto Rico 58, 483-488.
- LIU L.C. and CEDENO-MALDONADO A., 1979: A bioassay method for detecting herbicide concentrations in water. J. Agric. Univ. Puerto Rico 63(1), 80-83.
- LOATS K.V., NOBLE R. and TAKEMOTO B., 1981: Photosynthesis under low level SO₂ and CO₂ enhancement conditions in three duckweed species. Bot. Gaz. 142(3), 305-310.
- LOCK J.M., 1973: The aquatic vegetation of Lake George, Uganda. Phytocoenol. 1, 250-262.

- LOCKHART W.L. and BLOUW A.P., 1979: Phytotoxicity tests using the duckweed Lemna minor. Can. Spec. Publ. Fish. Aquat. Sci. 44, 112-118.
- LOCKHART W.L., BILLECK B.N., DE MARCH B.G.E. and WUIR D.C.G., 1983: Uptake and toxicity of organic compounds: studies with an aquatic macrophyte (Lemna minor). ASTM Spec. Tech. Publ. 802, 460-468.
- LOCKHART W.L., BILLECK B.N. and BUCHKO G.W., 1985: Phytotoxicity testing with common duckweed (Lemna minor). Can. Tech. Rep. Fish. Aquat. Sci. 1368, 297-298.
- LUKINA M.M., 1976: Peculiar features of pollen sac development in some species of Najadaceae Juss. and Lemnaceae S. Gray. (In Russian). Bot. Zh. (USSR) 61, 1536-1546.
- LOEFFELHARDT W. and KINDL H., 1979: Conversion of 4-hydroxyphenylpyruvic acid into homogentisic acid at the thylakoid membrane of Lemna gibba. FEBS Lett. 104, 332-334.
- LOEPFERT H., 1979: Evidence for electrogenic proton extrusion by subepidermal cells of Lemna paucicostata 6746. Planta 144(4), 311-315.
- LOEPFERT H., 1981: Energy coupling for membrane hyperpolarization in Lemna: evidence against an ATP-fueled electrogenic pump as the exclusive mechanism. Planta 151, 293-297.
- LOEPFERT H., 1983: Energy coupling for membrane hyperpolarization in Lemna: respiration rate, ATP level and membrane potential at low oxygen concentrations. Planta 159(4), 329-335.
- LOEPFERT H. and KOENIGSHOFER H., 1984: Die Energetik elektrogenen Transportprozesse bei Lemna-Protoplasten. Mitt. Bd. Bot.-Tag. Wien, 50 (0504).
- LOEPFERT H. and KRONBERGER W., 1979: Control of nitrate uptake by photosynthesis in Lemna paucicostata 6746. In: MARCELLE R., CLIJSTERS H. and VAN POUCKE M. (eds.), Photosynthesis and Plant Development, Proc. Conf. 1978. Junk, The Hague, 301-308.
- LOEPFERT H. and TROMBALLA H.W., 1984: K transport and membrane potential in Lemna. In: CRAM W.J. et al. (eds.), Membrane transport in plants. Symp. Abstr., 218-219.
- LOEPFERT H., KRONBERGER W. and KANDELER R., 1977: Correlation between nitrate uptake and alkalization by Lemna paucicostata 6746. In: THELLIER M. et al. (eds.), Transmembrane ionic exchanges in plants. Edit. CNRS et Univ. Rouen, Paris et Mont Saint Aignan, 283-288.
- LOEPFERT H., KRONBERGER W. and KANDELER R., 1978: Phytochrome-mediated changes in the membrane potential of subepidermal cells of Lemna paucicostata 6746. Planta 138, 133-136.
- LOETSCH B. and KINZEL H., 1971: Zum Calciumbedarf von Oxalatzpflanzen. Biochem. Physiol. Pflanzen 162(3), 209-219.
- LOEVE A., 1978: IOPB chromosome report LXI. Taxon 27, 375-392.
- LOEWENGART G., 1984: Aquatic toxicity and environmental fate studies - caprolactam. Proc. Symp. Ind. Approach Chem. Risk Assess. Conf. Proc. 36-52.
- LOENUS P. and LOENUS M.W., 1971: Isolation of glucose-6 phosphate cyclodolase from cell cultures of Acer pseudoplatanus and evidence for histidine at the active center of this enzyme. Plant Physiol. 47 (suppl.), 5.
- LOHMAN G., 1938: Wasserchemie und höhere Vegetation schwedischer Seen. Symb. Bot. Ups. 3/1, 252 pp.
- LOHMAN G., 1940: Die Verbreitung von Lemna trisulca in Fennoskandina-vien und Dänemark. Verh. Int. Ver. Theor. Angew. Limnologie 9, 204-209.
- LOOMAN J., 1983: Water meal, Wolffia arrhiza (Lemnaceae) in Saskatchewan. Can. Field-Nat. 97(2), 220-222.
- LOOKAN J., 1985: The vegetation of the Canadian Prairie Provinces. III.

- Aquatic and semi-aquatic vegetation., part 3. Aquatic plant communities. Phytocoenologia 14, 19-54.
- LOOS S. and LUETGE U., 1984: Effects of HgCl₂ on membranes of Lemna gibba in the energized and non-energized state. Physiol. Veg. 22(2), 171-179.
- LOOS W., 1962: Einfluss der Gibberellinsäure auf die vegetative Vermehrung und das Wurzelwachstum von Lemna minor L. Phytol. (Argentina) 18, 133-136.
- LOT A., NOVELO A. and QUIROZ A., 1979: The Chinampa: an agricultural system that utilizes aquatic plants. J. Aquat. Plant Manage. 17, 74-75.
- LOVORN J.R. and KIRKPATRICK C.M., 1982: Analysis of freshwater wetland vegetation with large-scale color infrared aerial photography. J. Wildl. Manage. 46, 61-70.
- LUDWIG F., 1909: Lemnaceen. In: VON KIRCHNER O., LOEW E. and SCHROETER C., Lebensgeschichte der Blütenpflanzen Mitteleuropas. Ulmer, Stuttgart. 1(3), 57-80.
- LUBOEND A., 1978: Unterschiedliche Nährstoffansprüche nahverwandter Arten und ihre ökologische Bedeutung. Diploma Thesis. Geobot. Inst. ETH, Stiftung Rübel, Zürich. 84 pp. (Polycopy).
- LUBOEND A., 1980: Effects of nitrogen and phosphorus upon the growth of some Lemnaceae. Veröff. Geobot. Inst. ETH, Stiftung Rübel, Zürich 70, 118-141.
- LUBOEND A., 1982: Effects of nitrogen and phosphorus upon the growth of some Lemnaceae. In: SYMOENS J.J., HOOPER S.S. and COMPERE P. (eds.), Studies on aquatic plants. Symp. Abstr., 143.
- LUBOEND A., 1983: Das Wachstum von Wasserlinsen (Lemnaceae) in Abhängigkeit des Nährstoffangebots, insbesondere Phosphor und Stickstoff. Veröff. Geobot. Inst. ETH, Stiftung Rübel, Zürich 80, 116 pp.
- LUETGE U., JUNG K.D. and ULLRICH-EBERHARDT C.I., 1981: Evidence for amino acid-H cotransport in Lemna gibba given by effects of fusococcin. Z. Pflanzenphysiol. 102(2), 117-125.
- LUKINA G.A., 1977: Flowering of the Lemnaceae. (In Russian). Byull. Inst. Biol. Vnutr. Vod. Akad. Nauk SSR 35, 29-31.
- LUKINA G.A., 1983a: Effect of UV irradiation on some representatives of the family Lemnaceae S. Gray. (In Russian). Gidrobiol. Zh. 19(1), 66-69.
- LUKINA G.A., 1983b: The effect of ultraviolet radiation on some representatives of the family Lemnaceae S. Gray. Hydrobiol. 19(1), 68-71.
- LUKINA G.A., 1984: Duckweeds under natural and laboratory conditions. (In Russian). Bot. Zh. 69, 81-83.
- LUTHER H., 1948: Die Funde von blühenden Lemnaceen in Finnland. Mem. Soc. Fauna Flora Fenn. 24, 161-170.
- LUTHER H., 1949: Vorschlag zu einer ökologischen Grundeinteilung der Hydrophyten. Acta Bot. Fenn. 44, 3-15.
- LUTHER H., 1951: Verbreitung höherer Wasserpflanzen im brackisch Wasser Finnlands. Acta Bot. Fenn. 50, 1-370.
- LYSENKO N.I. and GRODZINSKII D.M., 1974: Effect of small doses of previous irradiation on radio resistance of Spiridella polyrrhiza. (In Russian). Fiziol. Biokhim. Kul't. Rast. 6(4), 371-375.
- LYSENKO N.I., 1981: Effect of monofunctional alkylating mutagens on the radiation resistance of Spiridella polyrrhiza (L.) Schleid. and Pisum sativum. (In Russian). Nauk Dumka (Kiev) 1981, 43-46.
- MABRY T.J., MARKHAM K.R. and THOMAS M.B., 1970: The systematic identification of flavonoids. Books 1970, 354.
- MACDONALD I.R., 1975: Effect of vacuum infiltration on photosynthetic gas exchange in leaf tissue. Plant Physiol. 56, 109-112.

- MACCINA M.J. and SHIREMAN J.V., 1980: Effects of salinity on vegetation consumption and growth in grass carp *Ctenopharyngodon idella*. *Prog. Fish-Cult.* 42(1), 50-53.
- MACCINTIE H.D., LEOPOLD E.B. and ROHRER W.L., 1974: An early middle Eocene flora from the Yellowstone-Absaroka volcanic province, northwestern wind river basin, Wyoming. *Univ. Calif. Publ. Geol. Sci.* 108, 94.
- MACIEJEWSKA-POTAPCZYK W., KONORSKA L. and NARZYMSKA E., 1970: Proteins in duckweed (*Lemna minor* L.). *Acta Soc. Bot. Pol.* 39, 251-255.
- MACIEJEWSKA-POTAPCZYK W., KONORSKA L. and OLECHNIEWICZ K., 1975: Protein in *Lemna minor* L. *Biochem. Physiol. Pflanzen* 167, 105-108.
- MACIEJEWSKA-POTAPCZYK W., KONORSKA L. and RESZKA J., 1976: Influence of TAA and kinetin on *Lemna minor* L. proteins. *Acta Univ. Nicolai Copernici Nauk. Mat.-Przyr.* 37, 183-186.
- MACKENTHUN K.M., INGRAM W.M. and PORCES R., 1964: Limnological aspects of recreation lakes. *Pub. Health Serv. Publ. No.* 1167.
- MACNICOL P.K., DATKO A.H., GIOVANELLI J. and MUDD S.H., 1981: Homocysteine biosynthesis in green plants: physiological importance of the transsulfuration pathway in *Lemna paucicostata*. *Plant Physiol.* 68(3), 619-625.
- MADONI P. and VIAROLI P., 1985: Microfauna distribution in shallow macrophyte-covered basins. *Verh. Int. Verein. Limnol.* 22, 2353-2355.
- MADSEN J.D. and ADAMS M.S., 1985: The aquatic macrophyte communities of two streams in Wisconsin. *Wis. Acad. Sci., Arts Lett.* 73, 198-216.
- MAENG J., 1973: Biochemical and physiological studies on the flowering mechanism of two species of *Lemna*. Northeastern Univ. Diss. Abstr. Int. B 34, 2496-2497.
- MAENG J., 1976: Studies on inhibition factors and the role of phytochrome in the floral induction in short-day plants. (In Korean). *Korean J. Bot.* 19, 14-18.
- MAENG J., 1977: Studies on the effect of 2-chloroethylphosphoric acid [etephon] on the floral induction to photoperiodic plants: I. Inhibition of flowering in *Lemna perpusilla* 6746 by etephon. (In Korean). *Korean J. Bot.* 20, 77-82.
- MAENG J. and BAE J.M., 1984: Influence of salicylic acid and dimethylsulfoxide on flowering *Lemna gibba* G3. *Korean J. Bot.* 27(4), 215-222.
- MAENG J. and KHUAI A.K., 1973: Studies on the flowering mechanism in *Lemna*. I. Amino acid changes during flower induction. *Physiol. Plant.* 28, 264-270.
- MAENPAEAE P. and ARO E.M., 1986: Chlorophyll-protein complexes, chlorophyll a/b ratio and chloroplast ultrastructure in *Lemna minor* L. grown under different light conditions. *J. Plant Physiol.* 123(2), 161-168.
- MAGONE I.G. and GRODZINSKII D.M., 1973: Abscisic acid as a modifier of plant radiation damage. (In Russian). *Fiziol. Biokhim. Kul't. Rast.* 5(4), 427-430.
- MAGONE I.G. and TEIVANS A., 1985: Bioindication of emission products from a cement mill. (In Russian). *Zagryaz. Priir. Sredy Kal'siisoderzh. Pyl'yu* 90-95.
- MAHANTY H.K., 1975: A study on the effects of polychlorinated biphenyl (Aroclor 1242) on an aquatic plant *Spirodela oligorrhiza* (Kurz) Hegelm. *Bull. Environ. Contam. Toxicol.* 14(5), 558-561.
- MAHANTY H.K. and FINERAN B.A., 1976: Effects of a polychlorinated biphenyl (Aroclor 1242) on the ultrastructure of frond cells in the aquatic plant *Spirodela oligorrhiza* (Kurz) Hegelm. *New Zealand J. Bot.* 14(1), 13-18.
- MAHANTY H.K. and McWHIA J.A., 1976: Sensitivity of *Spirodela oligorrhiza*

- (Kurz) Hegelm. to a polychlorinated biphenyl (Aroclor 1242). *New Zealand J. Bot.* 14(1), 9-12.
- MAHAR H., 1974: The combined effects of ⁶⁰Co irradiation and selected herbicides on *Spirodela polyrrhiza*. *Int. Diss. Abstr.* B 35(5), 2489.
- MAHESHWARI J.K., 1960: The vegetation of marshes, swamps and riverside in Khandwa district (Madhya Pradesh). *J. Bombay Nat. Hist. Soc.* 57, 271-307.
- MAHESHWARI S.C., 1954: The embryology of *Wolffia*. *Phytomorphology* 4, 355-365.
- MAHESHWARI S.C., 1956a: Endosperm and seed of *Wolffia*. *Nature* 178, 925-926.
- MAHESHWARI S.C., 1956b: The endosperm and embryo of *Lemna* and systematic position of the Lemnaceae. *Phytomorphology* 6, 51-55.
- MAHESHWARI S.C., 1958a: The Lemnaceae. A contribution to their biology, morphology and systematics. Ph.D. Thesis. Univ. Delhi.
- MAHESHWARI S.C., 1958b: *Spirodela polyrrhiza*: The link between the aroids and the duckweeds. *Nature* 181, 1745-1746.
- MAHESHWARI S.C., 1959: Systematic position of the family Lemnaceae. *Abstr. Int. Congr. Montreal* 2, 246-247.
- MAHESHWARI S.C., 1963 see MAHESHWARI S.C. and CHAUHAN O.S., 1963
- MAHESHWARI S.C. and CHAUHAN O.S., 1963: In vitro control of flowering in *Wolffia microscopica*. *Nature* 198, 99-100.
- MAHESHWARI S.C. and GUPTA S., 1967: Induction of flowering in *Lemna paucicostata*, a short-day plant, by chelating agents and iron. *Planta* 77, 95-98.
- MAHESHWARI S.C. and KAPIL R.N., 1963a: Morphological and embryological studies on the Lemnaceae. I. The floral structure and gametophytes of *Lemna paucicostata*. *Am. J. Bot.* 50, 677-686.
- MAHESHWARI S.C. and KAPIL R.N., 1963b: Morphological and embryological studies on the Lemnaceae. II. The endosperm and embryo of *Lemna paucicostata*. *Am. J. Bot.* 50, 907-914.
- MAHESHWARI S.C. and KAPIL R.N., 1964: Morphological and embryological studies on the Lemnaceae. III. The seed and seedling of *Lemna paucicostata*. *J. Ind. Bot. Soc.* 43, 270-277.
- MAHESHWARI S.C. and KHURANA J.P., 1978: Floral induction in duckweeds by salicylic acid. In: SEN D.N. and BANSAL R.P. (eds.), *Environ. Physiol. Ecol. Plants*. B. and M.P. Singh Publ., Dehra Dun, India. 5-13.
- MAHESHWARI S.C. and MAHESHWARI N., 1963: The female gametophyte, endosperm and embryo of *Spirodela polyrrhiza*. *Beitr. Biol. Pflanzen* 39, 179-188.
- MAHESHWARI S.C. and SETH P.N., 1966a: Induction of flowering in *Wolffia microscopica* by the iron salt of ethylenediamine-di-o-hydroxyphenylacetic acid (Fe-EDDA). *Z. Pflanzenphysiol.* 55, 89-91.
- MAHESHWARI S.C. and SETH P.N., 1966b: Photoperiodic control of flowering in *Wolffia papulifera*. *Plant Cell Physiol.* 7, 163-165.
- MAHESHWARI S.C. and SETH P.N., 1967 see MAHESHWARI S.C. et al., 1967
- MAHESHWARI S.C. and VENKATARAMAN R., 1966: Induction of flowering in a duckweed - *Wolffia microscopica* - by a new kinin, zeatin. *Planta* 70, 304-306.
- MAHESHWARI S.C., VENKATARAMAN R. and SETH P.N., 1967: Effects of kinins and an iron chelate on the growth and flowering of *Wolffia microscopica*. *Proc. Int. Conf. Plant Growth Substances, Rostock. Wiss.-Z. Univ. Rostock, Math.-Natw. R.* 16, 655.
- MAHLOCH J.L. and McGRUFF E.C., Jr., 1974: Agricultural wastes. *J. Water Poll. Control Fed.* 46(6), 1280-1283.
- MAI D.H. and WALTHER H., 1978: Die Floren der Haselbacher Serie im Weis-

selster-Becken (Bez. Leipzig, DDR). Abh. Staatl. Mus. Min. Geol. Dresden 28, 149.

MAJID F.Z. and JAHAN M.A.A., 1984: Study on the effects of some aquatic weed extract solutions on seed germination and primary seedling vigour of some economic crops. Proc. 4th Nat. Bot. Conv., Bangladesh. 12 pp.

MAKAROVA Z.Y., 1968: The biomass of common duckweed in Kaliningrad waters and the dynamics of it during the growing period. (In Russian). Tr. Kaliningrad. Tekh. Inst. Rybn. Prom. Khoz. 20, 170-174.

MALECHA S.R., BUCK D.H., BAUR R.J. and ONIZUKA D.R., 1981: Polyculture of the freshwater prawn, *Macrobrachium rosenbergii*, Chinese and common carps in ponds enriched with swine manure. I. Initial trials. Aquaculture 25(2-3), 101-116.

MALEK L., 1981: The effect of drying on Spirodela polyrrhiza turion germination. Can. J. Bot. 59(1), 104-105.

MALEK L. and COSSINS E.A., 1982: Aspartate pathway activity in N- and S-deficient duckweed. Plant Physiol. 69(4 suppl.), 111.

MALEK L. and COSSINS E.A., 1983 see 1983a

MALEK L. and COSSINS E.A., 1983a: Senescence, turion development, and turion germination in nitrate- and sulfate-deficient Spirodela polyrrhiza. Relationships between nutrient availability and exogenous cytokinins. Can. J. Bot. 61(7), 1887-1897.

MALEK L. and COSSINS E.A., 1983b: Aminoacylation of tRNA and protein turnover in nitrate- and sulphate-deficient Spirodela polyrrhiza. Plant Cell Physiol. 24(8), 1353-1359.

MALEK L. and ODA Y., 1979: Effect of red and blue light on turion production. Plant Physiol. 63(5 suppl.), 156.

MALEK L. and ODA Y., 1980: Germination of Spirodela polyrrhiza turions: the role of culture conditions during turion development. Plant Cell Physiol. 21(2), 357-361.

MALENDE B., 1957: Die Zwerglinse, *Wolffia arrhiza* (L.) Wimmer, bei Steinhelm a. Main. Hess. Florist. Br. 6, 1-2.

MALIS G.P. and MUIR D.C.G., 1984: Fate of fenitrothion in shaded and unshaded ponds. ACS Symp. Ser. 238, 277-295.

MALKIN R. and POSNER H.B., 1978: On the site of function of the Rieske iron-sulfur center in the chloroplast electron transport chain. Biochim. Biophys. Acta 501, 552-554.

MANCINELLI A.L., 1977: Photocontrol of anthocyanin synthesis in Spirodela polyrrhiza. Plant Physiol. 59(6 suppl.), 49.

MANCINELLI A.L., 1980: Photoreceptors of the high irradiance responses of plant photomorphogenesis. Photochem. Photobiol. 32(6), 853-858.

MANCINELLI A., 1984: Photoregulation of anthocyanin synthesis. VIII. Effect of light pretreatments. Plant Physiol. 75(2), 447-453.

MANCINELLI A.L. and RABINO I., 1984: Photoregulation of anthocyanin synthesis. X. Dependence on photosynthesis of high irradiance response anthocyanin synthesis in *Brassica oleracea* leaf disks and Spirodela polyrrhiza. Plant Cell Physiol. 25(7), 1153-1160.

MANGI J., SCHMIDT K., PANKOW J., GAINES L. and TURNER P., 1978: Effects of chromium on some aquatic plants. Environ. Pollut. 16(4), 285-291.

MANISSEY J.K., KRISHNA R.D.S. and DEVARAJ K.V., 1981: Observations on

the feeding of the duckweed (*Lemna minor*) by an ostracod (*Cypris* sp.). Curr. Res., Univ. Agric. Sci., Bangalore 10(3), 52-53.

MANN H., 1967: The utilization of food by *Tilapia melanopleura* Dum. FAO Fish. Rep. 44(3), 408-410.

MANSOR M. and AHMAD M., 1986: The dominant aquatic weeds in peninsular Malaysia. Proc. 7th Int. Symp. Aquatic Weeds. European Weed Res. Soc., Wageningen. 207-212.

MARCIULIONIENE E.-D.P., 1980: Interactions of radionuclides and pollutants with phytocenoses in fresh-water ecosystems. Agrochem. Residue-Biota Interact. Soil Aquat. Ecosyst., Proc. Rep. Comb. Advis. Group Meet. Res. 113-124.

MARCIULIONIENE E.-D.P., DUSAUSKIENE-DUZ R.G., CIBIRAITIS N.A., NENINSKIENE V., SULIENE R. and TRAINAUSKAITIS I., 1979: Absorption of some radionuclides of metals by freshwater plants. (In Russian). Vzglimod. Vod. Zhiv. Vesh. 1, 90-97.

MARDER J.B., MATTOO A.K. and EDELMAN M., 1982: Photosynthesis and the rapidly-metabolized 32-kilodalton chloroplast membrane protein. Prog. Clin. Biol. Res. 102B, 91-100.

MARDER J.B., GOLOUBINOFF P. and EDELMAN M., 1984: Molecular architecture of the rapidly metabolized 32 kilodalton protein of photosystem. II. Indications for carboxyl terminal processing of a chloroplast membrane polypeptide. J. Biol. Chem. 259(6), 3900-3908.

MARIE-VICTORIN FRÈRE, 1931: Les Spadiciflores du Québec (Aracées, Lemnacees). Contr. Lab. Bot. Univ. Montréal 19, 60 pp.

MARKLUND F., 1936: Vergleichende Permeabilitätsstudien an pflanzlichen Protoplasten. Acta Bot. Fenn. 18, 1-110.

MARKS G.C., 1968: The flowering of *Lemna minor* and the establishment of *Centaurium pulchellum* in Northwestern Indiana. Proc. Indiana Acad. Sci. 78, 414-416.

MARQUES I.A. and ERISMANN K.H., 1984: Metabolism of glycolate by *Lemna minor* L. grown on different inorganic nitrogen sources. Plant Physiol. 75(1 suppl.), 156.

MARQUES I.A., OBERHOLZER M.J. and ERISMANN K.H., 1985: Metabolism of glycolate by *Lemna minor* L. grown on nitrate or ammonium as nitrogen source. J. Exp. Bot. 36(172), 1685-1697.

MARTI J., TROESCH L. and ERISMANN K.H., 1986: Gaswechsel-Test mit der Wasserlinse *Lemna minor* L. zur Untersuchung allelopathisch wirksamer Stoffe. Schweiz. Bot. Ges., Bull. 4(suppl.), 9-10.

MARTIN G., 1955: Action antitoxique des ions Mg^{++} à l'égard des ions Zn^{++} chez *Lemna minor*. C.R. Soc. Biol. 149, 2099-2102.

MARTIN G., 1963a: Différences spécifiques entre les besoins quantitatifs en chlore de *Lemna minor*, *L. perpusilla* et *Spirodela polyrrhiza*. Plant and Soil 18, 258-266.

MARTIN G., 1963b: Nécessité du chlore dans la nutrition de *Spirodela polyrrhiza* cultivé en conditions partiellement hétérotrophes. C.R. Acad. Sci., Paris 257, 294-296.

MARTIN G., 1964: Effets de la L-cystine sur la croissance de *Spirodela polyrrhiza* en conditions hétérotrophes. C.R. Acad. Sci., Paris 258, 5757-5759.

MARTIN G., 1965: Nécessité du chlore dans la nutrition de *Spirodela polyrrhiza* cultivé en conditions hétérotrophes. C.R. Acad. Sci., Paris 260, 5928-5930.

MARTIN G. and LAVOLLAY J., 1958a: Le chlore, oligo-élément indispensable pour *Lemna minor*. Experientia 14, 333-334.

MARTIN G. and LAVOLLAY J., 1958b: Sur la spécificité de la carence en chlore chez *Lemna minor*. C.R. Soc. Biol. 152(2), 241-244.

- MARTIN G. and LAVOLLEY J., 1959: Sur la spécificité de la carence en chlore chez Lemna minor. Recherches sur l'utilité de l'iode. C.R.Soc. Biol. 153, 1130-1133.
- MARTIN J.F., CAUDRON G., JAN E. and KOFMAN E., 1978: Le langage en traitement tertiaire. Expérimentation à partir de plantes aquatiques supérieures. Le cresson et les lentilles d'eau. I-III. Tech.Eau Assainissement 376, 23-34; 377, 21-37; 378/379, 19-25.
- MARTINSSON K., 1984: The flowering of Lemna minor. (In Swedish). Svensk. Bot.Tidskr. 78(1), 9-15.
- MARTONNE E. see DE MARTONNE E.
- MASCARO L.J., Jr., 1975: Characterization of apogalacturonans formed by a cell-free system from Lemna minor. Ph.D.Thesis. Univ.Microfilms Int., Ann Arbor, Mich., Order No. 75-27,298; 190 pp. Diss.Abstr.Int. B 36, 2777-2778.
- MASCARO L.J., Jr. and KINDEL P.K., 1973: Biosynthesis of cell wall apio-galacturonans by a cell-free system from Lemna minor. Fed.Proc. 32(3 part 2), 1257.
- MASCARO L.J., Jr. and KINDEL P.K., 1974: Biosynthesis of cell wall apio-galacturonans by a cell-free system from Lemna minor. Fed.Proc. 33(5 part 2), 1445.
- MASCARO L.J., Jr. and KINDEL P.K., 1977: Characterization of (¹⁴C) apio-galacturonans synthesized in a cell-free system from Lemna minor. Arch.Biochem.Biophys. 183, 139-148.
- MASON H.L., 1938: The flowering of Wolffia linguata (Hegelm.) Hegelm. Madroño 4(8), 241-251.
- MASON H.L., 1957: A flora of the Marshes of California. Lemnaceae. duckweed family. Univ.California Press,Berkeley/Los Angeles. 327-343.
- MASON R., 1950: Water plants of New Zealand. Hutchinson et al., Wellington. ton.
- MATHESON R., 1930: The utilization of aquatic plants as aids in mosquito control. Am.Nat. 64, 56-86.
- MATHESON R. and HINMAN E.H., 1929: Further studies on Chara spp. and other aquatic plants in relation to mosquito breeding. Am.J.Trop.Med. 9, 249-266.
- MATHEZ J., 1973: Nouveaux matériaux pour la flore du Maroc. II. Contribution à l'étude de la flore de la région d'Ifni. (Polycopie).
- MATHIS B.J., CUMMINGS T.F., GOWER M., TAYLOR M. and KING C., 1977: Dynamics of manganese, cadmium and lead in experimental power plant ponds. Res.Rep. Univ.Ill., Urbana-Champaign, Water Resour.Centre. 125, 62 pp.
- MATHIS B.J., CUMMINGS T.F., GOWER M., TAYLOR M. and KING C., 1980: Dynamics of manganese, cadmium and lead in experimental power plant ponds. Hydrobiologia 67(3), 197-206.
- MATHUR S.N. and YADAVA S.R., 1974: Hill-reactivity and some other metabolic processes of Spirodela polyrrhiza as affected by maleic hydrazide. In: PURI V., MURTY Y.S., GUPTA P.K. and BANERJI D. (eds.), Biology of the Land Plants. Symposium, Meerut Univ. 173-178.
- MATHUR S.N. and YADAV S.R., 1975: Effect of maleic hydrazide on the growth of Spirodela polyrrhiza: interaction with some purine and pyrimidine bases. Indian J.Plant Physiol. 18(1), 8-11.
- MATSUMOTO S., 1981: Absorption of nutritional salts by Spirodela polyrrhiza and its application to ecological salt control. (In Japan.). Kagaku to Seibutsu (Chemistry and Biology) 19(9), 594-600.
- MATSUURA R., NAGAOKA K., KUSABE K. and NAKATANI S., 1979: p-n-alkylbenzoic acid. (In Japan.). Ger.Offen.800807 Patent:Applic.No. 79/982526. 26 pp.

- MATTOO A.K. and CONLON C.A., 1985: Protein metabolism in higher plants: Enhanced breakdown of soluble and membrane proteins in Spirodela oligorhiza supplemented with copper sulphate. Plant Physiol. 77(4 suppl.), 131.
- MATTOO A.K. and EDELMAN M., 1985: Photoregulation and metabolism of a thylakoidal herbicide-receptor protein. In: ST.JOHN J.B., BERLIN E. and JACKSON P.C. (eds.), Symp.Agric.Res., Frontiers of membrane research in agriculture, Beltsville. 23-34.
- MATTOO A.K. and EDELMAN M., 1987: Intramembrane translocation and post-translational palmitoylation of the chloroplast 32-kDa herbicide-binding protein. Proc.Natl.Acad.Sci.USA. 84(6), 1497-1501.
- MATTOO A.K., EDELMAN M. and GRESSEL J., 1980: Dependence on light and differentiated thylakoids for synthesis and processing of 33.5 kD plastid membrane protein in Spirodela. Plant Physiol. 65(6 suppl.), 21.
- MATTOO A.K., PICK U., HOFFMAN-FALK H. and EDELMAN M., 1981: The rapidly metabolized 32,000-dalton polypeptide of the chloroplast is the "proteaceous shield" regulating photosystem II electron transport and mediating diuron herbicide sensitivity. Proc.Natl.Acad.Sci.USA. 78(3), 1572-1576.
- MATTOO A.K., MARDER J.B., GRESSEL J. and EDELMAN M., 1982: Presence of the rapidly-labelled 32,000-dalton chloroplast membrane protein in triazine resistant biotypes. FEBS Lett. 140(1), 36-40.
- MATTOO A.K., ST.JOHN J.B. and WERGIN W.P., 1983a: Is turnover of the 32,000-dalton thylakoid protein a signal for adaptation of the chloroplast to triazine herbicides and to light conditions. J.Cell Biochem. (suppl. 7 part B), 319.
- MATTOO A.K., WERGIN W.P. and ST.JOHN J.B., 1983b: Adaptive reorganization of thylakoid components of Spirodela oligorhiza cultured on sublethal doses of atrazine. Plant Physiol. 72(suppl.1), 56.
- MATTOO A.K., BAKER J.E. and LIEBERMAN M., 1983c: Copper induced ethylene production by Spirodela oligorhiza is a consequence of singlet oxygen mediated membrane damage. Plant Physiol. 72(suppl.1), 39.
- MATTOO A.K., MARDER J.B., ST.JOHN J.B., WERGIN W.P. and EDELMAN M., 1984a: D-threo-chloramphenicol-induced depletion of the 32 kDa protein in Spirodela: photosynthetic reactions, lipids and morphology. Plant Physiol. 75(1 suppl.), 22.
- MATTOO A.K., ST.JOHN J.B. and WERGIN W.P., 1984b: Adaptive reorganization of protein and lipid components in chloroplast membranes as associated with herbicide binding. J.Cell Biochem. 24(2), 163-176.
- MATTOO A.K., HOFFMAN-FALK H., MARDER J.B. and EDELMAN M., 1984c: Regulation of protein metabolism: coupling of photosynthetic electron transport to in-vivo degradation of the rapidly metabolized 32-kDa protein of the chloroplast membranes. Proc.Natl.Acad.Sci.USA 81(5), 1380-1384.
- MATTOO A.K., BAKER J.E. and MOLINE H.E., 1984d: Different mechanisms in Spirodela and Nicotiana of copper-induced ethylene production. Plant Physiol. 75(suppl.1), 185.
- MATTOO A.K., BAKER J.E. and MOLINE H.E., 1986a: Induction by copper ions of ethylene production in Spirodela oligorhiza: evidence for a pathway independent of 1-aminocyclopropane-1-carboxylic acid. Plant Physiol. 123(3), 193-202.
- MATTOO A.K., MARDER J.B., GABA V. and EDELMAN M., 1986b: Control of 32-kDa thylakoid protein degradation as a consequence of herbicide binding to its receptor. Plant Biol. 2, 607-613.
- MATTOO A.K., MEHTA R.A., CONLON C.A. and PORATH D., 1987: Turnover of

- ribulose-1,5-bisphosphate carboxylase-oxygenase involves its membrane association during copper-induced senescence of *Spirodela oligorrhiza*. *Plant Physiol.* 83(4 suppl.), 26.
- MATVEEV V.I., 1963: Blooming in *Lemna gibba* L. (In Russian). *Bot. Zh.* (USSR) 48, 272.
- MATVEEV V.I., 1977: On the flowering of duckweeds [Lemnaceae] in waterbodies of Trans-Volga territories. (In Russian). *Bot. Zh.* (USSR) 62, 1498-1500.
- MAUVE A.A., 1966: Flowering aquatic plants in South Africa. *Fauna and Flora* 17, 19-31.
- MAZUR B.J. and CHUI C.-F., 1985: Sequence of a genomic DNA clone for the small subunit of ribulose bis-phosphate carboxylase-oxygenase from tobacco. *Nucleic Acids Res.* 13(7), 2373-2386.
- MCALLA T.M. and PLUCKNETT D.L., 1981: Collecting, transporting, and processing organic fertilizers (in China). In: *Vegetable farming systems in China*. Westview Press, Boulder. 19-37.
- MCANN C., 1942: Observations on Indian duckweeds, Lemnaceae. *J. Bomb. Nat. Hist. Soc.* 43(2), 148-162.
- MCCLURE J.W., 1964: Taxonomic significance of the flavonoid chemistry and the morphology of Lemnaceae in axenic culture. *Ph.D. Thesis*. Univ. Texas, Austin. 235 pp. *Diss. Abstr.* 25, 4928-4929.
- MCCLURE J.W., 1967a: Flavonoid variation in *Lemna perpusilla* from different laboratories. *Plant Cell Physiol.* 8, 523-526.
- MCCLURE J.W., 1967b: Effects of light on the synthesis of flavonoids in *Spirodela intermedia*. *Plant Physiol.* 42(suppl. 44).
- MCCLURE J.W., 1968: Photocontrol of *Spirodela intermedia* flavonoids. *Plant Physiol.* 43, 193-200.
- MCCLURE J.W., 1969: Environmental control of flavonoid glycoflavone synthesis. 11th Int. Bot. Congr., Abstr., 134.
- MCCLURE J.W., 1970: Secondary constituents of aquatic angiosperms. In: *HARBORNE J.B. (ed.), Phytochemical Phylogeny*, *Proc. Phytochem. Soc. Symp.*, Bristol. Acad. Press, New York/London. 233-268.
- MCCLURE J.W., 1973: Regulation of growth and flavonoid accumulation in *Spirodela intermedia* by light, partial satisfaction by growth substances, and complications introduced by substances accumulating in the culture medium. *Am. J. Bot.* 60(4 suppl.), 28-29.
- MCCLURE J.W., 1975: The applicability of polyphenolic data to systematic problems in the Lemnaceae. *Aquat. Bot.* 1, 395-405.
- MCCLURE J.W. and ALSTON R.E., 1963: Chromatographic pattern of phenolic constituents in 18 duckweed spp. (Lemnaceae). *Am. J. Bot.* 50, 636.
- MCCLURE J.W. and ALSTON R.E., 1964a: Patterns of selected chemical components of *Spirodela oligorrhiza* formed under various conditions of axenic culture. *Nature* 201(4916), 311-313.
- MCCLURE J.W. and ALSTON R.E., 1964b: Glycoflavonoids of *Spirodela* species and their taxonomic implications. *Am. J. Bot.* 51, 685-686.
- MCCLURE J.W. and ALSTON R.E., 1966: A chemotaxonomic study of Lemnaceae. *Am. J. Bot.* 53, 849-860.
- MC COMBS P.J.A. and RALPH R.K., 1972a: Protein, nucleic acid and starch metabolism in the duckweed, *Spirodela oligorrhiza*, treated with cytokinins. *Biochem. J.* 129, 403-417.
- MC COMBS P.J.A. and RALPH R.K., 1972b: Cytokinin control of growth of *Spirodela oligorrhiza* in darkness. *Planta* 107(2), 97-109.
- MC COWEN M.C. et al., 1979: Fluridone, a new herbicide for aquatic plant management. *J. Aquat. Plant Manage.* 17, 27-30.
- MC CRADY J.W., WENTZ W.A. and LINDER R.L., 1986: Plants and invertebrates in a prairie wetland during duck brood-rearing. *Prairie Nat.* 18(1), 23-32.

- MCDONALD I.R., 1975: Effect of vacuum infiltration on photosynthesis gas exchange in leaf tissue. *Plant Physiol.* 56, 109-112.
- MC HARGUE J.S. and COFUE R.K., 1932: Manganese essential for growth of *Lemna major*. *Plant Physiol.* 7, 697-703.
- MC LAREN J.S. and SMITH H., 1976: The effect of abscisic acid on growth, photosynthetic rate and carbohydrate metabolism in *Lemna minor* L. *New Phytol.* 76, 11-20.
- MC LAREN J.S. and SMITH H., 1977: Effect of abscisic acid on photosynthetic products of *Lemna minor*. *Phytochemistry* 16, 219-221.
- MC LAY C.L., 1973: Wind-blown dust as a source of nutrients for aquatic plants. *Environ. Pollut.* 5, 173-180.
- MC LAY C.L., 1974: The distribution of duckweed *Lemna perpusilla* in a small southern Californian lake: An experimental approach. *Ecology* 55, 262-276.
- MC LAY C.L., 1976: The effect of pH on the population growth of three species of duckweed: *Spirodela oligorrhiza*, *Lemna minor* and *Wolffia arrhiza*. *Freshwater Biol.* 6, 125-136.
- MC MINN A., 1985: Structure of the vegetative shoots of *Spirodela polyrrhiza*. *Univ. Wisconsin.*
- MC NEIL M., DARVILL A.G., FRY S.C. and ALBERSHEIM P., 1984: Structure and function of the primary cell walls of plants. *Ann. Rev. Biochem.* 53, 625-633.
- MC VAUGH R., 1972: Nomenclature report of the committee for spermatophyta conservation of generic names. Part 15. *Taxon* 21(4), 531-535.
- MC WHA J.A. and JACKSON D.L., 1976: Some growth promotive effects of abscisic acid. *J. Exp. Bot.* 27(100), 1004-1008.
- MEIJER L. and SUTTON D.L., 1987: Influence of plant position on growth of duckweed. *J. Aquat. Plant Manage.* 25, 28-30.
- MELARAGNO J.E., 1974: Anatomical studies on the duckweed *Lemna minor* L. A light and electron microscope investigation with emphasis on ultrastructural details of sieve-element ontogeny. *Ph.D. Thesis*. Univ. of Pittsburgh, PA.
- MELARAGNO J.E. and WALSH M.A., 1974: Ultrastructural features of developing sieve elements in *Lemna minor*: the protoplast. *Am. J. Bot.* 61(5 suppl.), 60.
- MELARAGNO J.E. and WALSH M.A., 1976: Ultrastructural features of developing sieve elements in *Lemna minor*: The protoplast. *Am. J. Bot.* 63, 1145-1157.
- MELZER A., 1980: Oekophysiologische Aspekte der N-Ernährung submerser Wasserpflanzen. *Verh. Ges. Ökologie* 7, 357-362.
- MELZER A. and EXLER D., 1982: Nitrate- and nitrite-reductase activities in aquatic macrophytes. In: *SYMOENS J.J., HOOPER S.S. and COMPERE P. (eds.), Studies on aquatic vascular plants*. *R. Soc. Bot. Belg.* 128-135.
- MELZER A. and MUELLER M., 1984: In vivo-nitrate reductase activities of different species of Lemnaceae and comparison with endogenous nitrate depletion. *Proc. Int. Symp. Aquatic Macrophytes* 1983, Nijmegen, 139-144.
- MENDICINO J. and ABON-ISSA H., 1974: Conversion of UDP-D-glucuronic acid to UDP-D-apiose and UDP-D-xylose by an enzyme isolated from *Lemna minor*. *Biochim. Biophys. Acta* 364, 159-172.
- MENDICINO J. and HANNA R., 1970: The synthesis of isomers of D-apiofuranosyl 1-phosphate. *J. Biol. Chem.* 245(22), 6113-6124.
- MENDICINO J. and PICKEN J.M., 1965: Biosynthesis of the branched chain sugar-D-apiose in *Lemna* and *Parsley*. *J. Biol. Chem.* 240, 2797-2805.
- MENDIOLA N.B., 1919: Variation and selection within clonal lines of *Lemna minor*. *Genetics* 4, 151-182.

- MENSCHICK R., 1970: Untersuchungen über den durch die Stickstoffquelle beeinflussbaren Flavonoidgehalt der Sommerglieder und Winterknospen von *Spirodela polyrrhiza* (L.) Schleid. Ph.D.Thesis. Math.-Nat.Fak. Univ. Münster. 110 pp.
- MERIAUX J.-L., 1978: Etude analytique et comparative de la végétation aquatique d'étangs et marais du Nord de la France (Vallée de la Sennée et Bassin Houillier du Nord-Pas de Calais). Doc.Phytosoc.N.S. 3, 1-244.
- MERIAUX J.-L. and GEHU J.M., 1978: De l'analyse symphytosociologique des complexes de végétation à celles des végétations complexes. In: TUXEN R. (ed.), Assoziationskomplexe (Sigmerten). Ber.Int.Symp., Rinteln. 97-115.
- MERTEN O.W., 1979: Nachweis von Abscissinsäure im Zusammenhang mit der Bildung von Reservestoffen in Pflanzen. Diploma Thesis. Univ.Agric., Vienna, Austria. 120 pp.
- MESTAYER C.R., 1980: A study on the solar energy conversion efficiency and growth aspects of duckweed, *Spirodela oligorrhiza*. M.S.Thesis. Louisiana State Univ., Baton Rouge. 57 pp.
- MESTAYER C.R., CULLEY D.D., jr., STANDIFER L.C. and KOONCE K.L., 1984: Solar energy conversion efficiency and growth aspects of the duckweed, *Spirodela punctata* (G.F.W.Mey.) Thompson. Aquat.Bot. 19(1-2), 157-170.
- MEUSEL H., 1951: Die Bedeutung der Wuchsform für die Entwicklung des natürlichen Systems der Pflanzen. Feddes Rep. 54, 137-172.
- MEYER F.A., 1964: Aquatic plant control. Dept. of Fish and Game, Sacramento, CA. Inland Fish.Adm.Rep.No. 642.
- MEYER G.F.W., 1818: Primitiae Florae Essequeboensis adjectis descriptionibus centum circiter stirpium novarum, observationibusque criticis. H. Dieterich, Göttingen. 262.
- MICHELI P.A., 1729: Nova plantarum genera. Florentiae.
- MICHEWICZ J.E., SUTTON D.L. and BLACKBURN R.D., 1972: Water quality of small enclosures stocked with white amur. Hyacinth Contr.J. 10, 22-25.
- MIEHITZ G., 1956: Die Wasserlinsen auf Teichen. Allg.Fischereizeitung 81, 112.
- MIFLIN B.J. and LEA P.J., 1980: Ammonia assimilation. In: STUMPF P.K. and CONN E.E. (eds.), The biochemistry of plants. Acad.Press, New York. 5, 169-202.
- MIKI S., 1934: On fresh water plants new to Japan. Bot.Mag. 48(569), 326-337.
- MIKRAKOVA T.F., 1980: Effect of cadmium on the growth of *Lemna minor*. (In Russian). Biol.Vnstr.Vod. 48, 22-25.
- MIKRAKOVA T.F., 1983: Effect of salts of heavy metals on the growth and accumulative ability of *Lemna minor*. (In Russian). Trud.Akad.Nauk SSSR, Inst.Biol.Vnstr.Vod. 48, 31-37.
- MILDE C.A., 1853: *Wolffia Michellii* Hook. (*Lemna arhriza* L.). Bot.Z. 11, 896-897.
- MILLER A.T., 1978: Plant seeds in biological research in outer space. (In Russian). Latv.Psr.Zinat.Akad.Vestis. 7, 82-89.
- MILLER D.L., COMBATTI N., WAGNER G.J. and HILLMAN W.S., 1977: Protoplast isolation from *Lemna perpusilla* and *Lemna gibba*. Plant Physiol. 59(6 suppl.), 114.
- MILLERS A. and MAGONE I., 1978: Relation of the combined effect of vibration and ionizing radiation to the physiological state of the plant. (In Russian). Fiziol.Biohim.Issled.Rast. 135-140.
- MIQUEL F.A.W., 1853: Preliminary report on a new *Wolffia*. (In Dutch). Ned.Kruidk.Arch. 3, 425-429.

- MIRASHI M.V., 1954: Studies in the hydrophytes of Nagpur. J.Indian Bot. Soc. 33, 299-398.
- MIRASHI M.V., 1957: Studies in the hydrophytes of Umred. J.Indian Bot. Soc. 36, 396-407.
- MISRA G. and DAS N., 1969: Studies on the control of aquatic weeds of orissa-response of *Pistia stratiotes* L., *Spirodela polyrrhiza* Schleid. and *Ipomoea aquatica* Forsk to hormone herbicides. Hyacinth Contr.J. 8(1), 40-41.
- MITCHELL D.S., 1969: The ecology of vascular hydrophytes on Lake Kariba. Hydrobiol. 34(3-4), 448-464.
- MITCHELL D.S., 1978: Aquatic weeds in Australian inland waters. Austral. Govt.Publ.Serv., Canberra. 189 pp.
- MITRA E. and BANERJEE A.C., 1976: Utilization of higher aquatic plants in fishery waters. Proc.Reg.Sem. Noxious Aquat.Veg. 375-381.
- MIYABE K. and KUDO Y., 1932: Flora of Hokkaido and Saghalien II. Hokkaido Imp.Univ., Sapporo.
- MIYATA H., 1970: Endogenous light-on rhythm in respiration of a long-day duckweed, *Lemna gibba* G3. Plant Cell Physiol. 11, 293-301.
- MIYATA H., 1971a: Endogenous light-on rhythm in respiration of a long-day duckweed, *Lemna gibba* G3. II. On basic and rhythmic components of the rhythm. Plant Cell Physiol. 12, 517-524.
- MIYATA H., 1971b: Endogenous light-on rhythm in respiration of a long-day duckweed, *Lemna gibba* G3. III. Relationship to frond production. Plant Cell Physiol. 12, 969-977.
- MIYATA H. and YAMAMOTO Y., 1969: Rhythms in respiratory metabolism of *Lemna gibba* G3 under continuous illumination. Plant Cell Physiol. 10, 875-889.
- MIYAWAKI A. and TUXEN J., 1960: Ueber Lemnëta-Gesellschaften in Europa und Japan. Mitt.Florist.-Soz.Arb.gem. Stolzenau/Weser 8, 127-135.
- MOCKERIDGE F.A., 1920: The occurrence and nature of the plant growth producing substances in various organic material composts. Biochem J. 14, 432-450.
- MOCKERIDGE F.A., 1924: The formation of plant growth promoting substances by microorganisms. Ann.Bot. 38, 23-34.
- MODONI P. and VIAROLI P., 1985: Microfauna distribution in shallow macrophyte-covered basins. Verh.Int.Verein.Limnol. 22, 2353-2355.
- MOFFITT S. and BLACKMAN G.E., 1972: The uptake of growth substances. Part 15. The differential effects of progressive chlorination of phenoxacetic acid on entry into the epidermal and cut surfaces of stem segments. J.Exp.Bot. 23(74), 128-140.
- MOLNARI E. and HOFFMANN-OSTENHOF O., 1968: Untersuchungen über die Biosynthese der Cyclite, XXI. Ueber ein Enzymsystem, das myo-Inositol zu Phytinsäure phosphorylieren kann. Hoppe Seyler's Z.Physiol.Chem. 349(12), 1797-1799.
- MONACO P. and PREVITERA L., 1987: Studies on aquatic plants distributed in Italy. Part 5. Oxygenated fatty acids from *Lemna trisulca*. Phytochemistry 26(3), 745-748.
- MONINGER R., 1968a: Un mécanisme possible d'action de la lumière sur les biosyntheses de caroténoïdes chez les végétaux supérieurs chlorophylliens. Bull.Soc.Fr.Physiol.Vég. 14(4), 473-497.
- MONINGER R., 1968b: Sur la réalisation d'un dispositif ultrasensible permettant de séparer et de doser les caroténoïdes de petits échantillons de frondes étioilées de *Spirodela polyrrhiza* Schleid. C.R.Acad. Sci., Paris D 266(7), 672-675.
- MONINGER R., 1968c: Influence de la durée d'éclairement préalable de frondes étioilées de *Spirodela polyrrhiza* Schleid. sur les radio-acti-

- vités incorporées à l'obscurité dans leurs caroténoïdes, à partir d'acétate $2-14^{\circ}\text{C}$ de sodium. C.R.Acad.Sci., Paris D 267(7), 733-736.
- MONÉGER R., 1968d: Contribution à l'étude de l'influence exercée par la lumière sur la biosynthèse des caroténoïdes chez la *Spirodela polyrrhiza* (L.) Schleiden. *Physiol.Vég.* 6(2), 165-202.
- MONÉGER R., 1968e: Mise au point d'une méthode de micro-analyse pour l'étude des caroténoïdes de petits échantillons de végétaux étioles. *Physiol.Vég.* 6(4), 367-402.
- MONÉGER R., 1968f: Incorporations de radio-activité, à partir de bicarbonate $2-14^{\circ}\text{C}$ et d'acétate $2-14^{\circ}\text{C}$ de sodium, dans les caroténoïdes de frondes étioles de *Spirodela polyrrhiza* Schleiden. exposées à des radiations oligochromatiques. C.R.Acad.Sci., Paris D 267, 605-608.
- MONÉGER R. and JACQUES R., 1968: Action de radiations oligochromatiques sur les teneurs en caroténoïdes des frondes étioles de *Spirodela polyrrhiza* Schleiden. C.R.Acad.Sci., Paris D 267(3), 313-316.
- MONÉGER R., VERMEERSCH J., LECHEVALIER D. and RICHARD C., 1977: Micro-analyse du NADP et du NAD réduits et oxydés dans les tissus foliaires et dans les plastides isolés de *Spirodela* et de *Blé*. I. Problèmes posés par le dosage séparé du NADP et du NAD réduits et oxydés des extraits végétaux. *Physiol.Vég.* 15(1), 29-62.
- MONETTI P.G. and BASTELLI R., 1983: Caratteristiche chimiche e prove di conservabilità di piante acquatiche coltivate su reflui zootecnici. In: GHETTI P.F. (ed.), *Fitodepur. Impieghi Biomasse Prod., Atti Conv.* Int. 171-179.
- MONNIER A., 1971: Electrokinetic formulation of competitive interactions in cellular absorption processes. *Physiol.Vég.* 9(1), 107.
- MONNIER A. and THELLIER M., 1970: Interprétation électrocinétique des interactions ioniques phosphate-arsénate lors de l'absorption par la Lemna minor. C.R.Acad.Sci., Paris D 270(18), 2178-2181.
- MONOD T., 1949: Sur une Lemnace africaine: *Wolffella Welwitschii* (Hemelmayer 1865) comb. nov. *Mém.Soc.Hist.Nat.Afr.Nord* 2, 229-242.
- MONSELVE E.B.-I., PORATH D. and TAL M., 1984: Unusual 'tubular clusers' in the plastids of a duckweed (*Lemna paucicostata*) mutant incapable of photosynthesis and ammonium ion uptake. *New Phytol.* 98, 249-257.
- MONSELVE E.B.-I., PORATH D., TAL M. and ARZEE T., 1986: Partial recovery in plastids of Lemna aequinoctialis Welwitsch 1073 treated with L-methionine or L-cystine. *New Phytol.* 103, 283-291.
- MONTELVE see MONSELVE
- MOODY R.P., GREENHALGH R., LOCKHART L. and WEINBERGER P., 1978: The fate of fenitrothion in an aquatic ecosystem. *Bull.Environ.Contam.Toxicol.* 19(1), 8-14.
- MOORE J.R., 1962: The effect of a duckweed cover on the phyto-community structure of a pond. M.S.Thesis. Univ. of Pittsburgh, PA.
- MOORE J.R., 1965: Productivity and standing crop of vascular hydrophytes. Ph.D.Thesis. Univ. of Pittsburgh, PA.
- MOORE L.B. and EDGAR E., 1970: Flora of New Zealand. Shearer, Wellington, N.Z. 2, 53-55.
- MOORE P.D., 1983: Alien duckweed hiding quietly beside the Cam. *Nature* 302, 384.
- MORET L., 1943: Manuel de paléontologie végétale. Masson, Paris.
- MORI H., 1979a: Kinetics of induction and production of flowers in a short-day duckweed, *Lemna paucicostata* 6746, in darkness. *Plant Cell Physiol.* 20, 615-621.
- MORI H., 1979b: Effect of red light pulse on induction and production of flowers in a short-day duckweed, *Lemna paucicostata* 6746, in darkness. *Plant Cell Physiol.* 20, 623-630.

- MORI H., 1979c: Effect of temperature on induction and production of flowers in *Lemna paucicostata* 6746 in uninterrupted and interrupted darkness. *Plant Cell Physiol.* 20, 631-638.
- MORI H., 1979d: Effect of far-red light pulse on induction and production of flowers in *Lemna paucicostata* 6746 in darkness. *Plant Cell Physiol.* 20, 639-648.
- MORI H., 1984a: Induction and reproduction of flowers in a short-day duckweed, *Lemna paucicostata* 6746, in diluted Hutter's medium in uninterrupted and interrupted darkness. *Plant Cell Physiol.* 25(6), 923-928.
- MORI H., 1984b: Effects of ferrous and phosphate ions on flowering in *Lemna paucicostata* 6746 in diluted Hutter's medium. *Plant Cell Physiol.* 25(6), 929-933.
- MORRIS P.F. and BAKER W.G., 1977: Oxygen transport rates through mats of Lemna minor and Wolffia sp. and oxygen tension within and below the mat. *Can.J.Bot.* 55, 1926-1932.
- MORRIS R.O. and MACDONALD E.M.S., 1986: Immunoaffinity purification of plant hormones as an analytical tool. *J.Cell Biochem.* (suppl. 10 part B), 30.
- MORVAN C., DEMARTY M. and THELLIER M., 1979: Titration of isolated cell walls of Lemna minor L. *Plant Physiol.* 63, 1117-1122.
- MORVAN C., DEMARTY M., MONNIER A., MUQIBIL R. and THELLIER M., 1980: Titration of isolated cell walls of Lemna minor L. In: SPANSWICK R.M., LUCAS W.J. and DAINTY J. (eds.), *Plant membrane transport: Current conceptual issues*. Elsevier, North-Holland. Biomed.Press. 423-424.
- MORVAN C., MORVAN H., DEMARTY M. and THELLIER M., 1983: Réponses électriques des lentilles d'eau à différentes conditions expérimentales. *C.R.Soc.Biol. (Rouen)* 177(5/6), 679-688.
- MORVAN C., DEMARTY M. and THELLIER M., 1984: Influence des charges négatives des acides pectiques sur les propriétés d'échange d'ions des parois végétales. *C.R.Soc.Biol. (Rouen)* 178(6), 712-721.
- MORVAN C., DEMARTY M. and THELLIER M., 1985: Propriétés physicochimiques des parois de Lemna minor: échanges entre les ions calcium et lithium. *Physiol.Vég.* 23(4), 444-444.
- MOSS E.A., 1959: Flora of Alberta. Univ.Toronto Press. 546 pp.
- MOUTERDE P., 1966: Nouvelle Flore du Liban et de la Syrie. Beyrouth. 1, 194-196.
- MOYLE J.B., 1945: Some chemical factors influencing the distribution of aquatic plants in Minnesota. *Am.Midl.Nat.* 34(2), 402-420.
- MOYLE J.B. and HOTCHKINS N., 1945: The aquatic and marsh vegetation of Minnesota and its value to waterfowl. *Minn.Dept.of Conserv., Game and Fish Tech.Bull.* 3, 1-22.
- MUDD S.H. and DATKO A., 1986a: Methionine methyl group metabolism in Lemna. *Plant Physiol.* 81(1), 103-114.
- MUDD S.H. and DATKO A., 1986b: Phosphoethanolamine bases as intermediates in phosphatidylcholine synthesis by Lemna. *Plant Physiol.* 82(1), 126-135.
- MUDD S.H. and DATKO A., 1987: Patterns of methylation in phosphatidylcholine synthesis. *Plant Physiol.* 83(4 suppl.), 113.
- MUDROCH A. and CAROBIANCO J.A., 1979: Effects of treated effluent on a natural marsh. *J.Water Pollut.Contr.Fed.* 51(9), 2243-2256.
- MUEHL H. and LOEFELHARDT W., 1982: 4-Hydroxyphenylpyruvate dioxygenase is bound to the thylakoid membranes from the cyanobacterium *Anacystis nidulans*. *FEMS Microbiol.Lett.* 13(1), 9-12.
- MUELLER B. and ZIEGLER H., 1969: Die lichtinduzierte Aktivitätssteigerung der NADP-abhängigen Glycerinaldehyd-3-phosphat-Dehydrogenase.

- IX. Die Reaktion in isolierten Chloroplasten. *Planta* 85(1), 96-104.
- MUELLER E., 1973: Cyclic AMP as a signal carrier in biological systems. II. Is there a signal carrier function of cyclic AMP in higher plants. *Biol. Rundsch.* 11(6), 352-361.
- MUELLER M., 1983: Ökophysiologische Untersuchungen zum Stickstoffumsatz verschiedener Lemnaceen. Diploma Thesis, TU München. 118 pp.
- MUELLER P., 1975: Der Einfluss verschiedener CO₂-Konzentrationen auf das Wachstum und den Gehalt einiger Produkte des Photosyntheseabhängigen Stoffwechsels von Lemna minor L. bei Nitrat- und Ammoniumernährung. Lic.Thesis. Univ. Bern.
- MUELLER P., FELLER U. and ERISMANN K.H., 1977: Einfluss verschiedener CO₂-Konzentrationen auf Wachstum und stoffliche Zusammensetzung von Lemna minor L. bei Nitrat- und Ammoniumernährung. *Z. Pflanzenphysiol.* 85, 233-241.
- MUELLER T., 1977: Klasse: Lemnaceae. In: OBERDORFER E. (ed.), *Süddeutsche Pflanzengesellschaften*. (2nd ed.). Fischer, Stuttgart/New York. 1, 67-77.
- MUELLER T. and GOERS S., 1960: Pflanzengesellschaften stehender Gewässer in Baden-Württemberg. *Beitr. Naturk. Forsch. SW-Deutschl.*, Karlsruhe, 19, 60-100.
- MUELLER Z. and LAUTNER V., 1954: Feeding value of some water plants. II. (In Czech.). *Sb. Československé Akad. Zemedl. Ved.* 27A, 451-472.
- MUENSCHER W.C., 1944: Aquatic plants of the United States. Comstock Publ., Ithaca.
- MUONEN M., SHOWNAN J. and COUCH R., 1983: Nutrient absorption by Spirodela polyrrhiza. *J. Aquat. Plant Manage.* 21(2), 107-109.
- MUIR D.C.G. and GRIFF N.P., 1978: Disappearance of the experimental aquatic herbicide fluridone (EL-171) in small ponds. *PEST* 56.
- MUIR D.C.G., GRIFF N.P., BLOW A.P. and LOCKHART W.L., 1980: Persistence of fluridone in small ponds. *J. Environ. Qual.* 9, 151-156.
- MUIR D.C.G., PITZE M., BLOW A.P. and LOCKHART W.L., 1981: Fate of terbutryne in macrophyte-free and macrophyte-containing farm ponds. *Weed Res.* 21(2), 59-70.
- MUIR D.C.G., GRIFF N.P. and LOCKHART W.L., 1982: Comparison of laboratory and field results for prediction of the environmental behaviour of phosphate esters. *Environ. Toxicol. Chem.* 1(2), 113-119.
- MUIR D.C.G., RAWN G.P. and GRIFF N.P., 1985: Fate of the pyrethroid insecticide deltamethrin in small ponds: a mass balance study. *J. Agric. Food Chem.* 33, 603-609.
- MUKHOPADHYAY S.K., 1982: Noxious aquatic vegetation in West Bengal. Vishva-Bharati Coll., Sriniketan, India. *Abstr. Ann. Conf. Ind. Soc. Weed Sci.* 47.
- MUKHOPADHYAY S.K., 1986: Aquatic weed control in India. *Pesticides (Bombay)* 20(11), 15-22.
- MULLENDERS W., 1967: Flore de la Belgique, du Nord de la France et des Régions voisines. Desoer, Liège. 505-506.
- MURATA G., BEPPU T. and NOBUCHI T., 1981: On Mijinkoukikusa (water meal). (In Japan.). *Acta Phytotax. Geobot.* 32, 197.
- MURKOWSKI A., GRABIKOWSKI E., BRZOSTOWSKI A. and PROKOSKI Z., 1982: Luminiscence method for bioindication of phytotoxic environmental pollutants. (In Polish). *Postępy Fiz. Med.* 17(3-4), 85-91.
- MURKOWSKI G.S., ZOL'NIKOVA N.V. and BOROVKOV A.V., 1977: Duckweed as a test-object for the evaluation of phytotoxic activity of soil microorganisms. (In Russian). *S.-Kh. Biol.* 12, 227-230.
- MURPHY K.J. and EATON J.W., 1983: Effects of pleasure-boat traffic on macrophyte growth in canals. *J. Appl. Ecol.* 20(3), 713-729.

- MURPHY K.J., HANBURY R.G. and EATON J.W., 1981: The ecological effects of 2-methylthiothiazine herbicides used for aquatic weed control in navigable canals. I. Effects on aquatic flora and water chemistry. *Arch. Hydrobiol.* 91(3), 294-331.
- MUSIL C.F., 1973: Water plants of Natal. *Wildlife Protect. Conserv. Soc. South Africa*. 62 pp.
- MUSIL C.F., GRUNOW J.O. and BORMAN C.H., 1973: Classification and ordination of aquatic macrophytes in the Pongolo River Pans, Natal. *Botanica 11(1/2)*, 181-190.
- MUSTIN M., 1983: Dans les marais Poitevin: l'exemple des lentilles d'eau. *Biomasse Actual.* 3, 55.
- MUZAFFAROV A.M., TAUBAEV T.T. and ABDIEV M., 1968: Use of lesser duckweed (Lemna minor) in poultry feeding. (In Russian). *Uzb. Biol. Zh.* 12(3), 44-46.
- MUZAFFAROV A.M., TAUBAEV T.T. and ABDIEV M., 1971: Lesser duckweeds (Lemna minor L.) as an animal food and methods of its mass cultivation in the basins under the open sky. (In Russian). *Kult. Vod. Vysch. Vodn. Rast. Uzbekistane* 1971, 117-130.
- MUZTAR A.J., SLINGER S.J. and BURTON J.H., 1976: Nutritive value of aquatic plants for chicks. *Poult. Sci.* 55(5), 1917-1922.
- MUZTAR A.J., SLINGER S.J. and BURTON J.H., 1977: Metabolizable energy content of freshwater plants in chicken and ducks. *Poult. Sci.* 56(6), 1893-1899.
- MUZTAR A.J., SLINGER S.J. and BURTON J.H., 1978: Chemical composition of aquatic macrophytes. I. Investigation of organic constituents and nutritional potential. II. Amino acid composition of the protein and nonprotein fractions. III. Mineral composition of freshwater macrophytes and their potential for mineral nutrient removal from lake water. *Can. J. Plant Sci.* 58(3), 829-862.
- MUZTAR A.J., SLINGER S.J. and BURTON J.H., 1979: Chemical composition of aquatic macrophytes. IV. Carotenoids, soluble sugars and starch in relation to their pigmentation, and ensiling potential. *Can. J. Plant Sci.* 59, 1093-1098.
- MYERS R.W., 1977: A comparative study of nutrient composition and growth of selected duckweeds, Lemnaceae, on dairy waste lagoons. M.S. Thesis. Louisiana State Univ., Baton Rouge. 75 pp.
- NAKAMURA H., 1960: A study of Wolffia as a new food. *Rep. Microalgae Inst., Japan*, 1, 7-13.
- NAKASHIMA H., 1964: Effects of exogenous amino acids on the flower and frond production in duckweed, Lemna gibba G3. *Plant Cell Physiol.* 5, 217-225.
- NAKASHIMA H., 1965: Further studies on the action of free amino acids on flowering of duckweed, Lemna gibba G3. *Plant Cell Physiol.* 6, 441-452.
- NAKASHIMA H., 1966: The rhythmical change in sensitivity of a long-day duckweed, Lemna gibba G3, to dark-break. *Plant Cell Physiol.* 7, 11-24.
- NAKASHIMA H., 1967: Change in sensitivity to dark break and its relation to floral induction in long-day duckweed, Lemna gibba G3. *Plant Cell Physiol.* 8, 637-645.
- NAKASHIMA H., 1968: On the rhythm of sensitivity to light interruption in a long-day duckweed, Lemna gibba G3. *Plant Cell Physiol.* 9, 247-257.
- NAKASHIMA H., 1969: Effect of inhibitors of nucleic acid and protein synthesis on the reappearance of light interruption rhythm in a long-

- day duckweed, Lemna gibba G3. Plant Cell Physiol. 10(2), 259-270.
- NAKASHIMA H., 1973a: Reversal of the dark inhibition of flowering in a long-day duckweed, Lemna gibba G3, by thymidine and related nucleosides. Plant Cell Physiol. 14, 893-899.
- NAKASHIMA H., 1973b: Effect of light on the metabolism of thymidine in the long-day duckweed, Lemna gibba G3. Plant Cell Physiol. 14, 901-910.
- NAKASHIMA H., 1975: Diurnal change of light-dependent uridine incorporation into RNA in a long-day duckweed, Lemna gibba G3. Plant Cell Physiol. 16, 27-39.
- NAKASHIMA H., 1976: Diurnal rhythm of uridine incorporation into RNA regulated by two light-perceiving systems in a long-day duckweed, Lemna gibba G3. Plant Cell Physiol. 17, 209-217.
- NAKASHIMA H., 1978: Further studies on the diurnal rhythm of uridine incorporation into RNA in the long-day duckweed, Lemna gibba G3. Plant Cell Physiol. 19(3), 375-383.
- NAKASHIMA H., 1979a: Diurnal rhythm of nuclear RNA polymerase I activity in a duckweed, Lemna gibba G3, under continuous light conditions. Plant Cell Physiol. 20(11), 165-176.
- NAKASHIMA H., 1979b: Effects of incubation of nuclei from a long-day duckweed, Lemna gibba G3, under different ionic conditions on the activity of RNA synthesis. Plant Cell Physiol. 20, 715-724.
- NAKASHIMA H. and MORI H., 1970: DNA synthesis as related to the appearance of 'light interruption rhythm' in a long-day duckweed, Lemna gibba G3. Plant Cell Physiol. 11, 805-816.
- NAKASHIMA H. and TSUDZUKI T., 1976: Uptake of uridine by a long-day duckweed, Lemna gibba G3. Plant Cell Physiol. 17, 701-711.
- NAKASHIMA H. and TSUDZUKI T., 1977: The effects of calcium withdrawal on the uridine uptake process in a long-day duckweed, Lemna gibba G3. Z. Pflanzenphysiol. 84, 399-405.
- NAKAYAMA T., 1986: Treatment of sulfuric acid-containing wastewater by aquaculture. (In Japan.). Patent: Jp 86197098; Japan Kokai Tokyo Koho. 14 pp.
- NAPHADE M.S. and NITHUI G.F., 1969-1970: Duckweed as a poultry food. B.A. (Bansilal Amritlal), Agric. Coll. Mag. 22, 73-77.
- NARUKHAMEDOV K. and VASIGOV T., 1972: Mass culture of Lemna minor L. in the southwestern Kyzyl-Kum area. (In Russian). Kult. Vod. Vyssh. Vodn. Rast. Uzbekistane 1972, 136-140.
- NASKAR K., BANERJEE A.C., CHAKRABORTY N.M. and GHOSH A., 1986: Yield of Wolffia arrhiza from cement cisterns with different sewage concentrations and its efficacy as a carp feed. Aquaculture 51(3-4), 211-216.
- NASU Y., 1983: Basic studies of environmental monitoring using Lemna paucicostata. Heavy metal absorption and action. (In Japan.) Nippon Eiseigaku Zasshi 38(5), 839-852.
- NASU Y. and KUGIMOTO M., 1981: Lemna (duckweed) as an indicator of water pollution. I. The sensitivity of Lemna paucicostata to heavy metals. Arch. Environ. Contam. Toxicol. 10, 159-169.
- NASU Y. and KUGIMOTO M., 1986: Absorption of heavy metals by Lemna paucicostata. (In Japan.). Iden 40(8), 41-47.
- NASU Y., KUGIMOTO M., TANAKA O. and TAKIMOTO A., 1983: Comparative studies on the absorption of cadmium and copper in Lemna paucicostata. Environ. Pollut. A 32(3), 201-209.
- NASU Y., KUGIMOTO M., TANAKA O., YANASE D. and TAKIMOTO A., 1984: Effects of cadmium and copper co-existing in the medium on the growth and flowering of Lemna paucicostata in relation to their absorption. Environ. Pollut. A 33(3), 267-274.

- NASU A., KUGIMOTO M., TANAKA O. and TAKIMOTO A., 1985: Lemna as an indicator of water pollution and the absorption of heavy metals by Lemna. In: PASCOE D. and EDWARDS R.W. (eds.), Freshwater biological monitoring. Pergamon Press, New York. 113-120.
- NAUMANN H., 1964: Kleintierleben im Wasserlinseenteppich. I. Aqu. Terr. Z. 17, 377-379.
- NAUMANN H., 1965: Kleintierleben im Wasserlinseenteppich. II. Aqu. Terr. Z. 18, 346-348.
- NAVARRO ANDRES F., SANCHEZ ANTA M.A. and GALLEGO M.F., 1984: El aparato vascular de Lemnaceas como resultado de un proceso de hidromorfosis. Stud. Bot. 3, 331.
- NAZARENKO S.I. and LUTKOVA I.N., 1974: Advances in medicinal plants. (In Russian). Vliy. Fiz.-Khim. Fakt. Rast. Org. 1974, 110-118.
- NECHUSHTAI R., NELSON N., MATTOO A.K. and EDELMAN M., 1981: Site of synthesis of subunits to photosystem I reaction center and the proton-ATPase in Spirodela. FEBS Lett. 125(1), 115-119.
- NECHUSHTAI R., PETERSON C.C. and THORNBER J.P., 1987a: Photosystem I complex of higher plants: purification and characterization of its two chlorophyll-protein complexes. In: BIGGINS J. (ed.), Prog. Photosynth. Res., Proc. 7th Int. Congr. Photosynth. Nijhoff, Dordrecht, Netherlands, 2, 41-44.
- NECHUSHTAI R., PETERSON C.C., PETER G.F. and THORNBER J.P., 1987b: Purification and characterization of a light-harvesting chlorophyll-a/b-protein of photosystem I of Lemna gibba. Eur. J. Biochem. 164, 345-350.
- NEEL P.L. and SUTTON D.L., 1977: Growth of four species of plants as influenced by composted sewage sludge and duckweed (Lemna gibba L.) grown in sewage effluent. Proc. SNA Res. Conf. Annu. Rep. (South Nurserymen's Assoc.) 22, 41-43.
- NEGBI M., SHIRHAI E.D., BITHAN N. and PORATH D., 1972: Ethylene induced separation of daughter plants in Spirodela polyrrhiza. Isr. J. Bot. 21, 108-111.
- NEIFF J.J. and MARCHESI E., 1978: Caracterización sinoptica de la vegetación acuática anfibia en el área del futuro embalse de Salto Grande. Estimación del riesgo potencial del desarrollo de hidrofitos en el embalse. Proc. V Reunión sobre Aspectos de Desarrollo Ambiental, 6-10 November 1978, Centro de Ecología Aplicada del Litoral, Argentina. 35 pp.
- NEIFF A.P. and NEIFF J.J., 1984: Dynamics of aquatic floating plants and related fauna in rain pools in Chaco, Argentina. Physis (Buenos Aires) B 42, 53-67.
- NEIL J.H., 1976: The harvest of biological production as a means of improving effluents from sewage lagoons. Res. Rep., Res. Program Abatement Munic. Pollut. Provis., Canada, Ontario. Agreement Great Lakes Water Qual. 38, 35 pp.
- NELLES S. and PARTHIER B., 1969: Protein synthesis in sterile chloroplasts from Lemna minor L.: A contribution to the role of bacterial contamination. Exp. Cell Res. 58, 225-233.
- NELSON B., 1981: Duckweeds - nature's smallest flowering plants. Aquatics 3(1), 4-16.
- NEUFARHT A., HARTZ P. and KNAUF W., 1979: Significance of detergents in sewage for model organisms from natural ecosystems. J. Com. Esp. Deterg. 10th, 125-140.
- NEWBOLD A. and HOLT R.J., 1986: Additions to the checklist of the flora of Montgomery County, Pennsylvania, USA, IV. Biol. Abstr. RM. Bartonica (suppl. 52), 78.
- NEWTON R.J., 1972a: Effects of abscisic acid on the root of Lemna minor. Tex. J. Sci. 24(3), 388.

- NEWTON R.J., 1972b: Effects of abscisic acid on growth and metabolism of protein and ribonucleic acid in Lemna minor L. Ph.D. Thesis, Univ. Ann Arbor, Michigan. 82 pp.
- NEWTON R.J., 1974a: Abscisic acid effects on growth and metabolism in the roots of Lemna minor. *Physiol. Plant.* 30, 108-112.
- NEWTON R.J., 1974b: Dual pattern of DL-leucine absorption by duckweed root tips. *Plant Cell Physiol.* 15, 249-254.
- NEWTON R.J., 1977: Abscisic acid effects on fronds and roots of Lemna minor. *Am. J. Bot.* 64(1), 45-49.
- NEWTON R.J. and DUFFEY J., 1975: Turion induction in *Spirodela polyrrhiza*. *Plant Physiol.* 56(suppl.), 85.
- NEWTON R.J., SHELTON D.R., DISHARON S. and DUFFEY J.E., 1978: Turion formation and germination in *Spirodela polyrrhiza*. *Am. J. Bot.* 65(4), 421-428.
- NEWTON S.H., MARTIN J.M., FERGUSON J.N. and GRAY D.L., 1979: Grass carp aid removal of weeds in irrigation canals and reservoirs. *Arkansas Farm Res.* 28(4), 12.
- NEY L.F., 1960: Gas exchange by the duckweed family. *ONR Contract No. 2887*, Standford Res. Inst.
- NG Y.L. and THIMANN K.V., 1962: Studies on the biogenesis of anthocyanins. VIII. Identification of the anthocyanin of *Spirodela*. *Arch. Biochem. Biophys.* 96, 336-339.
- NG Y.L., THIMANN K.V. and GORDON S.A., 1961: The action spectrum of anthocyanin formation in *Spirodela oligorrhiza*. *Plant Physiol.* 36 (suppl.), 46.
- NG Y.L., THIMANN K.V. and GORDON S.A., 1964: The biogenesis of anthocyanins. X. The action spectrum for anthocyanin formation in *Spirodela oligorrhiza*. *Arch. Biochem. Biophys.* 107, 550-558.
- NGUYEN T.L., 1978: Studies of the biology of the trematode *Fasciolopsis buski* in Vietnam. (In Russian). *Med. Parazit. Parazit. Bolezni* 47(2), 81-84.
- NICKELL L.G., 1955: Effects of antigrass substances in normal and atypical plant growth. *Antimetabolites and cancer. Symp. Am. Ass. Adv. Sci.* 129-151.
- NICKELL L.G., 1956: Aseptic studies on metabolism of nitrogenous compounds in plants. *Bull. Torr. Bot. Club* 83(6), 421-427.
- NICKELL L.G., 1962: Some simple substituted pyrimidines and their effect on the growth of Lemna minor. *Phyton (Argentina)* 18, 59-63.
- NICKELL L.G. and FINLAY A.C., 1954: Antibiotics and their effects on plant growth. *Agr. Food Chem.* 2, 178-182.
- NIELSON E.S., 1944: Dependence of freshwater plants on quantity of carbon dioxide and hydrogen ion concentration. *Botan. Ark.* 11(8), 1-25.
- NIEVEYER R., 1975: Poly- and Metaphosphate in höheren Pflanzen (Lemnaeae). *Planta* 122, 303-305.
- NIKANOROV A.M. and ZHULIDOV A.V., 1986: Background levels of mercury in tissues of aquatic organisms in the eastern European subarctic. (In Russian). *Monit. Fonovogo Zagryaz. Prirod. Sred.* 3, 186-190.
- NIKITIN V.V. and SEIFULIN E.M., 1976: New species for the flora of the Turkmenistan. (In Russian). *Izv. Akad. Nauk Turkmen SSR, Ser. Biol. Nauk* 5, 89.
- NIKOLAEVA A.M., 1956: Duckweeds as a feedstuff for water birds. (In Russian). *Pticevodstvo* 6, 24-29.
- NIKOLSKIJ G.V. and VERIGIN B.V., 1966: The basic biological characteristics of white amur and bighead and their acclimatization in the water reservoirs of our country. (In Russian). *Piscevaja Promyslennost (Moskva)*, 30-40.

- NILSEN S. and DANIELSEN S., 1984: Photoinhibition of photosynthesis in Lemna gibba: The effect of O₂, CO₂ and the selective inhibition of photosystem II. *J. Plant Physiol.* 115, 39-48.
- NILSEN S. and DANIELSEN S., 1985: Photoinhibition in vivo. Does cultivation in high CO₂ protect plants against photoinhibition? *Plant Physiol.* 64(2), 11a.
- NINNEMANN H., 1979: Photoreceptors for circadian rhythms. *Photomed. Photobiol. Rev.* 207, 207-266.
- NISHIOKA H., NASU Y., KUGIMOTO M., KAIHARA S. and TAKIMOTO A., 1986: Flower-promoting effects of iron and EDDHA in Lemna paucicostata 151. *Plant Cell Physiol.* 27(7), 1369-1376.
- NISHIUCHI Y., 1974: Control effect of a pesticide to duckweeds. (In Japanese). *Noyaku Kensasho Hokoku* 14, 69-72.
- NISHIUCHI K., 1975: Effects of herbicides on Lemna paucicostata. (In Japanese). *Noyaku Tsushin* 93, 38-39.
- NISSSEN P. and BENSON A.A., 1982: Arsenic metabolism in freshwater and terrestrial plants. *Physiol. Plant.* 54(4), 446-450.
- NOIRFALISE A. and DETHIQUX M., 1977: Synopses des végétations aquatiques d'eau douce en Belgique. *Comm. Centre Ecol. For. Rur. N.S.* 14, 1-25.
- NOVACKY A. and ULLRICH-EBERIUS C.I., 1982: Evidence for dual uptake mechanism in the plasmalemma from H⁺/phosphate cotransport in Lemna gibba. *Plant Physiol.* 69(4 suppl.), 93.
- NOVACKY A. and ULLRICH-EBERIUS C.I., 1983: Proton/solute cotransport in higher plants case history: Lemna gibba Gl. *Curr. Top. Plant Biochem. Physiol., Proc. Inaug. Plant Biochem. Physiol. Symp.* 1, 136-144.
- NOVACKY A., FISCHER E., ULLRICH-EBERIUS C.I., LUETTGE U. and ULLRICH W. R., 1978a: Membrane potential changes during transport of glycine as a neutral amino acid and nitrate in Lemna gibba G3. *FEBS Lett.* 88 (2), 264-267.
- NOVACKY A., ULLRICH-EBERIUS C.I. and LUETTGE U., 1978b: Membrane potential changes during transport of hexoses in Lemna gibba Gl. *Planta* 138, 263-270.
- NOVACKY A., ULLRICH-EBERIUS C.L. and LUETTGE U., 1980: pH and membrane potential changes during glucose uptake in Lemna gibba Gl and their response to light. *Planta* 149, 321-326.
- NOWINSKA M., 1972: Edible duckweed. (In Polish). *Wszelchiat* 5, 123-124.
- NUSS R.F. and LOEWUS F.A., 1978: Further studies on oxalic acid biosynthesis in oxalate-accumulating plants. *Plant Physiol.* 61, 590-592.
- NYBERGH T., 1933: A new locality of Lemna gibba. *Mem. Soc. Faun. Flor. Fenn.* 8, 196-197.
- OBERDORFER E., 1960: Pflanzensoziologische Studien in Chile. Ein Vergleich mit Europa. *Flora et Vegetatio Mundi* 2, 208 pp.
- OBERDORFER E., 1970: Pflanzensoziologische Exkursionsflora für Süddeutschland. (3rd ed.). Fischer, Stuttgart. 987 pp.
- OBERDORFER E., 1977: Süddeutsche Pflanzengesellschaften. *Pflanzensoziologie* X. Fischer, Stuttgart. 1, 67-77.
- OBERMEYER-HAUVE A.A., 1966: A note on two rarely seen minute flowering plants, *Wolffia denticulata* and *W. Welwitschii* (Lemnaceae). *South Afr. J. Sci.* 62, 277-278.
- O'BRIEN M.C., 1985: The effect of herbicides on cell membrane permeability in Lemna minor L.; the influence of certain ions and hormones on herbicide activity. *Diss. Abstr. Int.* C 46(3), 656.
- O'BRIEN M.C. and PRENDEVILLE G.N., 1978: A rapid sensitive bioassay for determination of paraquat and diquat in water. *Weed Res.* 18, 301-303.
- O'BRIEN M.C. and PRENDEVILLE G.N., 1979: Effect of herbicides on cell membrane permeability in Lemna minor. *Weed Res.* 19, 331-334.

- O'BRIEN W.J., 1981: Use of aquatic macrophytes for wastewater treatment. *J. Environ. Eng. Div. (Am. Soc. Civ. Eng.)* 197, 681-698.
- ODA Y., 1962: Effect of light quality on flowering of *Lemna perpusilla* 6746. *Plant Cell Physiol.* 3, 415-417.
- ODA Y., 1969: The action of skeleton photoperiods on flowering in *Lemna perpusilla*. *Plant Cell Physiol.* 10(2), 399-409.
- ODEI M.A., 1973: Observations on some weeds of malacological importance in the Volta Lake. *Bull. Inst. Fond. Afrique Noire* A 35(1), 57-66.
- OREGREN E. and ORQUIST G., 1984a: Photoinhibition of photosynthesis in *Lemna gibba* as induced by the interaction between light and temperature. II. Photosynthetic electron transport. *Physiol. Plant.* 62(2), 187-192.
- OREGREN E. and ORQUIST G., 1984b: Photoinhibition of photosynthesis in *Lemna gibba* as induced by the interaction between light and temperature. III. Chlorophyll fluorescence at 77 K. *Physiol. Plant.* 62(2), 193-200.
- OREGREN E., ORQUIST G. and HALLGREN J.E., 1984: Photoinhibition of photosynthesis in *Lemna gibba* as induced by the interaction between light and temperature. I. Photosynthesis in vivo. *Physiol. Plant.* 62(2), 181-186.
- ORZENCIK S., KARATUEPENKCI M. and GUERESCI L.U., 1982: The effects of some pollutants on *Lemna gibba*. (In Turkish). *Oezel Sayi* 1, 461-467.
- OFFORD H.R., 1946: Rapid estimation of the phytocidal action of chemicals. *Science* 103, 474-476.
- OGAWA H., 1977: Modelling of the Heron pond ecosystem. *WRC Res. Rep.* 111. Univ. Water Resour. Cent. 132, 218-222.
- OGUNYEMI E.O., PITTMER F. and HOFMANN-OSTENHOF O., 1978: Studies on the biosynthesis of cyclitols XXXVI. Purification of myo-inositol-1-phosphate synthase of the duckweed, *Lemna gibba*, to homogeneity by affinity chromatography on NAD sepharose: Molecular and catalytic properties of the enzyme. *Hoppe-Seyler's Z. Physiol. Chem.* 359, 613-616.
- OHATA Y. and TACHIBANA Y., 1978: Effect of oxalate on the growth of *Lemna* and metabolic utilization of oxalate by *Lemna*. (In Japan.). *Nippon Dojo-Hiryogaku Zasshi*. 49, 16-20.
- OHATANI T. and ISHIGURI Y., 1979: Inhibitory action of blue and far-red light in the flowering of *Lemna paucicostata*. *Physiol. Plant.* 47, 255-259.
- OHATANI T. and KUMAGAI T., 1980a: Spectral sensitivity of the flowering response in green and etiolated *Lemna paucicostata* T-101. *Plant Cell Physiol.* 21, 1335-1338.
- OHATANI T. and KUMAGAI T., 1980b: Action spectra for the light inhibition of flowering and its reversal in *Lemna paucicostata* T-101. *Planta* 149, 332-335.
- OHATANI T. and KUMAGAI T., 1981: Phytochrome mediated effects of near UV radiation in the induction of flowering in etiolated *Lemna paucicostata* T-101, a short day-plant. *Planta* 153(6), 543-546.
- OHWY J., 1965: Flora of Japan. *Lemnaceae*. Smithsonian Inst., Washington D.C., 264-265.
- OKTYABR'SKAYA T.A. and SHISHKOV I.G., 1972: Experience in using herbicide simazine for controlling plants in water bodies. (In Russian). *Med. Parazit. Parazit. Bolezni*. 41(1), 64-66.
- OKUDA S., 1978: Pflanzensozioologische Untersuchungen über die Auenvegetation der Kanto-Ebene. *Bull. Inst. Environ. Sci. Tech. Yokohama Nat. Univ.* 4, 43-112.
- OLACZEK R., 1959: New locality of *Wolffia arrhiza* (L.) Wimm. in Mazowsze. (In Polish). *Zesz. Nauk. U.I. Ser.* 2, 5, 89-90.

- OLACZEK R. and KRZYWANSKI D., 1970: *Wolffia arrhiza* and the *Wolffia* arrhiza in Poland. (In Polish). *Zesz. Nauk. U.I. Ser.* 2, 36, 39-51.
- OLSEN C., 1930: On the influence of humus substances on the growth of green plants in water culture. *C.R. Trav. Lab. Carlsberg* 18, 1-16.
- OLSEN C., 1934a: The absorption of manganese by plants. *C.R. Trav. Lab. Carlsberg* 20, 1-34.
- OLSEN C., 1934b: Ueber die Manganaufnahme der Pflanzen. *Biochem. Z.* 269, 329-348.
- OLSEN C., 1936: Absorption of manganese by plants. II. Toxicity of manganese to various plant species. *C.R. Trav. Lab. Carlsberg* 21, 139-145.
- OLSEN C., 1950: Aquatic plants and hydrospheric factors. I. Aquatic plants in SW-Jutland. II. The hydrospheric types. *Svensk Bot. Tidskr.* 44, 1-34 and 332-374.
- ONO H., 1952a: The effect of growth substances and some physiological factors on the growth of roots of *Lemnaceae* plants. *Sieboldia (Fukuoka)* 1, 39-50.
- ONO H., 1952b: The effects of 2,4-dichlorophenoxyacetic acid upon the growth of roots. *Bot. Mag. (Tokyo)* 65, 224-227.
- OOTTA Y., 1965: Effects of growth substances on frond and flower production in *Lemna gibba* G3. *Plant Cell Physiol.* 6, 547-559.
- OOTTA Y., 1966: Light- and dark-growth in long-day duckweed, *Lemna gibba* G3, as affected by kinetin. *Plant Cell Physiol.* 7, 631-641.
- OOTTA Y., 1969: Frond and flower production in *Lemna gibba* G3 in presence of respiratory inhibitors. *Plant Cell Physiol.* 10(3), 621-633.
- OOTTA Y., 1970: Periodical growth response of *Lemna gibba* G3 to light-break. *Plant Cell Physiol.* 11, 417-425.
- OOTTA Y., 1971: Disappearance of rhythmicity in growth response to dark- and light-breaks in *Lemna gibba* G3 due to iron deficiency. *Plant Cell Physiol.* 12, 255-266.
- OOTTA Y., 1972a: A possible mechanism for sugar inhibition of duckweed flowering. *Plant Cell Physiol.* 13, 195-199.
- OOTTA Y., 1972b: The response of *Lemna gibba* G3 to a single long day in the presence of EDTA. *Plant Cell Physiol.* 13, 575-580.
- OOTTA Y., 1973: The length of the induction period vs. the minimum number of long-day cycles needed for floral induction in *Lemna gibba* G3. *Plant Cell Physiol.* 14, 307-317.
- OOTTA Y., 1974: Removal of the sugar inhibition of flowering in *Lemna gibba* G3 by catecholamines. *Plant Cell Physiol.* 15, 63-68.
- OOTTA Y., 1975a: Time measurement in photoperiodic floral induction in duckweeds. *Curr. Adv. Plant Sci.* 6, 665-674.
- OOTTA Y., 1975b: Photoperiodic requirements for flowering of the long-day duckweed, *Lemna gibba* G3. *Plant Cell Physiol.* 16, 885-894.
- OOTTA Y., 1975c: Short-day flowering of *Lemna gibba* G3 induced by salicylic acid. *Plant Cell Physiol.* 16, 1131-1135.
- OOTTA Y., 1976: Time measurement in photoperiodic floral induction in duckweeds. In: SMITH H. (ed.), *Commentaries in plant science*. Pergamon Press, Oxford. 199-208.
- OOTTA Y., 1977a: Removal of photoperiodic light requirements of *Lemna gibba* G3. *Plant Cell Physiol.* 18, 95-105.
- OOTTA Y., 1977b: Replacement by ionophores of the photoperiodic light requirement in *Lemna gibba* G3. *Plant Cell Physiol.* 18, 1363-1367.
- OOTTA Y., 1981: Bimodal floral response of *Lemna gibba* G3 to night interruption: photoperiodic time measurement. *Plant Cell Physiol.* 22(1), 99-113.
- OOTTA Y., 1983a: Floral inhibition in *Lemna paucicostata* 6746 due to night interruption. *Plant Cell Physiol.* 24(3), 327-332.

- OOTA Y., 1983b: Physiological structure of the critical photoperiod of Lemna paucicostata 6746. *Plant Cell Physiol.* 24(8), 1503-1510.
- OOTA Y., 1984: Physiological function of night interruption in Lemna paucicostata 6746: action of light as a phase on the photoperiodic clock. *Plant Cell Physiol.* 25(2), 323-331.
- OOTA Y., 1985: Measurement of the critical nyctiperiod by Lemna paucicostata 6746 grown in continuous light. *Plant Cell Physiol.* 26, 923-929.
- OOTA Y. and HOSHINO T., 1974: Diurnal change in temperature sensitivity of Lemna gibba G3 induced by acetylcholine in continuous light. *Plant Cell Physiol.* 15, 1063-1072.
- OOTA Y. and HOSHINO T., 1979: Spectral dependence of the critical photoperiod in the long-day duckweed Lemna gibba G3. *Plant Cell Physiol.* 20, 1531-1536.
- OOTA Y. and KONDO T., 1974: Removal by cyclic AMP of the inhibition of duckweed flowering due to ammonium and water treatment. *Plant Cell Physiol.* 15, 403-411.
- OOTA Y. and NAKASHIMA H., 1978: Photoperiodic flowering in Lemna gibba G3: time measurement. *Bot.Mag.(Tokyo), Spec. Issue 1*, 177-198.
- OOTA Y. and TSUDZUKI T., 1971: Resemblance of growth substances to metal chelators with respect to their actions on duckweed growth. *Plant Cell Physiol.* 12, 619-631.
- OOTA Y. and TSUDZUKI T., 1979: Evidence against involvement of circadian floral rhythm in the critical daylength measurement in Lemna gibba G3. *Plant Cell Physiol.* 20, 725-732.
- OPUSZYSKI K., 1972: Use of phytophagous fish to control aquatic plants. *Aquaculture* 5(1), 61-73.
- OREBAMJO T.O., 1973: Regulation of nitrate reductase in Lemna minor. Ph.D. Thesis. Univ. Manchester.
- OREBAMJO T.O. and STEWART G.R., 1974: Some characteristics of nitrate reductase induction in Lemna minor L. *Planta* 117, 1-10.
- OREBAMJO T.O. and STEWART G.R., 1975a: Ammonium repression of nitrate reductase formation in Lemna minor L. *Planta* 122, 27-36.
- OREBAMJO T.O. and STEWART G.R., 1975b: Ammonium inactivation of nitrate reductase in Lemna minor L. *Planta* 122, 37-44.
- ORLANDO J.A. and NEILANDS J.B., 1982: Ferrichrome compounds as a source of iron for higher plants. In: KEHL H. (ed.), *Chem.Biol.Hydroxamic Acids*; (Proc.Int.Symp.). Karger, Basel. 123-129.
- ORNES W.H., 1979: Effects of cadmium (II) on Azolla caroliniana Willd., Salvinia rotundifolia Willd., Spirodela polyrrhiza (L.) Schleid., Ceratophyllum demersum L., and Myriophyllum spicatum L. var. exalbes-cens (Fernald) Jepson grown in 10 or 50% Hoagland solution. Ph.D. Thesis. Iowa State Univ., Ames, IA. 199 pp. *Diss.Abst.Int.* B 39(110), 4712.
- ORNES W.H. and WILDMAN R.B., 1979: Effects of cadmium (II) on aquatic vascular plants. *Trace Subst.Environ.Health* 13, 304-312.
- ORON G. and PORATH D., 1981: Ammonia stripping of fish influent by duckweed in a circulating system. *Proc.Water Reuse Symp.* II, 2, 1638-1647.
- ORON G. et al., 1984 see 1985
- ORON G., WILDSCHUT L.R. and PORATH D., 1985: Wastewater recycling by duckweed for protein production and effluent renovation. *Water Sci. Tech.* 17(4-5), 803-817.
- ORON G., PORATH D. and WILDSCHUT L.R., 1986: Wastewater treatment and renovation by different duckweed species. *J.Environ.Eng.* 112(2), 247-263.

- ORON G., PORATH D. and JANSEN H., 1987a: Performance of the duckweed species Lemna gibba on municipal wastewater for effluent renovation and protein production. *Biotechn.Bioengine.* 29(2), 258-268.
- ORON G., DE-VECT A. and PORATH D., 1987b: The role of the operation regime in wastewater treatment with duckweed. *Water Sci.Technol.* 19(1-2), 97-106.
- OSTROW J.P., 1968: The effect of photoperiod on vegetative reproduction and senescence in Spirodela oligorrhiza. (Lemnaceae). M.S.Thesis. Univ. Miami.
- OSTROW J.P. and DIJMAN M.J., 1969: Photo-related aging in Spirodela oligorrhiza (Lemnaceae). *Quart.J.Fla.Acad.Sci.* 32(2), 119-126.
- OSTROW-SCHWEBEL J.P., 1973: Hormonal control of growth and development in Lemna minor with special emphasis on the role of abscisic acid (ABA). Ph.D.Thesis. College of A and M Univ., Texas. 143 pp. *Diss. Abstr.* B 34, 997.
- OSTROW-SCHWEBEL J.P., 1979: An abscisic acid-induced separation layer in Lemna minor (L.). *Fla.Sci.* 42, 172-176.
- OTAKI M., 1974: Distribution and description of Lemnaceae. (In Japan.). *Heredity* 28(8), 50-54.
- OTAKI S., 1978: Illustrated Japanese water plants. (In Japan.). Hokuryukan, Tokyo.
- OZIMEK T., 1983: The role of duckweeds in cycling of heavy metals in ponds supplied with post-sewage water. *Proc.Int.Symp. Aquatic Macrophytes*, September 18-23, Nijmegen, Netherlands, 172-176.
- OZIMEK T., 1985: Heavy metal content in macrophyte from ponds supplied with post-sewage water. In: SALANKI J. (ed.), *Symp.Biol.Hung.*, Heavy metals in water organisms. 29, 41-50.
- PALAVAN N., GOREN R. and GALSTON A.W., 1984: Effects of some growth regulators on polyamine biosynthetic enzymes in etiolated pea seedlings. *Plant Cell Physiol.* 25, 541-546.
- PAMPANINI R. and PROVASI T., 1921: La fioritura della Lemna minor L. nell'Orto Botanico di Firenze. *Bull.Soc.Bot.Ital.* 1921, 53.
- PAN S.-M. and CHEN S.-C., 1976: Scanning electron microscopy of chloroplasts from duckweed cells. *Taiwania* 21, 248-250.
- PAN S.M. and CHEN S.-C., 1979: The morphology of Wolffia arrhiza - a scanning electron microscopy study. *Bot.Bull.Acad.Sin.* 20(2), 89-96.
- PAN Y.-T., 1974: Characterization of a particulate UDP-D-apiose acceptor D-apiosyltransferase from Lemna minor. *Fed.Proc.* 33(5 part 2), 1445.
- PAN Y.-T. and KINDEL P.K., 1977: Characterization of particulate D-apiosyl- and D-xylosyltransferases from Lemna minor. *Arch.Biochem.Biophys.* 183, 131-138.
- PANCHO J.V., 1976: Philippine aquatic weeds. *Kalikaan* 5(1), 37-91.
- PANDIT A.K., KAUL V. and FOTEDAR D., 1978: A preliminary study of duckweed ecology and its natural control. *Int.J.Ecol.Environ.Sci.* 4(1-3), 107-115.
- PANTICKER K.N., SRINIVASAN R., VISWAM K. and RAJAGOPALAN P.K., 1985: Larvovorous potential of some cypriniformes fishes. *Ind.J.Med.Res.* 82(6), 517-520.
- PANKEY R.D., DRAUDT H.N. and DESROSIER N.W., 1965: Characterization of the starch of Spirodela polyrrhiza. *J.Food Sci.* 30, 627-631.
- PANKIN W., 1945: Zur Oekologie und Soziologie der Lemna-Standorte. *Arch.Hydrobiol.* 41, 225-232.
- PARDY R.L. and GLIDER W.V., 1984: The photic environment of the symbiotic hydroid, Hydra viridis. *Am.Midl.Nat.* 112(1), 196-197.
- PARK K.H., HAHN K.-W. and KANG B.G., 1987: Senescence in Lemna gibba G3:

- evidence against stomatal involvement in the regulation of frond senescence. 14th Int. Bot. Congr. Berlin, Abstr., 111.
- PARK W.C. and YATAZAWA M., 1979: Nitrogen fixation by Lemna-blue green algae association. (In Korean). J. Korean Soc. Soil Sci. Fert. 12(1), 47-49.
- PARKS C.R., SANDHU S.S. and MONTGOMERY K.R., 1972: Floral pigmentation studies in the genus *Gossypium*. IV. Effects of different growing environments on flavonoid pigmentation. Am. J. Bot. 59(2), 158-164.
- PASSARGE H., 1957: Ueber Wasserpflanzen- und Kleinhörschichtgesellschaften des Oberspreewaldes. Abh. Ber. Naturkd. Mus. G6rlitz, Leipzig, 35, 143-152.
- PASSARGE H., 1978: Zur Syntaxonomie mitteleuropäischer Lemnetae-Gesellschaften. Folia Geobot. Phytotax. 13, 1-17.
- PATIENCE R.L., STERRY P.R. and THOMAS J.D., 1983: Changes in chemical composition of a decomposing aquatic macrophyte, Lemna paucicostata. J. Chem. Ecol. 9(7), 889-911.
- PATNAIK S., 1976: Control of some aquatic vegetation in fish ponds at Cuttack. J. Ind. Fish. Soc. India 8, 221-226.
- PAUTOVA V.N. and GALIMULIN M.G., 1980: New findings of the rare higher aquatic plants in the East Siberia. (In Russian). Bot. Zh. 65, 1020-1022.
- PAWLOWICZ P. and SIEMINSKI A., 1987: Biotransformations. Part 20. Enantioselective hydrolysis of esters and the oxidation of aromatic-aliphatic alcohols obtained therefrom by *Spirodela oligorrhiza*. Phytocchemistry 26(4), 1001-1004.
- PEARSALL W.H., 1921: A suggestion as to factors influencing the distribution of free floating vegetation. J. Ecol. 9, 241-253.
- PECHENYUK E.V., 1982: Rare aquatic plants in the Khoperski State Reserve. (In Russian). Bot. Zh. 67(5), 647-651.
- PECHENYUK E.V., 1984: Lemna gibba (Lemnaceae) in Khoperski State Reserve. (In Russian). Bot. Zh. 69, 1101.
- PECHENYUK E.V., 1985: Flowering of Lemnaceae in the Khoperski State Reserve. (In Russian). Bot. Zh. 70(8), 1066-1070.
- PEKHOVA see PETKOVA
- PEDROTTI F., 1979: L'associazione *Ricciocarpetum natantis* (Segal 1965) Tx. 1972 dans le marais de Colfiorito (Italie Centrale). Doc. Phytosoc. N.S. 4, 795-802.
- PEKIC S., 1977: Endogenous hormones in Lemna perpusilla. Biol. Vestn. 25(2), 187.
- PEKIC S. and NESKOVIC H., 1982: Influence of phytochrome on the content of endogenous hormones in Lemna aequinoctialis during the long night period. Bull. Inst. Jard. Bot. Univ. Beograd (13), 15(1-3), 21-28.
- PENFOUND W.T., 1956: Primary production of vascular aquatic plants. Limnol. Oceanogr. 1(2), 92-101.
- PERLMAN D. and SEMAR J.B., 1967: Inhibition by purines of the production of fronds in Lemna perpusilla. Nature 215, 760-761.
- PERRING F.H. and WALTERS S.M., 1962: Atlas of the British Flora. Nelson, London/Edinburgh, 343-344.
- PERRY T.O., 1963: Differences in protein constituents in dormant and vegetative plant tissue (*Spirodela polyrrhiza* and *Pinus thunbergii*). AIBS Bull. 13(5), 73-75.
- PERRY T.O., 1968: Dormancy, turion formation and germination by different clones of *Spirodela polyrrhiza*. Plant Physiol. 43, 1866-1869.
- PERRY T.O. and BYRNE O.R., 1969: Turion induction in *Spirodela polyrrhiza* by abscisic acid. Plant Physiol. 44, 784-785.
- PESTER W., 1979: Biological determination of photosynthetic inhibitors

- in soils and water and application in bioassays to herbicide investigations. Z. Naturforsch. 34c, 964-964.
- PETER G. and THORBER J.P., 1984: Use of Lemna gibba to study light stress effects on higher plant photosynthetic apparatus. J. Cell. Biochem. (suppl. 8 part B), 232.
- PETKOVA L.M. and LUBJANOV I.P., 1969: Concentration of some microelements in macrophytes in the basins of the steppe zones of the Ukraine. (In Ukrainian). Ukr. Bot. Zh. 26, 90-96.
- PETKOVA L.R., 1980: Review of the research in the anatomy of reproductive organs of angiosperms in USSR for the last ten years. (In Russian). Bot. Zh. 65(12), 1681-1695.
- PEVERLY J.H., 1985: Element accumulation and release by macrophytes in a wetland stream. J. Environ. Qual. 14, 137-143.
- PFFERER W., 1986: Ueber die Aufnahme von Anilinfarben in lebenden Zellen. Unters. Bot. Inst. Tübingen 2, 179-329.
- PHILIPPI G., 1969: Laichkraut- und Wasserlinsengesellschaften des Oberrheingebietes zwischen Strassburg und Mannheim. Veröff. Landesst. Natursch. Landesst. pfl. Baden-Württemberg 37, 102-172.
- PHILIPPI G., 1971: Beiträge zur Flora der nordbadischen Rheinebene und der angrenzenden Gebiete. Beitr. Naturk. Forsch. Südw. Deutschl. 30, 9-47.
- PHILIPPI G., 1978: Veränderungen der Wasser- und Uferflora im Badischen Oberrheingebiet. Beitr. Veröff. Natursch. Landesst. pfl. Baden-Württemberg 11, 99-134.
- PHILIPPI R.A., 1957: Plantarum novarum Chilensium, centuria quarta. Linnaea 29, 1-47.
- PHILIPPI R.A., 1964: Plantarum novarum Chilensium. Linnaea 33, 239.
- PICCOLI F. and GERDOL R., 1981: Rice-field weed communities in Ferrara Province (Northern Italy). Aquat. Bot. 10, 317-328.
- PICKEN J.M. and MENDICINO J., 1967: The biosynthesis of D-apiose in Lemna minor. J. Biol. Chem. 242(7), 1629-1634.
- PIETERSE A.H., 1972: Studies on flowering and turion formation in Lemnaceae. Thesis. Univ. of Kentucky, Lexington. 124 pp.
- PIETERSE A.H., 1974a: On the morphology of Lemna minor in South Finland. Ann. Bot. Fenn. 11, 271-274.
- PIETERSE A.H., 1974b: Gibberellin-EDDHA interaction in flowering and gibbosity of Lemna gibba G3. Plant Cell Physiol. 15, 1125-1127.
- PIETERSE A.H., 1975a: Physiological, morphological and anatomical aspects of gibbosity in Lemna gibba. Aquat. Bot. 1, 333-344.
- PIETERSE A.H., 1975b: Induction of flowering in Lemna gibba G3 under short-day conditions by a combined treatment of EDDHA and BA. Proc. 12th Int. Bot. Congress, Leningrad, 308.
- PIETERSE A.H., 1976: Specific interactions in the physiology of flowering and gibbosity of Lemna gibba G3. Plant Cell Physiol. 17, 713-720.
- PIETERSE A.H., 1977: Control of tropical aquatic weeds. Bull. Dept. Agric. Res. Roy. Trop. Inst. 300, 20 pp.
- PIETERSE A.H., 1978a: Experimental control of flowering in Pistia stratiotes L. Plant Cell Physiol. 19, 1091-1093.
- PIETERSE A.H., 1978b: Interaction of naphthol with EDDHA/salicylic acid in flowering and gibbosity of Lemna gibba G3. Plant Cell Physiol. 19(7), 1307-1310.
- PIETERSE A.H., 1980: Specific interactions in the physiology of flowering in Lemna gibba G3 and Pistia stratiotes. Symp. Weed Sci. Soc. Toronto, 13 pp.
- PIETERSE A.H., 1981: Specific interactions in the physiology of flowering in Lemna gibba G3 and Pistia stratiotes. Int. Coll. Aquat. Vascular Plants, Brussels, Belgium. Abstr., 44.

- PIETERSE A.H., 1982: A review of chemically induced flowering in *Lemna gibba* G3 and *Pistia stratiotes*. *Aquat. Bot.* 13(1), 21-28.
- PIETERSE A.H. and MUELLER L.J., 1977: Induction of flowering in *Lemna gibba* G3 under short-day conditions. *Plant Cell Physiol.* 18, 45-53.
- PIETERSE A.H. and SABHARWAL P.S., 1969: Physiological studies with *Wolffia* under aseptic conditions. 11th Int. Bot. Congr., Abstr., 170.
- PIETERSE A.H., BHALLA P.R. and SABHARWAL P.S., 1970a: Induction of flowering in *Lemna gibba* G3 by Fe-EDHA. *Plant Cell Physiol.* 11, 675-676.
- PIETERSE A.H., BHALLA P.R. and SABHARWAL P.S., 1970b: Investigations on the effects of metal ions and chelating agents on growth and flowering of *Lemna gibba* G3. *Plant Cell Physiol.* 11, 879-889.
- PIETERSE A.H., BHALLA P.R. and SABHARWAL P.S., 1970c: Control of gibbosity in *Lemna gibba* G3 by ethylenediamine-di-o-hydroxyphenylacetic acid (EDDHA). *Acta Bot. Neerl.* 19(4), 521-524.
- PIETERSE A.H., BHALLA P.R. and SABHARWAL P.S., 1970d: Chemical induction of turions in *Wolffia* floridana (J.D. Smith) Thompson. *Acta Bot. Neerl.* 19(6), 901-905.
- PIETERSE A.H., BHALLA P.R. and SABHARWAL P.S., 1971a: Endogenous gibberellins in floating plants and turions of *Wolffia* floridana. *Physiol. Plant.* 24, 512-516.
- PIETERSE A.H., BHALLA P.R. and SABHARWAL P.S., 1971b: In vitro studies on turions of *Wolffia* floridana. *Am. J. Bot.* 58(5 part 2), 455.
- PIETSCH W., 1972: Ausgewählte Beispiele für Indikatoreigenschaften höherer Wasserpflanzen. *Arch. Naturforsch. u. Landschaftsforsch.* 12, 121-151.
- PIGNATTI S., 1957: La vegetazione delle risaie pavesi (studio fitosociologico). *Arch. Bot. Biogeogr. Ital.* 33, 4a Ser., 2, 1-68.
- PIISPANEN R. and LAEHDESMAKI P., 1983: Effect of vanadium on some water plants. *Geol. Foeren. Stockholm Forsh.* 105(1), 23-27.
- PIT P.E., 1979: Survey of the ecology of submerged aquatic macrophytes in central Canada. *Aquat. Bot.* 7, 339-357.
- PIT P.E. and SIMMONS K., 1986: Aquatic angiosperms at unusual depths in Shoal Lake. *Can. Field-Nat.* 100(3), 354-358.
- PIRSON A. and GOELLNER E., 1953: Zellphysiologische Untersuchungen an der Lemna-Wurzel bei verminderter Nitrat- und Phosphatversorgung. *Z. Bot.* 41, 147-176.
- PIRSON A. and GOELLNER E., 1954: Beobachtungen zur Entwicklungsphysiologie der Lemna minor L. *Flora* 140, 485-498.
- PIRSON A. and SCHAEFER G., 1957: Osmotischer Wasserentzug und Plasmolyse mit Polyäthylenoxyd. *Protoplasma* 48, 215-220.
- PIRSON A. and SEIDEL F., 1950: Zell- und stoffwechselphysiologische Untersuchungen an der Wurzel von Lemna minor L. unter besonderer Berücksichtigung von Kalium- und Kalziummangel. *Planta* 38, 431-473.
- PITTENDRIGH C.S., 1966: The circadian oscillation in *Drosophila pseudoobscura* pupae: A model for the photoperiodic clock. *Z. Pflanzenphysiol.* 54, 275-307.
- PIZZOLATO T.D. and FRICK H., 1979: Pyrimidine metabolism in Lemna minor. II. Specific inhibition of plastid replication in a higher plant by cytidine deoxyriboside. *Plant Physiol.* 63, 979-983.
- PODBIELKOWSKI Z., 1967: Rather rare species of vascular plants of the Warsaw Voivodship, Poland. *Fragmenta Florist. Geobot.* 13(3) 323-325.
- POIJAERVI L.A.P., 1928: Ueber die Basenpermeabilität pflanzlicher Zellen. *Acta Bot. Fenn.* 4, 1-102.
- POKORNY J. and REJMANKOVA E., 1983: Oxygen regime in a fish pond with duckweeds (Lemnaceae) and Ceratophyllum. *Aquat. Bot.* 17(2), 125-137.
- POLAR E. and KUECKECCZAR R., 1986: Influence of some metal chelators and light regimes on bioaccumulation and toxicity of Cd in duckweed (*Lemna gibba*). *Physiol. Plant.* 66, 87-93.

- POPPIUS B., 1901: Lemna trisulca. *Med. Soc. Fauna Flora Fenn.* 27, 45.
- PORATH D., 1973: Cultured duckweeds (Lemnaceae) as a model for studying effects of environmental conditions on the vegetative phase of higher plants. *Ph.D. Thesis. Univ. Tel Aviv.*
- PORATH D., 1979: Pathways of plastide differentiation in *Spirodela oligorrhiza*. *New Phytol.* 82, 733-737.
- PORATH D. and AGANI M., 1986: An alternative approach to study the avoidance of duckweed (*Lemna gibba* L. - Lemna minor L. complex) from arid zones. *Arch. Hydrobiol.* 108(1), 45-53.
- PORATH D. and BEN-SHAUL Y., 1971: Structural and physiological changes during heat bleaching in *Spirodela oligorrhiza*. *Isr. J. Bot.* 20(3), 157-168.
- PORATH D. and BEN-SHAUL Y., 1973: Growth, greening and phytochrome in etiolated *Spirodela* (Lemnaceae). *Plant Physiol.* 51(3), 474-477.
- PORATH D. and KOTON A., 1977: Enhancement of protein production in fish ponds with duckweed (Lemnaceae). *Isr. J. Bot.* 26(1), 51.
- PORATH D. and POLLOCK J., 1982: Ammonia stripping by duckweed Lemna gibba and its feasibility in circulating aquaculture. *Aquat. Bot.* 13(2), 125-132.
- PORATH D., HEPHER B. and KOTON A., 1979: Duckweed as an aquatic crop: Evaluation of clones for aquaculture. *Aquat. Bot.* 7, 273-278.
- PORATH D., EFRAI Y. and ARZEE T., 1980: Morphological patterns and heterogeneity in populations of duckweeds. *Aquat. Bot.* 9(2), 159-168.
- PORATH D., ORON G. and GRANTH G., 1986: Duckweed as an aquatic crop: Edible protein recovery, production and utilization. *Proc. 5th Int. Symp. Agric. Wastes, Chicago, 1985. Am. Soc. Agric. Engineers, St. Joseph, Michagan.* 680-687.
- POSNER H.B., 1961: Effects of x-irradiation on "handedness" of frond production in Lemna perpusilla 6746. *Plant Physiol.* 36(suppl.), 22.
- POSNER H.B., 1962a: Permanent and temporary effects of X-rays on the reproduction and aging of Lemna perpusilla. *Ph.D. Thesis. Yale Univ.*
- POSNER H.B., 1962b: Characteristics of X-ray-induced aberrants of Lemna perpusilla 6746. *Plant Cell Physiol.* 3, 275-284.
- POSNER H.B., 1966: Some effects of copper and EDTA on flowering in Lemna perpusilla, aberrant strain 1073. *Proc. Annu. Meetings, Am. Soc. Plant Physiol.* 29.
- POSNER H.B., 1967a: Aquatic vascular plants. In: WILT F. and WESSELS N. (eds.), *Methods in developmental biology*. Crowell, New York. 301-317.
- POSNER H.B., 1967b: Inhibitory effect of sucrose on flowering in Lemna perpusilla 6746 and mutant strain 1073. *Plant Cell Physiol.* 8, 535-539.
- POSNER H.B., 1969a: Inhibitory effect of carbohydrate on flowering in Lemna perpusilla. I. Interaction of sucrose with calcium and phosphate ions. *Plant Physiol.* 44, 562-566.
- POSNER H.B., 1969b: Further studies on the inhibitory effect of carbohydrates on flowering in Lemna perpusilla 6746. *Plant Physiol.* 44 (suppl.), 19-20.
- POSNER H.B., 1970: Inhibitory effect of carbohydrate on flowering in Lemna perpusilla. 2. Reversal by glycine and L-aspartate: Correlation with reduced levels of beta-carotene and chlorophyll. *Plant Physiol.* 45, 687-690.
- POSNER H.B., 1971: Inhibitory effect of carbohydrate on flowering in Lemna perpusilla. 3. Effects of respiratory intermediates, amino acids, and CO₂. Glucose 6-phosphate dehydrogenase activity. *Plant Physiol.* 48, 361-365.
- POSNER H.B., 1973a: Reversal of sucrose inhibition of Lemna flowering by adenine derivatives. *Plant Cell Physiol.* 14, 1199-1200.

- POSNER H.B., 1973b: Lack of flowering responses to DCMU in a mutant of *Lemma perpusilla* 6746. *Plant Cell Physiol.* 14, 1031-1033.
- POSNER H.B. and HILLMAN W.S., 1960: Effects of X irradiation on *Lemma perpusilla*. *Am.J.Bot.* 47, 506-511.
- POSNER H.B. and HILLMAN W.S., 1962: Aseptic production, collection and germination of seeds of *Lemma perpusilla* 6746. *Physiol.Plant.* 15, 700-708.
- POSNER H.B. and HILLMAN W.S., 1970: Sucrose ammonium interaction on the flowering *Lemma perpusilla*. *Plant Physiol.* 46(suppl.), 25.
- POSNER H.B. and ROSNER A., 1975: Effects of chloramphenicol on RNA synthesis in *Spirodela chloroplasts*. *Plant Cell Physiol.* 16, 361-365.
- POSNER H.B., GRESSEL J. and ROSNER A., 1974: Direct evidence for the lack of methylation of two pulse labeled plant RNAs. *Plant Cell Physiol.* 15, 807-811.
- POSNER H.B., POSNER R.S. and GOWER R.A., 1977: Effects of DCMU on long-day flowering of *Lemma perpusilla* 6746 and photosynthetic mutant strain 1073. *Plant Cell Physiol.* 18, 1301-1307.
- POSNO M., TORENVLIET D.J., LUSTIG H., VAN NOORT M. and GROOT G.S.P., 1985: Localization of three chloroplast ribosomal protein genes at the left junction of the large single copy region and the inverted repeat of *Spirodela oligorhiza* chloroplast DNA. *Curr.Genet.* 9(3), 211-219.
- POSNO M., VAN VLIET A. and GROOT G.S.P., 1986a: Localization of chloroplast ribosomal protein genes on *Spirodela oligorhiza* chloroplast DNA. *Curr.Genet.* 10(12), 923-930.
- POSNO M., VAN VLIET A. and GROOT G.S.P., 1986b: The gene for *Spirodela oligorhiza* chloroplast ribosomal protein homologous to *E. coli* ribosomal protein L16 is split by a large intron near its 5' end: structure and expression. *Nucleic Acids Res.* 14(8), 3181-3195.
- POSNO M., VERWEIJ W.R., DEKKER I.C., DE WARD P.M. and GROOT G.S.P., 1986c: The genes encoding chloroplast ribosomal proteins S-7 and S-12 are located in the inverted repeat of *Spirodela oligorhiza* chloroplast DNA. *Curr.Genet.* 11(1), 25-34.
- POST G.E., 1933: *Flora of Syria, Palestine and Sinai*. American Press, Beirut. 547-549.
- POTT R., 1980: Die Wasser- und Sumpfvegetation eutropher Gewässer in der westfälischen Bucht - Pflanzensoziologische und hydrochemische Untersuchungen. *Abh.Landesmus.Naturk. Münster* 42(2), 156 pp.
- POTT R., 1981: Ökologie und Indikatorwert von Wasserpflanzengesellschaften. *Mitt.Landesanst.Ökologie,Landschaftsentw.,Forstplanung Nordrhein-Westf., Sonderheft Landestagungen* 1980, 57-64.
- POTT R. and WITTIG R., 1985: Die Lemnetae - Gesellschaften nieder rheinischer Gewässer und deren Veränderungen in den letzten Jahren. *Tuexenia* 5, 21-30.
- POURRIOT R., 1972: Etude hydrobiologique de deux petits étangs de prairies: Observations sur la distribution de la température et du plancton et l'influence d'un couvert végétal à *Lemma minor*. *Ann.Hydrobiol.* 3(1), 33-46.
- PRASAD R., 1981: Management of aquatic weeds with chemicals impact of nonylphenol on macrophytes. *Proc.Int.Bot.Congr.* 13, 179.
- PRASAD R., 1984: Impact of glyphosate on macrophytes. *Plant Physiol.* 75 (1 suppl.), 139.
- PREVITERA L. and MONACO P., 1983: Fatty acid composition in *Lemma minor* - Characterization of a novel hydroxy C₁₆ acid. *Phytochem.* 22(6), 1445-1446.
- PREVITERA L. and MONACO P., 1984: A linear diterpene diol from *Lemma minor*. *Phytochem.* 23(1), 194-195.

- PRICE D.N. and WAIN R.L., 1976: Studies on plant growth-regulating substances. XII. Structure-activity relationships and metabolism of a group of nitrophenols capable of inhibiting chloroplast development. *Ann.Appl.Biol.* 83, 115-124.
- PRIDHAM J.B., 1964: The phenol glucosylation reaction in the plant kingdom. *Phytochem.* 3, 493-497.
- PRINTZ H., 1921: The vegetation of the Siberian-Mongolian frontiers. *Trondhejm.* 175 pp.
- PRISHCHEPOV G.P., 1974: Selectivity in the feeding of grass carp and the nutritive value to the fish of some plant species of Belorussian lakes. (In Russian). *Trudy Belorusskogo Nauchno-Issledovatel'skogo Inst. Rybnogo Khozayaistva* 10, 196-200.
- PRISZTER S., 1961: *Wolffia*, an adventitious plant. (In Hung.). *Bot.Közlem.* 49, 118-121.
- PROCTOR G.R., 1982: More additions to the flora of Jamaica. *J.Arnold Arboretum* 63, 207-210.
- PROSVIROVA L.V. and LEVANTIDOV L.Y., 1979: *Manganophyllia* in aquatic plants. (In Russian). *Khim.Biokhim.Okislennie Sist., Soderzh.d-Elem.* 7-10.
- PRYCE R.J., 1973a: Decomposition of aqueous solutions of gibberellic acid on autoclaving (*Lemma perpusilla*). *Phytochem.* 12(3), 507-514.
- PRYCE R.J., 1973b: Allogibberic acid: An inhibitor of flowering in *Lemma perpusilla*. *Phytochem.* 12, 1745-1754.
- PRYCE R.J., 1974: Inhibition of flowering by hexahydrofluorene-9-carboxylic acids related to allogibberic acid. *Phytochem.* 13, 2377-2382.
- PRYCE R.J., 1978: From natural to synthetic plant growth regulators. *Monogr.Br.Crop Prot.Counc.* 21, 181-186.
- PURVES W.K., 1961: Dark reactions in the flowering of *Lemma perpusilla* 6746. *Planta* 56, 684-690.
- QIAN K.C., 1982: A preliminary study on biology of *Tipula* aino Alexander. (In Chinese). *Insect Knowl.* 19(5), 9-11.
- QUIROZ F.A., 1981: Papel de algunas hidrofitas en la fertilidad del sistema chinampero. *Biotica* 5(4), 169-179.
- QUIROZ F.A., MIRANDA M.G. and HELGUERAS A.L., 1982: Uso potencial de algas hidrofitas como abono verde en la zona chinampera de Xochimilco. *Biotica* 7(4), 631-633.
- RAABE E.-W., 1970: *Wolffia arrhiza* in Schleswig-Holstein. *Kieler Notizen Pflk.Schl.-Holst.* 5, 13.
- RAABE U., 1984: *Lemma minuscula* Herter auch in Westfalen. *Gött.Florist. Rundbr.* 18(1-2), 42.
- RADENOVIC C., GRBOVIC M., VUCINIC Z. and DAMJANOVIC Z., 1980: A report on excitability phenomena of intact roots registered with microelectrodes. In: SPANSWICK R.M. et al. (eds.), *Plant membrane transport*. Elsevier, North-Holland. Biomed.Press. *Dev.Plant Biol.* 4, 617-618.
- RADNER B.S. and THIMMANN K.V., 1963: The biogenesis of anthocyanins. IX. The effect of ribonuclease on anthocyanin formation in *Spirodela oligorhiza*. *Arch.Biochem.Biophys.* 102, 92-95.
- RAFINESQUE SCHWALTZ C.S., 1808: Prospectus of two intended works on North-American Botany. Medical Repository, New York II/5, 350-356.
- RAGHAVENDRA A.S. and DAS V.S.R., 1976: Distribution of the C₄ dicarboxylic acid pathway of photosynthesis in local monocotyledonous plants and its taxonomic significance. *New Phytol.* 76(2), 301-305.
- RAGY H., PICKEN M. and MENDICINO J., 1973: Purification of a specific

- D-apitol dehydrogenase from a Micrococcus isolated from the surface of germinating parsley seeds. *Biochim. Biophys. Acta* 315, 259-271.
- RAHMAN A. and COX T.I., 1975: Bioassay techniques for the determination of herbicide residues. *Proc. 28th NZ Weed and Pest Conf. Soil Field Res. Org., Ruakura Agric. Res. Centre, Hamilton, New Zealand*, 96-100.
- RAKHIMOV A. and ABDIEV M., 1972: On the chemical composition of Lemna grown in various nutrient media. (In Russian). *Kult. Vod. Vyssh. Vodn. Rast. Uzbekistane* 1972, 112-115.
- RAKHIMOV A. and RAKHIMOVA S., 1975: Study of the biochemical characteristics of small duckweed (content of carotene and ascorbic acid in a small duckweed culture biomass). (In Russian). *Vod. Griby Sredn. Azii* 208-213.
- RAKHIMOV A. and RAKHIMOVA S., 1982: Effect of sewage from the swinery complex "Sergeli" and poultry farm "Uzbekistan" on the productivity and chemical composition of Lemna minor and Spirodela. (In Russian). *Uzb. Biol. Zh.* 1982(3), 18-20.
- RAKHIMOV A., 1983: On the possibility of growing aquatic macrophytes on certain sewage waters. (In Russian). *Uzb. Biol. Zh.* 1983(1), 21-24.
- RAKHIMOV A., RAKHIMOVA S. and ISMAILKHODZHAEV B., 1981: Effect of municipal sewage on the accumulation of organic substances, proteins and various fractions of carbohydrates in common duckweed biomass. (In Russian). *Uzb. Biol. Zh.* 1981(6), 14-17.
- RAMAYYA N. and RAJAGOPAL T., 1975: Pseudowolfia hyalina (Del.) Hart. and Plas (Lemnaceae) - A new record for Asia from Hyderabad (India). *Aquat. Bot.* 1, 71-73.
- RAMIREZ C. and BECK S.G., 1981: Makrophytische Vegetation und Flora in Gewässern der Umgebung von La Paz, Bolivien. *Arch. Hydrobiol.* 91(1), 82-100.
- RAMIREZ C., ROMERO M. and RIVEROS M., 1979: Habit, habitat, origin and geographical distribution of Chilean vascular hydrophytes. *Aquat. Bot.* 7(3), 241-253.
- RANE D.A. and TUKEY L.D., 1972: A rapid technique for comparing the effectiveness of various growth regulators. *Hortscience* 7(3), 329.
- RANJEVA R., GRAZIANA A., RANTY B., CAVALIE G. and BOUDET A.M., 1984: Phosphorylation of proteins in plants: a step in the integration of extra- and intracellular stimuli? *Physiol. Vég.* 22(3), 365-376.
- RAO A.S. and VERMA D.M., 1979: Materials towards a monocot flora of Assam, India 5, Lemnaceae. *Bull. Bot. Surv. India* 18, 30-36.
- RAO C.B., 1953: On the distribution of algae in a group of six small ponds. *J. Ecol.* 41, 62-71.
- RAO D.S.K., MANISSERY J.K., DEVARAJ K.V. and MURTHY G.N., 1982: Culture of Lemna minor in cement cisterns using different organic and inorganic manures. *Mysore J. Agric. Sci.* 16(4), 462-468.
- RAO K.V.N., RAO S.S.R., RAO K.N. and SRIMANNARAYANA G., 1980: Effect of some proanthocyanidins and catechins on the growth of Lemna paucicostata Hegelm. *Proc. Indian Acad. Sci. (Plant Sci.)* 89, 73-77.
- RAO R., 1969: Ultrastructure of Spirodela polyrrhiza (L.) Schleiden with specific reference to chloroplast development during turion germination. *Ph.D. Thesis*. NCSU, Raleigh. 121 pp.
- RAO R. 1982 see RAO R., 1969
- RAO S.S.R. and RAO K.V.N., 1984: Effect of certain proanthocyanidins and catechins on the nucleic acid and nitrogen contents of Lemna paucicostata Hegelm. *Proc. Indian Acad. Sci. (Plant Sci.)* 93(1), 1-5.
- RAO S.S.R. and RAO K.V.N., 1985a: Effect of certain proanthocyanidins on the chlorophyll content of Lemna paucicostata Hegelm. *Curr. Sci.* 54(18), 945-947.
- RAO S.S.R. and RAO K.V.N., 1985b: Mangiferin - a phenolic growth inhibitor. *Curr. Sci.* 54(20), 1075-1076.
- RAO S.V.R., 1984: Anaerobic digestion of two aquatic weeds for energy generation. *Resour. Manage. Optim.* 3(4), 343-352.
- RASOL M.K. and RENDIC L., 1977: The effect of some triazine derivatives on the growth and development of duckweeds. (In Serbo-Croat.). *Acta Bot. Croat.* 36, 75-82.
- RATHE W., 1952: Zur Physiologie des Kaliums. I. Mitt.: Parallelität von Kaliumgehalt und Säuregehalt lebender Zellen. *Z. Pflanzenernährung* 57, 151-163.
- RAU W. and SCHROTT E.L., 1979: Light mediated biosynthesis in plants. *Photochem. Photobiol.* 30(6), 755-765.
- RAVEN J.A., 1970: Exogenous inorganic carbon sources in plant photosynthesis. *Biol. Rev.* 45, 167-221.
- RAVEN J.A., 1981: Nutritional strategies of submerged benthic plants: the acquisition of C, N and P by rhizophytes and haptophytes. *New Phytol.* 88, 1-30.
- RAWN G.P., WEBSTER G.R.B. and MUIR D.C.G., 1980: Analysis and fate of ¹⁴C permethrin in an aquatic ecosystem. *Can. Plains Proc.* 9, 25-33.
- RAWN G.P., WEBSTER G.R.B. and MUIR D.C.G., 1982: Fate of permethrin in model outdoor ponds. *J. Environ. Sci. Health B* 17(5), 463-486.
- RAWN G.P., MUIR D.C.G. and WEBSTER G.R.B., 1983: Uptake and persistence of permethrin by fish, vegetation and hydrosol. *Can. Tech. Rep. Fish. Aquat. Sci.* 1151, 195-196.
- RAYNAL-ROQUES A., 1978: Les plantes aquatiques alimentaires. *Adansonia*, ser. 2, 18(3), 327-343.
- REAY P.F., 1972: The accumulation of arsenic from arsenic-rich natural water by aquatic plants. *J. Appl. Ecol.* 9, 557-565.
- RECHINGER K.H., 1943: Flora Aegaea. Springer, Wien. 846 pp.
- REDDY K.R., 1984a: Nutrient removal potential of aquatic plants. *Aquat. Bot.* 6(1), 15-16.
- REDDY K.R., 1984b: Nutrient transformations in aquatic macrophyte filters used for water purification. *Proc. Water Reuse Symp.* 2, 660-678.
- REDDY K.R. and DEBUSK W.F., 1985a: Nutrient removal potential of selected aquatic macrophytes. *J. Environ. Qual.* 14(4), 459-462.
- REDDY K.R. and DEBUSK W.F., 1985b: Growth characteristics of aquatic macrophytes cultured in nutrient-enriched water. II. Azolla, duckweed and Salvinia. *Econ. Bot.* 39(2), 200-208.
- REDDY K.R., SUTTON D.L. and BOWES G., 1983: Freshwater aquatic plant biomass production in Florida. *Proc. Soil Crop Sci. Soc. Florida* 42, 28-48.
- REGNAULT F., 1910: La culture des lentilles d'eau dans la lutte contre le paludisme. *Bull. Soc. Path. Exot.* 12, 735-736.
- REHMET J.B., 1974: A habitat-distribution study of the planarian flatworm (*Dugesia* sp.) in Dewart Lake, Indiana. *J. Sci. Lab. Derison Univ.* 55(4-7), 85-94.
- REID M.S., 1968: Response of Spirodela to phosphorus deficiency. *Ph.D. Thesis*. Univ. of Auckland.
- REID M.S. and BIELESKI R.L., 1970a: Response of Spirodela oligorrhiza to phosphorus deficiency. *Plant Physiol.* 46, 609-613.
- REID M.S. and BIELESKI R.L., 1970b: Changes in phosphatase activity in phosphorus-deficient Spirodela oligorrhiza. *Planta* 94(4), 273-281.
- REISFELD A. and EDELMAN M., 1979: In vitro translation and control of membrane protein synthesis in Spirodela chloroplasts. *Isr. J. Med. Sci.* 15(1), 109.
- REISFELD A., JAKOB K.M., GRESSEL J. and EDELMAN M., 1977: Two Spirodela

- chloroplast mRNAs and their discrete polypeptide products. *Isr.J.Bot.* 26(1), 49.
- REISFELD A., GRESSEL J., JAKOB K.M. and EDELMAN M., 1978a: Characterization of the 32,000 dalton membrane protein. I. Early synthesis during photoinduced plastid development of *Spirodela*. *Photochem.Photobiol.* 27, 161-165.
- REISFELD A., JAKOB K.M. and EDELMAN M., 1978b: Characterization of the 32000 dalton chloroplast membrane protein. II. The molecular weight of chloroplast messenger RNAs translating the precursor to P-32000 and full-size RUDP carboxylase large subunit. In: AKOYUNOGLU G. and ARGYROUDI-AKOYUNOGLU J.H. (eds.), *Chloroplast development*. Elsevier, North-Holland. Biomed.Press. 669-674.
- REISFELD A., JAKOB K.M. and EDELMAN M., 1978c: Molecular weight of chloroplast messenger RNA translating full-size RUDP carboxylase large subunit. *Plant Physiol.* 61(4 suppl.), 104.
- REISFELD A., JAKOB K.M. and EDELMAN M., 1978d: Characterization of the main membrane protein synthesized within the chloroplast of *Spirodela*. *Isr.J.Bot.* 27(1), 39.
- REISFELD A., MATTOO A.K. and EDELMAN M., 1982: Processing of a chloroplast-translated membrane protein in vivo. Analysis of the rapidly synthesized 32,000 dalton shield protein and its precursor in *Spirodela oligorrhiza*. *Eur.J.Biochem.* 124(1), 125-129.
- REJMANKOVA E., 1973: The influence of temperature and irradiance on the growth and production of duckweeds (*Lemna gibba* L., *Lemna minor* L. and *Spirodela polyrrhiza* [L.] Schleiden). (In Czech.). Thesis. Charles Univ., Praha. 100 pp.
- REJMANKOVA E., 1973a: Biomass, production and growth rate of duckweeds (*Lemna gibba* and *L. minor*). In: HEJNY S. (ed.), *Ecosystem study on wetland biome in Czechoslovakia*. Trebon. 101-106.
- REJMANKOVA E., 1973b: Seasonal changes in the growth rate of duckweeds (*Lemna gibba* L.) in the littoral of the Nesyt fishpond. In: KVET J. (ed.), *Littoral of the Nesyt fishpond*. *Ekol.Stud.Czech.Akad.Praha* 15, 103-106.
- REJMANKOVA E., 1973c: Seasonal changes in the growth rate of duckweed community (*Lemnetum gibbae*). *Folia Geobot.Phytotax.* 8, 1-13.
- REJMANKOVA E., 1973d: Biomass of submerged macrophytes growing in the Nesyt fishpond. In: KVET J. (ed.), *Littoral of the Nesyt fishpond*. *Ekol.Stud.Czech.Akad.Praha* 15, 107-110.
- REJMANKOVA E., 1973a: The growth of *Lemna* spp. under natural and semi-natural conditions. *Abstr.12th Int.Bot.Congr., Leningrad*. 164.
- REJMANKOVA E., 1975b: Comparison of *Lemna gibba* and *Lemna minor* from the production ecological viewpoint. *Aquat.Bot.* 1, 423-428.
- REJMANKOVA E., 1975c: Biology of duckweeds in a Pannonian fishpond. *Symp.Biol.Hung.* 15, 125-131.
- REJMANKOVA E., 1976: Germination of seeds of *Lemna gibba*. *Folia Geobot. Phytotax.* 11, 261-267.
- REJMANKOVA E., 1978: Growth, production and nutrient uptake of duckweeds in fishponds and in experimental cultures. In: DYKVOVA D. and KVET E. (ed.), *Pond littoral ecosystems. Ecol.Studies, Analysis and Synthesis*. Springer, Berlin/Heidelberg. 28, 278-291.
- REJMANKOVA E., 1979: The role of duckweeds in fishpond ecosystems. (In Czech.). Ph.D.Thesis. Univ. Pruhonice. 166 pp.
- REJMANKOVA E., 1981: On the production ecology of duckweeds. *Intern. Workshop on Aquatic Macrophytes, Illmitz, Austria*. 7 pp.
- REJMANKOVA E., 1982: The role of duckweeds (*Lemnaceae*) in small wetland water bodies of Czechoslovakia. In: GOPAL B. et al. (eds.), *Wetlands: ecology and management. Proc.1st Intern.Wetlands Conf.* (New Delhi 1980). 397-403.
- REJMANKOVA E., 1986: Relationship between competitive strength and fungal resistance in *Lemnaceae*. 8 pp. (Polycopy).
- REJMANKOVA E., KVET J. and REJMANEK M., 1979: The possibilities of the use of floating aquatic plants for biological water treatment. (In Czech.). 5th Conf.Limnology, Dum techniky CSVTS, Pardubice. 304-309.
- REJMANKOVA E., CULLEY D.D., KVET J. and REJMANEK M., 1983: Maximizing duckweed production by optimal strategy. 14 pp. (Polycopy).
- REJMANKOVA E., BLACKWELL M. and CULLEY D.D., 1986: Dynamics of fungal infection in duckweeds (*Lemnaceae*). *Veröff.Geobot.Inst.ETH, Stiftung Rübli,Zürich* 87, 178-189.
- REUTER L., 1948: Die Protoplasmatik der Schliesszellen von Schwimmpflanzen. I. Die Schliesszellen von *Lemna minor*. *Phyton(Austria)* 1, 76-79.
- REUVENY Z., 1974: ATP sulfurylase assay based on differential solubility of sulfate and adenosine phosphosulfate. *Fed.Proc.* 33(5/2), 1446.
- REVIER J.M., 1979: Ecological distribution of *Spirodela polyrrhiza* var. *masonii* in W. Nederland. (In Dutch). *Mimeogr.Rep.Hugo de Vries Lab.* 76.
- REZNÍK H. and MENSCHICK R., 1969: Flavonoide in Sommergliedern und Wintertknochen von *Spirodela polyrrhiza* (L.) Schleid. *Z.Pflanzenphysiol.* 61, 348-349.
- REZNÍK H. and NEUHAEUSEL R., 1959: Farblose Anthocyanine bei submersen Wasserpflanzen. *Z.Bot.* 47, 471-489.
- RHO J. and TAYLOR M., 1981: Composition, characterization and significance of phyllosphere and rhizosphere microflora of duckweed (*Lemna* sp.). *Abstr.Annu.Meet.Am.Soc.Microbiol.* 81, 180.
- RHODES D., 1976: The regulation of ammonia assimilation in *Lemna minor* L. Ph.D.Thesis. Univ. Manchester.
- RHODES D. and STEWART G.R., 1974: A procedure for the in-vivo determination of enzyme activity in higher plant tissue. *Planta* 118(2), 133-144.
- RHODES D., RENDON G.A. and STEWART G.R., 1975: The control of glutamine synthetase level in *Lemna minor* L. *Planta* 125, 201-211.
- RHODES D., RENDON G.A. and STEWART G.R., 1976: The regulation of ammonia assimilating enzymes in *Lemna minor*. *Planta* 129, 203-210.
- RHODES D., SIMS A.P. and STEWART G.R., 1979: Glutamine synthetase and the control of nitrogen assimilation in *Lemna minor*. In: HEWITT E.J. and CUTTING C.V. (eds.), *Nitrogen assimilation of plants*. Long Ashton Symp.(Proc.)1977. Acad.Press,London. 6, 501-520.
- RHODES D., SIMS A.P. and FOLKES B.F., 1980: Pathway of ammonia assimilation in illuminated *Lemna minor*. *Phytochem.* 19, 357-365.
- RHODES D., MYERS A.C. and JAMIESON G., 1981a: A rapid method for the determination of the N abundance of amino acids using electron impact gas chromatography mass spectrometry. *Plant Physiol.* 67(4 suppl.), 7.
- RHODES D., MYERS A.C. and JAMIESON G., 1981b: Gas chromatography mass spectrometry of N-heptafluorobutyryl isobutyl esters of amino acids in the analysis of the kinetics of ¹⁵N labeled ammonium assimilation. *Plant Physiol.* 68(5), 1197-1205.
- RHODES D., DEAL L., HAWORTH P., JAMIESON G.C., REUTER C.C. and ERICSON M.C., 1986: Amino acid metabolism of *Lemna minor* L. 1. Responses to methionine sulfoximine. *Plant Physiol.* 82(4), 1057-1062.
- RHODES L.W., 1968: The duckweeds: their use in the High School Laboratory. *Am.Biol.Teacher* 30(7), 548-551.
- RICHARD A., 1833a: *Reliquiae Richardianae ad analysin botanicam spectantes* *Hayadeae*. *Arch.Bot.* 1, 200-210.

- RICHARD A., 1933b: Observations sur le genre Lemna. Arch. Bot. 1, 210-212.
- RICHARD E.P., Jr., MILLER T.C. and BOWMAN D.H., 1980: Bifenox as an herbicide for the control of barnyardgrass in rice. MAFES Res. Highlights 43(4), 8.
- RICHARD J.M., RAVANEL P. and CANTIN D., 1987: Phytotoxicity of orellanine (a mushroom toxin). Toxicon 25(3), 350-354.
- RICHARDSON L.V., 1975: Water level manipulation: a tool for aquatic weed control. Hyacinth Control J. 1975, 8-11.
- RICHARDSON W.G., 1985: Bioassays for glyphosate. In: GROSSBARD E. and ATKINSON D. (eds.), The herbicide glyphosate. Butterworths, London. 286-298.
- RIDLEY H.N., 1930: The dispersal of plants throughout the world. Reeve, Ashford. 744 pp.
- RIEDL H., 1976: Lemnaceae. In: RECHINGER K.H. (ed.), Flora Iranica 119, 8 pp.
- RIEMER D.N. and TOTH S.J., 1968: A survey of the chemical composition of aquatic plants in New Jersey. New Jersey Agric. Exp. Stat. Bull. 820, 14.
- RIFAI S.A., 1979: The use of aquatic plants as feed for Tilapia nilotica in floating cages. Asian Aquaculture 2(6), 4-5.
- RIMES C.A. and GOULDER R., 1985: A note on the attachment rate of suspended bacteria to submerged aquatic plants in a calcareous stream. J. Appl. Bacteriol. 59(4), 389-392.
- RIMON D., 1964: Bud initiation in Spirodela oligorrhiza. Isr. J. Bot. 13, 24-29.
- RIMON D. and GALUN E., 1965 see 1968a
- RIMON D. and GALUN E., 1967: Morphogenesis in Spirodela oligorrhiza: Effects of pyrimidine base analogues on initiation, elongation and differentiation of fronds. Plant Cell Physiol. 8, 283-291.
- RIMON D. and GALUN E., 1968a: Morphogenesis of Spirodela oligorrhiza: ontogenesis of fronds. Bot. Gaz. 129, 138-144.
- RIMON D. and GALUN E., 1968b: Morphogenesis of Wolffia microscopica: frond and flower development. Phytomorphology 18, 364-372.
- RIMON D., GRESSEL J. and GALUN E., 1969: Pyrimidine analogues and nucleic acid synthesis in Spirodela. Isr. J. Bot. 18(3), 113-121.
- RIPL W., 1976: Processing lakes with damaged ecosystems. Vierteljahrsschr. Natf. Ges. Zürich 121(4), 301-308.
- RIVAS-MARTINEZ S., 1982: Vegetatio Matritensis. I. Datos sobre la vegetación flotante dulcícola de la clase Lemneta minoris. Lazaroa 4, 149-154.
- ROBERTS M.L., 1972: Wolffia in the bladders of Utricularia: an "herbivorous" plant? Mich. Bot. 11, 67-69.
- ROBERTS R.M. and LOEWUS F., 1968: Inositol metabolism in plants. VI. Conversion of myo-inositol to phytic acid in Wolffia floridana. Plant Physiol. 43, 1710-1716.
- ROBERTS R.M., SHAH R.H. and LOEWUS F., 1967: Inositol metabolism in plants. IV. Biosynthesis of apiose in Lemna and Petroselinum. Plant Physiol. 42(5), 659-666.
- ROBERTS R.M., CONNOR A.B. and CETORELLI J.J., 1971: The formation of glycoproteins in tissues of higher plants: Specific labelling with D-(1-C) glucosamine. Biochem. J. 125, 999-1008.
- ROBERTSON-CUNNINGHAM R.C. and BLACKMAN G.E., 1952: Effects of preliminary treatment on the subsequent variation in the resistance of Lemna minor to the phytotoxic action of 2,4-dichlorophenoxyacetic acid. Nature 170, 459.
- ROBINETTE R., 1980a: Use of duckweeds in catfish diets. M.S. Thesis. Dept. Wildlife Manage. Mississippi State Univ., State College.
- ROBINETTE H.R., BRUNSON M.W. and DAY E.J., 1980: Use of duckweed in diets of channel catfish. Proc. 13th Annu. Conf. SE Assoc. Fish Wildlife Age. 108-114.
- RODGERS J.H., Jr. and CHERRY D.S., 1975: Ash basin effluent impact on the aquatic flora of a stream and swamp drainage system. ASB Bull. 22(2), 76.
- RODGERS J.H. Jr., CHERRY D.S. and GUTHRIE R.K., 1978: Cycling of elements in duckweed (Lemna perpusilla) in an ash settling basin and swamp drainage system. Water Res. 12, 765-770.
- ROELOFS J.G.M., 1983: Impact of acidification and eutrophication on macrophyte communities in soft waters in the Netherlands. I. Field observations. Aquat. Bot. 17(2), 139-155.
- ROHWEDER H., 1937: Versuch zur Erfassung der mengenmäßigen Bedeckung der Darss und Zingst mit polyploiden Pflanzen. Ein Beitrag zur Bedeutung der Polyploidie bei der Eroberung neuer Lebensräume. Planta 27, 501-549.
- ROLA J., GABINSKA K. and GOUDEN B., 1975: Evaluation of biotests for determining herbicide effectiveness. (In Russian). Nauchn. Tsentr. Biol. Isal. Pushchino (USSR) 75(1), 205-211.
- ROMBACH J., 1961: Growth of Lemna minor as influenced by light and kine- tin. In: CHRISTENSEN B.C. and BUCHMANN B. (eds.), Progress in photo- biology. Elsevier, Amsterdam. 379-380.
- ROMBACH J., 1965: The influence of the phytochrome reaction on the growth of Lemna minor L. Meded. Landbouwhogeschool Wageningen 65(14), 1-11.
- ROMBACH J., 1966: The phytochrome reaction in Lemna minor L. Photochem. Photobiol. 5, 383-384.
- ROMBACH J., 1971: On the interaction of kinetin and phytochrome in Lemna minor growing in the dark. Acta Bot. Neerl. 20, 636-645.
- ROMBACH J., 1974a: Thiamine requirement and phytochrome in Lemna minor L. In: DE GREFF J.A. (ed.), Proc. Ann. Europ. Symp. Plant Morphogenesis, Antwerpen. 83-85.
- ROMBACH J., 1974b: Growth stimulation by cytokinins, thiamine and phyto- chrome in Lemna minor L. in darkness and in light. Acta Bot. Neerl. 23 (3), 348.
- ROMBACH J., 1976: Effects of light and phytochrome in heterotrophic growth of Lemna minor L. Meded. Landbouwhogeschool Wageningen 76(1), 1-114.
- ROMBACH J., 1978: Phytochrome variation and reversion in Lemna minor, L. gibba G3 and L. paucicostata 6746. Photochem. Photobiol. 27, 781-786.
- ROMBACH J. and SPRUIT C.J.P., 1968: On phytochrome in Lemna minor and other Lemnaceae. Acta Bot. Neerl. 17, 445-454.
- ROSENTHAL G.A., GULATI D.K. and SABHARWAL P.S., 1975: Studies on the growth effects of the canaline-urea cycle amino acids with Lemna minor L. Plant Physiol. 56, 420-424.
- ROSENTHAL G.A., GULATI D.K. and SABHARWAL P.S., 1976: Additive and synergistic growth-inhibiting properties of the canaline-urea cycle amino acid. Plant Physiol. 57, 493-496.
- ROSENER A., POSNER H.B. and GRESSEL J., 1973: Synthesis and processing of RNA in Lemna. Characterization by gel electrophoresis. Plant Cell Physiol. 14, 555-564.
- ROSENER A. and GRESSEL J., 1974a: The distribution of plastid ribosomes and the integrity of plastid ribosomal RNA during the

- greening and maturation of Spirodela fronds. Plant Cell Physiol. 15, 891-902.
- ROSNER A., EDELMAN M. and GRESSEL J., 1974b: Thermal denaturation of nucleic acids in polyacrylamide gels. Anal. Biochem. 58(2), 602-608.
- ROSNER A., JAKOB K.M., GRESSEL J. and SAGHER D., 1975a: The early synthesis and possible function of a 0.5×10^4 RNA after transfer of dark-grown Spirodela plants to light. Biochem. Biophys. Res. Commun. 67, 383-391.
- ROSNER A., POSNER H.B. and GRESSEL J., 1975b: Synthesis and processing of RNA in Lemna. Isr. J. Bot. 24(1), 46.
- ROSNER A., JAKOB K.M., GRESSEL J., SAGHER D. and EDELMAN M., 1976: Early synthesis and possible function of a 500000 molecular weight RNA after transfer of dark grown Spirodela plants to light. Isr. J. Bot. 25 (1-2), 98.
- ROSNER A., GRESSEL J. and JACOB K.M., 1977a: Discoordination of ribosomal RNA metabolism during metabolic shifts of Spirodela plants. Biochim. Biophys. Acta 474, 386-397.
- ROSNER A., HELLER D. and EDELMAN M., 1977b: Polyacrylamide gel as a medium for DNA dissociation and reassociation. Mol. Biol. Rep. 3(3), 207-212.
- ROSNER A., REISFELD A., JAKOB K.M., GRESSEL J. and EDELMAN M., 1977c: Shifts in the RNA and protein metabolisms of Spirodela (duckweed). In: BOGORAD I.B. and WEIL J. (eds.), Coll. Int. CNRS 261, 561-568.
- ROSS J.H., 1979: Structure-activity relationships in the toxicology of 4,4'-bipyridylum salts in the rat (Rattus norvegicus) and the duckweed (Spirodela oligorhiza). Diss. Abstr. Int. B 40(2), 699.
- ROSS J.H., KRIEGER R.I. and LIM L.O., 1979a: Herbicidal potency of paraquat homologues as a function of their physicochemical constants in duckweed. Toxicol. Appl. Pharmacol. 48(1/2), A194.
- ROSS J.H., LIM L.O. and KRIEGER R.I., 1979b: Herbicidal potency of 1,1'-alkyl-4,4'-bipyridylum salts as a function of their physicochemical constants in duckweed. Drug Chem. Toxicol. 2, 193-205.
- ROSS S.L., DOUGHTY C.R. and MURPHY K.J., 1986: Cause, effects and environmental management of a Lemna problem in a Scottish canal. Proc. 7th Intern. Symp. Aquatic Weeds. 277-283.
- ROSSI J.B., 1969: La germinación de semillas de Spirodela intermedia W. Koch. Soc. Argent. Bot. Bol. 11(4), 235-238.
- ROSTOVTEVA Z.P., 1967: The organogenesis and the characteristic structural features of Wolffia arhiza (L.) Wimm. during the vegetative period of life. (In Russian). Bot. Zh. (USSR) 52, 1177-1184.
- ROSTOWZEW S.I., 1901: On the flowering of duckweeds in the region of Moskwa. (In Russian). Ann. Inst. Agron. Moscou 7, 63-70.
- ROSTOWZEW S.I., 1905: On the biology and morphology of duckweeds. (In Russian). Ann. Inst. Agron. Moscou 11, 222-329.
- ROTH G., 1982: Vergleichende Untersuchungen über die Wirkung von Pentachlorphenol auf die Aktivitäten von Nitratreduktase und Nitratreduktase verschiedener Lemnaceen. Diploma Thesis. TU, München. (Polycopied).
- ROTTENBERG T., SCHARPETER E. and KANDELER R., 1981: Short period oscillation of starch levels in Lemna under light and darkness. Photosynth. Res. 1(4), 251-258.
- ROTMANN R., 1977: Management of weedy lakes and ponds with grass carp. Fisheries 2(5), 8-14.
- ROULET M.G., 1975: Wasserlinsen (Lemna minor L.) als Testpflanzen von Klärschlamm. Schweiz. Landwirtsch. Forsch. 14(1), 79-82.
- ROUSSEAU J., 1945: Le folklore botanique de Caughnawaga et le folklore botanique de l'île aux Coudres. Contrib. Inst. Bot. Univ. Montreal 55, 7-111.
- ROY G., 1912: Flore de France. Deyrolle, Paris. 13, 283-287.
- ROWE E.L., ZIOBRO R.J., WANG C.J.K. and DENCE C.W., 1982: The use of an alga Chlorella pyrenoidosa and a duckweed Lemna perpusilla as test organisms for toxicity bioassays of spent bleaching liquors and their components. Environ. Pollut. A 27(4), 289-296.
- ROWECK H., 1981: Die Gefäßspflanzen von Schwedisch Lappland. Flora Veget. Mundi 8. Cramer, Vaduz. 639.
- ROYBURGH W., 1832: Flora Indica. Mission Press, Calcutta. 3, 565-566.
- ROY R.P. and DUTT B., 1967: Cytology of Wolffia microscopica Kurz. Cytologia 32, 270-272.
- RUIZ-LEAL A., 1951: La floración de Lemna gibba L. Y Lemna Parodiana Giard. (Lemnaceae) en Mendoza. Rev. Fac. Cienc. Agr. 3(1), 1-8.
- RUNGE F., 1972: Die Flora Westfalens. (2nd ed.). Münster.
- RUNGE F., 1979: Neue Beiträge zur Flora Westfalens. Natur und Heimat 39, 69-103.
- RUSKIN F.R. and SHIPLEY D.W. (eds.), 1976: Making aquatic weeds useful: Some perspectives for developing countries. Appendix A: Duckweeds and their uses. Natl. Acad. Sci., Washington, D.C. 148-154.
- RUSOFF L.L., GANTT D.T., WILLIAMS D.M. and GHOLSON J.H., 1977: Duckweed - a potential feedstuff for cattle. J. Dairy Sci. 60(suppl. 1), 161.
- RUSOFF L.L., ZERINGUE S.P., ACHACOSO A.S. and CULLEY D.D., 1978: Feeding value of duckweed (an aquatic plant, family Lemnaceae) for ruminants. J. Dairy Sci. 61(suppl. 1), 186.
- RUSOFF et al., 1980 see 1980b
- RUSOFF L.L., ACHACOSO A.S. and GHOLSON J.H., 1980a: Nutritional studies with duckweeds. In: CRYE J.B. and CULLEY D.D. (eds.), U.S. Dept. Energy Final Report. Louisiana State Univ., Baton Rouge. 2 pp. (Polycopy).
- RUSOFF L.L., BLAKENEY E.W. and CULLEY D.D., 1980b: Duckweeds (Lemnaceae family) - a potential source of protein and amino acids. J. Agric. Food Chem. 28, 848-850.
- RUTTER J.C. and ERISMANN K.H., 1985: Influence of nitrogen availability on aminotransferases in Lemna minor L. J. Exp. Bot. 36(165), 583-589.
- RUTTNER F., 1947: Zur Frage der Karbonatassimilation der Wasserpflanzen. I. Die beiden Haupttypen der Kohlenstoffaufnahme. Oesterr. Bot. Z. 94, 265-284.
- RUTTNER F., 1948: Zur Frage der Kohlenstoffassimilation der Wasserpflanzen. II. Das Verhalten von Elodea canadensis und Fontinalis antipyretica in Lösungen von Natrium- bzw. Kaliumkarbonat. Oesterr. Bot. Z. 95, 208-238.
- RYANG H.S., KIM M.K. and JEON J.C., 1976: Control of perennial weeds in paddy rice in Korea. Proc. 5th Asian-Pac. Weed Sci. Soc. Conf. (Tokyo). 293-297.
- RYAZANOVA A.V. and LYSSENKO N.L., 1985: Dynamics of 2,4-DNA concentration in an aquatic environment and duckweed tissues in relation to pH. (In Russian). Yaroslavl Univ., 28-33. Ref. Biol. Zh. (1986), 106296.
- RYTHER J.H., ANDREWS D.A., DEBUSK T.A., FALKNER B.E., HANISAK M.D., STENBERG R.W., TUCKER C.S. and WILLIAMS L.D., 1980: Studies on biomass and biogas production by aquatic macrophytes. Proc. Bio-Energy (Washington D.C.), 130-133.
- RYTHER J.H., WILLIAMS L.D., HANISAK M.D., STENBERG R.W. and DEBUSK T.A., 1979: Biomass production by marine and freshwater plants. Proc. Bio-Energy, Golden, Colo., 13-23.

- SABAT S.C. and SAHA S., 1986: Photosynthetic metabolism of dark fixed carbon in some aquatic angiosperms. *Ind.J.Exp.Biol.* 24(8), 517-519.
- SABHARWAL P.S. and BHALLA P.R., 1973: Biological effects of tobacco smoke and its constituents on cells, tissues and chromosomes of plants. *Kentucky Univ.Tob.Health Res.Inst.Conf.Rep.* 4, 632-651.
- SABULARSE D.C. and ANDERSON R.L., 1982: Purification, properties and distribution of D-fructose-2,6-bisphosphate-activated inorganic pyrophosphatase: D-fructose-6-phosphate 1-phosphotransferase. *Plant Physiol.* 69(4 suppl.), 128.
- SÄGER A., 1925: The growth of duckweeds in mineral nutrient solutions with and without organic extracts. *J.Gen.Physiol.* 7, 517-526.
- SÄGER A., 1929: The flowering of the Lemnaceae. *Bull.Torr.Bot.Club* 56, 351-358.
- SÄGER A., 1930: A method of obtaining pure cultures of *Spirodela polyrrhiza*. *Bull.Torr.Bot.Club* 57, 117-121.
- SÄGER A., 1932 see 1934
- SÄGER A., 1933a: Manganese and the growth of Lemnaceae. *Am.J.Bot.* 20, 234-245.
- SÄGER A., 1933b: Gas injury to pure cultures of *Spirodela polyrrhiza*. *Bot.Club* 61, 479-480.
- SÄGER A., 1934: *Spirodela oligorrhiza* collected in Missouri. *Bull.Torr.Bot.Club* 61, 233-236.
- SÄGER A., 1937: The concentration of copper in nutrient solutions for *Spirodela polyrrhiza*. *Am.J.Bot.* 24, 640-643.
- SÄGESE E.J., FUGLIA T.A., LEATHER G., THOMPSON M.P., BILLIS D.D. and HOAGLAND P.D., 1985: Fractionation of allelochemicals from oilseed sunflowers and Jerusalem artichokes. In: *The chemistry of allelopathy*. *Am.Chem.Soc.(Washington D.C.)* 99-112.
- SAHAI R. and ROY P.S., 1977: Report of two natural enemies of *Spirodela polyrrhiza* (L.) Schleid. *Curr.Sci.* 46(23), 825.
- SAHAI R. and SINHA A.B., 1970: Contribution to the ecology of Indian aquatics. II. Studies on the growth rate of 'duck-weeds' (Lemna minor Linn. and *Spirodela polyrrhiza* (Linn.) Schleid.) under laboratory conditions. *Proc.Nat.Acad.Sci.India B* 39(1969), 143-144.
- SAHAI R., ROY P.S. and KAUR G., 1977a: A comparative study of the toxicity of synthetic detergents on two common duckweeds at Gorakhpur. *Indian J.Ecol.* 4, 242-245.
- SAHAI R., SINGH S.P. and GANGULEE K., 1977b: Role of synthetic detergent on the growth behaviour of *Spirodela polyrrhiza* (L.) Schleid. *Curr.Sci.* 46(19), 676-677.
- SAHAI R., ROY P.S. and WAJITH S.A., 1980: Effect of certain herbicides on the growth performance of *Spirodela polyrrhiza*. *Indian J.Weed Sci.* 12(1), 87-92.
- SAID M.Z.B.M., 1978: Growth of duckweed *Spirodela polyrrhiza* in a system utilizing dairy cattle manure. M.S.Thesis. Louisiana State Univ., Baton Rouge. 48 pp.
- SAID M.Z.B.M., CULLEY D.D. Jr., STANDIPER L.C., EPPS E.A., MYERS R.W., and BONEY S.A., 1979: Effect of harvest rate waste loading, and stocking density on the yield of duckweeds. *Proc.World Maricult.Soc.* 10, 769-780.
- SAJI H., FURUYA M. and TAKIMOTO A., 1982: Spectral dependence of night break effect on photoperiodic floral induction in Lemna paucicostata 441. *Plant Cell Physiol.* 23(4), 623-629.
- SAJI H., NAGATANI A., YAMAMOTO K.T., FURUYA M., FUKUMOTO T. and YAMASHITA A., 1984: Cross-reactivity of monoclonal antibodies against rye and pea phytochrome with phytochromes extracted from 8 different plant species. *Plant Sci.Lett.* 37(1-2), 57-62.
- SAKAI W.S., SHIROMA S.S. and NAGAO M.A., 1984: A study of raphide microstructure in relation to irritation. *Scann.Electr.Microsc.* 2, 979-986.
- SAKS Y., ILAN I. and NEGBI M., 1975: Production and release of abscisic acid in cultures of *Spirodela polyrrhiza* and its possible role in the regulation of morphogenetic events. *Plant Physiol.* 56(2 suppl.), 85.
- SAKS Y., NEGBI M. and ILAN I., 1980: Involvement of native abscisic acid in the regulation of onset of dormancy in *Spirodela polyrrhiza*. *Austr.J.Plant Physiol.* 7, 73-79.
- SAMPFORD M.R., 1952: Studies in the principles of phytotoxicity. II. Experimental designs and techniques of statistical analysis. *J. Exp.Bot.* 3, 28-46.
- SAWUELISSON G., 1934: Die Verbreitung der höheren Wasserpflanzen in Nord-europa (Fennoskandien und Dänemark). *Acta Phytogeogr. Suecica* 6, 211 pp.
- SANCHEZ ANTA A., NAVARRO ANDRES F. and ELENA-ROSELLO J.A., 1984: Adquisición de caracteres de adaptación de las Lemnaceas a la vida dulceacuicola. *Studia Bot.* 3, 281-286.
- SANDERHANN H., Jr., 1969: Specific and rapid determination of D-apiose. *Phytochem.* 8(8), 1571-1575.
- SANDERHANN H., Jr. and GRISEBACH H., 1970: Biosynthesis of D-apiose. V. NAD-dependent biosynthesis of UDP-apiose and UDP-xylose from UDP-D-glucuronic acid with an enzyme preparation from Lemna minor L. *Biochim.Biophys. Acta* 208, 173-180.
- SANDERHANN H., Jr., TISUE G.T. and GRISEBACH H., 1968: Biosynthesis of D-apiose. IV. Formation of UDP-apiose from UDP-D-glucuronic acid in cell-free extracts of parsley (*Apium petroselinum* L.) and Lemna minor. *Biochim.Biophys. Acta* 165, 550-552.
- SANKARAN U.K., 1977: Seasonal variation in chemical constituents of some aquatic plants. *J.Bombay Nat.Hist.Soc.* 69, 242-246.
- SANKHIA N. and HUBER W., 1979: Effect of abscisic acid on enzyme induction in Lemna minor L. *Z.Pflanzenphysiol.* 91, 7-15.
- SARAWEK S., 1977: The effect of pyridoxal phosphate on the activity of aldolase from Lemna minor. *J.Sci.Fac.Chiangmai Univ.* 4, 62-87.
- SARAWEK S. and DAVIES D.D., 1977a: The effect of pyridoxal phosphate on the activity of aldolase from Lemna minor L. *Planta* 137, 265-270.
- SARAWEK S. and DAVIES D.D., 1977b: The control of aldolase in Lemna minor L. in relation to nitrogen deficiency. *Planta* 137, 271-277.
- SARGENT J.A., 1957: Factors determining the pattern of vascular tissue in Lemna minor L. Ph.D.Thesis. Univ. London.
- SARGENT J.A. and WANGERMANN E., 1959: The effect of some growth regulators on the vascular system of Lemna minor. *New Phytol.* 58, 345-363.
- SAROSTEK J. and WOZAKOWSKA-NATKANIEC H., 1980: The populations of Lemna minor L. and *Spirodela polyrrhiza* (L.) Schleiden as bioindicators of heavy metals water pollutions in extent of foundry emission. *Proc. 3rd Int.Conf.Bioindic.Deterior.Reg.(Praha)* 25-31.
- SAROSTEK J. and WOZAKOWSKA-NATKANIEC H., 1984: The effect of calcium and magnesium on the radiosensitivity of *Wolffia arrhiza* (L.) Wimm. *Acta Soc.Bot.Pol.* 53(3), 399-410.
- SAROSTEK J., WOZAKOWSKA-NATKANIEC H. and NIEMIEC C., 1982: The influence of heavy metals on the dynamics of the Lemna minor population. (in Polish). *Acta Univ.Wratisl.* 530, Prace Bot. 25, 11-39.
- SASTROUTOMO S.S., 1982: Summer biomass of aquatic macrophytes in relation to sediment characteristics in Lake Aino-Numa, Miyaki. *Jap.J. Ecol.* 32(1), 45-55.
- SASTROUTOMO S.S., 1984: Classification and ordination of aquatic vegetation in Lake Aino-Numa, Miyaki. *Ecol.Rev.* 20(3), 205-222.

- SATAKE K. and SHIMURA S., 1983: Carbon dioxide assimilation from air and water by duckweed *Spirodela polyrrhiza* (L.) Schleid. Hydrobiol. 107 (1), 51-55.
- SATAKE K. and UHRO T., 1984: Selection of indicator plants for detection of heavy metal pollution. (In Japan.) Kok.Kog.Kenkyusho Kenkyu Hokoku 58, 11-20.
- SATO T. and ODA Y., 1977: Significance of timing and number of short-day cycles for initiation and subsequent development of floral buds in *Lemna perpusilla* 6746. Plant Cell Physiol. 18, 1041-1046.
- SATO T. and ODA Y., 1978: Effects of culture conditions on development in continuous light of floral buds in *Lemna perpusilla* 6746. Plant Cell Physiol. 19(4), 537-543.
- SAUNDERS G.W., 1957: Interrelations of dissolved organic matter and phytoplankton. Bot.Rev. 23, 389-409.
- SAUNDERS J.A. and McCURE J.W., 1976: The distribution of flavonoids in chloroplasts of twenty five species of vascular plants. Phytochem. 15(5), 809-810.
- SAUZE F., 1982: Cultures en alimentation subcontinue. Influence et optimisation de certains paramètres. Tech.Eau Assainiss. 422, 27-39.
- SAVAGE K.E., TRUELOVE B. and WIESE A.F., 1978: Herbicide movement from application sites and effects on non-target species. South.Coop.Ser. Bull. 234, 16 pp.
- SAVAGE S., 1945: A catalogue of the Linnaean Herbarium. Linn.Soc., London. 225 pp.
- SAVULESCU T., 1972: Flora Rep.Soc. Rom. Acad.Rep.Soc. Rom. 12, 780-787.
- SEBURLINO G., SCOPPOLA A. and MARCHIORI S., 1986: Contributo alla conoscenza degli ambienti umidi della Pianura Padana orientale: la Classe Lemneta minoris R.Tx. 1955 em. Schw. et R.Tx. 1981. Not.Soc.Ital. Fitosoc. 21, 61-70.
- SCHAEFER G., 1954: Beziehungen zwischen Stoffwechsel und Plasmazustand in der Wurzel von *Lemna minor* L. Ph.D.Thesis. Marburg.
- SCHAEFER G., 1955: Ein Versuch zur quantitativen Auswertung der Plasmolyseform und -zeitmethode. Protoplasma 44, 422-436.
- SCHAEFER G., 1956: Ueber die Wirkung von Stoffwechselfaktoren auf den Plasma-Wand-Kontakt in der Wurzel von *Lemna minor* L. Flora 143, 327-355.
- SCHAEFER G., 1958: Ueber Veränderungen des mechanischen und biologischen Kontaktes von Plasma und Zellwand in subplasmolytischen Konzentrationen von Anektrolyten. Planta 51, 414-439.
- SCHAEFER M., 1974: Der Einfluss des Schwefeldioxids (SO₂) auf die Sulfataufnahme durch *Lemna minor* L. unter standardisierten Bedingungen. Lic.Thesis. Univ. Bern.
- SCHAEFER M., BRUNOLD C. and ERISMANN K.H., 1975: Hemmung der Sulfataufnahme durch *Lemna minor* L. durch SO₂ in subletalen Konzentrationen. Experientia 31, 1414-1415.
- SCHAEFFNER M., 1905: Free-floating plants of Ohio. Ohio Natur. 6, 420-421.
- SCHARFETTER E., ROTTENBURG T. and KANDELER R., 1978: Die Wirkung von EDDHA und Salicylsäure auf Blütenbildung und vegetative Entwicklung von *Spirodela punctata*. Z.Pflanzenphysiol. 87, 445-454.
- SCHARFETTER E., BECK E. and KANDELER R., 1981: End-of-day far-red effect on short-period oscillations of starch content in *Lemna*. Europ. Symp., Light mediated plant development. Book of Abstracts 7, 14.
- SCHARFETTER E., FAERBER E. and KANDELER R., 1984: Blühinduktion und Äthylenhaushalt bei *Lemna*. Mitt.bd.Bot.-Tag.Wien, 42 (0406).
- SCHARFETTER E., FAERBER E. and KANDELER R., 1986: Involvement of ethylene in the effect of EDDHA on flowering and vegetative development in *Spirodela punctata*. Veröff.Geobot.Inst.ETH,Stiftung Rübel,Zürich 87, 168-177.
- SCHARFETTER E., LESEMANN Ch. and KANDELER R., 1987: Ethylene as a flower-promoting agent in *Lemna*. Phytol.(Austria), in press.
- SCHARFETTER E., MENKE K.H., GOETZ D., MEYER-SPASCHKE H. and DORFFLING K., 1982: Anzucht, Analyse und Nutzung des Stickstoff sammelnden Systems *Azolla-anabaena*, sp. Filicoides in Nährlösungen und Klärbrühen. Landw.Forsch. 35(3-4), 200-213.
- SCHAUMANN L., GALLIE P., ULLRICH W. and THELIER M., 1986: Application de la microscopie ionique analytique à l'utilisation des isotopes stables ¹⁵N et ¹³C comme traceurs, et pour faire l'image de la distribution de l'azote chez *Lemna gibba* L. C.R.Acad.Sci.Paris, III, 302(3), 109-115.
- SCHAEFFER F., RATHJE W. and SCHAPFAYER H., 1952: Zum Mechanismus der Stoffaufnahme pflanzlicher Zellen. (Ein Beitrag zur Kalium- und Natriumaufnahme lebender Zellen). Z.Pflanzenernährung 56, 139-151.
- SCHMID H.W., EHMEKE A. and HARTMANN T., 1980: Plant NAD-dependent glutamate dehydrogenase. Purification, molecular properties and metal ion activation of the enzymes from *Lemna minor* and *Pisum sativum*. Z.Naturforsch. 35C, 213-221.
- SCHNEIDER O., 1976: Zur biologischen Phosphorylierung von myo-Inositol und L-myo-Inositol-1-phosphat. Ph.D.Thesis. Univ. Vienna.
- SCHNEIDER O., PITNER F., BOLLMANN O. and KANDELER R., 1978: Die Wirkung von Stickstoffmangel und anderen Faktoren auf die Phytinsäureseigerung in *Lemna gibba* Gl. Z.Pflanzenphysiol. 88, 295-303.
- SCHENCK H., 1886: Vergleichende Anatomie der submersen Gewächse. Bibl. Bot. 1(1), 1-67.
- SCHER S. and AARONSON S., 1958: Nutritional factors in apochlorosis: comparative studies with algae and higher plants. In: The photochemical apparatus. Brookhaven Symp. in Biology 11, 343-347.
- SCHIFF J.A., 1980: Pathways of assimilatory sulphate reduction in plants and microorganisms. In: Excerpta medica, sulphur in biology. CIBA Foundation Symposium 72, 49-69.
- SCHLEIDEN M.J., 1839: Prodrum Monographiae Lemnacearum oder Conspectus generum atque specierum. Linnaea 13, 385-392.
- SCHLEIDEN M.J., 1844: Beiträge zur Botanik. Leipzig. 242 pp.
- SCHMIDT A., 1975: Distribution of APS-sulfofructosylase activity among higher plants. Plant Sci.Lett. 5, 407-415.
- SCHMIDT A., 1984: Occurrence of mercaptopyruvate sulfofructosylase activity in photosynthetic organisms. Z.Naturforsch. 39C(9-10), 916-921.
- SCHMIDT D., 1981: Pflanzensoziologische und ökologische Untersuchungen der Gewässer um Gützw. Natur u. Naturschutz Mecklenburg 17, 130 pp.
- SCHMIDT-CLAUSEN H.J. and ZIEGLER J., 1969: The influence of light quality on the activation of NADP-dependent glyceraldehyde-3-phosphate dehydrogenase. In: METZNER A. (ed.), Progress in photosynthesis research, Tübingen, 3, 1646-1652.
- SCHMIDT-CLAUSEN H.J., ZIEGLER I. and ZIEGLER H., 1969: Die lichtinduzierte Aktivitätssteigerung der NADP-abhängigen Glyceraldehyd-3-phosphat-Dehydrogenase. XI. Die Kinetik der reversiblen Reaktion bei *Lemna gibba*. Planta 86(3), 272-285.
- SCHMUTZ D. and BRUNOLD C., 1982: Rapid and simple measurement of ATP sulfurylase activity in crude plant extracts using an ATP-meter for bioluminescence determination. Anal.Biochem. 121(1), 151-155.
- SCHNYDER J., ROTTENBERG M. and ERISMANN K.H., 1975: The synthesis of threonine and thiothreonine from o-phospho-homoserine by extracts

prepared from higher plants. Biochem. Physiol. Pflanzen 167, 605-608.

SCHOLZ J.D., KERR R.I. and NUTTALL P.M., 1986: Treatment of wastewater by aquaculture systems. AWRC Res. Proj. No. 80/139, Australia. 104 pp.

SCHOLZ G., 1962: Versuche mit Bor an Lemnaceen: Die Wirkung des Bors auf den Kohlenhydratgehalt von Lemna minor L. Kulturpflanze 10, 63-71.

SCHOLZE U., 1986: Lemnetae in Vorarlberg. Diploma Thesis. Univ. Innsbruck. 146 pp. (Polycopy).

SCHOTT C.D. and WORTHLEY E.G., 1973: Effects of CS and its breakdown products on the growth of duckweeds. Edgewood Arsenal, Aberdeen Prov-ing Ground, Maryland Rep. No. EB-TR-73045. 26 pp.

SCHREINEMAKERS W.A.C., 1984: Effects of metal ions on growth of and on ion absorption by Spirodela polyrrhiza (L.) Schleid. Effects of iron, magnesium and zinc. Z. Pflanzenphysiol. 114(2), 123-129.

SCHREINEMAKERS W.A.C., 1986: The interaction between Cd-absorption and Cd-compartmentation in Wolffia gladiata. Acta Bot. Neerl. 35(1), 23-34.

SCHREINEMAKERS W.A.C. and DORHOUT R., 1985: Effects of copper ions on growth and ion absorption by Spirodela polyrrhiza (L.) Schleid. J. Plant Physiol. 121(4), 343-351.

SCHROEDER F., 1977: Die Mollusken der Pflanzengesellschaften in den Gewässern des Bremer Raumes. 1. Gastropoden der Lemnetae im Bremer Blockland. Abh. Naturw. Verein Bremen 38, 423-430.

SCHUIZ B., 1962: Wasserlinsen. Die neue Brehm-Bücherei, Ziemsen, Wittenberg. 95 pp.

SCHUSTER M., 1968: Die Bedeutung von Starklicht und Kupfer für die phytochromgesteuerte Morphogenese von Lemna perpusilla. Ph.D. Thesis. Univ. Würzburg. 102 pp.

SCHUSTER M. and KANDELER R., 1970: Die Bedeutung der Photosynthese für die Langtag-Blüte der Kurztagpflanze Lemna perpusilla 6746. Z. Pflanzenphysiol. 63(4), 308-315.

SCHUTH C.K., ISENSEE A.R., WOOLSON E.A. and KARNY P.C., 1974: Distribution of ^{14}C and arsenic derived from ^{14}C cacodylic acid in an aquatic ecosystem. J. Agric. Food Chem. 22(6), 999-1003.

SCHWABE-BRAUN A. and TUXEN R., 1981a: Zur Synsystematik der Klasse Lemnetae minoris in Europa. Ber. Int. Symp. Int. Ver. Vegk., Rinteln 1980, 181-205.

SCHWABE-BRAUN A. and TUXEN R., 1981b: Lemnetae minoris. Prodrömus der europäischen Pflanzengesellschaften 4. Cramer, Vaduz. 141 pp.

SCHWARZ R., FRIED W., PITTNER F. and HOFFMANN-OSTENHOF O., 1974: Untersuchungen über die Biosynthese der Cyclite. 32. Mitt. Anwendung der Affinitätschromatographie an NAD-Sepharose zum Nachweis und zur Reinigung von myo-Inositol-1-phosphat-Synthase in Erythrocyten von Hühnern und in Lemna gibba. Monatshefte Chem. 105, 445-451.

SCHWEBEL J.O., 1973: Hormonal control of growth and development in Lemna minor L., with special emphasis on the role of abscisic acid (ABA). Ph.D. Thesis. College Station, Texas. Univ. Microfilms Int., Ann Arbor, Mich., Order No. 73-19, 682; 129 + 14 pp. Diss. Abstr. Int. B 34, 1015.

SCHWEBEL J.O. and ANDERSON K., 1982: The effect of lead acetate and lead nitrate on the growth of Lemna minor L. Florida Sci. 45(suppl.1), 31.

SCHWEIGER G., GOLLE B., LOOS S. and RYGOL J., 1984: Effects of HgCl₂ and pentachlorophenol on passive and active membrane properties in tissue discs and intact duckweeds (Lemna gibba). In: CRAM W.J. et al. (eds.), Membrane transport in plants. Symp. Abstr. (Praha). 427-428.

SCHWIER H., 1953: Wasserlinsentrif in der Weser. Natur und Heimat 2, 1.

SCHWOBEL J. and TILLMANN G.C., 1972: Ammonium-Adaptation bei submersen Phanerogamen in situ. Arch. Hydrobiol. 42(suppl.1), 139-141.

SCOGGAN H.J., 1978: The Flora of Canada. Nat. Mus. Sci., Canada, 7(2), 456-457.

SCOPPOLA A., 1982: Considérations nouvelles sur végétations des Lemnetae minoris (R.Tx.1955) em. A. Schwabe et R.Tx.1981 et contribution à l'étude de cette classe en Italie centrale. Doc. Phytosoc. 6, 1-130.

SCOPPOLA A., 1983: Synthèse des Lemnetae minoris en Europe. Coll. Phytosoc. (Lille) 10, 513-520.

SCOTLAND M.B., 1934: The animals of the Lemna association. Ecology 15, 290-294.

SCOTLAND M.B., 1940: Review and summary of studies of insects associated with Lemna minor. J. New York Ent. Soc. 48, 319-333.

SCUTTHORPE C.D., 1967: The biology of aquatic vascular plants. Arnold, London. 610 pp.

SEAMAN D.E., 1964: Action spectrum PF paraquat toxicity in duckweed. Abstr. Weed Soc. Am. 1964, 102.

SEBACHER D.I., HARRISS R.C. and BARTLETT K.B., 1985: Methane emissions to the atmosphere through aquatic plants. J. Environ. Qual. 14(1), 40-46.

SEEWALDT E. and STACKEBRANDT E., 1982: Partial sequence of 16 S ribosomal RNA and the phylogeny of Prochloron. Nature 295, 618-620.

SEGAL S., 1966: Ecology of higher aquatic plants. (In Dutch). Vakbl. Biol. 46, 138-149.

SEKINE I., 1979: Alcohol and methane from Lemnaceae. (In Japan.). Jap. Kokai Tokyo Koho. Patent No. 56-78594.

SELISKAR A., 1983: A contribution to the knowledge of the classes Lemnetae and Potamogetonetea in Slovenia. (In Serbo-Croat.). Biol. Vestn. 31(1), 25-34.

SEMIKHINA K.A., 1979: New sites of rare aquatic species in water bodies of the Desna River flood plains, USSR. (In Russian). Ukr. Bot. Zh. 36, 214-218.

SETH P.N., VENKATARAMAN R. and MAHESHWARI S.C., 1970: Studies on the growth and flowering of a short-day plant, Wolffia microscopica. II. Role of metal ions and chelates. Planta 90, 349-359.

SETO M., TAKAHASHI Y., USHIJIMA T. and TAZAKI T., 1979: Chlorotic death of Lemna gibba by cadmium in different concentrations of nutritional minerals. (In Japan.). Jap. J. Limnol. 40(2), 61-65.

SEVERI A. and BARONI FORNASIERO R., 1982: Preliminary SEM observations of stomatal apparatus in Spirodela oligorrhiza (Kurz) Hegelm. in darkness and in light. Caryologia 35, 400-401.

SEVERI A. and BARONI FORNASIERO R., 1983a: Morphological variations in Lemna minor L. and possible relationships with abscisic acid. Caryologia 36(1), 57-64.

SEVERI A. and BARONI FORNASIERO R., 1983b: Effects of metabolic inhibitors and inophores on the stomatal apparatus in Spirodela oligorrhiza (Kurz.) Hegelm. and in Lemna minor L. Z. Pflanzenphysiol. 113(1), 39-46.

SEVERI A. and BARONI FORNASIERO R., 1983c: Stomatal responses in Spirodela oligorrhiza and in Lemna minor after treatments with some inhibitors and ionophores. G. Bot. Ital. 117(3-4), 178-179.

SEYMOUR F., 1969: The Flora of New England. C.E. Tuttle Co., Rutland, Vermont. 169-170.

SHAH K.L. and TYAGI B.C., 1985: A note on the role of Kachuga tectum Gray (Reptilia: Chelonia) in controlling aquatic weeds. Hydrobiologia 122, 243-245.

SHAHAK Y., POSNER H.B. and AVRON M., 1976: Evidence for a block between plastoquinone and cytochrome f in a photosynthetic mutant of Lemna with abnormal flowering behavior. Plant Physiol. 57, 577-579.

- SHALAR V.M., MOGYLYA V.M., KUMPERE A.G., RUDIK V.F. and OBUKH P.A., 1983: Cultivation of the higher aquatic plants and algae in media containing the waste water of livestock breeding complexes. (In Russian). Izv. Akad. Nauk SSSR Ser. Biol. Khim. Nauk 1983(6), 9-13.
- SHAMBHU M.B., DIGENTIS G.A., GULATI D.K., BOWMAN K. and SABHARWAL P.S., 1976: Chemical combinations of 2,4-dichlorophenoxyacetic acid (2,4-D) and polystyrene: Preparation and application for the control of duckweed. J. Agric. Food Chem. 24, 666-668.
- SHANNON E.L., 1953: The production of root hairs by aquatic plants. An. Midland Naturalist 50, 474-479.
- SHAPIRO J., 1957: Chemical and biological studies on the yellow organic acids of lake water. Limnol. Oceanogr. 2(3), 161-179.
- SHAPIRO J., 1958: Yellow acid-cation complexes in lake water. Science 127(3300), 702-704.
- SHAPIRO J., 1964: Effect of yellow organic acids on iron and other metals in water. J. Amer. Water Works Assoc. 56(8), 1062-1082.
- SHARITZ R.R. and LUVALL J.C., 1978: Growth of duckweed under constant and variable temperatures. In: THORP J.H. and GIBBONS J.W. (eds.), Energy and environmental stress in aquatic systems. DOE Symp. Ser. (CONF-771114), 48, 410-419.
- SHARMA K.P., 1979: An efficient device of harvesting fish in weed infested waters in Rajasthan. J. Ind. Fish Soc. India 11(1), 138-139.
- SHARP M., 1978: Study of effects of certain ammonium test solutions on Lemna minor, duckweed. J. Tenn. Acad. Sci. 53(4), 155.
- SHCHERBAKOV A.V., 1987: Lemna gibba L. in Moscow region. (In Russian). Biol. Mosk. Ispitat. Prirod. Biol. 92(3), 90-92.
- SHEPARD D.V. and THURMAN D.A., 1973: Effect of nitrogen sources upon the activity of L-glutamate dehydrogenase of Lemna gibba. Phytochem. 12, 1937-1946.
- SHELTON D.R., 1979 see NEWTON et al. 1978
- SHIBATA O. and TAKIMOTO A., 1975: Flowering response of Lemna perpusilla 6746 to a single dark period. Plant Cell Physiol. 16, 513-519.
- SHIBAYAMA H. and MIYAHARA M., 1978: Aquatic weeds in creeks of the paddy area on the lower region of Chikugo River. IV. Yearly and seasonal changes of cover degrees of weeds. (In Japan.). Weed Res., Japan, 23(3), 109-115.
- SHIH C.Y., 1979: SEM studies of the flowering of duckweed, Lemna perpusilla, 6746. SEM 1979(3), 479-486.
- SHIMOMURA H., SASHIDA Y. and NAKATA H., 1981: Plant growth regulating activities of crude drugs and medicinal plants. Shoyakugaku Zasshi 35(3), 173-179.
- SHIMOMURA H., SASHIDA Y., NAKATA H. and IMAMURA M., 1982: Plant growth inhibitors from methanol extracts of Gallae rhois. Shoyakugaku Zasshi 36(2), 132-133.
- SHIMURA Y. and VOGEL H.J., 1966: Diaminopimelate decarboxylase of Lemna perpusilla: Partial purification and some properties. Biochim. Biophys. Acta 118, 396-404.
- SHINOZAKI M. and TAKIMOTO A., 1983: Effects of some growth regulators and benzoic acid derivatives on flower initiation and root elongation of Pharbitis nil cultivar Kidachi. Plant Cell Physiol. 24(3), 433-440.
- SHINOZAKI M., HIKICHI M., YOSHIDA K., WATANABE K. and TAKIMOTO A., 1982: Effect of high-intensity light given prior to low-temperature treatment on the long-day flowering of Pharbitis nil. Plant Cell Physiol. 23(3), 473-477.
- SHIREMAN J.V., COLLE D.E. and ROTTMANN R.W., 1977: Intensive culture of grass carp Ctenopharyngodon idella in circular tanks. J. Fish. Biol. 11, 267-272.
- SHIREMAN J.V., COLLE D.E. and ROTTMANN R.W., 1978: Growth of grass carp fed natural and prepared diets under intensive culture. J. Fish. Biol. 12(5), 457-464.
- SHIREMAN J.V., ROTTMANN R.W. and ALDRIDGE F.J., 1983: Consumption and growth of hybrid grass carp fed four vegetation diets and trout chow in circular tanks. J. Fish. Biol. 22(6), 685-694.
- SHIROISHI T. and SHIMIZU R., 1984: Sensitivity of duckweed to Cr⁶⁺. (In Japan.). Toyama-ken Eisei Kenkyusho Nenpo 7, 264-266.
- SHISHKIN B.K., 1964: Lemnaceae. In: KOMAROV L. (ed.), Flora SSSR 3, 389-391.
- SHOAR-GHAFARI A., 1983: Etudes ultrastructurales de cellules oxalifères en vie latente ou en vie active dans deux plantes aquatiques (Spirodela polyrrhiza L. et Lemna minor L.). D.E.A. Univ. Paris VI. (P. and M. Curie). (Polycopy).
- SHREVE F. and WIGGINS I.L., 1964: Vegetation and flora of the Sonora Desert. Stanford Univ. Press, Stanford. 1, 327-329.
- SHUKLA A.C. and ANASTHI A.K., 1985: Influence of auxin-photoperiodic interrelationships of Wolffia arrhiza. Natl. Acad. Sci. Lett. (India) 8(3), 65-66.
- SHUKLA A.C. and PANDE S.N., 1975a: Correlative studies on auxin photoperiodic growth and moisture relationships of Spirodela polyrrhiza (Linn.) Schleid. Assoc. Adv. Plant Sci., Kalyani, India, 47-48.
- SHUKLA A.C. and PANDE S.N., 1975b: Changes in ascorbic acid accompanying auxin photoperiodic inductions in Spirodela polyrrhiza. Indian Sci. Congr. Assoc. Proc. 62, 104.
- SHUKLA A.C. and PANDE S.N., 1976: Correlative studies on auxin photoperiodic growth and moisture relationships of Spirodela polyrrhiza (Linn.) Schleid. Botanique (Nagpur) 7(1), 51-54.
- SHUKLA A.C., PANDE S.N. and SHUKLA P., 1972: Studies on hormonal photoperiodic interrelationships of Spirodela polyrrhiza. Indian Sci. Congr. Assoc. Proc. 59(3), 378-379.
- SHUKLA A.C., PANDE S.N. and SHUKLA P., 1973: Studies on hormonal photoperiodic interrelationships of Spirodela polyrrhiza (Linn.) Schleid. Plant Sci. 5, 60-63.
- SIBASAKI T. and ODA Y., 1979: Heterogeneity of dormancy in the turions of Spirodela polyrrhiza. Plant Cell Physiol. 20(3), 563-571.
- SIDOROFF S., 1910: La floraison de Lemna (une grande rareté).
- SIEFERMANN D., 1971: Ueber den Zusammenhang von Xanthophyllcyclus und Photosynthese bei Lemna gibba L. Ph.D. Thesis. Univ. Tübingen. 83 pp.
- SIEFERMANN D., 1972: Kinetic studies on the xanthophyll cycle of Lemna gibba L. - Influence of photosynthetic oxygen and supplied reductor. Proc. 2nd Int. Congr. Photosynth. 1971, 1, 629-635.
- STIEGEL S.M., 1961: Effect of reduced oxygen tension on vascular plants. Physiol. Plant. 14, 554-557.
- STIELING D.H., 1937: The influence of the phosphate-calcium ratio and of humates on chlorosis in Lemna. Iowa State Coll. J. Sci. 12, 151-154.
- STIK W.K., 1980: Growth and mineral uptake in the Lemnaceae. CSRS Calb Proj. No. Ca-D-LAW-3639-H. Rep. 0075135.
- SILVERTHORNE J. and TOBIN E.M., 1984: Demonstration of transcriptional regulation of specific genes by phytochrome action. Proc. Natl. Acad. Sci. USA 81(4), 1112-1116.
- SILVERTHORNE J., WINPEE C.F. and TOBIN E.M., 1985: Phytochrome regulation at the level of transcription. Photochem. Photobiol. 41(suppl.), 48 pp.

- SILVEY W.D., 1967: Occurrence of selected minor elements in the waters of California. U.S. Gov. Print. Off., Washington DC. Geol. Surv. Water-Supply 1535L, 1-25.
- SIMON E.W. and BLACKMAN G.E., 1953: Studies in the principles of phyto-toxicity. 4. The effects of the degree of nitration on the toxicity of phenol and other substituted benzenes. J. Exp. Bot. 4, 235-250.
- SIMON E.W., ROBERTS H.A. and BLACKMAN G.E., 1952: Studies on the principles of phytotoxicity. 3. The pH factor and the toxicity of 3,5-dinitro-o-cresol, a weak acid. J. Exp. Bot. 3, 99-109.
- SIMS A.P., FOLKES B.F. and BUSSEY A.H., 1968: Mechanisms involved in the regulation of nitrogen assimilation in micro-organisms and plants. In: HEWITT E.J. and CUTTING C.V. (eds.), Recent aspects of nitrogen metabolism in plants. Acad. Press, London. 91-114.
- SINCLAIR L.R. and FORBES R.B., 1980: Nutrient removal from drainage waters with systems containing aquatic macrophytes. Trans. Am. Soc. Agric. Engrs. 23, 1189-1194.
- SINGH R.P. and SHARMA V.K., 1975: Caloric values of hydrophytes at different vertical stratification. Geobios 2(1), 26.
- SINGH J.S. and SINGH K.P., 1967: Contribution to the ecology of ten noxious weeds. J. Indian Bot. Soc. 46(4), 440-451.
- SINHA A.B. and SHAI R., 1975: Effect of two herbicides on the control of duck-weeds. Sci. Cult. 41(6), 289-291.
- SIPKO L.L. and DIMITRIEVA S.A., 1971: Higher water vegetation and its fauna of Krotovaya Laga and Kuskan lakes (North Kulanda). (In Russian). Izv. Sib. Otd. Akad. Nauk SSSR Ser. Biol. Med. Nauk 3, 59-62.
- SJOGREN R.D., 1968: Experimental use of Lemna to prevent mosquito breeding in organically polluted waters. Proc. Pap. Annu. Conf. Calif. Mosquito Contr. Ass. Inc. 36, 31-33.
- SKAER H.B., FRANKS F. and ECHLIN P., 1979: Freeze-fracture studies of ice recrystallization in tissue quenched in aqueous solutions of polyvinylpyrrolidone. Cryo. Lett. 1971(1), 61-70.
- SKENDER A. and BERTIC B., 1986: A study on the role of submersed and floating macrophytes in the self-purification process of canals. (In Serbo-Croat.). Jug. Drust. Procv. Subz. Korava. 247-253.
- SKLAR F.H., 1980: A preliminary comparison of the uptake of chromium-51 and zinc-65 by three species of aquatic plants from Louisiana. Proc. La. Acad. Sci. 43, 46-51.
- SKLAR F.H., 1985: Seasonality and community structure of the backswamp invertebrates in a Louisiana cypress-tupelo wetland. Wetlands 5, 69-86.
- SKOGEN D., CHATURVEDI R., WEIDEMANN F. and NILSEN S., 1986: Photoinhibition of photosynthesis: effect of light quality and quantity on recovery from photoinhibition in Lemna gibba. J. Plant. Physiol. 126 (2-3), 195-205.
- SKOOG F. and THIMANN K.V., 1940: Enzymatic liberation of auxin from plant tissues. Science 92, 64.
- SLOOFF W. and CANTON J.H., 1983: Comparison of the susceptibility of 11 freshwater species to 8 chemical compounds. II. (Semi)chronic toxicity tests. Aquat. Toxicol. 4(3), 271-282.
- SLOVIN J.P., 1987: Characterization of a chlorophyll b-less mutant of Lemna gibba G3. Plant Physiol. 83(4 suppl.), 136.
- SLOVIN J.P. and COHEN J.D., 1985a: Selection of mutants in IAA conjugate hydrolysis. J. Cell Biochem. (suppl. 9 part C), 222.
- SLOVIN J.P. and COHEN J.D., 1985b: Analysis of indole-3-acetic acid levels in Lemna gibba G-3 by GC/MS using ^{13}C -IAA as an internal standard. Plant Physiol. 77(4 suppl.), 81.
- SLOVIN J.P. and COHEN J.D., 1985c: Generation of variant lines of Lemna gibba G-3 via tissue culture, for use in selecting auxin metabolism mutants. Plant Physiol. 77(4 suppl.), 11.
- SLOVIN J.P. and COHEN J.D., 1986: Production and mutant lines of Lemna gibba G-3 by cell culture and whole plant mutagenesis. Plant Physiol. 80(4 suppl.), 38.
- SLOVIN J.P. and COHEN J.D., 1987: Production of mutant lines of Lemna gibba G-3 by cell culture and whole plant mutagenesis. J. Cell Biochem. (suppl. 11 part B), 24.
- SLOVIN J.P. and TOBIN E.M., 1980: Effects of glyphosate on synthesis of chloroplast proteins in Lemna gibba L. G-3. Plant Physiol. 65(6 suppl.), 67.
- SLOVIN J.P. and TOBIN E.M., 1981: Glyphosate, a plant growth regulator, affects chloroplast membrane proteins. Biochim. Biophys. Acta 637(1), 173-184.
- SLOVIN J.P. and TOBIN E.M., 1982: Synthesis and turnover of the light-harvesting chlorophyll a/b-protein in Lemna gibba grown with intermittent red light: possible translational control. Planta 154(5), 465-472.
- SLOVIN J.P., BALDI B.G. and COHEN J.D., 1984: Hydrolysis of IAA-conjugates in Lemna gibba G-3. Plant Physiol. 75(1 suppl.), 107.
- SMART C.C. and ESSER R.P., 1968: Aphelenchoides Fragariae in aquatic plants. Plant Disease Reporter 52(6), 455.
- SMART C.C. and TREWAVAS A.J., 1983a: Abscissic-acid-induced turion formation in Spirodela polyrrhiza L. I. Production and development of the turion. Plant Cell Environ. 6(6), 507-514.
- SMART C.C. and TREWAVAS A.J., 1983b: Abscissic-acid-induced turion formation in Spirodela polyrrhiza L. II. Ultrastructure of the turion, a stereological analysis. Plant Cell Environ. 6(6), 515-522.
- SMART C.C. and TREWAVAS A.J., 1984a: Abscissic-acid-induced turion formation in Spirodela polyrrhiza L. III. Specific changes in protein synthesis and translatable RNA during turion development. Plant Cell Environ. 7(2), 121-132.
- SMART C.C. and TREWAVAS A.J., 1984b: Abscissic-acid-induced turion formation in Spirodela polyrrhiza L. IV. Comparative ion flux characteristics of the turion and the vegetative frond and the effect of ABA during early turion development. Plant Cell Environ. 7(7), 521-530.
- SMIRNOVA N.N., 1975a: Effect of exogenous amino acids on the productivity of some higher aquatic plants. (In Russian). Gidrobiol. Zh. 11(4), 47-53.
- SMIRNOVA N.N., 1975b: Effect of exogenous amino acids on the productivity of some higher aquatic plants. Hydrobiol. J. 11(4), 33-39.
- SMIRNOVA N.N., 1976: Amino acids as factors of the nutrition of some higher aquatic plants. (In Russian). Form. Kontrol. Kach. Poverkhn. Vod. 2, 143-148.
- SMIRNOVA N.N., 1981: Role of the higher aquatic plants in the cycles of organic substances of the water. (In Russian). Gidrobiol. Zh. 17(1), 100-101.
- SMITH D.H. and CASTLE J.E., 1960: Production of auxotrophs in a duckweed, Spirodela polyrrhiza. Plant Physiol. 35, 809-815.
- SMITH F.A. and WALTER N.A., 1980: Photosynthesis by aquatic plants. New Phytol. 86, 245-260.
- SMITH J.D., 1980: Wolfia (Wolffia) gladiata, var. Floridana. Bull. Torr. Bot. Club 7, 64-65.
- SMITH N.A. and RIPPINGALE R.J., 1983: Tertiary treatment of wastewater using aquatic plants. Tech. Pap. Fed. Conv., Conf. Proc. Aust. Water Wastewater Assoc., Sydney, 27, 1-9.

- SMITH R. and NIELSEN P.T., 1978: Hormonal effects on the growth and potassium content of *Lemna perpusilla*. *Va.J.Sci.* 29(2), 76.
- SMITH W.L., Jr., and ENNS W.R., 1967: Laboratory and field investigations of mosquito populations associated with oxidation lagoons in Missouri. *Mosq.News* 27(4), 462-466.
- SNELL K., 1907: Untersuchungen über die Nahrungsaufnahme der Wasserpflanzen. *Flora* 98, 213-249.
- SOLOSKI A., 1978: Biological evaluation of the toxicity of Gilow Reservoir waters. (In Polish). *Cuprum* 5(5), 27-29.
- SOLOSKI A., WERNIKOWSKA-UKLEJA A. and MOLTAN B., 1979: Toxicity of flotation reagents to aquatic organisms and their permissible concentration in surface waters. (In Polish). *Cuprum* 6, 32-35.
- SOUÈGES R., 1959: Embryogénie des Lemnacees. *Développement de l'embryon chez le Lemna minor* L. C.R.Acad.Sci.Paris, 248, 1896-1900.
- SOULEN T.K. and KOUKKARI W.L., 1977: Ammonium ion inhibition of flowering in *Lemna perpusilla*, a pH effect. *Plant Physiol.* 59(6 suppl.), 48.
- SPENCER-JONES D.H., 1981: Preliminary studies with PH40:62 - an experimental algalicide. *Proc.Conf.Aquatic Weeds and their Control*. Nat.Veg. Res.Sta., Wellesbourne, Warwick, UK; Assoc. of Applied Biologists. 177-182.
- SPILLER H., EISENMEIER S. and AHMED J., 1973: Activities of nitrate reductase and peroxidase in normal and iron deficient cultures of *Lemna minor*. *Hoppe-Seyler's Z.Physiol.Chem.* 354, 1245.
- SPOHR E., 1926: Ueber das Vorkommen von *Sium erectum* Huds. und *Lemna gibba* L. in Estland und über deren nordöstliche Verbreitungsgrenzen in Europa. *Acta Horti Bot.Univ.Tartuens.* 1(1), 22 pp.
- SPOONER W.E., 1967: Local population variation in *Spirodela polyrrhiza* (L.) Schleid. (Lemnaceae). M.S.Thesis. North Carolina State Univ., Raleigh. 32 pp. (Polycopy).
- SPRANKEL H., 1954: Blühende Wasserlinsen. *Kosmos* 6, 278-279.
- SRINIVASAN J., SOPORY S.K. and MUKHERJEE S., 1975: Photoregulation of nitrate reductase in *Lemna paucicostata*. In: MOTIL V.S. (ed.), Seminar on recent advances in plant sciences. Kalyani, India. Abstr., 38.
- SRINIVASAN R. and VISWAM K., 1986: Laboratory studies on the biology of *Mansonia annulifera* Theobald (1901). *Ind.J.Med.Res.* 83(4), 384-386.
- SRIVASTAVA V.C., 1978: Caloric values of floating plant community of Chilwa lake. *Geobios* 5(4), 174-175.
- STACKEBRANDT E. and WOESE C.R., 1981: The evolution of prokaryotes. In: Molecular and cellular aspects of microbial evolution. Cambridge Univ.Press. 1-31.
- STAFFORD H.A., 1964: Comparison of lignin-like products found naturally or induced in tissues of Phleum, Elodea, and Coleus, and in a paper peroxidase system. *Plant Physiol.* 39, 350-360.
- STAFFORD H.A., 1965: Factors controlling the synthesis of natural and induced lignins in Phleum and Elodea. *Plant Physiol.* 40, 844-851.
- STAHL E., 1983 see 1888
- STAHL E., 1988: Pflanzen und Schnecken. *Jenaische Z.Natw. Med.* 22, N.F. 15, 126 pp.
- STAKE E., 1968: Higher vegetation and phosphorus in a small stream in Central Sweden. *Schweiz.Z.Hydrol.* 30, 353-373.
- STANGENBERG M., 1967: Bacteriostatic effects of some Algae and Lemna minor extracts. *Hydrobiol.* 32, 88-96.
- STARIEWSKA-ZATEK W., 1964: New localities of *Wolffia arrhiza* Wimmer in the valley of the Sarna river near Poznan. (In Polish). *Przycz. Polski Zach.* 8, 86-89.
- STANLEY R.A., 1980: Methods of biological recycling of nutrients from livestock wastes: a literature review and systems analysis. Tennessee Valley Authority, National Fertilizer Development Center, 1-45. (Polycopy).
- STANLEY R.A. and MADEWELL C.E., 1975: Optimum dilution of swine wastes for growth of *Lemna minor* and *Euglena* sp. In: *Managing Livestock wastes*. Am.Soc.Agric.Engineers, St.Joseph, Michigan. 321-323.
- STANLEY R.A. and MADEWELL C.E., 1976a: Chemical tolerance of *Lemna minor* L. Tenn.Valley Authority, Muscle Shoals, Al. 17 pp. (Polycopy).
- STANLEY R.A. and MADEWELL C.E., 1976b: Thermal tolerance of *Lemna minor* L. Tenn.Valley Authority, Muscle Shoals, Al. 16 pp. (Polycopy).
- STANLEY T.D., 1979: Aquatic plants of Queensland part 3. The duckweeds of Queensland. *Queensland Agric.J.* 105(6), 570-572.
- STARFINGER U., 1983: Die Pleustophytenvegetation der Berliner Pfuhle in Berlin, Fachbereich Biologie. 104 pp. (Polycopy).
- STARFINGER U., 1985: Die flache Form von *Lemna gibba*. *Verh.Berl.Bot.Ver.* 4, 67-77.
- STATSENKO L.P., 1978: Control of mosquito hatching using the herbicide simazine. (In Russian). *Med.Parazitol.Parazit.Bolezni* 47, 93-95.
- STAVES R.P., 1980: Chromium uptake and its effects on the growth of duckweeds. M.S.Thesis. Louisiana State Univ., Baton Rouge. 101 pp. (Polycopy).
- STAVES R.P. and KNAUS R.M., 1985: Chromium removal from water by three species of duckweeds. *Aquat.Bot.* 23(3), 261-273.
- STEEMANN NIELSEN E., 1944: Dependence of freshwater plants on quality of carbon dioxide and hydrogen ion concentration. *Dansk Bot.Ark.* 11 (8), 25 pp.
- STEEMANN NIELSEN E., 1947: Photosynthesis of aquatic plants with special reference to the carbon sources. *Dansk Bot.Ark.* 12(8), 71 pp.
- STEEMANN NIELSEN E., 1954: On the reference of some freshwater plants in Finland for brackish water. *Bot.Tidskr.* 51, 242-247.
- STEINBACK K. and WATSON J., 1981: In vivo chloroplast thylakoid protein phosphorylation. *Plant Research Ann.Rep., Michigan State Univ., MSU/DOE. Plant Res.Lab.* '81, 32-33.
- STEINBACK K.E. and WATSON J., 1982: In vivo control of thylakoid protein phosphorylation. *Plant Physiol.* 69(4 suppl.), 29.
- STEINBERG R.A., 1941: Use of Lemna for nutrient studies on green plants. *J.Agric.Res.* 62, 423-430.
- STEINBERG R.A., 1943: Use of Lemna as a test organism. *Chronica Botanica* 7, 420.
- STEINBERG R.A., 1946: Mineral requirements of *Lemna minor*. *Plant Physiol.* 21, 42-48.
- STELZ T. and THELLIER M., 1971: Formulation électrocinétique de l'absorption du calcium par des végétaux calcicoles et calcifuges. *C.R.Acad.Sci.Paris*, D 272, 1101-1104.
- STELZ T., AYADI A., MONNIER A., DEMARTY M. and THELLIER M., 1975: Echange de calcium dans les espaces libres cellulaires de végétaux. *C.R.Soc.Biol.Rouen* 169, 1072-1076.
- STELZIG V. and BERNING A., 1984: Ein neues Vorkommen der Zwerglinse (Wolffia arrhiza [L.] Wimm.) im südlichen Emsland. *Natur und Heimat* 44(2), 54-55.
- STEPANOVA V.S., 1928: Influence of Lemna covering on a water basin. (In Russian). *Trav.Soc.Nat. Leningrad* 58, 63-82.
- STEPANOVA V.S., 1932: Die Geschwindigkeit des vegetativen Wachstums der Gattung Lemna. (In Russian). *Bot.Zh.(USSR)* 17, 524-529.

- STEPHENSON M., TURNER G., POPE P., COLT J., KNIGHT A. and TCHOBANOGLOUS G., 1980: The use and potential of aquatic species for wastewater treatment. Appendix A. The Environmental requirements of aquatic plants. California State Water Resources Control Board, 65, 291-440.
- STEPHENSON R.R. and KANE D.F., 1984: Persistence and effects of chemicals in small enclosures in ponds. Arch. Environ. Contam. Toxicol. 13, 313-326.
- STERKI V., 1910: Winter-buds of *Spirodela polyrrhiza* (L.). Ohio Naturalist 10(8), 181-182.
- STERN H.L., ARMSTRONG D.A., KNIGHT A.W. and CHIPPENDALE D.J., 1976: Survival and growth of juveniles of the giant Malaysian prawn, *Macrobrachium rosenbergii*, fed natural plant diets. Proc. 7th Annu. Meet. World Maricul. Soc. 1976, 667.
- STERRY P.R., THOMAS J.D. and PATIENCE R.L., 1983: Behavioral responses of *Biomphalaria glabrata* to chemical factors from aquatic macrophytes including decaying *Lemna paucicostata*. Freshwater Biol. 13(5), 465-476.
- STERRY P.R., THOMAS J.D. and PATIENCE R.L., 1985: Changes in the concentrations of short-chain carboxylic acids and gases during decomposition of the aquatic macrophytes, *Lemna paucicostata* and *Ceratophyllum demersum*. Freshwater Biol. 15(2), 139-153.
- STEBING L., RAMIREZ C. and ALBERDI M., 1980: Energy content of water- and bog-plant associations in the region of Valdivia (Chile). Vegetatio 43, 153-161.
- STEWART G.R., 1968: The effect of cycloheximide on the induction of nitrate and nitrite reductase in *Lemna minor* L. Phytochem. 7, 1139-1142.
- STEWART G.R., 1969: Abscisic acid and morphogenesis in *Lemna polyrrhiza* L. Nature 221, 61-62.
- STEWART G.R., 1972a: End-product repression of nitrate reductase in *Lemna minor* L. In: FARAS G.L. (ed.), Nucleic acids and proteins in higher plants. Akad. Kiado, Budapest. Symp. Biol. Hung. 13, 127-135.
- STEWART G.R., 1972b: The regulation of nitrite reductase level in *Lemna minor* L. J. Exp. Bot. 23, 171-183.
- STEWART G.R. and RHODES D., 1976: Evidence for the assimilation of ammonia via the glutamine pathway in nitrate-grown *Lemna minor* L. FEBS Lett. 64, 296-299.
- STEWART G.R. and RHODES D., 1977a: A comparison of the characteristics of glutamine synthetase and glutamate dehydrogenase from *Lemna minor* L. New Phytol. 79, 257-268.
- STEWART G.R. and RHODES D., 1977b: Control of enzyme levels in the regulation of nitrogen assimilation. In: SMITH H. (ed.), Regulation of enzyme synthesis and activity in higher plants. Acad. Press, London/ New York. 1-22.
- STEWART G.R. and SIMS A.P., 1967: Adaptive response of the enzyme malic dehydrogenase from cells of *Lemna minor* grown on different levels of calcium ions. SERAVAC.
- STEWART G.R. and SMITH H., 1972: Effects of abscisic acid on nucleic acid synthesis and the induction of nitrate reductase in *Lemna polyrrhiza*. J. Exp. Bot. 23, 875-885.
- STEWART G.R., LEE J.A., OREBAMJO T.O. and HAVILL D.C., 1974: Ecological aspects of nitrogen metabolism. In: BIELESKI R.L., FERGUSON A.L. and CRESSWELL M.M. (eds.), Mechanisms of regulation of plant growth. R. Soc. New Zealand, Wellington. Bull. 12, 41-47.
- STEWART R.R., 1972: Flora of West Pakistan. Fakhri Print. Press, Karachi, Pakistan. 36.
- STEWART W.C. and SERFLING S.A., 1979: The solar aquacell for primary, secondary or advanced treatment of wastewaters. Aquaculture systems for wastewater treatment. U.S. Environ. Prot. Agency, Publ. MCD-67 (EPA 430/9-80-006).
- STEYERMARK J., 1975: Flora of Missouri. (4th ed.). The Iowa State Univ. Press, Ames. 1, 386-391.
- STICKNEY R.R., ROWLAND L.O. and HESBY J.H., 1977: Water quality - *Tilapia* aurea interactions in ponds receiving swine and poultry wastes. Proc. 8th Annu. Meet. World Maricul. Soc. 1977, 55-71.
- STIEKEMA W.J., WINPEE C.F., SILVERTHORNE J. and TOBIN E.M., 1983a: Phytochrome control of the expression of two nuclear genes encoding chloroplast proteins in *Lemna gibba* L. G-3. Plant Physiol. 72(3), 717-724.
- STIEKEMA W.J., WINPEE C.F. and TOBIN E.M., 1983b: Nucleotide sequence encoding the precursor of the small subunit of ribulose 1,5-diphosphate carboxylase from *Lemna gibba* L. G-3. Nucleic Acids Res. 11(22), 8051-8061.
- STOECKLI B., BRAENDLE R. and ERISMANN K.H., 1975: Mitoseaktivität und Zellzyklus unter Begasung mit subletalen SO₂-Konzentrationen bei der Wasserlinse (*Lemna minor* L.). Experientia 31, 795-796.
- STOTANOFF N. and STEFANOFF B., 1924: Flora of Bulgaria. (In Bulgar.). Sofia. 204-206.
- STOM D.I. and ROTH R., 1981: Some effects of polyphenols on aquatic plants. I. Toxicity of phenols in aquatic plants. Bull. Environ. Contam. Toxicol. 27(3), 332-337.
- STOM D.I., GIL T.A. and BALAYAN A.E., 1986: Effect of individual phenolic compounds and of their mixtures on luminous bacteria. Part 3. Combined action of phenolic compounds on aquatic organisms. Acta Hydrochim. Hydrobiol. 14(5), 539-549.
- STONE W., 1911: The plants of southern New Jersey with a special reference to the flora of the Pine Barrens. Ann. Rep. New Jersey State Mus. 25, 310-319.
- STRACK D. and KRAUSE J., 1978: Reversed phase high performance liquid chromatographic separation of naturally occurring mixtures of flavone derivatives. J. Chromatogr. 156(2), 359-361.
- STRANSBAUGH P.D. and CORE E.L., 1952: Flora of West Virginia 1. West Virginia Univ. Bull. 52, 208-209.
- STRASBURGER E., NOLL F., SCHENCK H. and SCHIMPER A.F.W., 1978: Lehrbuch der Botanik. (31st ed.). Fischer, Stuttgart. 1178 pp.
- STRASBURGER K., 1981: Wasserpflanzengesellschaften im unteren Allertal. Ph.D. Thesis. Univ. Hannover. 207 pp.
- STRASBURGER K. and HOMANN J., 1982: Gesellschaften der Lemnetaalia im Meissendorfer Fischteichgebiet westlich von Celle. Tuexenia 2, 27-29.
- STRASSER R.J., 1971: Eine einfache Anlage zur kontinuierlichen Kultivierung von Lemnaceen mit automatischer Probenentnahme. Photosynthetica 5, 76-78.
- STRASSER R.J., 1974: Studies on the oxygen evolving system in flashed leaves. In: AVRON M. (ed.), Proc. 3rd Int. Congr. Photosynthesis (Rehovot, Israel). Elsevier, Amsterdam. 1, 497-503.
- STRASSER R.J., ERISMANN K.H. and METZNER H., 1970: Die Photooxidation von Sulfid durch Grana. Verh. Schweiz. Naturf. Ges. 150, 235-238.
- STRAUSS R., 1973: Influence de la concentration en alcalino-terreux du milieu de culture sur la croissance et le métabolisme de *Lemna minor* L. C. R. Soc. Biol. (Rouen) 167, 827-830.
- STRAUSS R., 1976: Effet de divers sels alcalins sur la croissance et la

- nutrition minérale de Lemna minor L. Int.Rev.Ces.Hydrobiol. 61(5), 673-676.
- STRAWINSKAJA E.A. (ed.), 1984: Conservation of aquatic ecosystems in the urbanized landscape. (In Russian). Nauka, Leningrad. 144 pp.
- STROTHER S., 1979: Hydrolysis of α -D-glucuronate-1-phosphate by extracts from Lemna minor L. Phytochemistry 18, 486.
- STROTHER S., 1981: Toxic effects of exogenous sorbose on Lemna minor and some other angiosperms. Ann.Bot. 47(4), 531-533.
- STROTHER S., 1984: Control of phosphatase activities of duckweed by senescence and phosphorus deficiency. Biol.Plant. 26(4), 309-311.
- STRUGGER S., 1934: Beiträge zur Physiologie des Wachstums. I. Zur protoplasmatisch-physiologischen Kausalanalyse des Streckungswachstums. Jb.Wiss.Bot. 79, 406-471.
- STUCKEY R.L., 1973: Bibliography of aquatic flowering plants of Ohio. Ohio Biol.Surv.Inform.Circ. 2, 11 pp.
- STUCKEY R.L., 1976: Aquatic vascular plants of the Sandusky River Basin. In: BAKER D.B., JACKSON W.B. and PRATER B.L. (eds.), Proc.Sandusky River Basin Int.Symp. Reference group Great Lakes pollution from land use activities. U.S.Gov.Printing Office. 295-333.
- STUCKEY R.L. and ROBERTS M.L., 1976: Rare and endangered vascular plants of Ohio. Sida 7(1), 32.
- SU K.L. and STABA E.J., 1972: Aquatic plants from Minnesota. I. Chemical survey. Univ.Minn.Water Res.Center Bull. 46, 50 pp.
- SU K.L. and STABA E.J., 1973: Toxicity, antineoplastic and coagulation effects of aquatic plants from Minnesota. Lloydia 36(1), 99-102.
- SU K.L., STABA E.J. and ABUL-HAJJ Y., 1973a: Preliminary chemical studies of aquatic plants from Minnesota. Lloydia 36, 72-79.
- SU K.L., ABUL-HAJJ Y. and STABA E.J., 1973b: Antimicrobial effects of aquatic plants from Minnesota. Lloydia 36, 80-87.
- SU M.T. and ASHBY E., 1929: The interaction of factors in the growth of Lemna. II. Technique for the estimation of dry weight. Ann.Bot. 43, 329-332.
- SUCKCHAROEN S., 1980: Mercury contamination of terrestrial vegetation near a caustic soda factory in Thailand. Bull.Environ.Contam.Toxicol. 24(3), 463-466.
- SUDZUKI M., WATANABE K., SUZUKI K. and NARITA K., 1983: Occurrence of Rotifera in the field under natural and intentionally changed conditions. Hydrobiologia 104, 341-347.
- SUN S., WANG H., LI Q., 1985: Preliminary studies on physiological changes and injury mechanism in aquatic vascular plants treated with cadmium. (In Chinese). Zhiwu Shengli Xuebao 11(2), 113-121.
- SUNESON S., 1959: Lemna gibba in Bohuslän. (In Swed.). Svensk.Bot.Tidskr. 53, 287-292.
- SUPNIEWSKA J.H., 1963: Observations on the action of trimethyl- β -chloroethyl-ammonium chloride on plants. I. Lemna minor L., Chlorella pyrenoidosa Prings., Riccia fluitans L. Bull.Acad.Pol.Sci.Ser.Sci.Biol. 11, 149-154.
- SUTANTO L., WIDYANTO L.S. and SOERJANI M., 1976: Tropical pest biology. 1. The effect of ametryne and cyanatryn on water hyacinth control. Biotrop Newslett. 15, 13.
- SUTER M., LAVANCHY P., VON ARB C. and BRUNOLD C., 1986: Regulation of sulfate assimilation by amino-acids in Lemna minor. Plant Sci. 44(2), 125-132.
- SUTHERLAND J.C. and BEVIS F.B., 1979: Reuse of municipal wastewater by volunteer freshwater wetlands. Rep.pref.Nat.Sci.Found., Washington. (Polycopy).
- SUTTON D.L., 1976: Utilization of duckweed by the white amur. In: FREE-MAN T.E. (ed.), Proc.4th Int.Symp.Biol.Control Weeds, 257-260.
- SUTTON D.L., 1981a: Who needs aquatic weeds? Weeds Today 1981, 6-7.
- SUTTON D.L., 1981b: Azolla. Aquatics 3(2), 4-5.
- SUTTON D.L. and ORNES W.H., 1975: Phosphorus removal from static sewage effluent using duckweed. J.Environ.Qual. 4, 367-370.
- SUTTON D.L. and ORNES W.H., 1977: Growth of Spirodela polyrrhiza in static sewage effluent. Aquat.Bot. 3, 231-237.
- SUTTON D.L., DURHAM A.D., BINGHAM S.W. and FOY C.L., 1969: Influence of simazine on apparent photosynthesis of aquatic plants and herbicide residue removal from water. Weed Sci. 17(1), 56-59.
- SUTTON D.L., BLACKBURN R.D. and STEWARD K.K., 1976: Control of aquatic plant growth. Annu.Res.Rep.IFAS, Fort Lauderdale, FL, USA, 365.
- SUTTON D.L., STEWARD K.K., PERKINS B.D. and VANDIVER V.V., 1977: Control of aquatic plant growth. Annu.Res.Rep.IFAS, Fort Lauderdale, FL, USA, 324-325.
- SUZUKI Y., YAMAGUCHI I. and TAKAHASHI N., 1985: Isolation and characterization of a flower inducing substance from Pharbitis purpurea. Abstr.12th Int.Conf.Plant Growth Substances, Heidelberg, FRG. 96.
- SWADER J.A., 1970: Nitrate uptake and reduction in Wolffia arrhiza. Ph.D.Thesis, Univ. Ann Arbor, Michigan. 92 pp.
- SWADER J.A. and STOCKING C.R., 1971: Nitrate and nitrite reduction by Wolffia arrhiza. Plant Physiol. 47, 189-191.
- SWADER J.A., STOCKING C.R. and LIN C.H., 1975: Light-stimulated absorption of nitrate by Wolffia arrhiza. Physiol.Plant. 34, 335-341.
- SYMONEIS J.J., HOPPER S.S. and COMPERE P. (eds.), 1982: Studies on aquatic vascular plants. R.Bot.Soc.Belgium, Brussels. 424 pp.
- SYTNYK K.M. and KORDYUM Y.L., 1980: Space botanical research. (In Russian). Ukr.Bot.Zh. 37(1), 1-10.
- SZABADOS M., HORVATH A. and CSEH E., 1983: Application of Lemna minor L. to testing the arsenic content of the soil. (In Hung.). Bot.Koezl. 70 (3-4), 171-177.
- SZALMA E. and BODROGKOZGY G., 1985: Phytocenology of Wolffia arrhiza Miyaw. et J.Tx.60. Element content of its species components as well as sediment- and water-samples. Tiscia 20, 45-53.
- SZUNAN J. and SKRZYDLEWSKI A., 1980: Land- und Wasserpflanzen in der Sumpfbibernalung. Deutsch.Pelztierzüchter 54(7), 107-111.
- TACHIMOTO M., KOBAYASHI M. and TAKAHASHI E., 1986: Bioassay of decomposed products of rapeseed cake using Lemna paucicostata 6746. Soil Sci.Plant Nutr. 32(2), 223-231.
- TAECKHOLM V., 1974: Students' Flora of Egypt. Lemnaceae. Cairo Univ., Beirut. 768-769.
- TAI-HSING H. et al., 1975: The growing of green manure. (In Chinese). Chiang-su jen min chiu pan she. 15 pp.
- TAKAOKI T., 1969: Measurement of osmotic quantities in higher plants. J.Sci.Hiroshima Univ., Ser.B, Div.2, 12, 199-210.
- TAKEMOTO B.K., 1986: Differential sensitivity of duckweeds to sulfite enrichment. Physiological and growth responses. Ph.D.Thesis. Bowling Green State Univ. 85 pp. Diss.Abstr.Int.B 47(2), 480.
- TAKEMOTO B.K. and NOBLE R.D., 1984: Differential sensitivity to sulfite enrichment by three species of duckweed (Lemnaceae). Plant Physiol. 75(1 suppl.), 38.
- TAKEMOTO B.K. and NOBLE R.D., 1986: Differential sensitivity of duckweeds (Lemnaceae) to sulphite. I. Carbon assimilation and frond replication rate as factors influencing sulphite phytotoxicity under low and high irradiance. New Phytol. 103(3), 525-539.

- TAKEMOTO B.K., NOBLE R.D. and HARRINGTON H.M., 1986: Differential sensitivity of duckweeds (Lemnaceae) to sulphite. II. Thiol production and hydrogen sulphide emission as factors influencing sulphite phytotoxicity under low and high irradiance. *New Phytol.* 103, 541-548.
- TAKIMOTO A., 1973: Flower initiation of *Lemna perpusilla* under continuous low-intensity light. *Plant Cell Physiol.* 14, 1217-1219.
- TAKIMOTO A., 1981: Duckweeds as system to investigate flowering. *Proc. Int. Bot. Congr.* 13, 50.
- TAKIMOTO A., 1987: Comparative studies on the photoperiodic and chemical induction of flowering in *Lemna*. 14th Int. Bot. Congr. Berlin, Abstr., 105.
- TAKIMOTO A. and KAIHARA S., 1986: The mode of action of benzoic acid and some related compounds on flowering in *Lemna paucicostata*. *Plant Cell Physiol.* 27(7), 1309-1316.
- TAKIMOTO A. and TANAKA O., 1973: Effects of some SH-inhibitors and EDTA on flowering in *Lemna perpusilla* 6746. *Plant Cell Physiol.* 14, 1133-1141.
- TAKIMOTO A. and TANAKA O., 1974: Effects of some sulphydryl inhibitors on floral initiation under various light conditions in *Lemna perpusilla*. In: *Plant growth substances*, 1973. Hirokawa Publ., Tokyo. 953-959.
- TAKIMOTO A. and TANAKA O., 1976: Long-day flowering of *Lemna perpusilla* 6746 in Mo-deficient medium. *Plant Cell Physiol.* 17, 299-303.
- TAKIMOTO A., KAIHARA S. and NISHIOKA H., 1987: A comparative study on the short-day and the benzoic acid-induced flowering in *Lemna paucicostata*. *Plant Cell Physiol.* 28(3), 503-508.
- TAMOT B.K. and MAHESHWARI S.C., 1987: Photoperiodic and chemical control of flowering in *Wolffia hyalina*, 7378. 14th Int. Bot. Congr. Berlin, Abstr., 130.
- TAMOT B.K., KHURANA J.P. and MAHESHWARI S.C., 1987: Obligate requirement of salicylic acid for short-day induction of flowering in a new duckweed, *Wolffia hyalina* 7378. *Plant Cell Physiol.* 28(2), 349-353.
- TAL S. and ZIV I., 1978: Culture of exotic species in Israel. *Bamidgeh* 30, 3-11.
- TAN Y., 1970: Composition and nutritive value of some grasses, plants and aquatic weeds tested as diets. *J. Fish. Biol.* 2, 253-257.
- TANAKA H., 1966: Response of *Lemna paucicostata* to boron as affected by light intensity. *Plant and Soil* 25, 425-434.
- TANAKA O., 1986: Flower induction by nitrogen deficiency in *Lemna paucicostata* 6746. *Plant Cell Physiol.* 27(5), 875-880.
- TANAKA O. and ASAGAMI K., 1986: Ferricyanide induces flowering by suppression of nitrate assimilation in *Lemna paucicostata* 6746. *Plant Cell Physiol.* 27(6), 1063-1068.
- TANAKA O. and CLELAND C.F., 1978: Influence of ferricyanide on flowering in *Lemna*. *Plant Physiol.* 61(4 suppl.), 52.
- TANAKA O. and CLELAND C.F., 1980: Comparison of the ability of salicylic acid and ferricyanide to induce flowering in the long-day plant, *Lemna gibba* G3. *Plant Physiol.* 65, 1058-1061.
- TANAKA O. and CLELAND C.F., 1981: Influence of ammonium on the ability of salicylic acid to induce flowering in the short-day plant *Lemna paucicostata* 6746. *Plant Cell Physiol.* 22(4), 597-602.
- TANAKA O. and TAKIMOTO A., 1975: Suppression of long-day flowering by nitrogenous compounds in *Lemna perpusilla* 6746. *Plant Cell Physiol.* 16, 603-610.
- TANAKA O. and TAKIMOTO A., 1977: Flower-promoting effect of some amino acids and amides in *Lemna paucicostata* 6746. *Plant Cell Physiol.* 18, 27-34.
- TANAKA O. and TAKIMOTO A., 1978: Effect of nitrate concentration in the medium on the flowering of *Lemna paucicostata* 6746. *Plant Cell Physiol.* 19, 701-704.
- TANAKA O., CLELAND C.F. and HILMAN W.S., 1979a: Inhibition of flowering in the long-day plant *Lemna gibba* G3 by Hutner's medium and its reversal by salicylic acid. *Plant Cell Physiol.* 20, 839-846.
- TANAKA O., TAKIMOTO A. and CLELAND C.F., 1979b: Enhancement of long-day flowering by Mo-deficiency and application of some amino acids and asparagine in the short-day plant *Lemna paucicostata* 6746. *Plant Cell Physiol.* 20, 267-270.
- TANAKA O., NASU Y., TAKIMOTO A. and KUGIMOTO M., 1982a: Absorption of copper by *Lemna* as influenced by some factors which nullify the copper effect on flowering and growth. *Plant Cell Physiol.* 23(7), 1291-1296.
- TANAKA O., NASU Y., YANASE D., TAKIMOTO A. and KUGIMOTO M., 1982b: pH dependence of the copper effect on flowering, growth and chlorophyll content in *Lemna paucicostata* 6746. *Plant Cell Physiol.* 23(8), 1479-1482.
- TANAKA O., CLELAND C.F. and BEN-TAL Y., 1983: Effect of ferricyanide, ferrocyanide and KCN on growth and flowering in the short-day plant *Lemna paucicostata* 6746. *Plant Cell Physiol.* 24(4), 705-711.
- TANAKA O., HORIKAWA W., NISHIMURA H. and NASU Y., 1986a: Flower induction by suppression of nitrate assimilation in *Lemna paucicostata* 6746. *Plant Cell Physiol.* 27(1), 127-133.
- TANAKA O., YOKOYAMA T., KONO Y. and NASU U., 1986b: Measurement of endogenous phosphorus levels in relation to flowering in the long-day plant *Lemna gibba* G3. *Plant Cell Physiol.* 27(7), 1277-1283.
- TANAKA O., NASU Y., SONOYAMA A., MAEHARA Y., KOBAYASHI T., NAWAFUNE H. and KUGIMOTO M., 1987: Effects of exogenous amino acids on iron uptake in relation to their effects on photoperiodic flowering in *Lemna paucicostata* 6746. *Plant Cell Physiol.* 27(7), 1277-1283.
- TANNAVACH I.T. and NEDELIC G.A., 1971: Concerning the dynamics of lake vegetation. *Commun. Bot.* 12, 289-295.
- TARVER D.P. et al., 1978: Aquatic and wetland plants of Florida. Tallahassee, FL: Florida Dept. Natural Resources. 127 pp.
- TASSERON-DE JONG J.G., 1968: Investigations on cytokinins. Effects of 6-benzylaminopurine on *Lemna minor* L. Thesis. Univ. Leiden.
- TASSERON-DE JONG J.G. and VELDSTRA J.H., 1971a: Investigations on cytokinins. 1. Effect of 6-benzylaminopurine on growth and starch content of *Lemna minor*. *Physiol. Plant.* 24, 235-238.
- TASSERON-DE JONG J.G. and VELDSTRA J.H., 1971b: Investigations on cytokinins. 2. Interaction of light and cytokinins as studied in *Lemna minor*. *Physiol. Plant.* 24, 239-241.
- TATKOWSKA E. and BUCZEK J., 1983: Effect of ammonium nutrition on the nitrate utilization, nitrate reductase activity and growth of *Spirodela polyrrhiza*. *Acta Soc. Bot. Pol.* 52(3-4), 241-252.
- TATKOWSKA E. and KOBYLANSKA D., 1978: The effect of sodium humate on cultures of *Spirodela polyrrhiza* (L.) Schleiden under aseptic conditions. *Ekol. Pol.* 26, 213-220.
- TATKOWSKA E. and TOPOROWSKA E., 1978: The effects of detergents on cultures of *Spirodela polyrrhiza* (L.) Schleiden under aseptic conditions. *Ekol. Pol.* 26, 221-229.
- TAUBAEV T.T. and ABDIEV M., 1971: The research on duckweeds under natural and cultivation conditions. (In Russian). *Kultiv. Vodn. Vyssh. Vodn. Rast. Uzbekistane*, 92-98.
- TAUBAEV T. and ABDIEV M., 1973: Duckweeds of water reservoirs in Uzbek-

kistan and their use in agriculture. (In Russian). Fan, Taskent. 87 pp.

TAUBAEV T.T., ABDIEV M. and KELDIBEKOV C., 1971a: The biological productivity of lesser duckweed (Lemna minor L.) in natural water bodies and in culture. (In Russian). Kultiv.Vodor.Vyssh.Vodn.Rast.Uzbekistane, 98-101.

TAUBAEV T.T., NESKUBO P.M., ABDIEV M. and NORMUCHAMEDOV C., 1971b: Use of Lemna minor as fodder for small cattle. (In Russian). In: TKACENKO I.P. (ed.), Cultivation of algae and water macrophytes in Uzbekistan. Taskent. 136-138.

TAUBAEV T.T., RAKHIMOVA L. and RAKHIMOVA S., 1982: Biological characteristics of cultured Spirodela polyrrhiza and Lemna minor. (In Russian). Sist.Ekol.Vodor.Gribov.Sred.Azii, Tashkent, 49-57.

TAYLOR J.S. and DUMEROFF E.B., 1975: Bud, root, and growth-regulator activity in Acer saccharum during the dormant season. Can.J.Bot. 53, 321-331.

TEIXEIRA A.R.N. and DAVIES D.D., 1974: The control of plant glutamate dehydrogenase by pyridoxal-5'-phosphate. Phytochem. 13, 2071-2080.

THANIKAIMONI G., 1969: Esquisse palynologique des Aracées. Trav.Sect. Sci.Tech.Inst.Fr.Pondichery 5(5), 31 pp.

THAYER D.D. and HALLER W.T., 1985: Effect of herbicides on floating aquatic plants. J.Aquat.Plant Manage. 23, 94-95.

THELLIER M., 1970: An electrokinetic interpretation of the functioning of biological systems and its application to the study of mineral salts absorption. Ann.Bot. 34, 983-1009.

THELLIER M., 1973: Electrokinetic formulation of ionic absorption by plant samples. In: ANDERSON W.P. (ed.), Ion transport in plants. Acad.Press, London/New York. 47-63.

THELLIER M. and AYADI A., 1967: Contribution à l'étude d'interactions ioniques phosphate-borate lors de l'absorption par la Lemna minor L. C.R.Acad.Sci.Paris, D 265, 1940-1943.

THELLIER M. and AYADI A., 1968: Interprétation électrocinétique de l'absorption du phosphate par la Lemna minor. C.R.Acad.Sci.Paris, D 267(22), 1839-1842.

THELLIER M. and AYADI A., 1971: Application of the electrokinetic formula to the interpretation of the regulation of overall cellular changes of potassium by calcium. In: BRODA E. et al. (eds.), First European Biophysics Congress, 14th to 17th September, 1971, Baden near Vienna, Austria. 211-215.

THELLIER M. and LE GUIEL J., 1967a: Etude, grâce à l'isotope stable ^{10}B , de l'absorption du borate par la Lemna minor L. C.R.Acad.Sci.Paris, D 264, 292-295.

THELLIER M. and LE GUIEL J., 1967b: Influence de la nutrition en bore sur l'incorporation de la leucine dans les protéines chez la Lemna minor L. C.R.Acad.Sci.Paris, D 264, 1848-1851.

THELLIER M. and TRONCOUR C., 1968: Contribution à l'étude d'interactions ioniques sulfate-borate lors de l'absorption par la Lemna minor L. C.R.Acad.Sci.Paris, D 266(2), 99-101.

THELLIER M., AYADI A. and TRONCOUR C., 1967: Contribution à l'étude d'interactions ioniques chlorure-borate et iode-borate lors de l'absorption par la Lemna minor L. C.R.Acad.Sci.Paris, D 265(22), 1687-1690.

THELLIER M., STELZ Th. and AYADI A., 1971a: Application des relations et de la loi d'Onsager à l'interprétation d'interactions entre processus cellulaires globaux. C.R.Acad.Sci.Paris, D 273, 2346-2349.

THELLIER M., THOIRON B. and THOIRON A., 1971b: Electrokinetic formula-

tion of overall kinetics of in vivo processes. Physiol.Vég. 9(1), 65-82.

THELLIER M., DUVAL Y. and DEMARTY M., 1979: Borate exchanges of Lemna minor L. as studied with the help of the enriched stable isotopes and of a (n, α) nuclear reaction. Plant Physiol. 63, 283-288.

THELLIER M., RIPOLL C., DEMARTY M. and LASSALLE J.-P., 1981: Thermodynamic model of ion transport: flux/force relationships and biological implications. 6th School on Biophys.Membr.Transp., Proc. 2, 31-57.

THERTOT R.F. and SANDERS D.R., Sr., 1975: Food preferences of yearling hybrid carp. Hyacinth Control J. 13, 51-53.

THIMANN K.V. and EDMONDSON Y.H., 1949: The biogenesis of the anthocyanins. I. General nutritional conditions leading to anthocyanin formation. Arch.Biochem. 22, 33-53.

THIMANN K.V. and RADNER B.S., 1955a: The biogenesis of anthocyanins. IV. The inhibitory effect of methionine and other sulfur-containing compounds on anthocyanin formation. Arch.Biochem.Biophys. 58, 484-497.

THIMANN K.V. and RADNER B.S., 1955b: The biogenesis of anthocyanins. V. Evidence for the mediation of pyrimidines in anthocyanin synthesis. Arch.Biochem.Biophys. 59, 511-525.

THIMANN K.V. and RADNER B.S., 1958: The biogenesis of anthocyanins. VI. The role of riboflavin. Arch.Biochem.Biophys. 74, 209-223.

THIMANN K.V. and RADNER B.S., 1960: Control of anthocyanin formation by purines and pyrimidines. Plant Physiol. 25(suppl.), 17.

THIMANN K.V. and RADNER B.S., 1962: The biogenesis of anthocyanins. VII. The requirement for both purines and pyrimidines. Arch.Biochem. 96, 270-279.

THIMANN K.V. and SKOOG F., 1940: The extraction of auxin from plant tissues. Am.J.Bot. 27, 951-960.

THIMANN K.V., SKOOG F. and BYER A.C., 1942: The extraction of auxin from plant tissues II. Am.J.Bot. 29, 598-606.

THIMANN K.V., EDMONDSON Y.H. and RADNER B.S., 1951: The biogenesis of anthocyanins. III. The role of sugars in anthocyanin formation. Arch.Biochem. 34, 305-323.

THOIRON A., THOIRON B., DEMARTY M. and THELLIER M., 1981: Compartmental analysis of sulfate transport in Lemna minor L. taking plant growth and sulfate metabolism into consideration. Biochim.Biophys.Acta 644 (1), 24-35.

THOIRON B., THOIRON A. and THELLIER M., 1969: Influence de la température sur l'absorption du sulfate par la Lemna minor L. C.R.Acad.Sci.Paris, D 269(21), 2103-2106.

THOMAS J.D., GREALY B. and FENNELL C.F., 1983: The effects of varying the quantity and quality of various plants on feeding and growth of Biomphalaria glabrata (Gastropoda). Oikos 41(1), 77-90.

THOMAS J.D., STERRY P.R. and PATIENCE R.L., 1984: Uptake and assimilation of short carboxylic acids by Biomphalaria glabrata (Say), the freshwater pulmonate snail host of Schistosoma mansoni (Sambon). Proc.R.Soc.Lond. B 222, 447-476.

THOMPSON C.H., 1896: The ligulate Wolffias of the United States. Ann. Rep.Miss.Bot.Gard. 7, 101-111.

THOMPSON C.H., 1898: A Revision of the American Lemnaceae occurring North of Mexico. Ann.Rep.Miss.Bot.Gard. 9, 21-42.

THOMPSON G.A., MUDD S.H., DATKO A.H. and GIOVANELLI J., 1980: Regulation of cystathionine γ -synthase and O-phosphohomoserine sulphydrylase in Lemna. Plant Physiol. 65(6 suppl.), 16.

THOMPSON G.A., DATKO A.H., MUDD S.H. and GIOVANELLI J., 1982a: Methionine biosynthesis in Lemna. Studies on the regulation of cysta-

- thionine- δ -synthase, O-phosphohomoserine sulphydrylase, and O-acetylserine sulphydrylase. *Plant Physiol.* 69(5), 1077-1083.
- THOMPSON G.A., DAYKO A.H. and MUDD S.H., 1982b: Methionine synthesis in Lemna. Inhibition of cystathionine δ -synthase by propargylglycine. *Plant Physiol.* 70(5), 1347-1352.
- THOMPSON G.A., DAYKO A.H. and MUDD S.H., 1983: Adaptation of Lemna paucicostata to sublethal methionine deprivation. *Plant Physiol.* 71(2) 241-247.
- THOMSON S., 1970: Wolfia. *Trail and Landscape* 4, 127-129.
- THORNER J.P., PETER G.F., NECHUSHAI R., CHITNIS P.R., HUNTER F.A. and TOBIN E.M., 1986: Electrophoretic separation of chlorophyll-protein complexes and their apoproteins. *Plant Biol.* 2, 249-258.
- THORLEY J.H.M., 1977: Root:shoot interactions. In: JENNINGS D.A. (ed.), *Integration of activity in the higher plants*. Symp. Soc. Experimental Biology. Cambridge Univ. Press. 31, 367-389.
- THORSTENSSON B., TILLBERG E. and TILLBERG J.-E., 1985: The effect of different growth limitations on assimilate partitioning in Lemna gibba. *Plant Physiol.* 64(2), 16A.
- TIKHONOV O.I., 1966: Flavonoids of L. minor. (In Ukrainian.). *Farm. Zh. (Kiev)* 21(1), 52-55.
- TIKHONOV O.I., KRIVENCHUK P.E. and LITVINENKO V.I., 1965: Flavonoids of the Lesser duckweed (Lemna minor). I. Preliminary studies. (In Ukrainian.). *Farm. Zh. (Kiev)* 20, 63-65.
- TIKHONOV O.I., KRIVENCHUK P.E., LITVINENKO V.I. and KOVALOV I.P., 1965: Flavonoids of Lemna minor II. (In Ukrainian.). *Farm. Zh. (Kiev)* 20, 53-55.
- TIKHONOV O.I. et al., 1966: The flavonoids of common duckweed. (In Ukrainian.). *Farm. Zh. (Kiev)* 21(3), 40-44.
- TILLBERG E., 1975: An abscisic acid-like substance in dry and soaked Phaseolus vulgaris seeds determined by the Lemna growth bioassay. *Physiol. Plant.* 34(3), 192-195.
- TILLBERG E., 1980: Glycolate and glyoxylate stimulation of growth in Lemna gibba. *Physiol. Plant.* 50, 158-160.
- TILLBERG E. and ELIASSON L., 1976: Indoleacetic acid enhancement of the inhibition of Lemna growth caused by abscisic acid. *Planta* 130, 53-55.
- TILLBERG E., HOLMVAL M. and ERICSSON T., 1979: Growth cycles in Lemna gibba cultures and their effects on growth rate and ultrastructure. *Physiol. Plant.* 46, 5-12.
- TILLBERG E., ELIASSON L. and ERICSSON T., 1980: Time course of uptake of C-abscisic acid by Lemna gibba in relation to growth. *Physiol. Plant.* 48, 584-587.
- TILLBERG E., DONS C., HAUGSTAD M. and NILSEN S., 1981: Effect of abscisic acid on CO₂ exchange in Lemna gibba. *Physiol. Plant.* 52(4), 401-406.
- TILLICH H.-J., 1985: Keimlingsbau und verwandtschaftliche Beziehungen der Araceae. *Gleditschia* 13, 63-73.
- TIMOFEVA S.S. and KRAEVA V.Z., 1983: β -cyanoalanine synthase of hydrophytes and its significance in detoxication of cyanide-containing waters. (In Russian). *Viniti (Irkutsk)* 5439-83. 35 pp.
- TIMOFEVA S.S. and MEN'SHIKOVA O.A., 1983: Development of biotechnology for hydrobotanical removal of thiocyanates from water by the multifactor experiment method. (In Russian). *Viniti (Irkutsk)* 2453-83. 34 pp.
- TIMOFEVA S.S. and MEN'SHIKOVA O.A., 1986: Use of macrophytes to improve biological treatment of thiocyanate-containing wastewater. (In Russian). *Vodn. Resur. (Irkutsk)* 6, 80-85.
- TISCHLER G., 1935: Die Bedeutung der Polyploidie für die Verbreitung der Angiospermen, erläutert an den Arten Schleswig-Holsteins mit Ausblicken auf andere Florenggebiete. *Bot. Jb.* 67, 1-37.
- TITOVA A.A., 1978a: Concentration of the herbicide 2,4-D (2,4-dichlorophenoxyacetic acid) by some higher water plants. *Hydrobiol. J.* 14(4), 96-97.
- TITOVA A.A., 1978b: Accumulation of the herbicide 2,4-D by some higher aquatic plants. (In Russian). *Gidrobiol. Zh.* 14, 110-111.
- TLOUAK E., CZERWINSKI W. and SIEWINSKI A., 1984: Biotransformation of testosterone to androstenedione and 5 α -androstane-3,17-dione by Lemnaceae. (In Polish). Patent: PL 126902 B2. 3 pp.
- TLOUAK E., PAWLOWICZ P., CZERWINSKI W. and SIEWINSKI A., 1986: Transformation of androstane derivatives by Spirodela oligorrhiza. *Phytochem.* 25(1), 61-64.
- TOBIN E.M., 1977: Polyadenylic-acid- plus RNA populations in Lemna gibba G3. *Plant Physiol.* 59(6 suppl.), 22.
- TOBIN E.M., 1978: Light regulation of specific mRNA species in Lemna gibba L. G3. *Proc. Natl. Acad. Sci. USA* 75, 4749-4753.
- TOBIN E.M., 1981a: Light regulation of the synthesis of two major nuclear-coded chloroplast polypeptides in Lemna gibba. In: AKOYUNOGLOU G. (ed.), *Photosynthesis*. Balaban Int. Sci. Serv., Philadelphia. 5, 949-959.
- TOBIN E.M., 1981b: Phytochrome-mediated regulation of messenger RNAs for the small subunit of ribulose 1,5-bisphosphate carboxylase and the light-harvesting chlorophyll a/b-protein in Lemna gibba. *Plant Mol. Biol.* 1(1), 35-51.
- TOBIN E.M., 1981c: White light effects on the mRNA for the light-harvesting chlorophyll a/b-protein in Lemna gibba G3. *Plant Physiol.* 67(6), 1078-1083.
- TOBIN E.M. and KLEIN A.O., 1974: Purification and translation of plant mRNA. *Plant Physiol.* 53(suppl.), 37.
- TOBIN E.M. and KLEIN A.O., 1975: Isolation and translation of plant messenger RNA. *Plant Physiol.* 56(1), 88-92.
- TOBIN E. and SILVERTHORNE J., 1984: Environmental regulation of gene expression in higher plants: light, temperature and the anaerobic response. In: CRESS D. (ed.), *Plant genetic engineering*. Dekker, New York.
- TOBIN E. and SILVERTHORNE J., 1985: Light regulation of gene expression in higher plants. *Ann. Rev. Plant Physiol.* 36, 569-593.
- TOBIN E.M. and SUTTIE L., 1980: Light effects on the synthesis of ribulose-1,5-bisphosphate carboxylase in Lemna gibba G-3. *Plant Physiol.* 65, 641-647.
- TOBIN E.M. and TURKALY E., 1982: Kinetic affects rates of degradation of mRNAs encoding two major chloroplast proteins in Lemna gibba L. G3. *J. Plant Growth Regul.* 1(1), 3-13.
- TOBIN E.M., STIEKEMA W.J., WIMPEE C.F. and SILVERTHORNE J., 1983: Light regulation of nuclear coded genes for chloroplast proteins ribulose-1,5-bisphosphate carboxylase small subunit and chlorophyll a/b-protein. *J. Cell Biochem. (suppl. 7 part B)*, 296.
- TOBIN E. and SILVERTHORNE J., STIEKEMA W.J. and WIMPEE C.F., 1984a: Phytochrome regulation of the synthesis of two nuclear-coded chloroplast proteins. *Sem.-Ser., Soc. Exp. Biol.* 21, 321-335.
- TOBIN E., WIMPEE C.F., SILVERTHORNE J., STIEKEMA W.J., NEUMANN G.A. and THORNER J.P., 1984b: Phytochrome regulation of the expression of two nuclear-coded chloroplast proteins. In: THORNER J.P., STAEHELIN L.A. and HALICK R.B. (eds.), *Biosynthesis of the photosynthetic apparatus*.

- tus. Molecular biology, development and regulation. UCLA Symp. Mol. Cell Biol. 14, 325-334.
- TOBIN E., WIMPEE C.F., KARLIN-NEUMANN G.A., SILVERTHORPE J. and KOHORN B.D., 1985: Phytochrome regulation of nuclear gene expression. In: STEINBACK K.E. et al. (eds.), Molecular biology of the photosynthetic apparatus. NATO Adv. Res. Workshop, Cold Spring Harbor, N.Y. 373-380.
- TODD G.W., MIDDLETON J.T. and BREWER R.F., 1956: Effects of air pollutants. Calif. Agr. 10, 7-8 and 14.
- TOIVONEN H., 1985: Changes in the pleustic macrophyte flora of 54 small Finnish lakes in 30 years. Ann. Bot. Fenn. 22(1), 37-44.
- TOMASZEWICZ H., 1977: Aquatic and bog vegetation in the reservoirs of the Ciechornice and Skwa drainage area on the Gostynin Lakeland. (In Polish). Monogr. Bot. 52, 144 pp.
- TOMIYAMA T., YONE Y. and ISHIO S., 1951: Biochemical studies on the liquefaction of fish body. II. On the effectiveness of "solubilized fish" to the growth of a plant, *Lemna paucicostata* Hegelm. (In Japanese). Sci. Bull. Fac. Agr. Kyushu Univ. 13, 306-312.
- TOMLISON P.B., 1982: Helobiae (Alismatidae). In: METCALFE C.R. (ed.), Anatomy of the monocotyledons VII. Clarendon Press, Oxford.
- TONAPI G.T. and VARGHESE G., 1983: Preliminary observations on the bioecology of the ectoprotectinatella burmanica Annadale. Curr. Sci. 52(13), 646-647.
- TORREY J., 1843: A flora of the state of New York. Albany. 2, 245.
- TOTH L., 1962: On some chemical properties of *Wolffia arrhiza* (L.) Wimm. Annal. Biol. Tihany 29, 275-282.
- TRAPEZNIKOV A.V. and TRAPEZNIKOVA V.N., 1979: 60 Co accumulation by freshwater plants under natural conditions. (In Russian). Ekologiya (Sverdlovsk) 2, 104-106.
- TRIEDEL S., 1974a: Einfluss von Morphaktinen auf den Stoffwechsel höherer Pflanzen. I. Die Wirkung auf die Atmung und auf einige glykolytische Enzyme. Biochem. Physiol. Pflanzen 166(5-6), 481-493.
- TRIEDEL S., 1974b: Einfluss von Morphaktinen auf den Stoffwechsel höherer Pflanzen. II. Die Wirkung auf die Photosynthese und den Gehalt an Stärke, ATP und Chlorophyll. Biochem. Physiol. Pflanzen 166(5-6), 495-509.
- TRELESE W., 1982: On the structure which favors cross-fertilization in several plants. Proc. Boston Soc. Nat. Hist. 21, 410-440.
- TRENT L. and MCARTHUR B., 1974: Results of testing Clean-Flo Lake cleaner in Florida lakes. Hyacinth Control J. 12, 44-45.
- TREMAVAS A., 1970: The turnover of nucleic acids in *Lemna minor*. Plant Physiol. 45, 742-751.
- TREMAVAS A., 1972a: Determination of the rates of protein synthesis and degradation in *Lemna minor*. Plant Physiol. 49, 40-46.
- TREMAVAS A., 1972b: Control of protein turnover rates in *Lemna minor*. Plant Physiol. 49, 47-51.
- TREMAVAS A., 1973: The phosphorylation of ribosomal protein in *Lemna minor*. Plant Physiol. 51, 760-767.
- TREMAVAS A.J., 1979: The dynamics of the meristem control by growth substances. Monogr. Brit. Plant Growth Regul. Group No. 3, 39-53.
- TRIDECH S. and ENGLAND A.J., Jr., 1980: Trace contaminant removal from secondary domestic effluent by vascular aquatic plants. Argonne Natl. Lab., Energy Environ. Syst., Tech. Rep. 96(2), 321-430.
- TRIDECH S., ENGLAND A.J., Jr., HERBERT M.J. and WILKINSON R.F., 1981: Tertiary wastewater treatment by the application of vascular aquatic plants. Chem. Water Reuse 2, 521-539.
- TRIVEDI B.S. and VERMA C.L., 1972: Spore and pollen assemblages of the Tertiary coal beds of Malaya. J. Palynol. 8, 27-36.
- TROTSCH G. and ULLRICH W., 1984: Der Einfluss von 6-Methylpurin auf die Nitrataufnahme bei *Lemna gibba* Gl. Mitt. Bot.-Tag. Wien, 55 (9513).
- TRUX R.E., CULLEY D.D., GRIFFITH M., JOHNSON W.A. and WOOD J.P., 1972: Duckweed for chick feed? Louisiana Agriculture 16(1), 8-9.
- TSAO T.H. and YIN L.M., 1985: Effects of polyamines (PA) and their biosynthesizing inhibitor MGBG on flowering of *Lemna paucicostata* 6746. Abstr. 12th Int. Conf. Plant Growth Substances, Heidelberg, FRG. 130.
- TSAO T.H., ZHONG H.W., JIAO S.P. and TAN Z.Y., 1985: Changes of endogenous ABA and GA contents during floral induction of *Lemna paucicostata* 6746. Abstr. 12th Int. Conf. Plant Growth Substances, Heidelberg, FRG. 96.
- TSAO T.H., ZHONG H.W., JIAO S.P. and TAN Z.Y., 1986: Changes of endogenous ABA and GA contents during floral induction of *Lemna aequinoctialis*. Acta Bot. Neerl. 35(4), 443-448.
- TSUCHIYA M., 1979: Control of aquatic weeds by grass carp (*Ctenopharyngodon idella* Val.). (In Russian). Jarq. 13(3), 200-203.
- TSUDZUKI T. and KONDO T., 1979: Further studies on potassium uptake rhythm in the long-day duckweed *Lemna gibba* G3 with special reference to vegetative growth. Plant Cell Physiol. 20, 1079-1086.
- TUCKER C.S., 1981: Relationships between culture density and the composition of 3 floating aquatic macrophytes. Hydrobiologia 85(1), 73-76.
- TUXEN R., 1971: Lemnetae. In: TUXEN R., Bibliographia phytosociologica syntaxonomica. 2. Cramer, Lehre.
- TUXEN R., 1974: Die Pflanzengesellschaften Nordwestdeutschlands. (2nd ed.). Lemnetae minoris. J. Cramer, Lehre. 35-83.
- TULGANOV A., 1972: Amino acid composition of protein isolated from *Lemna minor* (common duckweed). (In Russian). Dokl. Akad. Nauk Uzb. 29(7), 56-57.
- TURNER B.L., 1967: Plant chemosystematics and phylogeny. J. Pure Appl. Chem. 14, 189-213.
- TURNER P., SCHMIDT K., GAINES L., MANGI J. and PANKOW J., 1978: Effects of chromium on some aquatic plants. Environ. Pollut. 16(4), 285-291.
- TYKHONOV I.O. see TIKHONOV
- UBAYDULLAEV A.U., ABDIEV M. and NIEZOV B., 1971: Morphological and anatomical characteristics of lesser duckweed (*Lemna minor* L.). (In Russian). Kultiv. Vodor. Vyssh. Vodn. Rast. Uzb. 1971, 102-106.
- ULLMANN I. and VAETH R., 1978: Wasser- und Sumpfpflanzengesellschaften der verschiedenen Gewässertypen im Schweinfurter Raum (östliches Maindreieck). Ber. Bayer. Bot. Ges. 49, 137-163.
- ULLRICH W.R., 1979: Die Nitrataufnahme bei Grünalgen und ihre Regulation durch äussere Faktoren. Ber. Deutsch. Bot. Ges. 92, 273-284.
- ULLRICH W.R., 1987: Nitrate and ammonium uptake in green algae and higher plants: mechanisms and relationship with nitrate metabolism. In: Inorganic nitrogen metabolism. Springer, Berlin. In press.
- ULLRICH W.R. and NOVACKY A., 1981: Nitrate-dependent membrane potential changes and their induction in *Lemna gibba* Gl. Plant Sci. Lett. 22(3), 211-217.
- ULLRICH-EBERTUS C.I. and NOVACKY A., 1984: H^+ /phosphate cotransport and its interaction with arsenate and vanadate in transport and metabolism in *Lemna gibba* Gl. In: CRAM W.J. et al. (eds.), Membrane transport in plants. Symp. Abstr., 433.
- ULLRICH W.R., SCHMITT H.-D. and ARNTZ E., 1981: Regulation of nitrate uptake in green algae and duckweeds. Effect of starvation and induction. In: BOTHE H. and TREBST A. (eds.), Biology of inorganic nitrogen and sulfur. Springer, Berlin/Heidelberg/New York. 244-251.

- ULLRICH W.R., LARSSON M., LARSSON C.-M., LESCH S. and NOVACKY A., 1984a: Ammonium-induced membrane depolarization and inhibition of nitrate and phosphate uptake in *Lemna gibba* Gl. In: CRAM W.J. et al. (eds.), Membrane transport in plants. Symp. Abstr., 432.
- ULLRICH W.R., LARSSON M., LARSSON C.-M., LESCH S. and NOVACKY A., 1984b: Ammonium uptake in *Lemna gibba* Gl., related membrane potential changes, and inhibition of anion uptake. *Physiol. Plant* 61(3), 369-376.
- ULLRICH-EBERIUS C.I., NOVACKY A. and LUETTIGE U., 1978: Active hexose uptake in *Lemna gibba* Gl. *Planta* 139, 149-153.
- ULLRICH-EBERIUS C.I., NOVACKY A. and LUETTIGE U., 1980: Extracellular pH changes during glucose uptake in *Lemna gibba*. In: SPANSWICK R.M., LUCAS W.J. and DAINY J. (eds.), North-Holland. Biomed. Press. 551-552.
- ULLRICH-EBERIUS C.I., NOVACKY A., FISCHER E. and LUETTIGE U., 1981: Relationship between energy-dependent phosphate uptake and the electrical membrane potential in *Lemna gibba* Gl. *Plant Physiol.* 67(4), 797-801.
- ULLRICH-EBERIUS C.I., NOVACKY A. and BALL E., 1983: Effect of cyanide in dark and light on the membrane potential and the ATP level of young and mature green tissues of higher plants. *Plant Physiol.* 72(1), 7-15.
- ULLRICH-EBERIUS C.I., NOVACKY A. and VAN BEL A.J.E., 1984: Phosphate uptake in *Lemna gibba* Gl.: Energetics and kinetics. *Planta* 161(1), 46-52.
- UNEMOTO T., 1971: Effect of chlorogenic acid on flower production in long-day duckweed, *Lemna gibba* G3. *Plant Cell Physiol.* 12, 165-169.
- UNEMURA K. and OOTA Y., 1965a: Effects of nucleic acid and protein-antimetabolites on frond and flower production in *Lemna gibba* G3. *Plant Cell Physiol.* 6, 73-85.
- UNEMURA K. and OOTA Y., 1965b: Flowering in *Lemna paucicostata* as compared with that in *Lemna perpusilla* 6746. *Plant Cell Physiol.* 6, 793-798.
- UNEMURA K., INOKUCHI H. and OOTA Y., 1963: Flowering in *Lemna gibba*. *Plant Cell Physiol.* 4, 289-292.
- UNDERWOOD J.L., HESTAND R.S. and THOMPSON B.Z., 1986: Utilization of triploid grass carp for control of floating vegetation in wastewater retention ponds. Proc. 26th Ann. Meet. Aquat. Plant Manage. Soc., Sarasota, Florida, 13-16.
- UNNI K.S., 1970: Seasonal variation in chemical constituents of some aquatic plants. *J. Bombay Nat. Hist. Soc.* 69(1), 242-246.
- UOTILA P. et al., 1983 see 1984
- UOTILA P., BAYTOP A. and LANDOLT E., 1984: Duckweeds (Lemnaceae) in Turkey. *Webbia* 38, 839-844.
- URBAN E., 1983: Energy balance and nitrogen changes of grass carp *Ctenopharyngodon idella* larvae and fry under different food conditions. *Ecol. Pol.* 30(3-4), 327-392.
- URBANSKA see URBANSKA-WORYTKIEWICZ
- URBANSKA-WORYTKIEWICZ K., 1975: Cytological variation within *Lemna L.* *Aquat. Bot.* 1, 377-394.
- URBANSKA-WORYTKIEWICZ K., 1980: Cytological variation within the family of Lemnaceae. *Veröff. Geobot. Inst. ETH, Stiftung Rübel, Zürich* 70, 30-101.
- URBANSKA-WORYTKIEWICZ K., 1982: Cytological variation within the family Lemnaceae. In: SYMOENS J.J., HOOPER S.S. and COMPARE P. (eds.), Studies on aquatic vascular plants. *R. Bot. Soc. Belgium, Brussels*. 1-15.
- USTYENKO P.M., 1984: Floristic findings on the territory of the planned Mezim National Park. (In Ukrainian). *Ukr. Bot. Zh.* 41(4), 64-67.
- UYGUNLAR S., GUERESCI L.U. and BUETVEN G., 1981: Farklı kirleticilerin *Lemna gibba* da radioaktiv etiketli CO₂ alımına etkisi. (In Turk.; not translated). *Atom Enerji Kom. Yay.*, Baskıda.
- VAILLANT F., 1982: Sur une cohabitation entre des Lépidoptères et des Trichoptères comprise par des Mollusques. *Entomologiste* 38(2), 58-64.
- VALLENTINE J.R. and WHITTAKER J.R., 1956: On the presence of free sugars in filtered lake water. *Science* 124, 1026-1027.
- VALLISNERI A., 1710: De arcanis Lenticulae palustris, semine ac admiranda vegetatione, in Prima raccolta d'osservazioni e d'esperienze. Padova.
- VAN BEUSEKOM C.F., 1967: Ueber einige Apiose-Vorkommnisse bei den Helobiae. *Phytochem.* 6, 573-576.
- VAN BRUGEN T., 1976: The vascular plants of South Dakota. Iowa State Univ. Press, Ames, USA. 123.
- VAN COILLIE R., THELLEN C. and DOL J.C., 1981: Detection of retarded toxicity in water pollution. *Water Pollut. Res. J. Can.* 15(3), 203-216.
- VANDENBERGHE A., CHEN M.-W., DAMS E., DE BAERE R., DE ROECK E., HUYSMANS E. and DE WACHTER R., 1984: The corrected nucleotide sequences of 5S RNAs from six angiosperms. With some notes on 5S RNA secondary structure and molecular evolution. *FEBS Lett.* 171(1), 17-23.
- VAN DER PLAS F., 1971: Flora Malesiana Ser. I. *Spermatophyta* 7, 219-237.
- VAN DER VALK A.G. and DAVIS C.B., 1978: The role of seed banks in the vegetation dynamics of prairie glacial marshes. *Ecology* 59, 322-335.
- VAN DER VELDE G., 1979: Nymphaeoides peltata (Gmel.) O.Kuntze (Menyanthes) as a food plant for *Catalysta lemnata* (L.) (Lepidoptera, Pyralidae). *Aquat. Bot.* 7(3), 301-304.
- VAN DER WERFF M., 1981: Ecotoxicity of heavy metals in aquatic and terrestrial higher plants. Ph.D. Thesis. Vrije Univ. Amsterdam. Elinkwijk BV, Utrecht. 126 pp.
- VAN DER WERFF M. and PRUYT M.J., 1982: Long-term effects of heavy metals on aquatic plants. *Chemosphere* 11(8), 727-739.
- VAN DONSELAAR J., 1968: Water and marsh plants in the artificial Brokoppo Lake (Surinam, South America) during the first three years of its existence. *Acta Bot. Neerl.* 17(3), 183-194.
- VAN DYKE G.D., 1965: Incorporation of ¹⁴C-aromatic amino acids into C-glycosyl flavones of *Lemna perpusilla*. M.S. Thesis. Univ. of Texas, Austin.
- VAN DYKE J.M., 1973: Nutritional study of the white amur (*Ctenopharyngodon idella* Val.) fed on duckweed (*Lemna minor* L.). M.S. Thesis. Univ. of Florida. 34 pp.
- VAN DYKE J.M. and SUTTON D.L., 1977: Digestion of duckweed (*Lemna* spp.) by the grass carp (*Ctenopharyngodon idella*). *J. Fish. Biol.* 11, 273-278.
- VAN EE J.H. and PLANTA R.J., 1982: Isolation and characterization of polyribosomes and non-ribosomal ribonucleoprotein particles from the duckweed *Spirodela oligorhiza*. *Plant Sci. Lett.* 25(3), 337-344.
- VAN EE J.H., VAN IN'T VELD W.A. and PLANTA R.J., 1980a: Isolation and characterization of chloroplast DNA from the duckweed *Spirodela oligorhiza*. *Plant Physiol.* 66, 572-575.
- VAN EE J.H., VOS Y.J. and PLANTA R.J., 1980b: Physical map of chloroplast DNA of *Spirodela oligorhiza*; analysis by the restriction endonucleases PstI, XhoI and SacI. *Gene* 12(3-4), 191-200.
- VAN EE J.H., VOS Y.J., BOHNET H.J. and PLANTA R.J., 1982: Mapping of genes on the chloroplast DNA of *Spirodela oligorhiza*. *Plant Mol. Biol.* 1(2), 117-131.

- VAN EYK J., 1963: Investigation on the mode of action of kinetin with Lemna minor L. Ph.D. Thesis. Univ. Leiden.
- VAN EYK J. and VELDSTRA H., 1966: A comparative investigation of kinetin (6-furfurylaminopurine) and some similarly substituted purines and pyrimidines with Lemna minor (L.). Phytochem. 5, 457-462.
- VAN HOREN F., 1969: Observations sur la physiologie des Lemnacees. Bull. Soc. R. Bot. Belg. 8, 15-88.
- VAN HOREN F., 1970: On the hibernation of Lemnaceae. J. Bot. 8, 36-40.
- VAN LOON L.C., TREWAVAS A. and CHAPMAN K.S.R., 1975: Phosphorylation of chromatin-associated proteins in Lemna and Hordeum. Plant Physiol. 55, 288-292.
- VAN MAZIK M.E., 1975: An inverse relation between the multiplication rate and the starch content of the fronds of Spirodela polyrrhiza (L.) Schleiden cultivated in the presence or absence of branched chain amino acids. Acta Bot. Neerl. 24(2), 241-242.
- VAN OVERBEEK J., LOEFFLER J.E. and MASON M.I.R., 1968: Mode of action of abscisic acid. In: WIGHTMAN F. and SETTERFIELD G. (eds.), Biochemistry and physiology of plant growth substances. Runge Press, Ottawa. 1593-1607.
- VAN OVERBEEK J. and MASON M.I.R., 1968: Dormin and cytokinin: growth regulation of Lemna. Acta Bot. Neerl. 17, 441-444.
- VAN STADEN J. and BORNWANN C.H., 1969: Inhibition and promotion by abscisic acid of growth in Spirodela. Planta 85, 157-159.
- VAN STADEN J. and BORNWANN C.H., 1970a: Spirodela growth test. A possible bioassay for abscisic acid. J. S. Afr. Bot. 36(1), 9-12.
- VAN STADEN J. and BORNWANN C.H., 1970b: Cytokinin and gibberellin effects on abscisic acid induced inhibition of growth in Spirodela. J. S. Afr. Bot. 36, 207-213.
- VAN VIERSEN W. and VERHOEVEN J.T.A., 1983: Plant and animal communities in brackish supra-littoral pools ('dobben') in the northern part of the Netherlands. Hydrobiologia 98(3), 203-221.
- VARELA M.E., CORRALES M.A., TELL G., NEIFF J.J., 1978: Limnological studies in the Riachuelo River Basin (Corrientes, Argentina). V. Aquatic Biota of the floating islands in La Brava pond in relation to its environment. Ecosyst. 5(9), 97-118.
- VARELY see VARELA
- VARENKO N.I. and CHUIKO V.T., 1971: Role of higher aquatic plants in the migration of manganese, zinc, copper and cobalt in the Dneprodzerzhinsk Reservoir. Hydrobiol. J. 7(3), 45-48.
- VARENKO N.I. and LUBJANOV I.P., 1973: On accumulation of trace elements by some hydrophytes of the Dneprodzerzhinsk and Zapozozhiye reservoirs. (In Ukrainian). Ukr. Bot. Zh. 30, 165-170.
- VARPOLONEVA T.A., 1976: The major plant formations of the Izhevsk Reservoir (USSR) and their productivity. (In Russian). Bot. Zh. 61(6), 896-900.
- VARGHESE T.J., DEVARAJ K.V. and SHANTARAM B., 1976: Relative growth of the grass carp Ctenopharyngodon idella fed on Utricularia and a mixture of Azolla and Lemna. J. Ind. Fish. Soc. India 8, 206-211.
- VARSNEY C.K. and SINGH K.P., 1976: A survey of aquatic weed problem in India. In: VARSNEY C.K. and RZOSKA J. (eds.), Aquatic weeds in Southeast Asia. Junk, The Hague. 31-41.
- VAUGHAN D. and ORD B.G., 1982: An in vitro effect of soil organic matter fractions and synthetic humic acids on the generation of superoxide radicals. Plant Soil 66(1), 113-116.
- VAUGHAN D., DEKOCK P.C. and ORD B.G., 1982: The nature and localization of superoxide dismutase in fronds of Lemna gibba and the effect of copper and zinc deficiency on its activity. Physiol. Plant. 54(3), 253-257.
- VAUGHAN D., DEKOCK P.C. and ORD B.G., 1983: Effects of benzyldene and abscisic acid on superoxide dismutase in fronds of the duckweed Lemna gibba. Physiol. Plant. 58(2), 239-242.
- VAVRUSKA A., 1966: Determination of nutrients in the most widely spread aquatic, littoral and swamp plants with regard to their utilization for composting. (In Czech.). Prace Vyzk. Ust. Ryb. Hydrol. 6, 41-68.
- VEDENINA Y.A. and ZAVARZIN G.A., 1977: Biological elimination of nitrous oxide under oxidative conditions. (In Russian). Mikrobiologiya 46(5), 898-903.
- VEEN J., 1975: Preliminary studies of the flavonoid pattern of Lemna gibba L. and Lemna minor L. Aquat. Bot. 1, 417-421.
- VELENOVSKY J., 1907: Vergleichende Morphologie der Pflanzen. Prag. 2, 340.
- VELUTHAMBI K., MAHADEVAN S. and MAHESHWARI R., 1981: Trehalose toxicity in Cuscuta reflexa. Correlation with low trehalase activity. Plant Physiol. 68(6), 1369-1374.
- VELUTHAMBI K., GIOVANELLI J., THOMPSON G.A., MUDD S.H. and DATKO A.H., 1983: Regulation of threonine synthase of Lemna paucicostata 6746. Plant Physiol. 72(1 suppl.), 33.
- VENKATARAMAN R., SETH P.N. and MAHESHWARI S.C., 1970: Studies on the growth and flowering of a short-day plant, Wolffia microscopica. I. General aspects and induction of flowering by cytokinins. Z. Pflanzenphysiol. 62, 316-327.
- VERIGIN B.V., 1962: Foreign experiments on cultivation, acclimatization and distribution of white amur and bighead. (In Russian). Piscevaia Promyslenost (Moskva).
- VERMAAK J.F., SWANPOEL J.H. and SCHOONBEE H.J., 1976: Absorption and accumulation of phosphorus-32 by Oedogonium and some aquatic macrophytes. Water SA 2(1), 7-12.
- VERMEERSCH J., 1976: Mise au point d'une méthode de microanalyse des nucléotides pyridiniques dans les cellules de Spirodèle et de Blé. Application à l'étude des effets du saccharose et de l'obscurité. Ph.D. Thesis. Univ. P. et M. Curie, Paris.
- VERMEERSCH J., LECHEVALLIER D. and MONEGER R., 1977a: Sur les nucléotides pyridiniques du Blé et de la Spirodèle cultivés dans diverses conditions (obscurité, lumière rouge de faible puissance, dichlorophényl-diméthylurée). C.R. Acad. Sci. Paris, D 284(18), 1785-1787.
- VERMEERSCH J., MONEGER R. and LECHEVALLIER D., 1977b: Sur le microdosage séparé des nucléotides pyridiniques réduits et oxydés dans les feuilles et les plastides de Blé et de Spirodèle. C.R. Acad. Sci. Paris, D 284(15), 1529-1531.
- VERHADSKY W. and VINOGRADOFF A., 1931: Sur la composition chimique des Lemna comme caractéristique des espèces. C.R. Acad. Sci. Paris, 193, 560-561.
- VESTER H., 1940: Die Areale und Arealtypen der Angiospermen-Familien. 2. Teil. Bot. Arch. 41, 295-356.
- VICKERY R., 1983: Lemna minor and Jenny Greenteeth. Folklore 94, 247-250.
- VIERSEN see VAN VIERSEN
- VIJAYARAGHAVAN S.J., GUPTA A., GUHA-MUKHERJEE S. and SOPORY S.K., 1982: Stimulation of nitrate reductase by light and ammonium in Spirodela oligorhiza. J. Exp. Bot. 33(135), 705-716.
- VINCENT G.A. and BERGERON Y., 1983: La caractérisation d'herbiers aquatiques du lac des Deux-Montagnes (Québec) à partir de paramètres physiques de l'eau. Can. J. Bot. 61, 400-411.

- VINCE-PRUE D., 1976: Phytochrome and photoperiodism. In: SMITH H. (ed.), Light and plant development. Proc. 22nd Univ. Nottingham Easter School in Agricultural Science, Sutton, England. 347-369.
- VINCE-PRUE D., 1981: Daylight and photoperiodism. In: SMITH H. (ed.), Plants and the daylight spectrum. Acad. Press, London. 223-242.
- VINCE-PRUE D., 1983: Photomorphogenesis and flowering. In: SHROPSHIRE W., Jr. and MOHR H. (eds.), Encyclopedia of plant physiology. Photo-morphogenesis. Springer, Berlin. N.S. 16B, 457-490.
- VINTEJOUX C., 1958: Recherches sur la racine de Lemna minor L. (Lemnaceae). Ann. Sci. Nat. Bot. 11e sér., 19, 211-261.
- VINTEJOUX C., 1969: Sur le mode de formation asymétrique des jeunes frondes de Spirodela polyrrhiza. C.R. Acad. Sci. Paris, D 269 (1), 44-47.
- VINTEJOUX C., 1978: Répartition à l'échelle ultrastructurale de composés phénoliques dans les cellules différenciées des turions de Spirodela polyrrhiza L. (Lemnaceae). Actes Soc. Sav. 1, 257-268.
- VINTEJOUX C., 1982a: Particularités physiologiques, cytologiques et cytochimiques des hibernacles de quelques plantes aquatiques. In: SYMOENS J.J., HOOPER S.S. and COMPÈRE P. (eds.), Studies on aquatic plants. R. Bot. Soc. Belg., Brussels. 17-18.
- VINTEJOUX C., 1982b: Caractères du réticulum, des dictyosomes et de l'appareil vacuolaire dans les cellules des turions d'Utricularia neglecta L. (Lentibulariaceae) et de Spirodela polyrrhiza (L.) Schield. (Lemnaceae), en rapport avec l'évolution saisonnière. In: SYMOENS J.J., HOOPER S.S. and COMPÈRE P. (eds.), Studies on aquatic vascular plants. R. Bot. Soc. Belg., Brussels. 29-34.
- VINTEJOUX C. and SHOAR-GHAFARI A., 1985: Répartition et ultrastructure comparées des cellules oxalifères en vie latente et en vie active de Spirodela polyrrhiza L. (Lemnaceae). Bull. Soc. Bot. Fr. Lett. Bot. 1, 25-39.
- VINTEJOUX C. and TOMILIN Y.A., 1984: Indication of radioactive contamination of water systems by the radionuclide content in immersed macrophytes. (In Russian). Gig. Sanit. (Moscow) 5, 87-89.
- VISSER S.A., 1971: Investigations on the nitrogen flow in a model ecosystem. Proc. 4th Intern. Coll. Soil Zoology, Dijon. Ann. Zoologie: Ecologie Animale (hors série), 119-129.
- VOGEL H.J., 1959: On biochemical evolution: lysine formation in higher plants. Proc. Nat. Acad. Sci. 45, 1717-1721.
- VOGEL H.J. and HIRVONEN A.P., 1971: Diaminopimelate decarboxylase (Spirodela oligorrhiza). In: TABOR H. and WHITE-TABOR C. (eds.), Methods in enzymology. Metabolism of amino acids and amines. Acad. Press, London. 17B, 146-150.
- VOGEL R.L. and EISINGER J.E., 1979: Frequency of aquatic macrophytes in East Central Illinois, USA. Trans. Ill. State Acad. Sci. 72 (1), 37-41.
- VOLKER R. and SMITH S.G., 1965: Changes in aquatic vascular flora of Lake East Okoboji in historic times. Iowa Acad. Sci. 72, 65-72.
- VON ARB C. and BRUNOLD C., 1980: Analysis of the regulation of adenosine 5'-phosphosulfate sulfotransferase activity in Lemna minor L. using ³⁵S-density labeling. Planta 149, 355-360.
- VON ARB C., SUTER M. and BRUNOLD C., 1985: Regulation of adenosine 5'-phosphosulfate sulfotransferase activity by amino acids in Lemna minor L. Plant Physiol. 77 (4 suppl.), 160.
- VON HALACSY E., 1904: Conspectus Florae Graecae III. Engelmann, Leipzig. 295.
- VON HALACSY E., 1908: Conspectus Florae Graecae Supplementum. Engelmann, Leipzig. 107.
- VON HAVEK A., 1933: Prodr. Fl. Pen. Balc. 3, Lemnaceae. Rep. Spec. Nov. Regn. Veg. Beth. 30 (3), 424-425.
- VON HUMBOLDT A., BONPLAND A. and KUNTH C.S., 1816: Nova genera et species plantarum. Luteiae Parisiorum. 1, 371-372.
- VON LINNE C., 1753: Species Plantarum Stockholm. 2, 970.
- VON LINNE C., 1771: Mantissa Plantarum Stockholm. 2, 294.
- VON LOON see VAN LOON
- VOSKRESENSKAYA N.P., 1979: Effect of light quality on carbon metabolism. In: GIBBS M. and LATZKE E. (eds.), Photosynthesis II. Encycl. Plant Physiol., N.S. Springer, Berlin/Heidelberg/New York. 6, 174-180.
- VOVK P.S., 1976: The use of white amur as a biocontrol agent for canals and industrial reservoirs. (In Russian). Kiew, Ukrainian SSR, Nauk Dumka, 183-206.
- VRHOVEC B. and WITSCHER M., 1970: The effect of amitrole on the fine structure of developing chloroplasts. Acta Bot. Croat. 29, 43-49.
- VRUCHINSKIY K.K., GRIB I.V. and GRIB A.V., 1970: Content of chlororganic pesticides in water plants. (In Russian). Gidrobiol. Zh. 6 (6), 107-109.
- VRUCHINSKIY K.K., GRIB I.V. and GRIB A.V., 1971: The content of organochlorine insecticides in aquatic plants. Hydrobiol. J. 6 (6), 91-93.
- VUILLOT M., BARBE J. and STEINER B., 1983: Conditions de production de biomasse dans les lagunes d'épuration. Biomasse Actualité, No. Spéc. 3 (Les végétaux aquatiques), 44-46.
- VOYCK L., 1895a: The flowering of Lemna. (In Dutch) Bot. Jaarboek 7, 60-72.
- VOYCK L., 1895b: Sur la floraison de quelques espèces de Lemna. Nederl. Kruidk. Arch. 2/6, 755-756.
- VYAS L.N., 1964: A study of hydrophytes and marsh plants of Alwar and environs. J. Indian Bot. Soc. 43, 17-30.
- WAGNER D.T., 1969: A monocentric holocarpic fungus in Lemna minor L. (Ressia amoeboides). Nova Hedwigia 18 (1), 203-208.
- WALBRIDGE C.T., 1977: A flow-through testing procedure with duckweed (Lemna minor L.). US Environ. Prot. Agency, Environ. Res. Lab. Duluth, Minnesota. EPA-600/3-77-108. 19 pp.
- WALKER E.H., 1976: Flora of Okinawa and the southern Ryukyu Islands. Smithsonian Inst. Press, Washington. 290-291.
- WALKER J.R.L. and EVANS S., 1978: Effect of quaternary ammonium compounds on some aquatic plants. Mar. Pollut. Bull. 9 (5), 136-137.
- WALLACE J.W., Jr., 1967: Investigations of flavone biosynthesis in the Lemnaceae. Ph.D. Thesis. Univ. of Texas, Austin. 118 pp.
- WALLACE J.W., 1969: Evidence against an in vivo equilibrium between 6-carbon and 8-carbon glycosylflavones. Abstr. 11th Int. Bot. Congr. Seattle. 231.
- WALLACE J.W., 1973: Aspects of flavone and C-glycosylflavone B-ring biosynthesis. Am. J. Bot. 60 (4 suppl.), 29.
- WALLACE J.W., 1975: Biosynthetic studies on flavones and C-glycosylflavones: B-ring oxidation patterns. Phytochem. 14 (8), 1765-1768.
- WALLACE J.W. and ALSTON R.E., 1966a: Incorporation and utilization of flavonoid aglycones in Spirodela polyrrhiza. Plant Physiol. 41 (suppl.), 73.
- WALLACE J.W. and ALSTON R.E., 1966b: C-glycosylation of flavonoids. Plant Cell Physiol. 7, 699-700.
- WALLACE J.W. and GRISEBACH H., 1973: The in vivo incorporation of a flavone into C-glycosylflavones. Biochim. Biophys. Acta 304 (3), 837-841.
- WALLACE J.W. and MABRY T.J., 1970: The conversion of the 8-C-glycosylflavone vitexin to the 6-isomer, isovitexin, in Lemna minor. Phytochem. 9, 2133-2135.

- nutrition, rate of respiration and rate of ageing in fronds of Lemna minor. New Phytol. 54, 182-198.
- WARD C.H., WILKS S.S. and CRAFT H.L., 1963: Use of algae and other plants in the development of life support systems. Am.Biol.Teach. 25 (7), 512-521.
- WARD C.H., WILKS S.S. and CRAFT H.L., 1970: Effects of prolonged near weightlessness on growth and gas exchange of photosynthetic plants. In: CORUM C.J. (ed.), Developments in industrial microbiology. Am. Inst. of Biol.Sci. 11, 276-295.
- WARD C.H., KLAINE S.J. and GRANT G.J., 1981: Toxicity bioassays with aquatic plants. Proc.13th Int.Bot.Congr., Abstr. 179.
- WARKENTIN M.J., 1968: Observations on the behavior and ecology of the nutria in Louisiana. Tulane Studies in Zoology and Botany 15(1), 10-17.
- WARNSTORF C., 1896: Blütenbiologische Beobachtungen bei Neu Ruppin im Jahre 1896. Z.Natw.Ver.Harz, Werningerode 11, 9-20.
- WATANABE K. and TAKIMOTO A., 1977: Effects of some metabolic inhibitors on flowering of Lemna gibba G3, a long-day duckweed. Plant Cell Physiol. 18, 1369-1372.
- WATANABE K. and TAKIMOTO A., 1979: Flower-inducing effects of benzoic acid and some related compounds in Lemna paucicostata 151. Plant Cell Physiol. 20, 847-850.
- WATANABE K., FUJITA T. and TAKIMOTO A., 1981: Relationship between structure and flower inducing activity of benzoic acid derivatives in Lemna paucicostata 151. Plant Cell Physiol. 22(8), 1469-1479.
- WATANABE K., TAKIMOTO A., IWAMURA H. and FUJITA T., 1983: Flower inducing activity of benzoic acid derivatives for Lemna minor. Plant Cell Physiol. 24(5), 889-897.
- WATSON R.R. and ORENSTEIN N.S., 1975: Chemistry and biochemistry of apiose. Adv.Carbohydr.Chem.Biochem. 31, 135-184.
- WATT J.M. and BREYER-BRANDWILK M.G., 1962: Medicinal and poisonous plants of Southern and eastern Africa. Livingstone, Edinburgh/London. 669.
- WATTENDORFF J., 1985: Calcium oxalate crystals in plants (Spermatophyta): Occurrence and formation. Conf.Crystal Deposition and Dissolution in Tissues. Lugrin. 6 pp.
- WATZEL R., 1933: Die Wasserlinsen. Blätter Aqu.Terr.k. 43, 90-94. 238 pp.
- WAYMAN C.B., BARLETT G.E. and ALBERTS J.J., 1977: Distribution of pu in aquatic macrophytes from midwestern watershed. In: WHITE M.G. and DUNAWAY P.B. (eds.), Transuramics in natural environments. Nevada Applied Ecology Group Symp.Gatlinburg, Tennessee. 505-516.
- WCISLO H., 1963: Some observations on clones of Lemna trisulca L. grown under aseptic conditions. Acta Biol.Crac.Ser.Bot. 6, 161-176.
- WCISLO H., 1970: Karyological studies in Polish representatives of Spadiciflorae. Acta Biol.Crac.Ser.Bot. 13, 79-88.
- WEAVER C.I. and WETZEL R.G., 1980: Carbonic anhydrase levels and internal lacunar CO₂ concentrations in aquatic macrophytes. Aquat.Bot. 8 (2), 173-186.
- WEBER-OLDECOOP D.W., 1969: Die Wasserpflanzengesellschaften im östlichen Niedersachsen. Ph.D.Thesis. Univ. Hannover.
- WEBER-OLDECOOP D.W., 1971: Wasserpflanzengesellschaften im östlichen Niedersachsen (I and II). Int.Rev.Gesamten Hydrobiol. 55, 913-967.
- WEBER-OLDECOOP D.W., 1973: Anmerkung zur Entdeckungsgeschichte von Wolffia arrhiza in Niedersachsen. Gött.Flor.Rundbr. 7, 29-30.
- WEDDELL H.A., 1849: Observations sur une espèce nouvelle du genre Wolffia. Ann.Sci.Nat. 12, 155-173.
- WALLACE J.W., MABRY T.J. and ALSTON R.E., 1969: On the biogenesis of flavone O-glycosides and C-glycosides in the Lemnaceae. Phytochem. 8 (1), 93-99.
- WALLSTEN M., 1981: Changes of lakes in Uppland Central Sweden during 40 years. Acta Univ.Ups..Symb.Bot.Ups. 23(3), 88 pp.
- WALSH G.E., 1965a: Studies on dissolved carbohydrate in Cape Cod waters. I. General survey. Limnol.Oceanogr. 10(4), 570-576.
- WALSH G.E., 1965b: Studies on dissolved carbohydrate in Cape Cod waters. II. Diurnal fluctuations in Oyster Pond. Limnol.Oceanogr. 10(4), 577-582.
- WALSH M.A. and MELARAGNO J.E., 1974: Ultrastructural features of developing sieve elements in Lemna minor L.: The sieve plate and lateral walls. Am.J.Bot. 61(5 suppl.), 67.
- WALSH M.A. and MELARAGNO J.E., 1976: Ultrastructural features of developing sieve elements in Lemna minor L.: Sieve plate and lateral sieve areas. Am.J.Bot. 63, 1174-1183.
- WALSH M.A. and MELARAGNO J.E., 1981: Structural evidence for plastid inclusions as a possible sealing mechanism in the phloem of monocotyledons. J.Exp.Bot. 32(127), 311-320.
- WALSH M.A. and PALMER J.M., 1979: Duckweed (Lemna minor) not just for the birds. Utah Sci.Agric.Exp.Sta. 40(3), 82-85.
- WALTER H. and BRECKLE S.W., 1983: Oekologie der Erde. Oekologische Grundlagen in globaler Sicht. URB. Fischer, Stuttgart. 1, 238 pp.
- WALTER H. and LIETH H., 1967: Klimadiagramm-Weltatlas. Fischer, Jena.
- WALTER H., HARNICKEL E. and MUELLER-DOMBOIS O., 1975: Klimadiagramm-Karten der einzelnen Kontinente und die ökologische Klimagliederung der Erde. Vegetationsmonographie der einzelnen Grossräume. Fischer, Stuttgart. 10 (9 maps with text).
- WALTERS M.B., 1950: Wolffia papulifera und Lemna minima in Ohio. Ohio J.Sci. 50, 266.
- WANG H., SUN S., KUANG P. and LIU Z., 1984: Uptake, distribution, accumulation and effect of cadmium on aquatic vascular plants. (In Chinese). Huanjing Kexue Xuebao 4(3), 248-257.
- WANG W., 1986a: Toxicity tests of aquatic pollutants by using common duckweed. Environ.Pollut.B 11(1), 1-14.
- WANG W., 1986b: The effect of river water on phytotoxicity of Ba, Cd and Cr. Environ.Pollut.B 11(3), 193-204.
- WANGERMANN E., 1952: Studies in the morphogenesis of leaves. 8. A note on the effects of length of day and of removing daughter fronds on ageing of Lemna minor. New Phytol. 51, 355-358.
- WANGERMANN E., 1965: Longevity and ageing in plants and plant organs. In: RUHLAND W. (ed.), Encyclopedia of plant physiology XV, 2. Springer, Berlin/Heidelberg/New York. 1026-1057.
- WANGERMANN E. and ASHBY E., 1950: Morphogenesis in Lemna minor. Proc. Linn.Soc.London 162, 10-13.
- WANGERMANN E. and ASHBY E., 1951: Studies in the morphogenesis of leaves. VII, 1. Effects of light intensity and temperature on the cycle of ageing and rejuvenation in the vegetative history of Lemna minor. New Phytol. 50, 186-199.
- WANGERMANN E. and LACEY H.J., 1952: Some effects of ultra-violet radiation on Lemna minor. Nature 170, 126-127.
- WANGERMANN E. and LACEY H.J., 1953: Studies in the morphogenesis of leaves. IX. Experiments on Lemna minor with adenine, triiodobenzoic acid and ultra violet radiation. New Phytol. 52, 298-311.
- WANGERMANN E. and LACEY H.J., 1955: Studies in the morphogenesis of leaves. X. Preliminary experiments on the relation between nitrogen

- WEDDELL H.A., 1954: Note sur le *Wolffia Michellii* Schleiden. Bull.Soc. Bot.Fr. 1, 54.
- WEDGE R. and BURRIS J.E., 1979: Effects of temperature and light intensity on photosynthesis in *Lemna minor* and *Spirodela oligorhiza*. Plant Physiol. 63(5 suppl.) 64.
- WEDGE R.M. and BURRIS J.E., 1982: Effects of light and temperature on duckweed photosynthesis. Aquat.Bot. 12(2), 133-140.
- WEGLER R., 1977: Chemie der Pflanzenschutz- und Schädlingsbekämpfungsmittel. Herbizide. Springer, Berlin. 5.
- WEGOREK W., 1983: Pesticide cycle in the agroecosis. (In Russian). Agrokhiimiya 1, 93-101.
- WEGOREK W. and LIPA J.J. (eds.), 1980: The circulation of pesticides in the agroecosis. Proc.19th Conf.Sci.Inst.Plant Protection, Poznan, Poland. 185-204.
- WEIK K.L. and MOHLENBROCK R.H., 1968: Contributions to a flora of Illinois. 3. Lemnaceae. Trans.Ill. State Acad.Sci. 61, 382-399.
- WEIMER W.C. and ARMSTRONG D.E., 1979: Naturally occurring organic phosphorus compounds in aquatic plants. Environ.Sci.Technol. 13, 826-829.
- WEINBAUM S.A., GRESSEL J., REISFELD A. and EDELMAN M., 1979a: Characterization of the 32,000 d chloroplast membrane protein. III. Probing its biological function in *Spirodela*. Plant Physiol. 64, 828-832.
- WEINBAUM S.A., GRESSEL J., REISFELD A. and EDELMAN M., 1979b: Specific depletion of the 32000 d protein from thylakoids by chloramphenicol. Plant Physiol. 63(5 suppl.), 99.
- WEINBERGER P. and CAUX P.Y., 1985: Effect of the solvent carrier dioxanol on some growth parameters of the aquatic angiosperm *Lemna minor* L. Can.Tech.Rep.Fish.Aquat.Sci. 1368, 265-287.
- WEINBERGER P. and IYENGAR S., 1983: Effects of aminocarb, fuel oil 585 and nonylphenol on the growth and development of *Lemna minor* L. In: SHUVAL H.I. (ed.), Dev.Ecol.Environ.Qual. Proc.Int.Meet.Isr.Ecol.Soc., 595-607.
- WEINBERGER P. and GREENHALGH R., 1983: Review of ecotoxicity of metacil in freshwater environment: chemical and phytoecological impact studies. In: NRIAGU J.O. (ed.), Aquatic toxicology. Wiley, New York. 438-448.
- WEISENSEL M., 1968a: Vergleichende Untersuchungen zum Einfluss der Temperatur auf lichtinduzierte Chloroplastenverlagerungen. I. Die Wirkung verschiedener Lichtintensitäten auf die Chloroplastenanordnung und ihre Abhängigkeit von der Temperatur. Z.Pflanzenphysiol. 59, 56-69.
- WEISENSEL M., 1968b: Vergleichende Untersuchungen zum Einfluss der Temperatur auf lichtinduzierte Chloroplastenverlagerungen. II. Die statische Bewegungsgeschwindigkeit der Chloroplasten und ihre Abhängigkeit von der Temperatur. Z.Pflanzenphysiol. 59(2), 153-171.
- WEISENSEL M.H., 1987: Ionic currents at the surface of growing leaves of *Lemna trisulca*. 14th Int.Bot.Congr.Berlin, Abstr., 95.
- WEJNAR R. and MICHEL D., 1983: Untersuchungen zum Einfluss des Nitrofens auf die Chloroplastenstruktur von *Lemna gibba* L. Beitr.Biol.Pflanzen 58, 435-441.
- WEJNAR R. and TAIS S., 1983: Untersuchungen zum Einfluss des Nitrofens auf Wachstum und Pigmentgehalt von *Lemna gibba* L. Beitr.Biol.Pflanzen 58, 427-434.
- WELBORN T.L., Jr., 1979: Water hyacinth. Information sheet, Coop.Extension Serv., Mississippi State Univ. 2 pp.
- WELLMANN E. and GRIEBACH H., 1971: Purification and properties of an enzyme preparation for *Lemna minor* L. catalyzing the synthesis of UDP-apiose and UDP-D-xylose from UDP-D-glucuronic acid. Biochim.Biophys.Acta 235, 389-397.
- WELSH R.P.H. and DENNY P., 1978: The vegetation of Nyumba ya Mungu reservoir, Tanzania. Biol.J.Linn.Soc. 10, 67-92.
- WELSH R.P.H. and DENNY P., 1980: The uptake of lead and copper by submerged aquatic macrophytes in two English lakes. J.Ecol. 68, 443-455.
- WELWITSCH F., 1859: Apontam. Phytogeogr. sobre a flora da prov. de Angola. Annaes Conselho Ultram. 55, 578.
- WENTSEL R.S. and BERRY J.W., 1975: Cadmium and lead levels in Palestine Lake, Palestine, Indiana. Proc.Indiana Acad.Sci. 84, 481-490.
- WERNER D., 1967: Untersuchungen über die Rolle der Kieselsäure in der Entwicklung höherer Pflanzen. Planta 76, 25-36.
- WETSTEYN L.P.M.J., 1983: The distribution of flat forms of *Lemna gibba* L. in the Netherlands. (In Dutch). Int.Rapp.Hugo de Vries Lab., Univ. Amsterdam. 81 pp.
- WETSTEYN L.P.M.J., DE LANGE L. and PIETERSE A.H., 1984: The influence of mould-infection of the nutrient medium on the dimensions of *Lemna gibba* L. Acta Bot.Neeri. 33(4), 559-560.
- WETZEL R.G., 1969: Factors influencing photosynthesis and excretion of dissolved organic matter by aquatic macrophytes in hardwater lakes. Ver.Int.Ver.Limnol. 17, 72-85.
- WETZEL R.G. and MANNY B.A., 1972: Secretion of dissolved organic carbon and nitrogen by aquatic macrophytes. Verh.Int.Ver.Limnol. 18, 162-170.
- WHITE H.L., 1936a: The interaction of factors in the growth of *Lemna*. VII. The effect of potassium on growth and multiplication. Ann.Bot. 50, 175-196.
- WHITE H.L., 1936b: The interaction of factors in the growth of *Lemna*. VIII. The effect of nitrogen on growth and multiplication. Ann.Bot. 50, 403-417.
- WHITE H.L., 1936c: The interaction of factors in the growth of *Lemna*. IX. Further observations on the effect of light intensity on growth and multiplication. Ann.Bot. 50, 827-848.
- WHITE H.L., 1937a: The interaction of factors in the growth of *Lemna*. XI. The interaction of nitrogen and light intensity in relation to growth and assimilation. Ann.Bot.(n.s.) 1, 623-648.
- WHITE H.L., 1937b: The interaction of factors in the growth of *Lemna*. XII. The interaction of nitrogen and light intensity in relation to root length. Ann.Bot.(n.s.) 1, 649-654.
- WHITE H.L., 1938: The interaction of factors in the growth of *Lemna*. XIII. The interactions of potassium and light intensity in relation to root length. Ann.Bot.(n.s.) 2, 911-917.
- WHITE H.L., 1939: The interaction of factors in the growth of *Lemna*. XIV. The interaction of potassium and light intensity in relation to growth and assimilation. Ann.Bot.(n.s.) 3, 619-648.
- WHITE H.L., 1940: The interaction of factors in the growth of *Lemna*. XV. On a rhythmic growth cycle of *Lemna* colonies associated with transference to a potassium free nutrient solution. Ann.Bot.(n.s.) 4, 495-504.
- WHITE H.L. and TEMPLEMAN W.G., 1937: The interaction of factors in the growth of *Lemna*. X. The interaction of nitrogen and light intensity in relation to respiration. Ann.Bot.(n.s.) 1, 191-204.
- WHYTE R.J. and SILVY N.J., 1981: Effect of cattle on duck food plants in Southern Texas (USA). J.Wild Manage. 45(2), 512-515.
- WIEGLEB G., 1976: Untersuchungen über den Zusammenhang zwischen Chemismus und Makrophytenvegetation stehender Gewässer in Niedersachsen. Ph.D.Thesis. Univ. Göttingen. 113 pp.

- WIEGLEB G., 1978a: Der soziologische Konnex der 47 häufigsten Makrophyten der Gewässer Mitteleuropas. Vegetatio 38(3), 165-174.
- WIEGLEB G., 1978b: Untersuchungen über den Zusammenhang zwischen hydrochemischen Umweltfaktoren und Makrophytenvegetation in stehenden Gewässern. Arch. Hydrobiol. 83(4), 443-484.
- WIEGLEB G., 1978c: Vergleich ökologischer und soziologischer Artengruppen und Makrophyten des Süßwassers. Verh. Ges. Oekol., Kiel 1977, 243-249.
- WIEWIORKA Z. and SAKOSIEK J., 1986: Effect of aluminium, lithium and strontium on the ecological populations of Lemna minor and Spirodela polyrrhiza. Proc. 7th Int. Symp. Aquatic Weeds, Wrocław, Poland. 401-406.
- WIGGERS F.H., 1980: Primitive floras Holartic. Killae.
- WIGHTMAN F. and SETTERFIELD G., 1968: Biochemistry and physiology of plant growth substances. Ottawa. 1593-1607.
- WIKOFF S.D., 1949: Lemna trisulca, a southward extension of range on the Atlantic coastal plain. Bartonica 25, 72-73.
- WILKINSON R.E., 1963: Effects of light intensity and temperature on the growth of waterstar grass, coontail and duckweed. Weeds 11, 287-290.
- WILKINSON R.E., 1964: Effects of red-light intensity on the growth of waterstar grass, coontail and duckweed. Weeds 12(4), 312-313.
- WILKS S.S., 1962: Preliminary report on the photosynthetic gas exchange potentialities of the family Lemnaceae (duckweeds). Biologistics for space-systems. Symp. AMRL-TDR-62-116, 265-278.
- WILLEY R.G., DOSKOCIL M.J. and LEWIS C.A., 1974: Potential of the white amur (Ctenopharyngodon idella Val.) as a biological control for aquatic weeds in Indiana. Proc. Indiana Acad. Sci. 83, 173-178.
- WILLIAMS N., 1978: A study of the effects of a diet supplement of Lemnaceae on growth, egg production and chemical content by Gallus domesticus. Louisiana State Univ., Fish. Sect. (Polycopy).
- WILLIS J.H., 1962: A handbook to plants in Victoria. Melbourne Univ. Press. 1, 268-271.
- WILLOMITZER J., LUCKY Z. and KOLAR Z., 1972: Molluscicidal effectiveness of fluorinated derivatives of salicylic acid. Acta Vet. Brno 41, 31-37.
- WILSON W., 1830: Lemna gibba. Remarks on the structure and germination. Hooker Bot. Miscellany 1, 145-149.
- WILSON A.J. and ROBERTS A.W., 1982: Experience in the use of a polymeric cryoprotectant in the freezing of plant tissue. J. Microsc. 125(3), 287-298.
- WILSON L.G. and KANG J., 1972: Sulfurylase activity in Lemna. Plant Research, Ann. Rep. MSU/AEC Plant Res. Lab., Michigan State Univ. 87-88.
- WILSON L.G. and KANG J., 1973: Sulfurylase activity in Lemna gibba and other higher plants. Plant Research, Ann. Rep. MSU/AEC Plant Res. Lab., Michigan State Univ. 113-115.
- WIMPEE C.F., 1984: Organization and expression of light-regulated genes in Lemna gibba L. G-3. Ph.D. Thesis. UCLA. 184 pp.
- WIMPEE C.F. and TOBIN E.M., 1982: Cloning and characterization of two light regulated nuclear genes encoding chloroplast proteins. Plant Physiol. 69(4 suppl.), 139.
- WIMPEE C.F. and TOBIN E.M., 1983: Characterization of light regulated nuclear genes in Lemna gibba. J. Cell Biochem. (suppl. 7 part B), 276.
- WIMPEE C.F. and TOBIN E.M., 1986: Cloning and expression of genes for the small subunit of ribulose biphosphate carboxylase. Methods Enzymol., Plant Mol. Biol. 118, 396-409.
- WIMPEE C.F., STIEKEMA W.J. and TOBIN E.M., 1983: Sequence heterogeneity in the RuBP carboxylase small subunit gene family of Lemna gibba. Plant Mol. Biol. 12, 391-401.
- WINTER E.J., 1937: Growth of Lemna minor. Nature 139, 1070.
- WISE D.L., AUGENSTEIN D.C. and RYHER J.H., 1979: Methane fermentation of aquatic biomass. Resour. Recovery Conserv. 4, 217-238.
- WISTRAND G. and LUNDQVIST J., 1977: New plant localities in Pite (Lappmark, Sweden) and adjacent areas. Sven. Bot. Tidskr. 71(3), 225-238.
- WITTE K.E., 1985: Zur Kultur von Wolffia arrhiza L. Monod. Aquaplanta 10(3), 7.
- WITZTUM A., 1966: A descriptive and experimental study of symmetry in Lemna. Ph.D. Thesis, Cornell Univ., Ithaca, N.Y. 104 pp.
- WITZTUM A., 1974a: Ultraviolet irradiation and pigment cell idioblasts in Spirodela oligorhiza (Lemnaceae). Am. J. Bot. 61(7), 713-716.
- WITZTUM A., 1974b: Abscission and the axillary frond in Spirodela oligorhiza (Lemnaceae). Am. J. Bot. 61(8), 805-808.
- WITZTUM A., 1977: An ecological niche for Lemna gibba L. that depends on seed formation. Israel J. Bot. 26, 36-38.
- WITZTUM A., 1979: Morphogenesis of asymmetry and symmetry in Lemna perpusilla Torr. Ann. Bot. 43, 423-430.
- WITZTUM A. and KEREN O., 1978a: Factors affecting abscission in Spirodela oligorhiza (Lemnaceae). I. Ultraviolet radiation. New Phytol. 80(1), 107-110.
- WITZTUM A. and KEREN O., 1978b: Factors affecting abscission in Spirodela oligorhiza (Lemnaceae). 2. Sucrose. New Phytol. 80(1), 111-115.
- WITZTUM A. and SHAPIRA Z., 1977: Exudation and chloroplast fragmentation as a result of ultraviolet irradiation in Spirodela oligorhiza. Israel J. Bot. 26(3), 109-114.
- WITZTUM A. and KEREN O., 1978: The effect of ultraviolet radiation and sucrose on IAA levels in Spirodela oligorhiza. Ann. Bot. 42(179), 595-598.
- WITZTUM A., POSNER H.B. and GOWER R.A., 1979: Phototactic chloroplast displacement in the photosynthetic mutant, Lemna paucicostata strain 1073. Ann. Bot. 44, 1-4.
- WOHLER J.R., 1966: Productivity of the duckweeds. M.S. Thesis. Univ. Pittsburgh, Pa. 69 pp.
- WOHLER J.R., WOHLER I.M. and HARTMAN R.T., 1965: The occurrence of Spirodela oligorhiza in Western Pennsylvania. Castanea 30, 230-231.
- WOJKE S., 1968: Wolffia arrhiza (L.) Wimm., die Zwerglinse, auch heute noch am Niederrhein. Niederrhein. Jb. 10, 35-38.
- WOJKE S., 1969: Die kleinste Blütenpflanze Europas (Wolffia arrhiza). Mikrokosmos 59(7), 193-194.
- WOLEK J., 1974a: Kritische Uebersicht der Pleustongesellschaften Polens (Klasse Lemnetae). Fragm. Florist. Geobot. 20(3), 365-379.
- WOLEK J., 1974b: A preliminary investigation in interactions (competition, allelopathy) between some species of Lemna, Spirodela, and Wolffia. Ber. Geobot. Inst. ETH, Stiftung Rübel, Zürich 42, 140-162.
- WOLEK J., 1974c: Experimental control of flowering in Spirodela polyrrhiza (L.) Schleid., strain 7401 - a preliminary report. Ber. Geobot. Inst. ETH, Stiftung Rübel, Zürich 42, 163-170.
- WOLEK J., 1979: Experimental investigations on competition and allelopathy between Spirodela polyrrhiza (L.) Schleid. and Wolffia arrhiza (L.) Wimm. Fragm. Flor. Geobot. 25(2), 281-350.
- WOLEK J., 1981: Assessment of the possibility of exoornithochory of duckweeds (Lemnaceae) in the light of researches into the resistance of these plants to desiccation. Ekol. Pol. 29(3), 405-419.
- WOLEK J., 1983: Determinants of community structure for the pleustonic plants (the Lemnetae class). Ekol. Pol. 31(1), 173-200.
- WOLEK J., 1984: Intraspecific variation and the competitive abilities of Spirodela polyrrhiza (L.) Schleiden. Ekol. Pol. 32(4), 637-649.

- WOLF-BLUMER M., 1983: Wasserlinsen-Blüten. Br. Bot. Garten Zürich 17(3), 4 pp.
- WOLFE H.S., 1926: The auximon question. Bot. Gaz. 81(2), 228-231.
- WOLFF J.F., 1801: Commentatio de Lemna. Altdorffii et Norimbergae. 32 pp.
- WOLFF J.F., 1804: Extrait d'une dissertation sur les lenticules. Bull. Sci. Soc. Philomat., Paris 3(13), 142-143.
- WOLVERTON B.C., 1979: Engineering design data for small vascular aquatic plant wastewater treatment systems. Seminar on aquaculture systems for wastewater treatment, Davis, Ca. EPA 430/9-80006, 179-192.
- WOLVERTON B.C., 1984: A review of aquaculture wastewater treatment systems. Workshop Low-Cost Water. Treatm. Syst. Comm. Municip. Albuquerque, Baton Rouge and Dallas, August 1984. 28 pp.
- WOLVERTON B.C. and McDONALD R.C., 1979: Upgrading facultative wastewater lagoons with vascular aquatic plants. J. Water Pollut. Contr. Fed. 51(2), 305-313.
- WOLVERTON B.C. and McDONALD R.C., 1980: Energy from aquatic plant wastewater treatment systems. Sci. Tech. Aerosp. Rep. 18(8), Abstr. No. N80-17545.
- WOLVERTON B.C. and McDONALD R.C., 1981: Energy from vascular plant wastewater treatment systems. Econ. Bot. 35(2), 224-232.
- WONG K.F. and COSSINS E.A., 1976: Control of methionine synthesis by lysine in Lemna minor. Phytochem. 15, 921-925.
- WONG K.F. and DENNIS D.T., 1973: Aspartokinase in Lemna minor L.: Studies on the in vivo and in vitro regulation of the enzyme. Plant Physiol. 51, 327-331.
- WOODFORD E.K., 1950: Assessment of relative toxicity and evaluation of selective herbicides. Proc. Int. Bot. Congr. Stockholm 7, 186-190.
- WORTHLEY E.G. and SCHOTT C.D., 1971: The comparative effects of CS and various pollutants on freshwater phytoplankton colonies of Wolffia papulifera Thompson. Edgewood Arsenal DM. Techn. Rep. EATR 4594. 17 pp.
- WOZAKOWSKA-NATKANIEC H., 1977a: Ecological radiosensitivity of Lemna minor L. and Spirodela polyrrhiza (L.) Schleiden. Monogr. Bot. 55, 53-106.
- WOZAKOWSKA-NATKANIEC H., 1977b: Ecological differentiation of Lemna minor L. and Spirodela polyrrhiza (L.) Schleiden populations. Acta Soc. Bot. Pol. 46, 201-229.
- WRIGHT C.H., 1909: Wolffia denticulata Hegelm. Kew Bull. 1909, 394.
- WRIGHT D.M., 1973: The Fly-fisher's plants. Their value in trout waters. David and Charles, Newton Abbot, England.
- WROBLEWSKI R., 1973: A fine structural investigation of the chloroplasts from the root of Lemna minor L. J. Submicrosc. Cytol. 5, 97-105.
- WROBLEWSKI R., 1976: Identification of raphide and tannin idioblasts in Lemna minor L. root cells by means of analytical electron microscopy. J. Ultrastruct. Res. 57(2), 234-235.
- WUESTLING W. and BOEHM H., 1979: Der Vertikaltest - ein Versuch des parallelen Einsatzes von Organismen verschiedener Organisationshöhe beim Test von Phytoeffektoren. Wiss. Beitr. Martin-Luther-Univ., Halle-Wittenberg, DDR 79(4), 45-50.
- WUESTLING W. and BOEHM H., 1980: Die Submerskultur der Lemnaceae Wolffia arrhiza (L.) Horkel et Wimmer im Unterwasserlicht-Fermenter. Abh. Akad. Wiss. DDR, Math. Natw. Tech. 79(3N), 199-204.
- WUNDER W., 1947: Verschiedenartige Nutzung von Karpfenteichen. Allg. Fischereiztg. 72, 300-306.
- WYLIE G.D. and JONES J.R., 1986: Limnology of a wetland complex in the Mississippi alluvial valley of southeast Missouri, USA. Arch. Hydrobiol. Suppl. 74(3), 288-314.
- WYNNE-EDWARDS V.C., 1941: Wolffia punctata Griseb. in Quebec. Can. Field-Nat. 55, 110.
- YAKOVLEV N.V., 1980: Sensitivity of plants to ammonia and sulfur dioxide. (In Russian). Gaz. Rast. 80, 153-154.
- YAMANI K.A.O., SALIB A.G., NOWAR M.S., MABROUK N.S. and ABOU EL NAGA K.M., 1978: Biological value of water lentil plants and microscopic algae as rich protein sources in poultry rations. Egypt. J. Anim. Prod. 18(2), 177-185.
- YAMASHITA S. and GOTO Y., 1968: Aseptic culture of Lemna and its application to the analysis of the flowering process in higher plants. Books, 400-409.
- YANG C.C., 1978: Preparation of aseptic culture of local species (Spirodela, Lemna paucicostata, Wolffia arrhiza) of Lemnaceae in Taiwan. Bull. Inst. Chem. Acad. Sin. 25, 13-17.
- YANG Y.P., 1978: Lemnaceae of Taiwan. In: LI H.L. et al. (eds.), Flora of Taiwan. V. Angiospermae. Epoch. Publ., Taipei. 816-818.
- YATAZAWA M.H., SUSILO H. and PARK W.C., 1979: Nitrogen fixing capacity in tropical aquatic plants. Biotrop. Bogor, Indonesia. 19 pp.
- YOKOTA R. and SHIMADA J., 1958: Physiological action of kojic acid on green plants. A growth-promoting action on Spirodela polyrrhiza. (In Japan.). Kagaku (Tokyo) 28(10), 531.
- YONE Y. and TOMIYAMA T., 1952: Studies on antianemic ingredients of the liver. VI. The growth promoting effect of liver extract and its purified fractions upon Lemna paucicostata Hegelm. VII. The growth promoting effect of several antianemic substances upon Lemna paucicostata Hegelm. (In Japan.). Bull. Jap. Soc. Sci. Fish. 17, 659-668.
- YONG K.L. and THO Y.L., 1976: Cardiac glycoside-like substance in Malaysia oleander and Lemna minor plant extracts. J. Sains Farm. Malays. 1(1), 12-13.
- YOON J.S. and NOBLE R.D., 1982: Genetic effects of sulfur dioxide in Lemna minor. Bot. Soc. Am. Meet., Abstr.
- YOON J.S., BULLION P., MASTERS S. and NOBLE R.D., 1981: Cytogenetic effects of sulfur dioxide. Genetics 97(1 suppl.), 117.
- YORK H.H., 1905: The hibernacula of Ohio water plants. Ohio Naturalist 5, 291-293.
- YOSHIOKA Y., OSE Y., GOTO M. and HIKINO H., 1987: Relations between the test methods for ecotoxicity. (In Japan.). Eisei Kagaku 33(1), 11-19.
- YOSHIMURA F., 1941: On the minimum concentration of manganese necessary for the growth of Lemnaceae plants. (In Japan.). Bot. Mag. (Tokyo) 55, 163-175.
- YOSHIMURA F., 1943: The significance of molybdenum for the growth of Lemnaceae plants. (In Japan.). Bot. Mag. (Tokyo) 57, 371-386.
- YOSHIMURA F., 1944: Heterotrophic culture of some Lemnaceae plants with sugars. (In Japan.). Bot. Mag. (Tokyo) 58, 15-26.
- YOSHIMURA F., 1950: Physiological studies in Lemnaceae plants. III-VI. (In Japan.). Bot. Mag. (Tokyo) 63(743-744), 63-69.
- YOSHIMURA F., 1952: Influence of the light on the consumption of nitrate and ammonia in Lemnaceae plants. (In Japan.). Bot. Mag. (Tokyo) 65(769-770), 176-185.
- YOUNG M., 1970: The regulation of potassium uptake in Lemna minor L. Ph.D. Thesis. Univ. East Anglia, Norwich.
- YOUNG M. and SIMS A.P., 1972: The potassium relations of Lemna minor L. I. Potassium uptake and plant growth. J. Exp. Bot. 23, 958-969.
- YOUNG M. and SIMS A.P., 1973: The potassium relations of Lemna minor L. II. The mechanism of potassium uptake. J. Exp. Bot. 24(79), 317-327.
- YOUNG M., JEFFERIES R.I. and SIMS A.P., 1970: The regulation of potassium uptake in Lemna minor L. Abh. Deutsch. Akad. Wiss., Berlin, Med. 67-82.

- YOUNG et al., 1978 see YANG Y.P., 1978
- YUI S., 1956: Duckweeds in rice fields and their removal in an easy manner. (In Japan.). *Nogyo Oyobi Engei* 31, 1113-1116.
- YUI S. and KOIKE F., 1955: Studies on P.C.O. (Pentachlorophenol). I. Test of removal of duckweeds in rice fields. (In Japan.). *Nogyo Oyobi Engei* 30, 1107-1108.
- YUKAWA I. and TAKIMOTO A., 1976: Flowering response of *Lemna paucicostata* in Japan. *Bot. Mag. (Tokyo)* 89, 241-250.
- YUSUF R. and OMER S., 1986: Lemnaceae. In: ALI S.I. (ed.), *Flora of Pakistan*. 17 pp. (Polycopy).
- ZAWADZKI B., 1971: Comparison of the prometryn absorption in marshy and in humus-rich soils using the biotests with *Spirodela polyrrhiza* Schleiden and *Wolffia arrhiza* (L.) Wimm. (In Dutch). Report Proefstation voor de Groenteteelt in de Vollegrond. Alkmaar, Netherlands.
- ZAWADZKI B., 1974: *Wolffia arrhiza* (L.) Wimm. as a biotest for determination of prometryn residues in the soil. In: ANTOSZEWSKI R., HARRISON L. and ZYCH C.C. (eds.), *Proc. 19th Int. Horticult. Congr., Warsaw*, 240.
- ZAWADZKI B., 1975: Biological test for trace amounts of prometryn in water using an aquatic plant *Wolffia arrhiza* (L.) Wimm. (In Polish). *Acta Agrobot.* 28(1), 57-66.
- ZAWADZKI B., 1976: Absorption of prometryn by humic-alluvial soil and peat soil. *Biul. Warzyw.* 19, 251-267.
- ZAWADZKI B. and REJMAN S., 1978: Evaluation of the serviceableness of *Wolffia arrhiza* (L.) Wimm. for quantitative determinations of herbicides. *Biul. Warzyw.* 21, 193-200.
- ZENNIE T.M. and McCURE J.W., 1977: The flavonoid chemistry of *Pistia stratiotes* L. and the origin of the Lemnaceae. *Aquat. Bot.* 3, 49-54.
- ZERINGUE H.R., Jr., 1987: A possible relationship between phytoalexin production in the cotton leaf and a phytotoxic response. *Phytochem.* 26(4), 975-978.
- ZERNALOV Y., 1978: How many photosystems take part in green plant photosynthesis? (In Bulg.). *Fiziol. Rast. (Sofia)* 4(2), 90-97.
- ZIEGLER H. and ZIEGLER I., 1965: Der Einfluss der Belichtung auf die NADP-abhängige Glycerinaldehyd-3-phosphat-Dehydrogenase. *Planta* 65, 369-380.
- ZIEGLER H. and ZIEGLER I., 1966: Die lichtinduzierte Synthese der NADP⁺-abhängigen Glycerinaldehyd-3-phosphat-Dehydrogenase. III. Die Bedeutung des Chlorophylls und der Einfluss von Stoffwechsellinhibitoren. *Planta* 69, 111-123.
- ZIEGLER H., ZIEGLER I. and SCHMIDT-CLAUSEN H.J., 1968: The light induced increase in the activity of NADP dependent glyceraldehyde-3-phosphate dehydrogenase. VIII. The dependence of the enzyme activity in *Lemna* on the intensity and the length of illumination. *Planta* 81(2), 181-192.
- ZIEGLER H., ZIEGLER I., SCHMIDT-CLAUSEN H.J., MUELLER B. and DOERR I., 1969: Activation of NADP⁺-dependent glyceraldehyde-3-phosphate dehydrogenase in relation to photosynthetic electron transport. In: METZNER H. (ed.), *Progress in photosynthesis research*. Tübingen. 3, 1636-1645.
- ZIMMERMANN M.-A., 1980: Der Einfluss von Calcium und Magnesium auf das Wachstum von mitteleuropäischen Lemnaceen-Arten. Diploma Thesis. Geobot. Inst. ETH, Stiftung Rübel, Zürich. 92 pp. (Polycopy).
- ZIMMERMANN M.-A., 1981: Einfluss von Calcium und Magnesium auf das Wachstum von mitteleuropäischen Lemnaceen-Arten. *Ber. Geobot. Inst. ETH, Stiftung Rübel, Zürich* 48, 120-160.
- ZOGG H., 1985: Brandpilze Mitteleuropas. *Cryptogamica Helv.* 16, 277 pp.
- ZOLLINGER J., 1956: *Lemna minor*, die kleine Wasserlinse. *Leben und Umwelt* 12(8).
- ZUBERER D.A., 1981: Nitrogen fixation (C_2H_2) associated with duckweed mats in Texas and Florida. *Abstr. Ann. Meet. Am. Soc. Microbiol.* 1981, 179.
- ZUBERER D.A., 1982: Nitrogen fixation (acetylene reduction) associated with duckweed (Lemnaceae) mats. *Appl. Environ. Microbiol.* 43(4), 823-828.
- ZUBERER D.A., 1984: Microbial colonization of some duckweeds (Lemnaceae): Examination by scanning and transmission electron and light microscopy. *Aquat. Bot.* 18(3), 275-285.
- ZUKOWSKI E., ECKSTEIN Z. and DOMANSKA H., 1967: Recherches sur l'extension de l'activité biologique des dérivés d'acide 6-aminopénicillanique (6-APA). *Meded. Rijksfac. Landbouwwet. Gent* 32(3/4), 1050-1061.
- ZURZYCKA A., 1951: The influence of the wave length of light on the movements of chloroplasts in *Lemna trisulca*. *Acta Soc. Bot. Pol.* 21, 17-37.
- ZURZYCKA A. and ZURZYCKI J., 1950: The influence of temperature on phototactic movements of chloroplasts. I. *Acta Soc. Bot. Pol.* 20, 665-680.
- ZURZYCKA A. and ZURZYCKI J., 1951: The influence of some metallic ions on the phototactic movements of chloroplasts. *Acta Soc. Bot. Pol.* 21, 113-124.
- ZURZYCKA A. and ZURZYCKI J., 1953: Studies on the phototactic movements of chloroplasts. I. *Acta Soc. Bot. Pol.* 22, 667-678.
- ZURZYCKA A. and ZURZYCKI J., 1957: Cinematographic studies on phototactic movements of chloroplasts. *Acta Soc. Bot. Pol.* 26, 177-206.
- ZURZYCKI J., 1953: Arrangements of chloroplasts and light absorption in plant cell. *Acta Soc. Bot. Pol.* 22, 299-320.
- ZURZYCKI J., 1955a: Chloroplast arrangement as a factor in photosynthesis. *Acta Soc. Bot. Pol.* 24, 27-63.
- ZURZYCKI J., 1955b: Photosynthesis in polarized light. *Acta Soc. Bot. Pol.* 24, 539-547.
- ZURZYCKI J., 1955c: The dependence of photosynthesis on the arrangement of chloroplasts. *Experientia* 11, 263.
- ZURZYCKI J., 1957a: The destructive effect of intense light on the photosynthetic apparatus. *Acta Soc. Bot. Pol.* 26, 157-175.
- ZURZYCKI J., 1957b: Formative effects of various spectral regions of light on *Lemna trisulca* L. *Med. Landbouwhog. Wageningen* 57, 1-14.
- ZURZYCKI J., 1960: Studies on the centrifugation of chloroplasts in *Lemna trisulca*. *Acta Soc. Bot. Pol.* 29, 385-393.
- ZURZYCKI J., 1961: The influence of chloroplast displacements on the optical properties of leaves. *Acta Soc. Bot. Pol.* 30, 503-527.
- ZURZYCKI J., 1962: The action spectrum for the light dependent movements of chloroplasts in *Lemna trisulca* L. *Acta Soc. Bot. Pol.* 31, 489-538.
- ZURZYCKI J., 1964: The effects of simultaneous action of short and long wave parts of spectrum on the movements of chloroplasts. *Acta Soc. Bot. Pol.* 33, 133-139.
- ZURZYCKI J., 1965: The energy of chloroplast movements in *Lemna trisulca*. *Acta Soc. Bot. Pol.* 34, 637-666.
- ZURZYCKI J., 1969: Experimental modification of the reaction pattern of *Lemna* leaf cells to polarized light. *Protoplasma* 68, 193-207.
- ZURZYCKI J., 1970: Light respiration of *Lemna trisulca*. *Acta Soc. Bot. Pol.* 39, 485-495.
- ZURZYCKI J., 1971: Effect of linear polarized light on the O_2 uptake in leaves. *Biochem. Physiol. Pflanz.* 162, 310-317.

ZURZYCKI J., 1972: Primary reactions in the chloroplast rearrangements. Acta Protozool. 11, 189-200.
 ZURZYCKI J. and LEIATKO Z., 1969: Action dichroism in the chloroplast rearrangements in various plant species. Acta Soc.Bot.Pol. 38, 493-506.
 ZURZYCKI J. and ZURZYCKA A., 1953: Kinematographic method of chloroplast movements analysis. Acta Soc.Bot.Pol. 22, 679-687.
 ZURZYCKI J. and ZURZYCKA A., 1955: Influence of some catalyst poisons on phototactic movements of chloroplasts. Acta Soc.Bot.Pol. 24, 663-674.
 ZURZYCKI J., WALCZAK T., GABRYŚ H., KAJFOSZ J., 1983: Chloroplast translocations in Lemna trisulca L. induced by continuous irradiation and by light pulses. Kinetic analysis. Planta 157(6), 502-510.
 ZUTSHI D.P. and KAUL V., 1963: Lemna trisulca L. - a new record for Kashmir. Trop.Ecol. 4, 95-96.

SURVEY OF SUBFAMILIES, GENERA, SECTIONS AND SPECIES IN THE LEMNACEAE

L e m n a c e a e

Lemnoideae	Wolffioideae
Spirodela	Wolffiella
sect. Spirodela	sect. Stipitatae
S. intermedia	W. hyalina
S. polyrhiza	W. repanda
sect. Oligorrhizae	sect. Rotundae
S. punctata	W. rotunda
Lemna	sect. Wolffiella
sect. Lemna	W. neotropica
L. gibba	W. Welwitschii
L. disperma	W. lingulata
L. minor	W. oblonga
L. japonica	W. gladiata
L. obscura	W. denticulata
L. ecuadoriensis	Wolffia
L. turionifera	sect. Pseudorrhizae
sect. Hydrophylla	W. microscopica
L. trisulca	sect. Elongatae
sect. Alatae	W. elongata
L. perpusilla	sect. Pigmentatae
L. aequinoctialis	W. brasiliensis
sect. Biformes	W. borealis
L. tenera	sect. Wolffia
sect. Uninerves	W. australiana
L. valdiviana	W. angusta
L. minuscula	W. arrhiza
	W. columbiana
	W. globosa

ABBREVIATIONS

(especially chemical substances; however, abbreviations of chemical elements, chemical symbols, and chemical formulas are not included)

ABA	abscisic acid
ACC	1-aminocyclopropane-1-carboxylic acid
ADH	alcohol dehydrogenase
ADP	adenosine-5'-diphosphate
ANAS	acetylcholine
AMP	adenosine-5'-monophosphate
AOA	aminooxyacetic acid
APS	adenosine-5'-phosphosulfate
ASA	acetylsalicylic acid
ATP	adenosine-5'-triphosphate
AVG	aminoethoxyvinyl glycine
BA, BAP	6-benzyladenine, 6-benzylamino purine
BEA	benzoic acid
BOD	biochemical oxygen demand
BZ	benzimidazole
CA	transcinnamic acid
CAMP	cyclic adenosine-5'-monophosphate
Case	carboxylase
CCC	2-chloroethyltrimethyl ammonium chloride
CCCP	carbonyl cyanide m-chlorophenyl hydrazine
Cdr	cytidine deoxyriboside
CIPC	isopropyl m-chlorocarbamate
CMU	3-(4-p-chlorophenyl)-1,1'-dimethylurea (= monuron)
COD	chemical oxygen demand
CS	o-chlorobenzilidene malonitrile
2,4-D	2,4-dichlorophenoxyacetic acid
DAD	diaminodurene
DBSS	dodecylbenzene sodiumsulfate
DBST	trimethylamine alkylbenzenesulfate
DCA	2,4-dichloroanisole
DCCD	N,N'-dicyclohexyl carbodiimide
DCMU	3-(3,4-dichlorophenyl)-1,1'-dimethylurea (= diuron)
DDT	dichlorodiphenyl-trichloromethylmethane
DMC	N,N-dimethylmorpholinium chloride
DMHM	N-dodecylmorpholine-N-methylmorpholinium chloride
DMNP	N-dodecylmorpholine-N-methylmorpholinium chloride
DMSO	dimethylsulfoxide
DNA	desoxyribonucleic acid
DNOC	4,6-dinitro-ortho-cresol
DNP	2,4-dinitrophenol
DOC	dissolved organic carbon
DOM	dissolved organic matter
DON	dissolved organic nitrogen
EAA	essential amino acid index
EDDHA	ethylenediamine-di-o-hydroxyphenylacetic acid
EDTA	ethylenediaminetetraacetic acid
EGTA	ethylene-bis(oxyethylenetrinitrile) tetraacetic acid
EHOPP	2-ethylhexyldiphenyl phosphate
ENF	nonylphenolethoxy sulfate
Ethephon	see ethrel

Ethrel	2-chloroethane phosphonic acid
FCCP	p-trifluoromethoxy carbonylcyanide phenylhydrazine
FDP	fructose diphosphate
FDU	5-fluorodeoxyuridine
FMN	flavine mononucleotide
FU	5-fluorouracil
GA	gibberellin
GDH	glutamate dehydrogenase
GMP	guanosine monophosphate
GOGAT	glutamate synthase
GOT	glutamate-oxaloacetate transaminase
GPD	glyceraldehyde-3-phosphate dehydrogenase
GS	glutamine synthetase
HC	homocysteine
HCCH	hexachlorocyclohexane
α -HPMS	2-pyridylhydroxymethane sulfonic acid
8-HQ	8-hydroxyquinoline
I	ionic strength
IAA	indoleacetic acid
IBA	indolebutyric acid
IMP	inosine 5'-monophosphate
2-IP, IP	N-(Δ^1 -isopentenyl) adenine
LDP	long-day plant
MACC	1(malonylamino)cyclopropane-1-carboxylic acid
MBAS	methylene blue active substances
MCPA	2-methyl-4-chlorophenoxyacetic acid
MDH	malate dehydrogenase
MGBG	methylglyoxal bis(guanylhydrazine)
MIBA	methylhydroxybutanoic acid
MP	6-methylpurine
mRNA	messenger ribonucleic acid
MSMA	monosodium methanearsenate
MSO	methionine sulfoximine
MTA	5'-methylthioadenosine
MTR	methylthioribose-1-phosphate
NAA	naphthaleneacetic acid
NAD	nicotinicamide-adenine-dinucleotide (oxidized)
NADH	nicotinicamide-adenine-dinucleotide (reduced)
NADP	nicotinicamide-adenine-dinucleotide-phosphate (oxidized)
NADPH	nicotinicamide-adenine-dinucleotide-phosphate (reduced)
NIR	nitrite reductase
NR	nitrate reductase
OAS	o-acetylserine
OPH	o-phosphohomoserine
PAA	phenylacetic acid
PABA	p-aminobenzoic acid
PAG	propargylglycine
PAL	phenylalanine ammonia lyase
PAR	photosynthetically active radiation
PCB	polychlorinated biphenyls
PCP	pentachlorophenol
PCFBA, PCIB	p-chlorophenoxyisobutyric acid
PEP	phosphoenolpyruvate
PER	peroxidase
PI	inorganic phosphate
PMG	n-phosphonomethyl glycine (= glyphosate)

LIST OF THE FIGURES

PPA	phenylpropionic acid	14
PS	photo-system	
QAC	quaternary ammonium compound	
RNA	ribonucleic acid	
rRNA	ribosomal ribonucleic acid	
RUDP	ribulose-1,5-diphosphate	
RuDPase	ribulose-1,5-diphosphate carboxylase	
SA	salicylic acid (= 2-hydroxybenzoic acid)	15
SAH	S-adenosylhomocysteine	54
SAM	S-adenosylmethionine	
SDP	short-day plant	
SHAM	salicylhydroxamic acid	72
2,4,5-T	2,4,5-trichlorophenoxyacetic acid	80
TAL	tyrosine ammonia-lyase	
TCDD	2,3,7,8-tetrachlorodibenzo-p-dioxine	
2,4,6-TCP	2,4,6-trichlorophenol	
TIBA	2,3,5-triiodobenzoic acid	88
TO	tetrazolium oxidase	
TPM	tripropylene-glycol-methylether	89
TPP	triophenylphosphate	
trNA	transfer ribonucleic acid	94
TU	2-thiouracil	95
UDP	uridine diphosphate	100
UMP	uridine monophosphate	
XMC	3,5-xylyl methylcarbamate	107
	Fig. 1. 1. The N and P content of Lemna gibba in relation to the N and P content of the nutrient solution	119
	Fig. 1. 2. The Cd, Cr, Cu, Pb, and Zn content of Spirodela polyrrhiza in relation to the concentration of the metals in the nutrient solution	136
	Fig. 1. 3. Banding patterns for 6 different enzyme systems of 11 Lemnaceae species	137
	Fig. 2. 1. Light response curves for photosynthetic rates of Lemna gibba measured in air and in CO ₂ enriched air at 5000 ppm CO ₂	143
	Fig. 2. 2. Growth rate of Lemna gibba and Wolffia arrhiza in relation to the concentration of H ₂ O ₂ solution	176
	Fig. 2. 3. The effect of the concentrations of potassium, calcium, magnesium, nitrate, phosphate, and sulfate on the growth rate of Spirodela polyrrhiza	177
	Fig. 2. 4. The effect of the concentrations of iron, manganese, zinc, and chloride on the growth rate of Spirodela polyrrhiza	178
	Fig. 2. 5. Requirement and tolerance of Spirodela intermedia for Ca and Mg	179
	Fig. 2. 6. Requirement and tolerance of Wolffia angusta for Ca and Mg	180
	Fig. 2. 7. Relation between growth rate and nitrogen supply in Lemna gibba	181
	Fig. 2. 8. Depletion of ammonium and nitrate in the medium by Lemna gibba adapted to a relative nitrogen addition rate of 0.25 per day	182
	Fig. 2. 9. The relative distribution of dissolved inorganic carbon among its various chemical forms as a function of pH	189
	Fig. 2.10. Growth rate of Spirodela polyrrhiza in relation to sucrose concentration	190
	Fig. 2.11. Growth rate of Spirodela polyrrhiza in the dark with 1% sucrose, at 1000 lux without sucrose and at 1000 lux with 1% sucrose	191
	Fig. 2.12. The effect of EDTA on the growth of Spirodela polyrrhiza	
	Fig. 2.13. Growth rates of Lemna minor at different temperatures and different light intensities	
	Fig. 2.14. Growth rate of different Lemnaceae species in relation to temperature	
	Fig. 2.15. Mean dry weight per frond of Lemna minor at different temperatures and different light intensities	
	Fig. 2.16. Mean net assimilation rates of Lemna minor plotted against temperature	
	Fig. 2.17. Relation between growth rate and assimilation rate of Lemna minor at different temperatures	
	Fig. 2.18. Turion formation of Spirodela polyrrhiza and Lemna turionifera in relation to temperature	
	Fig. 2.19. Germination of turions in relation to temperature	
	Fig. 2.20. Relation between light intensity and growth rate in Wolffia arrhiza	
	Fig. 2.21. Growth rates of Lemna minor at different light intensities and different temperatures	
	Fig. 2.22. Growth rate of different species in relation to light intensity	

Fig. 2.23. Mean dry weight per frond of <i>Lemna minor</i> at different light intensities and different temperatures	192	Fig. 2.53. Flavonoid nucleus, numbering system, and substitutions of the flavone, glycoflavone, and flavonols and of the anthocyanidins detected in <i>Lemnaceae</i>	354
Fig. 2.24. Dry weight per frond in relation to light intensity	193	Fig. 2.54. Metabolic pathways leading to several subgroups of flavonoids in <i>Lemnaceae</i>	355
Fig. 2.25. Growth rate of <i>Lemna gibba</i> at three light intensities in solutions with and without sucrose in relation to temperature	195	Fig. 2.55. Intermediate steps in the light action on anthocyanin biosynthesis in <i>Spirodela punctata</i>	358
Fig. 2.26. Growth rate of <i>Lemna gibba</i> and in relation to the length of day in solution with sucrose and without sucrose	198	Fig. 2.56. Hypothetical scheme for a self-sustaining and light-entrainable rhythm working in <i>Lemna gibba</i> G3	364
Fig. 2.27. Growth rate of <i>Wolffia arrhiza</i> in relation to the length of day in solution with sucrose and without sucrose	199	Fig. 3.1. Activity of some enzymes of <i>Wolffia arrhiza</i>	404
Fig. 2.28. The effect of red light and kinetin on the development of intercellular air spaces in <i>Lemna minor</i>	213		
Fig. 2.29. The vascular system in fronds of <i>Lemna minor</i> as influenced by four concentrations of triiodobenzoic acid	214		
Fig. 2.30. The length of life of <i>Lemna minor</i> fronds, the number of daughter fronds produced during the life-span of one mother frond, and the length of daughter fronds as influenced by the N- and P- content of nutrient medium	221		
Fig. 2.31. Senescence and frond development in <i>Lemna aequinoctialis</i>	227		
Fig. 2.32. Arrangement of chloroplasts in the same cells induced by polarized light	233		
Fig. 2.33. Effect of abscisic acid on growth and turion formation of <i>Spirodela polyrrhiza</i>	237		
Fig. 2.34. The contrary influence of mother frond senescence on flower induction in daughter fronds of <i>Lemna gibba</i> G1 and <i>L. aequinoctialis</i> 6746	249		
Fig. 2.35. Movability of stomata in <i>Lemnaceae</i>	271		
Fig. 2.36. Scheme of the xanthophyll cycle	277		
Fig. 2.37. Photorespiration in <i>Lemna minor</i> as shown by the course of CO ₂ -evolution during repeated light-dark changes (20:20 min) under pure oxygen	280		
Fig. 2.38. ATP content of intact <i>Lemna aequinoctialis</i> plants as a function of respiration rate	281		
Fig. 2.39. Transmembrane potential changes in single subepidermal cells of <i>Lemna aequinoctialis</i> fronds induced by short red and far-red light pulses	287		
Fig. 2.40. Velocity of K ⁺ -absorption plotted against K ⁺ -concentration (<i>Lemna minor</i>)	288		
Fig. 2.41. Accumulation of released ABA in the medium of a <i>Spirodela polyrrhiza</i> culture	296		
Fig. 2.42. The effects of continuous light, DCMU, and sucrose on malate accumulation in <i>Lemna aequinoctialis</i> 6746	305		
Fig. 2.43. Biosynthesis of myo-inositol, apiose, and xylose in <i>Lemna</i>	309		
Fig. 2.44. Location of phosphatase in control and phosphorus-deficient <i>Spirodela punctata</i> plants	313		
Fig. 2.45. The glutamine pathway of ammonium assimilation	317		
Fig. 2.46. Summary of the free and bound pathways of sulfate assimilation in plants	321		
Fig. 2.47. Methionine biosynthesis in <i>Lemna</i>	325		
Fig. 2.48. Biosynthetic pathway of branched-chain amino acids	329		
Fig. 2.49. Methionine metabolism	330		
Fig. 2.50. Possible secondary structure of <i>Lemna minor</i> chloroplast 5 S rRNA	341		
Fig. 2.51. Restriction fragment map of <i>Spirodela punctata</i> chloroplast DNA	345		
Fig. 2.52. (4R)-4-hydroxyisophytol, a novel diterpene isolated from <i>Lemna minor</i>	350		

LIST OF THE TABLES

Table 1. 1. Variation of content of elements in % of the dry weight	10	Table 2.13. The effect of low nitrogen supply on characteristics of Lemnaceae	103
Table 1. 2. Mineral ratios in Lemnaceae, according to many authors	11	Table 2.14. pH values of culture solutions with nitrate, ammonium-nitrate, and ammonium after cultivation of 14 and 40 days	110
Table 1. 3. Variation of chemical composition of <i>Wolffia globosa</i> from Japan between summer and winter	17	Table 2.15. Phosphorus requirement and phosphorus tolerance for growth of some Lemnaceae species	113
Table 1. 4. Differences in mineral content between <i>Spirodela polyrrhiza</i> , <i>Lemna minor</i> , and <i>L. trisulca</i>	17	Table 2.16. The effect of low phosphorus supply on characteristics of Lemnaceae	115
Table 1.5a. Mineral content of three Lemna species cultivated under identical conditions	17	Table 2.17. Tolerance of <i>Lemna minor</i> towards different micro-nutrients	123
Table 1.5b. Differences of minimum and maximum content of N and P between different species	18	Table 2.18. Influence of cytokinins on growth rates of Lemnaceae	151
Table 1. 6. Mineral content of Lemnaceae species grown at concentrations of minerals to accomplish half of the maximum growth rate	18	Table 2.19. Effects of some herbicides on Lemnaceae	162
Table 1. 7. Variations of organic components in Lemnaceae	20	Table 2.20. Effects of some algicides, fungicides, insecticides, and other toxic substances on Lemnaceae	169
Table 1. 8. Amino acid composition of the proteins of Lemnaceae	24	Table 2.21. Growth rate of different species of Lemnaceae at two light intensities	196
Table 1. 9. Carbohydrate content in Lemnaceae	26	Table 2.22. Grouping of different species and clones of Lemnaceae according to their behaviour towards yeast extracts and casamino acids in darkness	207
Table 1.10. Percentage of sugars in Lemnaceae	26	Table 2.23. Meaning of the categories A), B), C), and D) in table 2.22.	208
Table 1.11. Content of different sugars within the 3 carbohydrate groups of pectins, hemicelluloses and celluloses in <i>Lemna gibba</i>	27	Table 2.24. Radio sensitivity of different Lemnaceae species	209
Table 1.12. Chlorophyll and carotenoid content in Lemnaceae	32	Table 2.25. Substances influencing flower initiation in Lemnaceae	244
Table 1.13. Flavonoids of Lemnaceae	37	Table 2.26. Effects of blue, red, and far-red irradiation on flowering in <i>Lemna gibba</i> (LDP) and <i>L. aequinoctialis</i> (SDP)	262
Table 1.14. Content of lignin components in Lemnaceae	41	Table 2.27. Concentration of some ions in cell compartments of <i>Spirodela polyrrhiza</i>	299
Table 1.15. Relative chemical composition of raphe idioblasts in <i>Lemna minor</i>	43	Table 3. 1. Productivity of Lemnaceae in different regions of the world	372
Table 1.16. Enzymes determined in members of Lemnaceae	48	Table 3. 2. Number of fronds of Lemnaceae per m ² water surface in nature	373
Table 1.17. Enzyme reactions to 7 different stainings of 11 Lemnaceae species	53	Table 3. 3. Energy content of different Lemnaceae	374
Table 1.18. Polypeptid composition of large and small subunits of Fraction-1-protein for different species of Lemnaceae	55	Table 3. 4. Content of essential amino acids of Lemnaceae, rice, soybean, <i>Chlorella</i> , and egg compared to FAO reference pattern	376
Table 1.19. Relative chemical composition of tannin idioblasts in <i>Lemna minor</i>	56	Table 3. 5. Content of protein, carbohydrate and fat in different crop plants from northern Thailand	377
Table 2. 1. Composition of some nutrient solutions used for Lemnaceae growth	60	Table 3. 6. Daily removal of N, P, and K by Lemnaceae during warm seasons	391
Table 2. 2. Comparison of the molar nutrient content of 7 different nutrient solutions	62	Table 3. 7. Accumulation factors for different elements in Lemnaceae	396
Table 2. 3. CO ₂ uptake of <i>Lemna minor</i> at different light intensities and 5% CO ₂	73		
Table 2. 4. CO ₂ uptake of <i>Spirodela punctata</i> , <i>Lemna minor</i> and <i>L. valdiviana</i> at a light intensity of 240 E m ² and second and at different CO ₂ concentrations	74		
Table 2. 5. Carbon uptake from water in <i>Spirodela polyrrhiza</i> in mg C per g dry weight and hour	75		
Table 2. 6. Carbon uptake from water in <i>Spirodela polyrrhiza</i>	76		
Table 2. 7. Comparison of growth rates of 15 species of Lemnaceae grown in Hutner solution 1/3 and 1/100	80		
Table 2. 8. Lower pH limits for growth of Lemnaceae	84		
Table 2. 9. Nutrient requirements of <i>Spirodela polyrrhiza</i>	86		
Table 2.10. The effect of low potassium content on characteristics of Lemnaceae	88		
Table 2.11. Maximum and minimum requirement of Ca and Mg for long-lasting growth of Lemnaceae	96		
Table 2.12. Nitrogen requirement and nitrogen tolerance of different species of Lemnaceae	101		

INDEX OF SUBJECTS

The most detailed information is given under the figure in fat print

A
 ABA 15,45,97,147,154,156,213,219,220,222,223,224,225,226,235,237,238,
 242,244,248,249,250,251,256,266,268,270,271,273,274,276,284,287,290,
 295,296,297,298,300,306,315,336,337,338,340,352,366,400
 ABA concentration 157
 ABA uptake 294
 abbreviations 582
 abscisic acid see ABA
 abscission 219
 ACC 46,145,160,212,218,219,244,250,330,331,332
 accumulation factor 300,394,299
 Acer pseudoplatanus 175
 acetate 136,321,323,324,351,354
 acetic acid 175
 acetohydroxy acid synthetase see AHAS
 2-acetohydroxybutyrate 327,328
 2-acetolactate 328
 acetylcholine 56,246,264,265
 acetylcholine esterase 264
 acetylsalicylic acid see ASA
 o-acetylserine see OAS
 o-acetylserine sulphydrylase 326
 Achyranthis radix 175
 acknowledgements 410
 acryl acids 172
 actinomycin 267,278
 Adanson 402
 adenine 141,150,260,330
 adenosine 154,260,330
 adenosine-5'-diphosphate see ADP
 adenosine-5'-monophosphate see AMP
 adenosine-3'-phosphate 344
 adenosine-5'-phosphosulfate see APS
 adenosine-5'-phosphosulfate cyclase see APS cyclase
 adenosine-5'-phosphosulfate kinase see APS kinase
 adenosine-5'-phosphosulfate sulfotransferase see APS sulfotransferase
 adenosine-5'-triphosphate see ATP
 adenosine-5'-triphosphatase see ATPase
 S-adenosylhomocysteine see SAH
 S-adenosylmethionine see SAM
 adenylyl cyclase 266
 adenylylates 260,318
 adenylate kinase 260
 ADH 47,48,53,54
 ADP 245,259,260,268,277,318
 Aedes 409
 aeration 143
 aerenchyma 343,373
 aerobic digestion 402
 afaion 274

aflatoxin 169
 Ag 10,134,245,256,257,396
 agar medium 65,227,246,254,267,268,269
 agmatine 46,112,225,320,330,331
 AHAS 327,328,329
 air chambers 146,212
 air drying 239
 air pollution 236
 air spaces 144,157,204,212,213,235,310
 Al 10,124,300,395,396
 alachlor 162
 Al₂(SO₄)₃ 82
 alanine 23,24,112,113,139,140,141,205,245,257,258,278,285,316,318,324
 alanine aminotransferase 48,55,324,404
 alanine dehydrogenase 48
 alanine glyoxylate aminotransferase 48
 alanine uptake 293
 albumine 21
 alcohol dehydrogenase see ADH
 alden 344
 aldohexoses 138,141,294
 aldolase 48,223
 aldolase activity 104,283
 aldrin 175
 algae 388,393,402,406
 algicidal effect 403
 algicides 169,170
 aliphatic carbonic acids 162
 Alismatis rhizoma 175
 alkali metals 87,92
 alkaline cations 287
 alkaline-earth cations 92,290
 alkaloids 56
 alkanes 29
 alkenes 29
 alkylbenzene sodium sulfonate 171,172
 alkylbenzene sulfonic acid 172
 allanthoin 319
 allelopathic substances 400
 allelopathy 297
 allogibberic acid 156,212,244,248
 allosteric effector 279
 alkylphenoxypolyethoxy ethanol see ENF
 Alternaria 173, 402
 aluminum see Al
 ametryne 161,162,167,274
 amides 112
 9-amine-9-fluorenephosphonic acid 163
 amines 112
 amino acid(s) 12,22,99,103,104,109,112,117,139,140,141,205,206,223,
 224,257,278,285,293,304,315,316,318,320,331,338,347,375,376
 amino acid biosynthesis 324
 amino acid content 103,159,376,381
 amino acid incorporation 197
 amino acid transport systems 134
 amino acid uptake 139,141,284,285,293

p-aminobenzoic acid see PABA
 2-aminobutyrate 327,329
 α-aminobutyric acid 23
 aminocarb 171
 1-aminocyclopropane-1-carboxylic acid see ACC
 L-aminoethoxyvinyl glycine see AVG
 9-aminofluorene-9-yl-phosphine oxides 163
 aminooxyacetic acid see AOA
 aminopimelate decarboxylase 48
 aminotriazole 173,229
 amitrole 170,293
 ammonia see NH₃
 ammonium see NH₄⁺
 AMP 260,277,314,318,321,327
 β-amylase 48
 amyloplasts 204
 mylose 25
 Anabaena 383,407
 Anacyctis nidulans 341
 Anas platyrhynchos 386
 anaerobic conditions 239
 Anastatica 220
 5α-androstane-3,17-dione 405
 androstenedione 405
 Angelica dahurica radix 175
 anions 290
 pranic acid 174
 Anopheles 408
 antheraxanthin 33,276,277
 anthocyanidin 354
 anthocyanins 35,36,37,38,40,117,144,355,356
 anthocyanin accumulation 116,15,126,353,357
 anthocyanin biosynthesis 202,203,358
 anthocyanin content 103,145,154,203,357
 anthocyanin formation 69,100,106,138,145,175,201,356,357,359,400
 antibiotics 142
 antimetabolites 167
 antimony see Sb
 AOA 145,212,217,219,244,250
 apigenin 35,36,354,355,356
 apigenin-7-O-glucoside 356
 apiobiose 310
 apiofuranosyl-1,6-glucopyranoside 254
 apiogalacturonan 27,308
 apiose 26,27,307,308
 D-aposyltransferase 48
 apoproteins 334
 appendage (of pouch) 81,105
 application 369
 APS 321,322
 APS cyclase 321
 APS kinase 321
 APS sulfohydrolase 321
 APS sulfotransferase 48,321,322
 Ar 281
 arabinose 26,27,28,136,139
 Araceae 27
 Araliae cordatae foliae 175
 aralphatic carbonic acids 162,165
 aresine 170
 arginine 23,24,99,105,107,112,139,140,141,211,245,316,320,322,328,330,
 343,376
 arginine uptake 294
 argon 281
 arochlor 170
 aromatic sulfonates 282
 arsenate 133,246,260,268,291
 arsenate detoxification 133
 arsenate influx 133
 arsenic see As
 arsenite 133
 arsoniumphospholipids 133
 As 10,133,396
 As concentration 133
 ASA 144,244,252
 ascorbate 108,147,307,353
 ascorbic acid see vitamin C
 ascorbic acid oxidase 48
 aseptic culture 120,142,369
 aseptic culture technique 59
 ash content 20,105,377
 Asiaticum radix 175
 asparagine 23,24,99,105,112,139,141,205,245,257,316,322,332
 aspartate 112,141,245,254,257,258,278,293,318,324,325,326,327,361
 aspartate aminotransferase 48,55,324,405
 aspartate kinase 48,324,326
 3-aspartate semialdehyde 324
 aspartic acid 23,24,99,105,111,112,113,139,140,278,320,324
 aspartokinase see aspartate kinase
 3-aspartylphosphate 324
 aspirin 144
 assimilates 260
 assimilation 178,371
 assimilation rate 71,179,180,218,272
 Astragali radix 175
 atebirin 260,268
 ATP 89,233,245,259,260,268,273,274,277,278,286,309,317,318,321,322,330,
 336
 ATP level 165,278,281,282,285,287
 ATP synthesis 273
 ATPase 48,133,285,286,287,333,335,347
 ATPase activity 89
 ATP-phosphohydrolase 48,404
 ATP-protein phosphokinase 336
 ATP-sulfurylase 48,321,322
 atrataf 170
 atrazine 29,162,167,229,275,349,350
 Aucubae japonicae foliae 175
 autoclaving 136
 autoradiographic methods 269
 auxin(s) 43,148,215,254
 auxin antagonists 148

auxin level 145,212,250,238
 auxin metabolism effectors 148
 auxin uptake 149,294
 AVG 145,212,217,219,244,250,325,326
 azaguanine 173,229,357
 azaserine 316,317
 azauracil 173,229
 azide 172,244,251,362,365
 Azolla 132,392,407

 B 10,12,15,123,276,282,395,396,398
 B concentration 123,299
 B deficiency 124,293,304,359
 B requirements 86,123
 B shortage 124
 B tolerance 123
 B toxicity 124
 BA 83,97,150,160,206,211,219,220,223,224,225,230,237,241,150,266,267,
 290,333,344,349,352
 Ba 10,12,98,396
 back-hybridization 347
 bacteria 220,381,383,402,407
 bacteriostatic effect 403
 BAP see BA
 banding pattern 54
 barium see Ba
 bathing 370
 BEA 46,145,147,211,244,249,252,253,254,266,294,353
 benzanthrane 78,175,244,253
 benzene 168
 benzimidazole see BZ
 benzoanthracene see benzanthrane
 benzoic acid see BEA
 p-benzoquinone 174
 6-benzyladenine see BA
 6-benzylamino purine see BA
 bicarbonate (ions) see HCO₃⁻
 biomass 372,373
 biotechnology 403
 birds 382
 bird food 386
 bleaching 135,186,104,231
 bleaching effect 173,351
 bleaching substances 169,173,229
 blue light 189,201,202,203,229,232,233,234,236,237,261,262,263,267,276,
 278,279,283,324,333,339,342,343,357,359
 BOD 392
 Bonner-Devirian solution 60,61,62,132,261
 borate 12,124,292
 borate diester 12,13,292,299
 borate monoester 13,299
 boric acid 134,299
 boron see B
 Br, Br⁻ 10,122,396
 branching pattern 215,216

bromacil 161,162
 bromate 172
 bromine see Br
 building material 135
 buoyancy 155
 burns 402
 butyric acid 175
 BZ 148

 C 10,118
 C assimilation 75
 C content 371
 C fixation 106
 C-glycosyl flavanone 355
 C-glycosyl flavones 355,356,360
 C metabolism 304
 C uptake 75,76,118,172
 C/N ratio 11,12,15,102
 C, plants 74,278,304
 C, plants 74
 C₄ plants 74
 C₄- 125,173,246,265,282,284,286
 CO 78,282
 CO₂ 71,119,138,245,268,303,304,307,319,330,359
 CO₂ aeration 72,138
 CO₂ assimilation 74,278
 CO₂ assimilation rate 104
 CO₂ compensation point 74
 CO₂ concentration 71,105,241,272,279,332
 CO₂ content 75,120
 CO₂ diffusion 64,73,75,120
 CO₂ enrichment 238,258,259,272
 CO₂ fixation 74,75,76,91,159,178,273,275,278,279
 CO₂ level 197
 CO₂ output 130,173,280,361,362
 CO₂ production 400
 CO₂ shortage 183
 CO₂ source 138
 CO₂ supply 71,80,138,188
 CO₂ uptake 73,74,129,159,173,194,195,272
 CO₃²⁻ 64,118,119,120,138
 Ca, Ca²⁺ 10,12,13,43,56,83,86,89,92,124,142,144,186,209,225,234,246,
 265,276,288,289,290,295,298,301,302,319,363,364,391,396
 Ca chloride see CaCl₂
 Ca concentration 88,92,158,290
 Ca content 10,16,17,158,197,226,290,298
 Ca deficiency 222,223,303,307,340
 Ca exchange 158
 Ca fixation 299
 Ca nitrate see Ca(NO₃)₂
 Ca oxalate 43,307
 Ca oxalate crystals 98
 Ca requirements 86,92,94,95,96
 Ca tolerance 92,94,95
 Ca uptake 97,142
 Ca/Mg ratio 11,92,93

CaCl₂ 82,288,302
 Ca(NO₃)₂ 105
 cadmium see Cd
 caesium see Cs
 caffeic acid 57,297,353
 caffeine 209
 5-caffeoylquinic acid 57,353
 Cairina scutulata 386
 calcicole species 97
 calcium see Ca
 Callitriche 12
 calmodulin 363,364
 callus tissue 66,227,228
 Calthrix 407
 Cambarus 389
 CAMP 246,266,321,337
 campesterol 29,350,351
 canaline 140,343
 canaline-urea cycle 140
 Candida albicans 403
 canavanine 140,328,343
 caprolactam 174
 Carassius auratus 388
 N-carbonyl putrescine 330
 carbaryl 171
 carbofurane 171
 carbohydrate(s) 20,25,135,153,238,258,277,283
 carbohydrate content 17,20,26,124,377,382
 carbohydrate/protein ratio 205
 carbohydrate synthesis 124
 carbohydrate transport 124
 carbon see C
 carbon dioxide see CO₂
 carbon monoxide see CO
 carbonate see CO₃²⁻
 carbonic acid 164
 carbonic acid amide 161
 carbonic anhydrase 48,278
 carboxylase see Case
 carbuncles 403
 cardenolide 28
 cardiac glycoside-like substances 56,403
 cardinal points (of temperature) 176
 β -carotene 33,43,164,202,351,386
 carotinoid(s) 31,156,203,351,357
 carotinoid content 32,154,159,164,171,192,386
 carotinoid production 202
 carp 388
 carrier transport 109,285,290,291,293,295
 casamino acids 99,113,139,205,207,208,242,245,257,320
 casein hydrolysate see casamino acids
 Casuarina 175
 catalase 48,353
 catechin 174
 catfish 388
 catechol 169,173
 catecholamines 246,266
 cations 13,287
 cattle 381,384,393
 CCC 155,243,244,248,268,344,352
 CCCP 89,287,365
 Cd 10,12,14,15,19,129,131,219,395,396,398
 Cd absorption 132
 Cd compartmentation 132,300
 Cd concentration 132
 Cd toxicity 132
 Cd uptake 132
 Cdr 204,229,230
 Ce 10,208,300,396
 cell compartments 117
 cell differentiation 212
 cell division 156,167,200,343
 cell elongation 148,156
 cell length 87,88,157
 cell membrane permeability 159,166
 cell membranes 78,98,111,129,154,170
 cell multiplication 148,150
 cell number 157
 cell proteins 174
 cell size 151
 cell wall 27,41,57,98,197,299,303,307,310,312
 cell wall fragments 175
 cellular multiplication rate 69
 cellulase 48,220,308
 celluloses 17,26,27
 Ceratophyllum 12,120,124,132,299
 Cerium see Ce
 Chamomilla flos 175
 channel catfish 388
 chelating agents 81,82,122,129,135,139,142,251
 chelating effect 83
 chemical composition 9
 chemostat cultures 65,189,191
 chickens 376,384,386
 chilling 180,239,240,241,242,272
 chinampa 406
 Chlamydomonas 335
 chloramphenicol 173,205,229,246,267,275,278,306,335,340
 Chloranthi glabri foliae 175
 chlorate 172
 Chlorella 376,383
 chloride see Cl
 chlorine see Cl
 o-chlorobenzilidene malononitrile see CS
 chlorobromuron 162
 2-chloroethane phosphonic acid see ethrel
 chlorofluorene 162,165,273,274,283
 chlorogenic acid 174,244,253,353
 p-chlorophenoxyisobutyric acid see PCPIB
 3-(4-p-chlorophenyl)-1,1'-dimethylurea see CMU
 chlorophyll a/b apoprotein 201,226,332,333,334,343
 chlorophyll content 17,31,32,77,99,100,103,104,129,130,145,154,156,
 159,164,165,166,168,170,171,172,174,192,274,278,349

chlorophyll decomposition 156
 chlorophyll degradation 225,352,357
 chlorophyll methyl esters 331
 chlorophyll synthesis 133,172,173,186
 chlorophyll system 188
 chlorophyllide 351
 chloroplast(s) 28,29,31,91,118,130,138,154,166,168,194,203,235,270,273,
 276,283,304,316,317,319,326,327,332,333,335,336,340,341,349,362,364,
 367
 chloroplast development 228,351,357
 chloroplast DNA 344,347
 chloroplast formation 164
 chloroplast membranes 171
 chloroplast movement 202,231,270
 chloroplast position 201,270
 chloroplast proteins 34,334,342
 chloroplast ribosomes 335,341,342
 chloroplast RNA 339,340,341,346
 chloroplast translocation 201,202
 chloroplast ultrastructure 228,231
 6-chloropurin 163
 chlorosis 76,111,142,166,170,211
 chlorotic symptoms 87,116,117,125,132,167
 chlorpromazine 363
 choline 141,294,331
 chromatine 336
 chromatography 34,38,360
 chromium see Cr
 chromosomal nondisjunction 77
 chromosome changes 209
 chymotrypsin 43
 cinnamic acids 57,297,352,354,366
 Cinnamon cortex 175
 CIPC 164,165
 circadian clock see clock
 circadian rhythms 265,289,353,361
 citrate 43,122,136,142,143,295,344
 citrulline 139,140,141,328,343
 Cl⁻ 10,13,43,56,91,121,298,396
 Cl⁻ absorption 122
 Cl⁻ accumulation 157
 Cl⁻ concentration 88,121,122
 Cl⁻ content 91,121
 Cl⁻ requirements 86,121
 Cl⁻ uptake 292
 clock 263,265,361,363,367,368
 CMU 270,271,357,362
 CN⁻ 125,173,246,265,282,284,286
 Co 10,12,86,130,219,244,250,300,337,396
 Co concentration 130
 coal distillate 175
 cobalt see Co
 COD 390,392
 cold-blooded animals 387
 coliforms 394
 coloured light 260
 companion cell 298
 compartment 117
 compartmental analysis 299
 compensation point 190,191
 complex salts 142
 concentration factor see accumulation factor
 condensed phosphates 311,312
 coniferyl alcohol 42
 continuous light 198,229,230,239,240,242,249,251,253,255,263,264,266,
 267,272,277,287,289,304,305,324,339,342,347,353,362,363,365,366,368
 copper see Cu
 Coptidis rhizoma 175
 corn silage 384
 cortex 298
 Cortinarius speciosissimus 166
 Corydalis tuber 175
 cosmonautic plant 406
 cough 402
 p-coumaric acid 174,352
 coumarin 147,174
 5-p-coumaroylquinic acid 57,353
 p-coumaryl alcohol 42
 cow(s) see cattle
 cow-dung 402
 CPS 162
 Cr 10,15,19,130,300,396,398
 Cr concentration 130,131
 Cr tolerance 130,131
 crayfish 389
 o-cresol 169
 crop plant 382
 croton aldehyde 174
 crude fibre content 20
 crude oil 175
 Crustaceae 388
 cryptochrome 201,202
 crystal cells 43
 CS 169,174
 Cs 10,90,208,289,300,396
 Ctenopharyngodon idella 387
 CTS 162
 Cu 10,12,14,15,19,86,128,130,134,144,145,146,150,159,211,212,219,245,
 252,253,256,257,258,260,261,264,266,300,301,359,370,395,396,398
 Cu concentration 128,129,332
 Cu requirements 86,128
 Cu tolerance 123
 Cu toxicity 128,332
 Cu uptake 129,212,256,257
 CuSO₄ 130,147,164,169,226,297
 Culex 408,409
 cultivation 59,379
 cultivation plants 380
 culture technique 65
 cutrine plus 166
 cyanatryn 161,162,168
 cyanide see CN

cyanidin 354,355
 cyanidin glucosides 35,36,38,39,202
 cyanoacrylate 168
 β -cyanoalanine synthase 48
 cyanobacterin 170
 Cyanophyta 407
 cyclic adenosine-3'-5'-monophosphate (cyclic AMP) see cAMP
 cyclic adenosine-2'-3'-phosphate 344
 cyclic nucleotides 30,337
 cyclic photophosphorylation 260
 cycloheximide 173,229,246,267,335
 p-cymenes 173
 Cyperus 407
 Cyprinus carpio 388
Cypripis 389
 cystathionine 322,325,326,330
 cystathionine- δ -synthase 48,326
 cysteic acid 117,322
 cysteine 23,24,77,117,245,257,321,322,323,325,326
 cysteine synthase 48,321,322,323,326
 cystine 117,118,125,141,228,245,291,292,322,323
 cytidine 230
 cytidine deoxyriboside see Cdr
 cytochrome 31,34,232,275,333
 cytochrome oxidase 48,282
 cytochrome reductase 48
 cytochrome(s) 45,142,145,147,150,160,205,206,222,2245,225,237,238,240,
 241,244,250,256,337
 cytokinin level 203,338
 cytokinin uptake 154
 cytoplasm 57,117,158,235,298,299,303,311,362,364
 cytoplasmatic ribosomes 333,334,335,336,341,342,343
 cytoplasmatic RNA 339,340
 cytosin 167
 cytosin-thymine dimers 348
 cytosol 283,341

 D
 2,4-D 66,79,83,148,160,165,216,218,227,293,294,295
 DAD 275
 dairy heifers 384
 dalapon 293
 Daphnia 388
 dark conditions (darkness) 202,204,205,206,207,208,218,231,232,239,242,
 251,263,270,277,278,284,287,302,303,318,324,332,340,343,353,359,361,
 364,365,368
 dark growth 204
 dark-red light 114
 Datko 4 solution 61,62
 daughter fronds 67,149
 daylength 198,199,260,261,267,268
 day-neutral plants 144,248,249,252,257,260,261
 DBSS 142
 DBST 169,171,315
 DCCD 286
 DCMU 109,161,162,164,218,234,245,259,260,267,268,274,275,278,279,283,
 286,287,305,314,362

 DDT 169,170,399
 decarboxylation 106
 deep freezing 65
 deltamethrine 399
 deplasmolysis time method 295
 desiccation 239,241,302
 desoxyribonucleic acid see DNA
 detergents 169,171,282
 deuterium 87
 deuterium oxide 87,224
 diacyldigalactosyl glycerol 349
 diacylgalactosyl glycerol 29,249
 diacylglycerol 349
 diaminodurene see DAD
 diaminopimelate decarboxylase 324
 diaminopimelic acid 324
 diazinon 174
 1,4-diazobicyclo[2,2,2]octane 332
 dibutyltinphthalene sodiumsulfonate 171
 2-dichloroacetamido-3-chloro-1,4-naphthaquinon 170
 2,4-dichlorophenoxyacetic acid see 2,4-D
 3-(3,4-dichlorophenyl)-1,1'-dimethylurea see DCMU
 dichlorodiphenyl-trichloromethylmethane see DDT
 2,4-dichlorophenyl-4-nitrophenylether see nitrofen
 2,4 dichloropyrimidine 162,167
 dicoumarol 244,253
 N,N'-dicyclohexyl carbodiimide see DCCD
 dieldrin 175
 diethyldithiocarbamate 245
 di(2-ethylhexal)phthalate 174
 diethylsulfate 209
 diffusion potential 173,284,287
 digalactosyl diglyceride 349,350
 digitoxose 28,403
 digoxin 403
 2,3-dihydroxy flavanone 356
 dihydroxy xanthophyll 33
 2,5-dimethyldodecanoic acid 161
 dimethylhydroquinone 174
 N,N-dimethylmorpholinium chloride see DMMC
 dimethylnitrosamine 78,175
 dimethyl pyrocatechol 174
 dimethylurea 295
 2,4-dinitrophenol see DNP
 dinoseb 293
 Dioscorides 402
 diphenoxylon 162
 α,α' -dipyridyl 244,251
 diquat 162,166,173,370
 disinfectants 169
 dissolved organic carbon see DOC
 dissolved organic matter see DOM
 dissolved organic nitrogen see DON
 diterpenes 57
 dithionate 117
 diuretic 403

diurnal oscillation 289,347
diuron see DCMU
DNMC 344
DNMM 169
DNMP 169
DNMSO 265
DNA 153,269,339,343
DNA content 167,174
DNA formation 153
DNA functions 344
DNA precursors 230
DNA repair 347
DNA segments 345
DNA structure 344
DNA synthesis 164,172,224,343
DNA transcription 347
DNase 348
DNOC 171,274
DNP 164,172,240,246,259,260,362,365
Docauer solution 61,63
dodecylbenzene sodiumsulfate see DBSS
N-dodecylcyclomethylene-N-methylmorpholinium chloride see DNMM
N-dodecylloxymethylene-N-methylpiperidinium chloride see DNMP
dominance 215,217
dominant daughter frond 215
dormancy 235
dowanol 170
driselase 308
dropsy 403
drugs 175,400
druses 43
dry weight see frond weight
drying 380
ducks 382,386,393
duration of light 198

E
EAA 375,377
ectoplasm 234
economic importance 369
EDDHA 143,144,158,212,217,218,219,220,244,249,250,251,252,253,254
EDTA 92,93,122,129,143,233,245,251,252,253,254,256,290,307,319,320,359
Egeria densa 374
egg 376,384
EGTA 363
EHDPP 399
Eichhornia 12,170,381,389,392,398,399,402,407
Einstein 188
electrochemical gradient 290
electroconductivity 363
electrolytes 287,363
electron flow 168
electron transport 159,194,260,272,273,275,276,278,334,357
electrophoresis 34
Elodea 12,41,120,398
elongation zone 302,303

embryo 217,242,269
Emerson enhancement effect 274
endodermis 298
endogenous factors 211
endogenous rhythms 231,236,263
endonucleases 344,345
endoplasmatic reticulum 235,337,350
endothall 166,399
endrin 166
energy budget 245
energy charge 277,278,318
energy content 374
energy conversion efficiency 188,375
energy generating processes 258
energy metabolism blockers 109
energy production 402
energy suppliers 135
energy units 188
ENF 169,171,315
environmental factors 211,372
enzymatic activity 104,177
enzymatic processes 140
enzyme(s) 47,98,164
enzyme content 55
enzyme production 403
enzyme systems 54,103
epiallogibberic acid 212,248
epicatechin 173
epidermis 79,302,303,312,313
epidermis cells 182,302
epigallocatechin 173
epinephrine 266
epoxylutein 202
epoxystearic acid 169
erg 188
Escherichia coli 347,348,403
esculetin 174
esserine 246,264
essential amino acid index see EAA
ethalfluralin 162
ethanol 136,175,240,365,402
ethanolamine 141,294,331
ethephon 160,211,212,217,219,244,250,268
ethionine 246,267
ethrel see ethephon
ethyl alcohol see ethanol
ethylene 46,156,160,212,215,219,220,225,250,251,254,268,330
ethylene formation 212,254,330,331
ethylene glycol 174
ethylene level 145
ethylene production 129,220,226,252,331,332
ethalensamine 209
ethylenediamine-di-o-hydroxyphenylacetic acid see EDDHA
ethylenediaminetetraacetic acid see EDTA
ethylhexyldiphenyl phosphate 399
ethyl-N-phenylcarbamate 174

etiolated plants 228,335,356
 etioplasts 228
 eutrophy of water 216
 evaporation 302,407
 evapotranspiration 302,407
 evapotranspiration/evaporation ratio 302,407
 excision of frond primordia 214
 extensin 22,226
 eye diseases 403
 Eyster solution 61,62

F
 F, F⁻ 10,122,140
 false dawn 368
 false dusk 368
 FAO reference 375,376
 far-red light 200,202,211,218,222,229,236,241,258,261,262,263,268,286,
 287,295,304,306,334,356,357,369
 fat(s) 17
 fat content 17,377
 fatty acid(s) 28,154,167,175,273,349
 fatty acid content 156
 FCCP 270,271
 FDP 273
 FDU 167,173,229,246,266
 Fe 10,12,15,81,86,125,142,145,146,150,152,211,225,244,251,252,300,353,
 396,398
 Fe availability 142
 Fe concentration 88
 Fe content 10,17,77,126,132,142,252,253,300,301
 Fe deficiency 212,315,359
 Fe effects 126
 Fe humate 142
 Fe requirements 86,126
 Fe supply 126
 Fe tartrate 142
 Fe tolerance 123
 Fe uptake 212,253
 Feigel's test 57
 female flower part 153,161
 feminization 268
 fenithrothion 170
 ferric citrate 253,254
 ferricyanide 126,246,256,257,265,266
 ferritin 12
 ferrocyanide 246,265
 ferulic acid 41,57,353
 1-feruloylglucose 57,353
 fibres 377
 ficin 43
 fish 382,384,387,393,394
 fish effluent 388,394
 fish ponds 370
 fishing 370
 flavacoumarine 173
 flavanone 354,355,356
 flavanoneol 355
 flavin(s) 203,232
 flavone-O-glucoside 256
 flavones 35,36,37,354,355,356,360
 flavonoid(s) 34,35,202,297,354
 flavonoid content 154,156
 flavonoid formation 156
 flavonol-converting enzyme 48
 flavonols 35,36,37,39,40,354,355,356
 floral bud 269
 flower development 152,267
 flower differentiation 267
 flower inducing activity 129,146,147
 flower induction (initiation) 116,144,158,174,179,243
 flower inhibition 243
 flower primordia 153,156,160,267,268,269
 flower production 90
 flower promotion 126,134,141,144,147,243
 flowering 77,103,106,120,125,126,127,129,133,134,138,141,144,145,147,
 149,152,155,156,160,167,175,182,200,203,222,243,367,368
 fluometuron 162,165,274
 fluoridone 162
 fluorine see F
 5-fluorodeoxyuridine see FDU
 fluorodifen 161,162
 p-fluorophenylalanine 285
 5-fluorouracil see FU
 FWN 234,306
 folic acid 146,147
 food conversion rate 387,388,393
 food plant 382
 foot-candle 188
 formate 279
 fowl 382
 fractures 402
 fresh weight (of fronds) 69,99
 frond age 197
 frond area see frond size
 frond clusters 93,114,117,126,158
 frond colonies 145
 frond colour 92,157
 frond counting 67
 frond development 152,155,157
 frond differentiation 167
 frond disconnection 138,152,155,157,160
 frond elongation 167,200,202
 frond form 150
 frond length 148,177,211,221,227
 frond length/width ratio 69,93,102,103,114,115
 frond morphogenesis 211
 frond multiplication 200,219,220,282,283,304,320
 frond nerves 149,160
 frond number 373,374
 frond primordia 215,230,238
 frond production 193

frond regeneration 228
 frond separation 144,220
 frond size 69,87,88,90,93,102,103,114,115,117,130,138,144,148,150,
 155,157,160,177,199,200,211,219,226,400
 frond thickness 69,93,157
 frond weight 69,71,77,82,87,88,90,92,102,103,104,105,114,115,117,142,
 144,150,155,165,176,178,192,193,270,272,371,373,400
 β -D-fructofuranosidase see invertase
 fructose 28,109,136,138,215,231,245,258,278,285
 D-fructose-6-phosphate 1-phosphotransferase 48
 fruit development 267
 FU 167,246,266,343
 fucose 26,27
 fuel oil 175
 fuel production 402
 fumarate 43
 Funaria 234
 fungicides 169,170
 Fusarium roseum 403
 fusicoccin 173,285,287,344

 G
 GA 39,45,146,155,160,202,239,240,241,248,250,256,268,295
 GA uptake 156,248,294,295
 GA₁ 147,155,156,211,215,216,219,220,238,239,240,242,243,244,248,266,
 268,344,350,351,352,359
 Ga 10,86,396
 galactolipids 28,29,153,349
 galactosamine 27
 galactose 26,27,136,138,225,231,283
 D-galacturosonyl transferase 48,310
 Galen 402
 gallium see Ga
 Gallia rhois 175
 gas (illuminating) 78
 gas exchange 71,182,283,304,400,406
 gaseous substances 71
 GDH 47,48,53,54,91,105,138,140,223,318,319
 Ge 134
 GeO₂ 134
 GeO₃ 134
 gel electrophoresis 47,338,339,341,344
 generation of energy 270
 germanium see Ge
 germination of seeds see seed germination
 gibberellins see GA
 gibberic acid 212,248
 gibbosity 102,103,115,144,145,155,160,212
 Ginseng radix 175
 globuline 21
 glucosamine 27,337
 glucose 26,27,28,108,109,136,215,225,231,238,245,258,278,283,285,287,
 289,302,303,307,319,332,350,362,366
 glucose transport 134
 glucose uptake 173,293
 glucose-1-phosphate 309,310,314

glucose-6-phosphate 30,246,260,307,309,310,311
 glucose-6-phosphate dehydrogenase 48,98,223,283
 glucuronic acid 136,308,309
 glutamate 141,245,257,258,317,319
 glutamate dehydrogenase see GDH
 glutamate glyoxylate aminotransferase 49
 glutamate oxaloacetate transaminase see GOT
 glutamate pathway 318,319
 glutamate pyruvate aminotransferase 49
 glutamate synthase see GOGAT
 glutamic acid 23,24,99,105,111,112,113,139,140,141,205,278,316,320
 glutamine 23,24,105,107,111,112,115,139,141,205,245,257,316,318,322,361
 glutamine pathway 316,319
 glutamine synthetase see GS
 glutamine synthetase/glutamate synthase cycle 91,316
 glutaric acid 136
 glutathione 23,322
 glutathione peroxidase 49
 gluteline 21
 glyceraldehyde 3-phosphate dehydrogenase see GPD
 glycerol 295
 glycine max 175
 glycine 23,24,112,139,140,234,245,257,258,278,285,287,306,316,318,332
 glycine oxidase 234
 glycine uptake 141,173,293
 glycoflavones 35,36,37,38,40,354
 glycolate 43,169,279,280,306,307
 glycolate oxidase 43,49,106
 glycolyase 49
 glycolysis 103,319
 glycoproteins 337
 α -glycosidase see maltase
 β -glycosidase 404
 glycoside-like substances 56,403
 glycosylation 39
 glycyl-alanine 112
 glycyl-aspartic acid 112,320
 glycylglycyl peptides 165
 glycyrrhiza radix 175
 glyoxylate 106,169,234,280,306,307
 glyphosate see PMG
 glyphosine 168,334
 GMP 154
 GOGAT 49,91,317,318,319
 goldfish 388
 Gomori procedure 313
 GOT 47,49,53,54
 GPD 49
 gramicidine 246,265
 gramoxone 274
 grana 167,228,229,230,349
 grana lamellae 349
 grana structure 164
 Granati cortex 175
 grass-carp 377,387
 green light 202,205,216,263,357

growth factors 135,400
 growth inhibition 129,132,164,205
 growth processes 156
 growth promoting substances 142
 growth rate 67,71,77,79,80,81,82,83,87,88,90,91,92,93,99,100,105,117,
 118,126,128,129,130,134,136,137,142,147,148,155,160,165,168,173,175,
 176,177,180,181,183,189,190,191,194,195,196,198,200,203,210,237,270,
 371,372,400
 growth stimulation 130,136,139,142,148,157,280
 growth substances 43,204,294
 GS 49,91,106,140,273,316,317,318,319
 guaiacol 169,173
 guanidine derivatives 112
 β -guanido propionic acid 112
 guanine 167
 guanosine deoxyriboside 230
 guanosine monophosphate see GMP
 H
 H⁺ 10
 H⁺ cotransport 116,134,141,284,290,293,294
 H⁺ extrusion pump 105,109
 HCCH 170
 HCN 173
 HCO₂ 76,118,119,120,138
 HF 78
 H₂S 77,117,320,322,323
 H₂S content 77
 H₂S emission 118,320
 H₂S fumigation 118,291,292,323
 H₂SeO₃ 133
 halogens 121
 handedness 216,217
 harvest 380
 HC see homocysteine
 HCCH 399
 heat tolerance 186
 heavy metals 14,123,212,219,251,300,301,394,400
 Helodea see Elodea
 hemicelluloses 26,27
 hepatitis 402
 Heracleum lanatum 175
 herbicides 130,133,166,216,273,274,292,370,399,400
 heterodromous branching 216
 heterocyclic compounds 162,164,167
 heterotrophic conditions 218,219,334,361,367
 heterotrophic growth 146,204
 heterotrophy 120
 hexachlorobiphenyl 170,399
 hexachlorocyclohexane see HCCH
 hexahydrofluorene-9-carboxylic acid 156
 hexametaphosphates 13
 hexene 77,273,274
 hexose(s) 138
 hexose uptake 138,284,293,294
 Hg 10,130,134,245,396,398
 HgCl₂ 134,285
 high-intensity light 211,216,218,232,238,242,255,258,259,266,270,272,
 276,278,283,307,320,352,362,363,365
 high-performance liquid chromatography 360
 Hill reaction 91,203,229,273,275,276
 Hillman solution see M solution
 histidine 23,24,139,245,258,376
 histones 336,337
 Hoagland solution 21,60,61,62,82,93,98,121,132,142,174,206,259,261
 HOE 2977 169
 homocysteine 322,330
 homocysteine biosynthesis 324
 homodromous branching 215,216
 homogentisic acid 353
 homo-orientin 36
 homoserine 324,326,327
 Hordeum vulgare 41
 hormonal state 243
 hormones 43,142,148,244
 hormone-like substances 135
 horses 385
 α -HPNS 307
 8-HQ 129,146,244,245,253
 human nutrition 382,389
 humate 142
 humic acid 142
 humus substances 142
 Hutter solution 21,60,61,62,79,80,82,120,121,259
 hybridization technique 346
 hydrocarbons 29
 Hydrocharitaceae 27,308
 hydrogen see H
 hydrogen fluoride see HF
 hydrogen sulfide see H₂S
 hydroquinone 169,174
 hydrothol-47 168
 p-hydroxybenzaldehyde 41,142
 2-hydroxybenzoic acid see SA
 p-hydroxy-benzoic acid 254,353
 8-hydroxyquinoline see 8-HQ
 hydroxycinnamic acid 41,353,360
 4-hydroxycoumarin 147,244,253
 (12S)hydroxyhexadeca-8Z,10E,14Z-trienoic acid 29
 (4R)-4-hydroxy-isophytol 29,57,350
 hydroxyllysine 24
 hydroxymethyl transferase 140
 4-hydroxy-3-nitrobenzaldehyde 230
 4-hydroxy-3-nitrobenzoic acid 230
 4-hydroxy-3-nitrobenzyl alcohol 230
 2-hydroxy-1-phenyl-1,4-pentadione 244,253
 4-hydroxyphenylpyruvic acid 353
 4-hydroxyphenylpyruvate dioxygenase 49,353
 hydroxyproline 24,245
 hydroxypyruvate reductase 49
 8-hydroxyquinoline see 8-HQ

- I
- IAA 43,146,147,148,174,202,215,217,219,220,228,240,244,250,254,264,266,268,338
- IAA content 138,220
- IAA level 203,219,225
- IAA oxidases 145,254
- IAA uptake 295
- IAA-L-alanine 66
- IBA 148
- Ictalurus punctatus 388
- illuminating gas 78
- indicator plant 400
- indoleacetic acid see IAA
- indolyl-3-acetic acid see IAA
- indolylbutyric acid see IBA
- inner cortex 298
- infrared irradiation 236
- inosine 5'-monophosphate see IMP
- inorganic substances 86
- inositol 307
- inositol hexaphosphate see phytic acid
- inositol phosphates 13,30,300,312
- insecticides 169,170,399
- insulin-related substances 47
- intercellular spaces 57,144
- inuline 136
- invert sugar 136
- invertase 48,49,294,404
- iodate 172
- iodide see J⁻
- iodine see J
- iodineacetate 172
- iodoacetamide 245
- ion transport 284
- ionophores 265,270
- 2-IP (IP) 66,150,227,237,241
- Ipecacuanhae radix 175
- Ipomoea 407
- iron see Fe
- isocitrate 254
- isocitrate dehydrogenase 49
- isoelectric point 56
- isoleucine 23,24,140,245,258,324,327,376
- isoleucine synthesis 140,327,328,329
- isoleucine uptake 293
- isoorientin 354,355,356
- isoorientin-3'-methylether 355
- isopentenyl adenine see 2-IP
- isopentenyl aminopurine see 2-IP
- isoprenoids 57,350
- 2-isopropylmalate synthetase 328,329
- isoproterenol 246,266
- isosaponarin 35,36
- isovitexin 354,355,356
- J
- J, J⁻ 10,122,292,396
- Joule 188
- Juncus procerus 374
- K
- K, K⁺ 10,12,87,98,234,298,302,363,391,392,396
- K⁺ absorption 89,288
- K⁺ accumulation 158
- K⁺ concentration 88,284,287,288,364
- K content 10,15,16,17,88,89,93,159,196,197,202,225,298,300
- K deficiency 218,304
- K⁺ level 225
- K⁺ proton exchange carrier 134
- K⁺ removal 391
- K⁺ requirements 86
- K⁺ shortage 114
- K⁺ uptake 89,97,111,287,288,289,365
- K⁺ uptake rate 89,91,173
- K⁺ uptake rhythm 90,172,173,202,265,363,364
- K/Ca ratio 159
- K/Na ratio 11
- KCl 11
- KCN
- K, (Fe(CN)₆) 147,172
- KHCO₃ 128
- KH₂PO₄ 114
- KNO₃ 105
- K₂SO₄ 118,320
- kaempferol 36,202,354,355,356,359
- kanomycin 173,229
- kerosene 168
- α-ketoglutaric acid 205
- 4-keto-intermediates 308,309
- khair-nam 382
- kinetin 39,81,150,160,202,205,206,212,213,215,216,224,225,228,231,238,240,241,242,250,268,334,338,351,400
- Klebsiella 407
- Knop solution 82,143
- kojic acid 142
- Krantz leaf anatomy 74,278
- Krebs cycle 319
- krenite 399
- krezamon 274
- L
- La 10,300,396
- lactose 81,136,283
- lanolin 79
- lanthanum see La
- laser radiation 210
- LDP 90,144,200,243,248,249,250,251,252,255,257,258,260,267,368
- leakage 224,265,292,296,363
- lead see Pb
- left-handed 215,217
- Lemnetum 373

Lemno-Utricularietum 373

lenacil 162
length/width ratio 69,93,102,103,114,115,211
leucine 23,24,113,139,140,141,245,293,328,376
leucine content 140
leucine synthesis 328
leucine uptake 293
leucocyanidin 173
Li 10,86,90,234,246,265,289,365
LiCl 265
life-span 104,154,199,203,221,222,226,227,248
light 188,314,332,335,356,357,358,265
light compensation point 190,191,195
light dependence 242
light duration 198
light effect 194,195,196,197,351,359
light energy 188
light influence 188
light intensity 72,73,74,87,102,118,124,137,150,176,178,183,188,189,
196,197,200,212,242,259,261,267,270,271,272,273,277,282,264,365
light periods 198
light pulses 201
light saturation 189,194,195,196,197,198
light stimulation 116
light-dark cycle 261,264,266
lignins 41
linoleic acid 28,154,167,332,349,350
linolenic acid 28,167,349,350
linuron 161,162,293
lipid(s) 20,28,98,331
lipid composition 153
lipid content 20,28,154,167
lipid droplets 235
lipid metabolism 167
lipid oxidation 332
lipochroms 98
lithium see Li
long-day conditions 120,144,218,222,226,227,236,239,240,242,243,248,
249,250,252,253,254,255,257,258,259,260,262,263,264,265,267,268,
286
long-day plants see LDP
long-short-day plant 249,250,260
Loniceræ flos 175
low-intensity light 231,233,262
lucenin 36
lutein 33,146,164,202,351
lutein-epoxide 33
luteolin 36,354,355,356
luteolin-7-O-glucoside 38,356
lutonarin 36
lux 188
lycopene 32
lycopersene 29,57,351
lyophilization 65,403
lysine 23,24,140,245,257,324,325,326,327,335,376

M
M solution 60,61,62,132
MACC 330,331,332
Macrobrachium 389
magnesium see Mg
Magnoliæ cortex 175
maize 33,346
malate 43,74,254,305,316
malate accumulation 305,306
malate content 105,301
malate dehydrogenase see MDH
malathion 174
male organs 153
maleic hydrazide 172
malic acid see malate
malic enzyme 49
malonic acid 136
l (malonylamino)cyclopropane-l-carboxylic acid see MACC
malonyl-CoA 354
malonylcyanidin monoglucoside 35,36
maltase 504
mammalian feed 384
manganese see Mn
mangiferin 174
mannitol 81,136,223,303
mannose 26,27,138
manure 142,406
masculinization 268
matatril 169,171
maximum growth rate 68,101,114,137,138,185,194,196,197,371,380
maximum temperature 186
maximum yield 379
Mayas 383
Mazerozyme R-10 308
MBAS 392
MCPA 161,162
MDH 46,49,53,54,55,98,223
measurements 67
mechanical damage 282
mecoprop 166
Medicago sativa 33
medicinal plants 175
membrane 124,129
membrane potential 109,173,200,284,293
membrane protein 334
membrane transport 264,284
menstruation 402
Menthae herba 175
mercaptopyruvate sulfotransferase 49
mercury see Hg
meristem 209,216,220,228,302
messenger ribonucleic acid see mRNA
metabolic activity 118
metabolic inhibitors 240
metabolites 167
metal ions 123,142

metaphosphates 13
methane 57,394,402
methionine 23,24,113,117,139,140,141,167,245,322,324,332,333,334,357,358,375,376
methionine biosynthesis 140,324
methionine metabolism 330
methionine pathway 324,326,327
methionine sulfoximine see MSO
methionine-5'-methylthioadenosine cycle 225
methoxone 172
methyl group transfer 330,331
methyl groups 140,330
methylbenzopyrene 78,175
2-methyl-4-chlorophenoxyacetic acid see MCPA
methylcoumarins 172
5,10-methylenetetrahydrofolate 327
methylglyoxal bis(guanyldrazone) see MGBG
3-O-methylglucose 285,293
methylhydroxybutanoic acid see MHBA
5-methylmethionine sulfonium 322
methylparathion 169
6-methylpurine see MP
methyltetrahydrofolate 326,327,330
methylthio recycling 330,331
5'-methylthioadenosine see MTA
methylthioribose-1-phosphate see MTR
methyleurea 295
Mg 10,17,43,56,83,89,92,127,142,209,222,223,225,234,288,298,301,318,337,340,347,363,391,396
Mg concentration 88
Mg content 87,88,197,290
Mg requirements 86,92,94,95,96
Mg tolerance 92,94,95
Mg/Ca ratio 12
MgCl₂ 82,288
MgSO₄ 338
MGBG 244,251
MHBA 307
mice 385,403
Microchaete 407
micronutrients 123
microsomes 29
microsporogenesis 269
mineral accumulation 300
mineral composition 159
mineral content 9,17
mineral deficiency 223,231,235,236,239,259
mineral elements 356,359
mineral location 298
mineral ratios 11
minimum temperature 84
mitochondria 138,235,281,283,318,319,364
mitochondrial DNA 346
mitotic activity 118
mitotic index 77
mixotrophic conditions 277,362

mixotrophic growth 197
Mn 10,12,15,17,81,86,98,123,126,132,142,145,212,219,244,251,252,274,275,300,301,307,310,337,352,395,396
Mn concentration 88,127
Mn requirements 86,127
Mn tolerance 9,123,127
Mo 10,15,81,86,109,117,122,123,125,219,257,258,291,300,359,396
Mo content 117
Mo toxicity 125
Mo tolerance 123
MoO₃ 117,125,245,256
MoO₄²⁻ deficiency 125
molluscicides 171
molybdate see MoO₄²⁻
molybdenum see Mo
monogalactosyl diglyceride 350
mono- β -D-glucopyranoside 353
monohydroxy xanthophyll 33
mononucleotides 314
monophenolase activity 282
monosodium methanearsenate see MSNA
Mori 402
morphactins 165,273,274,283
mosquito breeding 408
mosquito larvae 408
motor-boating 370
MP 109,161,163,168
mRNA 30,203,333,334,335,336,342
MSA 145
MSNA 163
MSO 174,273,316,317
MTA 330
MTR 330
Mugil cephalis 388
mullet 388
multiplication rate 67
muskrat 385
mutagenesis 66,173,209
mutagenous effects 77
mutagens 209
Mycobacterium smegmatis 403
mycotoxins 173
Myocaston 385
myo-inositol 307,309,310,312
myo-inositol kinase 49,312
myo-inositol hexaphosphate see phytic acid
myo-inositol-1-phosphate 307,308,310,312
myo-inositol-1-phosphate kinase 49,312
myo-inositol-1-phosphate synthase 49,307
myo-inositol-triphosphate 312
myo-inositol-triphosphate kinase 49,312
Myricae cortex 175
Myriophyllum spicatum 120
N
N 78,99,391,392,396,398

N accumulation 407
 N assimilation 311
 N balance 116
 N concentration 100,104,139
 N content 10,14,17,18,21,77,83,103,104,130,167,196,375
 N deficiency 22,102,104,222,223,231,235,236,237,239,240,303,306,312,319,359
 N limitation 191,273
 N removal 223,391
 N requirement 86,99,101
 N shortage 114,235,240
 N source 83,99
 N starvation 108,224,283
 N supply 99,103,218,221,273
 N tolerance 101
 N utilization 106,111
 N₂ 10,12,99
 N₂ aeration 78,279
 N₂ atmosphere 272
 NH₃ 82,211,245,257,260,314,316,326,388
 NH₄⁺ 12,86,99,104,105,107,110,129,132,139,145,164,179,206,211,265,278,279,289,301,304,306,315,316,317,322,332,337,338
 NH₄⁺ absorption 83
 NH₄⁺ assimilation 91,125,316
 NH₄⁺ concentration 22,361
 NH₄⁺ content 107,394
 NH₄⁺ depletion 107
 NH₄⁺ ions 82,104
 NH₄⁺ carbamate 171
 NH₄⁺ removal 393,394
 NH₄⁺ sulfamate 162,164
 NH₄⁺ thiocyanate 162,164
 NH₄⁺ uptake 109,111,141,197,228
 NH₄⁺ utilization 109
 NH₄⁺-N 100
 NH₄Cl 105
 NH₄NO₃ 105,106,110
 (NH₄)₂SO₄ 105,150
 NO₂⁻ 99,257,314,316
 NO₂⁻ reductase see Nir
 NO₂⁻ reduction 316,317
 NO₂⁻ 18,19,45,71,86,99,104,109,110,114,139,179,245,256,278,279,285,289,298,301,304,306,317,319,320,337,363,364
 NO₂⁻ absorption 83,290
 NO₂⁻ assimilation 109,111,124,125,138,256,289
 NO₂⁻ carrier 290,291
 NO₂⁻ concentration 88,104,108,109,186,255,282,290,291,339,361
 NO₂⁻ deficiency 218,222,223,241,340
 NO₂⁻ depletion 107
 NO₂⁻ efflux 290
 NO₂⁻ level 366
 NO₂⁻ pollution 401
 NO₂⁻ reductase see NR
 NO₂⁻ reduction 125,159,291,305,314
 NO₂⁻ starvation 290
 NO₂⁻ storage 109
 NO₃⁻ supply 302
 NO₃⁻ uptake 104,108,109,111,112,124,141,193,196,197,284,289,290,291,364
 NO₃⁻/H⁺ cotransport 104
 NO₃⁻-N 99,100
 Na 10,12,17,86,90,98,121,225,287,299,302,365,391,396
 Na acetate 273
 Na arsenite 164,172
 Na azide 172,293
 Na chlorate 162,164
 Na content 91
 Na fluoroacetate 171
 Na laurylsulfate 264
 Na phytate 174
 Na proton exchange carrier 134
 NaAsO₂ 133
 NaCl 91,121,273,306,319
 NaCN 365
 Na₂CO₃ 91,120
 NaF 122
 NaHCO₃ 120
 NaN₃ 172,365
 NaNO₂ 105
 NaNO₃ 111
 NAA 145,148,212,215,218,219,220,254
 NAD 30,276,308,309,363,364
 NAD/NADP cycle 366
 NAD glutamate dehydrogenase 49,98,106,316
 NAD kinase 363,364
 NAD glyceraldehyde-3-phosphate dehydrogenase see NAD GPD
 NAD GPD 49,278,362,363,364
 NAD nitrate reductase 50,314
 NADH 245,259,309,318,319
 NADH cytochrome-c-reductase 49
 NADP 30,276,362,363,364
 NADP glucose-6-phosphate dehydrogenase 362
 NADP glutamate dehydrogenase 49
 NADP glyceraldehyde-3-phosphate dehydrogenase see NADP GPD
 NADP GPD 49,278,362,363,364
 NADP isocitric dehydrogenase 50,98,362
 NADP malate dehydrogenase 50,306
 NADP nitrate reductase 50,315
 NADP phosphatase 363,364
 NADP 6-phosphogluconate dehydrogenase 50
 NADPH 245,259,273,279,318
 NADPH/NADP ratio 276
 Najadaceae 27
 nalidixic acid 173,229
 naphthaleneacetic acid see NAA
 β-naphthol 145,212,244,253
 naringenin 354,355,356
 Nasturtium 392
 Nb 10,300,396
 NDP phosphohydrolase 321
 Nkal 282
 neochrome 34

neoxanthin 33,164,202,351
 nerves 149,160
 Ni 10,12,86,129,130,300,319,320,395,396,398
 Ni concentration 130
 nickel see Ni
 nick-translation 347
 nicotinamide-adenine-dinucleotide-phosphate (oxidized) see NADP
 nicotinamide-adenine-dinucleotide-phosphate (reduced) see NADPH
 nicotinic acid 43,246,266
 nigericin 270,271
 night temperature 213,264
 niobium see Nb
 NiR 50,108,314,315,316
 NiR activity 315
 nitrate see NO₃
 nitrate reductase see NR
 nitriles 162,165
 nitrite see NO₂
 nitrofen 162,164,352
 nitrogen see N
 nitrophenols 162,164,230
 NMU 66
 N-nitroso-N-ethylurea 209
 N-nitroso-N-methylurea 209
 nonylphenol 171
 nonylphenolethoxy sulfate see ENF
 norflurazon 357
 N-phosphonomethylglycine see PMG
 NR 91,108,109,256,258,314,315,316,319
 NR activity 78,104,106,107,108,109,125,129,147,167,168,199,200,315,
 366
 nuclear genes 347
 S₁ nuclease 50
 nucleic acid(s) 30,224,229,312,314,331
 nucleic acid synthesis 124,173,229,266
 nucleotide(s) 30,153,318
 nucleotide sequences 346
 nucleus 336,347,367
 Nutria 385
 nutrient assimilation 177
 nutrient concentration 79
 nutrient needs 64
 nutrient recycling 390
 nutrient removal 391
 nutrient requirements 86
 nutrient solution 60,61,79
 nutrient supply 195
 nutrient transport 177
 nutrient uptake 177,392
 nutritive substances 375
 nutritive value 375,382
 Nylon-6 174
 O
 O₂ concentration 281,282,286
 O₂ consumption 281
 O₂ content 74
 O₂ evolution 178,186,191,194,272,274,275,279,283
 O₂ fumigation 279
 O₂ pressure 197
 O₂ production 71,164,272
 O₂ supply 387
 O₂ uptake 172,282,283,362
 o-acetylserine see OAS
 OAS 48,321,322,323,325
 OAS sulfhydrylase 326
 octamers 342
 n-octanol 294
 Oedogonium 116
 oil 175
 oligonucleotides 314
 oligophosphates 13,311
 oligosaccharides 254
 Ondatra 385
 OPH 325,326,327
 OPH sulfhydrylase 50,326
 o-phenanthroline see phenanthroline
 o-phosphohomoserine see OPH
 o-phosphoric acid see phosphoric acid
 optimum temperature 183
 orellanine 166,173
 Oreochromis 388
 organic acid 305
 organic components 20
 organic anions 42
 organic nitrogen 112
 organic substances 56,83,120,135,293,398
 orientin 36,39,202,354,355,356,359
 ornithine 139,140,343
 ornithine-urea cycle 343
 Oronectes 389
 orotic acid 167
 orthophosphate 311,314
 oscillations 289,347,363
 oscillators 111
 osmotic pressure 302
 osmotic stress 219,223,240,303
 o-succinyl-L-homoserine see under succinyl
 ouabaine 264
 oxalate(s) 42,142,290,306,307,383
 oxalate accumulation 125,306
 oxalate content 115
 oxalate oxidase 42
 oxalate synthesis 306,307
 oxalic acid 43,98,316,383
 oxidation/carboxylation ratio 106,179
 2-oxoglutarate 316,317,318
 2-oxobutyrate 327,329
 oxyfluorfen 293
 oxygen see O₂
 oxyproline 23

ozonated hexene 77
 ozone 22,77,204,273,274,352
 ozone sensibility 77
 ozone-hexene mixture 274

P
 P 10,12,13,113,142,298,391,392,396,398
 P carrier 133
 P compounds 30,163
 P concentration 114,116,218
 P content 10,14,15,17,18,114,116,336
 P deficiency 115,116,222,235,303,304,312,313,314,344,349,359,404
 P incorporation 116
 P limitation 196
 P removal 391,393
 P requirement 86,113,114,116
 P shortage 114,115,116
 P starvation 116
 P supply 115,116,221,241
 P tolerance 113
 P uptake 116
 P/Fe ratio 159
 PO₄ 13,17,18,19,30,45,113,246,260,265,285,291,300,306,326,327,349
 PO₄ acquisition 312
 PO₄ assimilation 311
 PO₄ buffer 233
 PO₄ concentration 88
 PO₄ deficiency 222,223,273,339,340
 PO₄ esters 30,311
 PO₄ ions 401
 PO₄ pollution 401
 PO₄ solubility 81
 PO₄ storage 307,311
 PO₄ supply 251,273,302
 PO₄ uptake 108,116,284,290,291,292,363
 PAA 83,148
 PABA 147,215
 PAG 325,326
 PAL 50,127,203,352,353,366
 palisade parenchyma 228
 palmitic acid 28,350
 P-aminobenzoic acid see PABA
 papillae 69,144,146,211
 PAPS 321
 papules see papillae
 PAR 188
 paraquat 161,162,166,173
 parathion 169
 parenchyma 228
 parthenocarpic development 269
 pathogens 383,394
 patulin 169,173
 Pb 10,12,15,19,131,300,394,396,398
 Pb acetate 131
 Pb concentrations 131
 Pb toxicity 131

Pb uptake 131
 PbNO₂ 131
 PCB 164,170,198,399
 p-chlorophenoxyisobutyric acid see PCPIB
 PCIB 148,212,254
 p-coumaric acid see under C
 p-coumaroylquinic acid see under C
 p-coumaryl alcohol see under C
 PCP 170
 PCPIB(A) 146
 peat 142
 pectic polysaccharides 307
 pectin(s) 27
 pectin fraction 27
 pectin methyl esters 331
 pectinase 50,154,220,308
 pentachlorophenol see PCP
 pentamers 342
 pentose-phosphate pathway 283,362
 PEP carboxylase 50,74,223,306
 peptides 320,347
 PER 47,53,54,55
 permeability 159,166
 permethrine 171
 peroxidase 50
 peroxisomes 306
 pesticides 170
 petunidin 35,354,355
 petunidin glucosides 35,36
 Pfr see phytochrome
 PH 40:62 169,170
 PH 63,81,90,105,106,109,110,111,119,120,122,129,130,132,140,147,149,
 197,215,216,255,284,285,291,294,304,305,319,320
 pH effects 83
 pH growth limits 82,84
 pH optimum 81
 pH tolerance 82
 phaeophytin 34
 Phorbis purpurea 253
 Pharmaceutical substances 402
 phasionids 386
 Phellodendri cortex 175
 o-phenanthroline 244,251
 phenol(s) 164,169,173,174,353,399
 phenol derivatives 162,164
 phenol test 41
 phenol uptake 83
 phenolases 283
 phenolic substances 244,251
 phenoxacetic acid 294
 phenyl groups 140
 phenylacetic acid see PAA
 phenylalanine 23,24,245,330,352,376
 phenylalanine ammonia lyase see PAL
 phenylenediamine 275
 phenylglyoxal 244,253

- phenylpropanes 353,354
- phenylpropanoids 352
- phenylpropionic acid see PPA
- phenylthiocarbamide 359
- phenylthiourea 172
- phlobaphene-like substances 35,56,203,219
- phloem 298
- phloroglucinol-HCl technique 42
- phosphate see PO₄
- phosphatase(s) 50,308,312,313,314,362,363
- phosphatase activity 115
- phosphatase location 313
- phosphatidic acid 30,349
- phosphatidyl choline 30,311,330,331,349
- phosphatidyl ethanolamine 30,167,311,349
- phosphatidyl glycerol 30,167,311,349
- phosphatidyl inositol 30,167,311,349
- phosphatidyl serine 30,349
- phosphodiesterase 50
- phosphoenolpyruvate carboxylase see PEP carboxylase
- phosphofructokinase 50
- 6-phosphogluconate dehydrogenase 223,362
- phosphoglyceric acid 30,274,278,311
- o-phosphohomoserine see OPH
- phospholipid(s) 28,30,153,311,349
- phospholipid content 29,230
- phospholipid synthesis 349
- N-phosphonomethylglycine see PMG
- N,N-bis [phosphonomethyl]glycine see glyphosine
- o-phosphoric acid-monoester phosphohydrolase 50,404
- phosphorus see P
- phosphorylase 50
- phosphorylcholine 331
- photoassimilation 178
- photoinhibition 194,271,276
- photokinetic activity 233
- photon flux density 118,188
- photoperiodic sensitivity 129
- photoperiodism 260,361
- photoperiods 339
- photophosphorylation 260
- photoreactivation 348
- photoreceptor 232,283
- photoregulation 201
- photorespiration 278,279,280,306
- photosynthates 357
- photosynthesis 90,112,118,138,141,159,167,174,177,190,194,196,197,228,233,239,259,270,287,291,304,305,317,320,323,333,352,357,358,406
- photosynthesis activity 78,91,171
- photosynthesis rate see photosynthetic rate
- photosynthetic area 71
- photosynthetic electron flow 168
- photosynthetic mutant 34,232
- photosynthetic pigment(s) 166,202,233,356,358
- photosynthetic process 274
- photosynthetic rate 71,72,74,77,87,88,100,102,103,114,115,116,120,124,165,167,178,186,191,197
- photosystem I 274,275,279,335
- photosystem II 131,272,274,275,276,334,336
- phragmites 373
- phthalate 174
- phycobilliproteins 346
- p-hydroxybenzaldehyde see under H
- p-hydroxybenzoic acid see under H
- p-hydroxycinnamic acid see under H
- physiological characteristics 58
- physiological tests 379
- phytadiene 29
- phytate see phytic acid
- phytic acid 30,308,311,312
- phytochemical characteristics 9
- phytochrome 31,34,38,45,206,218,229,236,241,261,263,284,287,333,356,361,362,366
- phytochrome action 201
- phytochrome content 201
- phytohormones 43,148,243,294
- phytoplankton 388,393,408
- phytol 29,57,350
- PI (inorganic phosphate) 13
- Picrin test 57
- pigment cells 35,40,56
- pigment content 164
- pigment systems 188
- pigs 376,384,394
- Pirson-Seidel solution 21,60,61,62,260
- pisciculture 394
- Pistia 398,407
- pistil 268,269
- pistil development 158
- plasma damage 186
- plasma membrane permeability 224
- plasmalemma 57,89,105,109,133,134,141,283,284,285,287,289,290,291,293,312
- plasmalemma flux 158
- plasmid DNA 348
- plasmid uptake 347,348
- plasmolysis 302,303
- plasmolysis time 87,88,303
- plastid(s) 141,201
- plastid division 228,230
- plastid polypeptides 333,335
- plastoquinone 232,275,353
- PMG 163,166,168,293,328,330
- podagra 402
- polarized light 232,233,272,283
- pollen cells 269
- pollination 269
- pollution 370,390
- polyacrylamide gel electrophoresis 338,339,344
- polyamine(s) 211,225,244,251,254,331,343
- polyamine biosynthesis 330
- poly(A)RNA 333,334,336,342

polychlorinated biphenyls see PCP
 polyethylene glycol 28,240,350,352
 polyethylene oxide 303
 polygalacturonic acid 310
 Polygalae radix 175
 polyhydroxy xanthophyll 33
 polypeptid clusters 55
 polypeptid composition 55
 polyphenolase 50
 polyphenolic substances 57
 polyribosomes 30,342
 polysaccharides 307
 poly(U)sepharose 342
 polyuridylic acid 342
 polyuronic acid 299
 Potamogeton 210
 Potamogetonaceae 27,108
 potassium see K
 potassium cyanide see KCN
 poultry 386
 PPA 148
 Pr 10,300,396
 praeodinium see Pr
 prawns 388
 prenylquinones 353
 preservation technique 65
 primordial cells 228
 proanthocyanidin 173
 Procambius 389
 processing 380
 procyanidin 173
 productivity 371
 Prochloron 346
 projection (of lower surface) 81,105
 prolamellar body 228
 prolamine 21
 proline 24,139,141,245,258
 prolongation effect 81
 prometryne 161,162,167,274,293
 propargylglycine see PAG
 propelargonidin 173
 propionic acid 175
 proplastids 228
 propylgallate 332
 prostaglandin-like fatty acid 29
 protein(s) 12,20,21,53,54,104,140,197,239,323,330,331,380,386
 protein amino acids 112
 protein bands 337
 protein content 17,10,21,77,106,115,124,149,153,159,167,168,174,192,
 199,204,223,224,278,306,337,338,375,377,381,382,384
 protein degradation 22,87,103,165,167,220,222,223,224,338
 protein efficiency ratio 377
 protein extraction 383
 protein kinase 50,336,337
 protein metabolism 153,224,337
 protein nitrogen 98
 protein nutrition 386
 protein phosphokinase 336
 protein phosphorylation 336
 protein synthesis 22,87,117,153,160,173,205,220,222,223,224,229,267,
 332,338,343,359,366
 protein turnover 222,224,332
 protein yield 377
 proteinases 284,287
 proton(s) 284,287
 proton co-transport processes 116,134,141,284,290,293,294
 proton electrochemical gradient 141,285
 proton extrusion pump 105,109,284
 protoplast 308
 Pruni cortex 175
 Pseudomys scripta 389
 p-trifluoromethoxy carbonylcyanide phenylhydrazine see trifluoromethoxy
 Pu 396
 purine 112,150,163
 purine analogs 357,358
 purine bases 141,294
 puromycin 351
 putrescine 112,251,320,330,331
 Pyrethri flos 175
 pyridine analogs 358
 pyridine nucleotides 30,276
 2-pyridylhydroxymethane sulfonic acid see α HPMS
 pyridoxal phosphate 283,319
 pyridoxamine phosphate 319
 pyrimidine(s) 112,167,205
 pyrimidine base analogs 167,357
 pyrimidine nucleotides 230
 pyrocatechol 174
 α -pyrone fluorescence method 295
 pyrophosphate 50,115
 pyrophosphate 113,314
 pyrophosphate phosphohydrolase 50
 pyruvate 205,326,328,329
 pyruvate kinase 50,362
 pyruvic acid see pyruvate
 Q
 QAC 162,164,166,169,170
 quantum yield 195
 quaternary ammonium compound see QAC
 quercetin 36,202,354,355,356,359
 quinol 353
 R
 Ra 10
 rabbits 385
 radiation 188
 γ -radiation 208
 radioactive contamination 210
 radioactive substances 209
 radionucleids 209
 radiosensitivity 98,209,406

radium see Ra

ram 385

raphide idioblast 43

raphides 43,307

Rb 10,12,90,365

Rb uptake 287,289

recycling 391,394

red light 189,200,201,202,203,205,206,211,212,213,218,222,228,229,233,
235,236,237,241,261,262,263,276,278,286,287,324,333,334,342,343,356,
357,361,364,368

redox pump 286

rejuvenation 220

removal (of metals and toxicants) 391,394

resorcinol 174,353

respiration 159,164,172,177,233,279,281,362,363,365,367,406

respiration rate 100,103,115,124,129,165,174,177,186,192,281,282

respiratory inhibitors 246,365

respirometer 270

resting fronds 30,31,158,184,200,235

restriction fragment map 345,346

rhamnose 26,27,283

Rhei rhizoma 175

rheumatism 403

riboflavin 201,357,358,359

ribonuclease see RNase

ribonucleic acid see RNA

ribonucleosides 229

ribonucleoprotein 30

ribose 136,283

ribosomal ribonucleic acid see rRNA

ribosomes 319,336,341

ribulose-1,5-diphosphate see RuDP

ribulose-1,5-diphosphate carboxylase see RuDPCase

ribulose-5-phosphate kinase 51

rice 376,377

Rieske center 275

right-handed 215,217

RNA 30,201,311,314,338

RNA biosynthesis 159,160,357

RNA content 77,153,174,224

RNA degradation 225

RNA metabolism 339

RNA polymerase 50,98,347,367

RNA synthesis 153,205,223,224,225,347,357,358,366

RNA synthetic activity 98

RNase 50,312,313,314,343,357,358

RNase activity 344

root(s) 79,81,89,113,180,204,298,312

root cells 103,115,302

root elongation 149,303

root growth 125,127,128,129,132,200,218,219,400

root length 69,71,81,88,89,90,93,102,103,104,105,114,115,116,144,145,
148,157,159,165,174,192,200,218,219,400

root morphogenesis 217

root number 144,145,217

root primordia 217

rRNA 223,314,339,366,367

rRNA degradation 223,225

rRNA genes 345,346

rRNA synthesis 147,223,225

rubidium see Rb

Ru 210,396

RuDP 273,279

RuDPCase 51,55,201,226,332,333,334,335,336,342,346,347

ruthenium see Ru

S

S 10,12,13,43,56,116,133,140,298

S²⁻ 117,320,321,322,323,325,326

S assimilation 320

S concentration 117

S deficiency 22,117

S fixation 118

S requirements 86,117

S-uptake 118

SH groups 134

SO₂ 76,273,320,323

SO₂ concentration 77

SO₂ fumigation 274,292,304,323

SO₄²⁻ 117,320,321,322,323

SO₄²⁻ detoxification 118,320

SO₄²⁻ reductase 321,323

SO₄²⁻ tolerance 118

SO₄²⁻ 12,116,125,274,285,291,314,320,321,322,323,325,326

SO₄²⁻ absorption 117,124

SO₄²⁻ assimilation 321,322

SO₄²⁻ concentration 88,292

SO₄²⁻ content 117

SO₄²⁻ deficiency 222,235,237,241

SO₄²⁻ efflux 292

SO₄²⁻ metabolism 292

SO₄²⁻ uptake 77,117,118,284,290,291,292

SA 129,142,144,171,211,212,219,220,244,252,253,254,265,267,353

saccharose see sucrose

s-adenosylhomocysteine see SAH

s-adenosylmethionine see SAM

SAH 322,330

salicylaldehyde 144,245,282

salicylhydroxamic acid see SHAM

salicylic acid see SA

Salmonella 383,394

salt stress 91,273

Salvinia 132

SAM 322,327,330,331

SAM-WTA-NTR cycle 330,331

SAM-SAH-HC cycle 330

Sandoz 6706 351

saponaretin 36

saponarin 35

saponins 56

Sb 10,300,396

Schizonepetae herba 175

Sclerids 170,399
 sclerids 42
 sclerenchymatic tissue 41
 scopolin 174
 scorbut 403
 Scutellariae radix 175
 SDP 90,144,200,248,249,250,251,252,255,258,260,261,263,264,267,368
 Se 10,133,140,396,398
 seasonal effects 68
 seasonal factor 236
 seed(s) 184,235,269
 seed coat 242
 seed dormancy 241
 seed germination 180,217,241
 selenate 133,323
 selenium see Se
 selenite 133,323
 selenocystine 133,323
 selenomethionine 133,323,332
 semipermeability 77
 Senegae radix 175
 senescence 28,104,115,220,235,236,238,248,249,251,256,303,314,350,352
 serine 23,24,141,245,257,258,278,318,321,323
 serine amide 112
 serine glyoxylate aminotransferase 51
 serine hydroxymethyltransferase 140,326
 serine phosphate 336
 serine proteinases 254
 serine transacetylase 321
 sequestrene 143
 sevin 175
 sewage 390,391,392,393
 SHAM 211,282,284
 shark teeth phosphorus 113
 sheep 385
 Shigella 383,394
 short-day conditions 120,172,200,216,218,222,226,227,235,236,238,240,
 248,250,251,252,255,257,258,259,261,262,263,264,265,266,267,277,286,
 312,339
 short-day plant see SDP
 shrimp 389
 Si 10,12,134
 sieve elements 298
 silage 381
 silicic acid 134
 silicium see Si
 silver see Ag
 simazine 161,162,167,274,293
 Sinapis semen 175
 I-sinapoylglucose 57,353
 sinapoyl alcohol 42
 β -sitosterol 29,57,350
 skeleton photoperiods 264
 smog (synthetic) 77,273
 Sn 10,300,396
 sodium see Na
 sodium acids see NaN,
 soil extract 142
 solar energy conversion 188,375
 somatostatin-like substances 46
 sorbose 136,225,231
 soybean 376,377
 space flight 406
 space-labor conditions 406
 Spaeotillus natans 403
 spermidine 46,225,251,330,331
 spermine 251,330,331
 spinach 47,346
 spongy parenchyma 228
 squalene 29,351
 Sr 10,43,86,98,208,396
 stalk 148,151,202
 stamen 268
 stamen development 158
 Staphylococcus aureus 403
 starch 136,138,178,180,272,304
 starch accumulation 116,153,218,230,274,304
 starch content 20,25,26,71,76,83,103,115,124,153,165,204,235,304,375
 steady-state culture 65
 Stellaria media 403
 Stegomyia 408
 stele 298
 sterols 56
 sterol(s) 29
 sterol content 156,350,351
 stigmasterol 29,350,351
 stipe 114,149,157,211,219
 stipe development 155
 stipe length 211
 stocking density 374,379,380
 stomata 159,182,212,213,270,271,302
 stomata number 199
 stomata size 199
 stomatal index 212,213
 Streptococcus 383,394
 streptomycin 66,173,229,357
 stress factors 220,223,224,226,319,339,352
 stress hormone 273
 stroma 164,229,235,342
 stroma thylakoids 274
 strontium see Sr
 submerged species 64,74,75,120
 submerged culture 65
 succinate 136
 o-succinyl-L-homoserine 326
 sucrose 26,44,81,109,136,137,138,176,183,186,195,198,199,204,205,208,
 215,216,218,219,220,225,238,240,242,245,257,258,259,260,265,266,268,
 269,276,277,278,282,284,286,289,305,314,319,332,349,350,357,358,366
 sucrose concentration 136,137
 sucrose supply 82,239,312,315,339,340,352,357,365
 sucrose uptake 129,138,294
 sugar(s) 25,26,27,65,71,106,205,215,218,225,238,239,278,293

sugar content 26,27,103,115,153,168,193,304
 sugar metabolism 153
 sugarphosphate esters 278
 sulfaguanidine 150
 sulfanil amide 117,150
 sulfanilic acid 117
 sulfapal 282
 sulfate see SO₄²⁻
 sulfathiazole 150
 sulphydryl blockers 245,258
 sulphydrylase 51
 sulfide see S²⁻
 sulfite see SO₃²⁻
 sulfocyanate 240
 sulfolipids 28,322,323
 sulfonamides 172
 sulfonyl group 322
 sulfur see S
 sulfur dioxide see SO₂
 sulfuric acid see H₂SO₄
 sulph- see sulf-
 super surf 169,172
 superoxide dismutase 51,128,142,154,159
 survival temperatures 184
 swellings 403
 syphilis 403
 syringaldehyde 41,42

T
 2,4,5-T 165,399
 tafapon 170
 tafazine 170
 TAL 51,352,353,366
 tannic acid 174,244,253
 tannin(s) 56
 tannin idoblasts 56
 tartrate 122,142,143
 taste 383
 taurine 23
 TCDD 165,399
 2,4,6-TCP 169,170
 TD-1929 168
 temperature 106,176,194,222,226,229,230,232,238,239,242,264,267,272,
 291,303,307,312,361,365,366
 temperature dependence 176,192,194,195
 temperature extremes 239
 temperature influence 176
 temperature tolerance 184,186,187
 temporal compartmentation 361
 tentoxin 173
 terbutryn 162,168
 terpenes 56
 test plant 400
 testosterone 405
 2,3,7,8-tetrachlorodibenzo-p-dioxine see TCDD
 tetracycline 173,229

tetrametaphosphates 13
 tetraphophylene benzolsulfonate 172
 tetrathionate 117
 tetrazolium 275
 tetrazolium oxidase see TO
 thermal treatment 230
 thiamine see vitamin B₁
 thiazole 146,205
 thio compounds 12
 thiocarbonic acid derivatives 164
 thiokinetic 150
 thiol(s) 322
 thiol content 118
 thiol production 118,320
 thiosulfate 117
 thiosulfonate reductase 321
 2-thiouracil see TU
 thiourea 117,295
 D-threo-chloramphenicol
 threonine 23,24,140,245,257,258,324,325,326,327,329,376
 threonine deaminase 329
 threonine phosphate 337
 threonine synthase 51,327
 threonyl peptides 165
 thylakoid(s) 138,164,167,225,228,229,230,235,335,341,353
 thylakoid degradation 225
 thylakoid membranes 167,334,350
 thylakoid protein 226,275
 thylakoid structure 352
 thylakoid superoxide dismutase 51
 thymidine 167,224,230,269,343
 thymine-thymine dimers 348
 Ti 10,12,300,396
 TIBA 83,148,214,215,216,294,295
 Tilapia 388
 tissue cultures 66,173
 tissue differentiation 212
 titanium see Ti
 TO 47,51,53,54
 tobacco 175
 tobacco smoke 78,175
 p-toluol sulfonic acid 172
 tonoplast 57,289
 tonoplast permeability 224,300
 toxic gases 71
 toxic substances 135,161,172
 toxicants 398,400
 TPM 170
 TPP 399
 trace elements 122
 transaminases 224,324
 trans-cinnamic acid 366
 transcription 347
 transmembrane potential 173,286
 trans-1,3-phytadiene 57,350
 transpiration 177,302

Trapa 407
 trehalase 51
 triacylglycerol 29
 triazine 167,275
 2,4,6-trichlorophenol see TCP
 2,3,5-triiodobenzoic acid see TIBA
 2,4,5-trichlorophenoxyacetic acid see 2,4,5-T
 trimers 342
 trimetaphosphates 13
 trimethylamine alkylbenzenesulfate see DBST
 trimethyl- β -chloroethylammonium chloride see CCC
 triphenylphosphate 399
 tripolyphosphate 113,314
 tris 143
 triterpenes 350
 tRNA 338,339
 tRNA genes 345,346
 tryad 166
 tryptin 43
 tryptophane 24,139,140,245,257,330,375,376
 TU 162,167,246,266,267,343
 tungstate see WO
 tungsten see W
 turgor pressure 302
 turion(s) 25,30,31,184,186,235,296,298,299,312,405
 turion dormancy 239
 turion formation 100,101,103,104,114,115,117,125,152,154,158,160,179,
 181,237,338,400,406
 turion germination 104,152,179,182,239
 turion induction 160,235,359
 turion-like fronds 235,239,312
 turion size 114
 turion weight 114
 turtles 389
 Typha 407
 tyramine 141,294
 tyrosine 23,24,141,245,330,352,353
 tyrosine ammonia-lyase see TAL

 U
 UDP
 UDP-D-apiose 308,309,310
 UDP-apiose/UDP-xylose synthase 51,308
 UDP-galacturonic acid 310
 UDP-glucose 309,310
 UDP-D-glucuronate carboxylase 51,310
 UDP-D-glucuronic acid 308,309,310
 UDP-D-glucuronic acid carboxy-lyase 51
 UDP-D-glucuronic acid cyclase 51
 UDP-D-xylose 308,309,310
 uracil 167,343
 Uranotaenia 409
 urea 12,99,112,113,141,245,257,294,295,316,332
 urea assimilation 316
 urea derivatives 161,162,164
 urease 51,319,320

 o-ureido-L-homoserine 343
 uridine 98,141,147,167,224,225,230,339,340,343,347,366
 uridine deoxyriboside 229
 uridine diphosphate see UDP
 uridine kinase 51
 uridine monophosphate see UMP
 uridine uptake 295
 UTP 309
 UV irradiation 34,44,66,203,219,220,222,226,348,351,352,353,357

 V
 V 10,133,300,396
 vacuole 117,298,299,300,311
 valine 23,24,113,139,140,245,258,327,328,376
 valine synthesis 140
 valine-leucine biosynthesis 140,327
 valine-leucine pathway 328
 valine tRNA 346
 valine uptake 293
 valinomycin 246,265,270,271
 vanadate/H⁺ cotransport 133
 vanadium see V
 vanillic acid 174
 vanillin 41,42
 vascular cells 41
 vascular system 212,214,215,343
 vegetative growth 67
 vibrations 406
 vicinin 36
 vinylglycine 326
 violaxanthin 33,202,276,277,351
 violet light 263
 virus 383
 vitamin(s) 43,135,146,204,382
 vitamin A 43,146,378
 vitamin B₁ 43,146,200,205,206,378
 vitamin B₂ 43,146,147,378
 vitamin B₃ 147
 vitamin B₆ 43,146,378
 vitamin B₁₂ 146
 vitamin C 43,146,147,245,259,296,307,378
 vitamin E 43,378
 vitamin K 43,147,246,266
 vitamin PP 43,378
 vitexin 35,36,39,202,354,355,356,359

 W
 W 125
 WO 125,245,256,257,258,305,315
 W7 363
 waste water 390,402
 waste water systems 392,398,399
 water content 9,167,239
 water deficit 270
 water flow 177
 water fowl 386

water loss	407,408	
water permeability	177	
water purification	370,390,392,393,398	
water relations	302	
water transport	79	
weight	see frond weight	
weightlessness	406	
whewellite	43	
white anur	see grass-carp	
white light	189,202,203,231,241,263,276,324,334,335,342,343,356,259,362	
X		
X irradiation	208	
xanthine-xanthine oxidase system	142	
Xanthium	352	
xanthone group	174	
xanthophyll(s)	33,46,386	
xanthophyll cycle	276,277	
xanthoxin	46	
Xim Ha	383	
XMC	171	
X-ray microanalysis	298	
X-rays	216,217,269,406	
xylem	298	
xylene	168	
xylose	26,27,136,309,310	
xylosyltransferase	51	
3,5-xylyl methylcarbamate	see XMC	
Y		
Y	10,196,300	
yeast extract	65,205,206,207,208,242	
yellow light	202	
yttrium	see Y	
Z		
Zannichelliaceae	27,108	
Zea mays	33,346	
zeatin	45,146,147,150,217,244,266,349	
zeaxanthin	33,276,277	
zinc	see Zn	
zirconium	see Zr	
Zn	10,12,15,17,19,86,98,122,127,130,132,142,159,219,300,301,395,396,398	
Zn concentration	88,127	
Zn deficiency	128,359	
Zn requirement	86	
Zn tolerance	123,128	
Zn toxicity	128	
zooplankton	388,402	
Zr	10,300,396,398	

CONTENTS OF VOLUME 1 (E. LANDOLT, 1986.		
Veröff.Geobot.Inst.ETH,Stiftung Rübel, Zürich, 71)		
1. Introduction - Acknowledgements	7	
2. Morphological characteristics: variability and function	13	
2.1. General	13	
2.2. Structure and organization of Lemnaceae plants; interpretation of different organs	13	
2.2.1. Description of the frond	13	
2.2.2. Interpretation of the frond	20	
2.3. Growth and ageing of fronds; formation of daughter fronds	27	
2.3.1. Mode of growth; growth rate	27	
2.3.2. Lifespan and ageing	28	
2.3.3. Number of daughter fronds	31	
2.3.4. Position of daughter fronds	31	
2.4. Characteristics of normal vegetative fronds	36	
2.4.1. Form, size and weight	36	
2.4.2. Elements of the cells	41	
2.4.3. Colour and brightness; pigment cells	46	
2.4.4. Epidermis, stomata and papillae (papules)	48	
2.4.5. Air spaces and parenchymatic tissue	54	
2.4.6. Nerves and tracts of elongated cells in the basal section of fronds; vascular and sieve cells	58	
2.4.7. Oxalate-crystal idioblasts	61	
2.4.8. Pouches (pockets) and cavities protecting daughter fronds	64	
2.4.9. Connection between mother and daughter frond (stipe)	66	
2.4.10. Prophyllum in Spirodela	67	
2.4.11. Roots	69	
2.4.12. Organs of stabilization	77	
2.5. Characteristics of resting fronds and turions	80	
2.5.1. Different kinds of resting fronds	80	
2.5.2. Factors causing formation of turions	84	
2.5.3. Factors causing germination of turions	86	
2.6. Characteristics of flowering and fruiting fronds	89	
2.6.1. Aspects of flowering fronds	89	
2.6.2. Morphology and interpretation of flowering organs	92	
2.6.3. Development of flowers and fruits	105	
2.6.4. Ripening of flowers	113	
2.6.5. Fruits and seeds	115	
2.6.6. Germination of seeds	123	
2.7. Use of morphological features in taxonomy of Lemnaceae; preparation of transparent slides	127	
3. Karyology		
3.1. Chromosome numbers	129	
3.2. Morphological characteristics and DNA content of chromosomes	129	
4. Habitat demands and ecological behaviour	137	
4.1. General	138	
4.2. Vegetative growth	138	
4.2.1. Physical characteristics of the water body	139	
4.2.2. Temperature conditions	142	

4.2. 3. Light condition	147	6.1.3.28. <u>W. brasiliensis</u>	357
4.2. 4. Chemical composition of the water	149	6.1.3.29. <u>W. borealis</u>	362
4.3. Sexual reproduction	167	6.1.3.30. <u>W. australiana</u>	363
4.3. 1. Flowering conditions	167	6.1.3.31. <u>W. angusta</u>	365
4.3. 2. Pollination and fruit setting	178	6.1.3.32. <u>W. arnhila</u>	366
4.4. Dispersal and survival	181	6.1.3.33. <u>W. columbiana</u>	371
4.4. 1. Dispersal	181	6.1.3.34. <u>W. globosa</u>	377
4.4. 2. Survival during unfavourable conditions	183	6.2. Altitudinal distribution	381
5. Position within the ecosystem	186	6.2. Relationship between ecological demands, climatic factors, and geographic distribution	392
5.1. Ecosystems with Lemnaceae	186	6.3. 1. Local factors interfering with the general distribution	392
5.2. Specific abiotic factors in ecosystems with Lemnaceae	189	6.3. 2. General climatic factors limiting the distribution of Lemnaceae	395
5.3. Relationship of Lemnaceae to other organisms	192	6.3. 3. Distribution within the climate zones	402
5.3. 1. Herbivores and parasites	192	6.4. Dispersal and threat by man	406
5.3. 2. Mutualists, commensalists, and amensalists	195	7. Systematic position, taxonomical relations, species differentiation	410
5.3. 3. Competitors	201	7.1. Paleobotanical records	410
5.4. Relationship between members of the family of Lemnaceae and other small, free-floating, vascular plant species	204	7.2. Position within the plant kingdom	412
5.5. Plant communities with Lemnaceae	207	7.3. Taxonomical relationship within the family	417
5.5. 1. Characteristics of Lemnaceae communities and principles of classification	207	7.3. 1. Differentiation of the genera	417
5.5. 2. Survey of Lemnaceae communities	215	7.3. 2. Differentiation within the genera	421
5.5. 3. Influence on Lemnaceae communities by human activities	228	7.3. 3. Differentiation within the species	429
6. Geographic distribution; distribution patterns	229	8. Nomenclature	436
6.1. Geographic distribution	229	8.1. Typification and validity of names	436
6.1. 1. Procedure and methods	229	8.1. 1. General remarks	436
6.1. 2. Distribution of the family; species diversity	233	8.1. 2. Names of family, subfamilies, genera and species	437
6.1. 3. Distribution of the different species	238	8.2. List of names and synonyms	455
6.1.3. 1. <u>Spirodela intermedia</u>	238	9. Description of family, genera, and species of Lemnaceae with key to the	461
6.1.3. 2. <u>S. polyrrhiz</u>	243	9.1. Description of the family of Lemnaceae with key to the subfamilies	461
6.1.3. 3. <u>S. punctata</u>	255	9.2. Subfamily of Lemnoideae	463
6.1.3. 4. <u>Lemna gibba</u>	262	9.2. 1. Description of the subfamily of Lemnoideae with key to the genera	463
6.1.3. 5. <u>L. disperma</u>	273	9.2. 2. Genus <u>Spirodela</u> Schleiden	466
6.1.3. 6. <u>L. minor</u>	275	9.2. 3. Genus <u>Lemna</u> L.	471
6.1.3. 7. <u>L. japonica</u>	283	9.3. Subfamily of Wolffioideae	530
6.1.3. 8. <u>L. ecuadoriensis</u>	284	9.3. 1. Description of the subfamily of Wolffioideae with key to the genera	530
6.1.3. 9. <u>L. obscura</u>	284	9.3. 2. Genus <u>Wolffiella</u> Hegelmaier	532
6.1.3.10. <u>L. turionifera</u>	287	9.3. 3. Genus <u>Wolffia</u> Horkel	544
6.1.3.11. <u>L. trisulca</u>	292	References. All the references and a complete survey of the Lemnaceae bibliography are given in vol. 2	553
6.1.3.12. <u>L. perpusilla</u>	301	Abbreviations (especially chemical substances)	555
6.1.3.13. <u>L. aequinoctialis</u>	303	List of the figures in vol. 1	558
6.1.3.14. <u>L. tenera</u>	320	List of the tables in vol. 1	559
6.1.3.15. <u>L. valdiviana</u>	321	Index of subjects and names of plants	565
6.1.3.16. <u>L. minuscula</u>	327	Contents of vol. 2 (E. LANDOLT and R. KANDELER, 1987. Veröff. Geobot. Inst. ETH, Stiftung Rüböl, Zürich, 95)	
6.1.3.17. <u>Wolffiella hyalina</u>	335		
6.1.3.18. <u>W. repanda</u>	336		
6.1.3.19. <u>W. rotunda</u>	337		
6.1.3.20. <u>W. neotropica</u>	337		
6.1.3.21. <u>W. Welwitschii</u>	339		
6.1.3.22. <u>W. lingulata</u>	343		
6.1.3.23. <u>W. oblonga</u>	347		
6.1.3.24. <u>W. gladiata</u>	352		
6.1.3.25. <u>W. denticulata</u>	353		
6.1.3.26. <u>Wolffia microscopica</u>	355		
6.1.3.27. <u>W. elongata</u>	355		

VERÖFFENTLICHUNGEN DES GEBOTANISCHEN INSTITUTES DER
EIDG. TECHN. HOCHSCHULE, STIFTUNG RÜBEL, IN ZÜRICH

- 1a. (Beiblatt). *Curvuletum*
Von E. RÜBEL 1922 (vergr.)
- 2a. (Beiblatt). Betrachtungen über einige pflanzensoziologische Auffas-
sungs-differenzen. Verständigungsbeitrag Schweden-Schweiz
3a. (Beiblatt). Vorschläge zur Untersuchung von Buchenwäldern
Von E. RÜBEL. 1925 (vergr.)
- 4a. The present state of geobotanical research in Switzerland
Von E. RÜBEL. 1928. Fr. 1.-
1. Ergebnisse der internationalen pflanzengeographischen Exkursion
durch die Schweizer Alpen 1923
Redigiert von E. RÜBEL. 1924 (vergr.).
2. Vegetationsstudium im Limmattal
Von M. SCHERRER. 1925 (vergr.)
3. Festschrift Carl Schröter
Redigiert von H. BROCKMANN-JEROSCH. 1925 (vergr.)
4. Ergebnisse der internationalen pflanzengeographischen Exkursion
durch Schweden und Norwegen 1925
Redigiert von E. RÜBEL. 1927. Fr. 8.-
5. Pollenanalytische Untersuchungen an Schweizer Mooren und ihre
florensgeschichtliche Deutung
Von P. KELLER. 1928. Fr. 5.-
6. Ergebnisse der internationalen pflanzengeographischen Exkursion
durch die Tschechoslowakei und Polen 1928
Redigiert von E. RÜBEL. 1930. Fr. 7.-
7. Die Flora Graubündens
Von J. BRAUN-BLANQUET und E. RÜBEL. 1932 (vergr.)
8. Die Buchenwälder Europas
Redigiert von E. RÜBEL. 1932. Fr. 10.-
9. Die postglaziale Entwicklungsgeschichte der Wälder von Norditalien
Von P. KELLER. 1931. Fr. 4.-
10. Ergebnisse der internationalen pflanzengeographischen Exkursion
durch Rumänien 1931
Redigiert von E. RÜBEL. 1933. Fr. 3.-
11. Das Grosse Moos im westschweizerischen Seeland und die Geschichte
seiner Entstehung
Von W. LÜDI. 1935. Fr. 8.-
12. Ergebnisse der internationalen pflanzengeographischen Exkursion
durch Mittelitalien 1934
Redigiert von E. RÜBEL. 1936. Fr. 4.-
13. Anton Schneeberger (1530-1581), ein Schüler Konrad Gesners in Polen
Von B. HRYNIEWIECKI. 1938. Fr. 2.-
14. Ergebnisse der internationalen pflanzengeographischen Exkursion
durch Marokko und Westalgerien 1936
Redigiert von E. RÜBEL und W. LÜDI. 1939 (vergr.)
15. Die Geschichte der Moore des Sihltals bei Einsiedeln
Von W. LÜDI. 1939. Fr. 3.-
16. Zur Geschichte des Waldes im Oberhasli (Berner Oberland)
Von E. HESS. 1940. Fr. 3.-
17. Stratigraphie und Waldgeschichte des Wauwilermooses
Von H. HÄRRI. 1940. Fr. 3.-
18. Klimaverhältnisse im Albisgebiet
Von W. LÜDI und B. STÜSSI. 1941. Fr. 2.-

Addresses of the authors: Prof. Dr. Elias Landolt
Geobotanisches Institut ETH
Stiftung Rübel
Zürichbergstrasse 38
CH-8044 Zürich
Switzerland

Prof. Dr. R. Kandeler
Botanisches Institut
Universität für Bodenkultur
Gregor Mendel-Strasse 33
A-1180 Wien
Austria

19. Pollenanalytische Untersuchung der Gletscherbewegung
Von V. VARESCI. 1942. Fr. 4.-
20. Die Ostgrenze Fennoskandiens in pflanzengeographischer Beziehung
Von A. KALELA. 1943. Fr. 2.-
21. Pollenanalytische, stratigraphische und geochronologische Untersuchungen aus dem Faulenseemoos bei Spiez.
Von M. WELTEN. 1944. Fr. 5.-
22. Les associations végétales de la vallée moyenne du Niger
Par G. ROBERTY. 1946. Fr. 5.-
23. Die Pflanzengesellschaften der Schynige Platte bei Interlaken und ihre Beziehungen zur Umwelt
Von W. LÜDI. 1948. Fr. 12.-
24. Die Geschichte der Moore und Wälder am Pilatus
Von P. MÜLLER. 1949. Fr. 5.-
25. Die Pflanzenwelt Irlands (The flora and vegetation of Ireland)
Redigiert von W. LÜDI. 1952. Fr. 15.-
26. Ueber die spät- und postglaziale Vegetationsgeschichte des Simmentals
Von M. WELTEN. 1952. Fr. 6.-
27. Die Pflanzenwelt des Eiszeitalters im nördlichen Vorland der Schweizer Alpen
Von W. LÜDI. 1953. Fr. 10.-
28. Die Arten der *Bromus erectus*-Wiesen des Schweizer Juras
Von H. ZOLLER. 1954. Fr. 14.-
29. Aktuelle Probleme der Pflanzensoziologie
Herausgegeben von W. LÜDI. 1954. Fr. 6.-
30. Verbreitungsbiologie der Blütenpflanzen
Von P. MÜLLER. 1955 (vergr.)
31. Die Pflanzenwelt Spaniens (I. Teil)
Redigiert von W. LÜDI. 1956. Fr. 16.-
32. Die Pflanzenwelt Spaniens (II. Teil)
Von R. TÜXEN, unter Mitarbeit von E. OBERDORFER. 1958 (vergr.)
33. Festschrift Werner Lüdi
Redigiert von M. WELTEN und H. ZOLLER. 1958. Fr. 15.-
34. Verhandlungen der vierten internationalen Tagung der Quartärbotaniker in der Schweiz 1957
Herausgegeben von W. LÜDI. 1959. Fr. 9.-
35. Ergebnisse der internationalen pflanzengeographischen Exkursion durch die Ostalpen 1956
Redigiert von W. LÜDI. 1959. Fr. 15.-
36. Die Pflanzenwelt der Tschechoslowakei
Ergebnisse der internationalen pflanzengeographischen Exkursion (IPE) durch die Tschechoslowakei 1958
Redigiert von W. LÜDI. 1961. Fr. 11.-
37. Festschrift Franz Firbas
Redigiert von W. LÜDI und O. L. LANGE. 1962. Fr. 25.-
38. Qualität und Quantität der Rehäsung in Wald- und Grünlandgesellschaften des nördlichen Schweizer Mittellandes
Von F. KLÖTZLI. 1965. Fr. 25.-
39. Vegetations- und bodenkundliche Methoden der forstlichen Standortskartierung - Ergebnisse eines internationalen Methodenvergleichs im Schweizer Mittelland
Herausgegeben von H. ELLENBERG. 1967 (vergr.)
40. Natürliche Waldgesellschaften und Waldstandorte der Umgebung von Zürich
Von N. KUHN. 1967 (vergr.)
41. Stickstoff- und Wasserversorgung von Feuchtwiesen (Balance d'azote et d'eau dans les prairies humides - The nitrogen and water relations of wet meadows)
Herausgegeben von H. ELLENBERG. 1968. Fr. 25.-
42. Die Taraxacum-Arten der Schweiz
Von J. L. VAN SOEST. 1969. Fr. 25.-
43. Zur Vegetation und Flora der Westalpen (Contribution à l'étude phytogéographique des Alpes occidentales) - Ergebnisse der 14. internationalen pflanzengeographischen Exkursion (IPE) durch die Westalpen
Redigiert von P. OZENDA und E. LANDOLT. 1970. Fr. 38.-
44. Ecologie comparée des prairies marécageuses dans les Préalpes de la Suisse occidentale
Par M. YERLY. 1970. Fr. 39.-
45. L'azote dans quelques forêts, savanes et terrains de culture d'Afrique tropicale humide (Côte d'Ivoire)
Par P. DE RHAM. 1970. Fr. 25.-
46. Uebersicht der Pflanzengesellschaften der rätischen Alpen im Rahmen ihrer Gesamtverbreitung. III. Teil: Flachmoorgesellschaften (Scheuchzerio-Caricetea fuscae)
Von J. BRAUN-BLANQUET. 1971. Fr. 29.-
47. Vergleichend-ökologische Untersuchungen an Plateau-Tannenwäldern im westlichen Aargauer Mittelland
Von K. PFADENHAUER. 1971. Fr. 21.-
48. Vergleich alpiner Rasen auf Silikat- und Karbonatboden. Konkurrenz- und Stickstoffformenversuche sowie standortkundliche Untersuchungen im Nardetum und im Seslerietum bei Davos
Von A. GIGON. 1971. (vergr.)
49. La végétation du bassin de l'One (Pyrénées centrales). Quatrième note: les forêts
Par R. NEGRE. 1972. Fr. 35.-
50. Versuche einer vergleichend-ökologischen Analyse der Buchen-Tannen-Wälder des Schweizer Jura (Weissenstein und Chasseral)
Von J. PFADENHAUER. 1973. Fr. 28.-
51. Gesellschaften nasser Standorte in den Alpen und Dinariden (Ergebnisse der 13. Tagung der Ostalpin-Dinarischen Gesellschaft für Vegetationskunde 1972)
Redigiert von E. LANDOLT. 1973 (vergr.)
52. Morphologisch-ökologische Untersuchungen an *Scabiosa columbaria* L.s.l.
Von F. GROSSMANN. 1975. Fr. 28.-
53. Zytotaxonomische Untersuchungen an *Campanula patula* L.s.l. und *Campanula Rapunculus* L. in der Schweiz und in Oesterreich
Von M. HAUSER. 1975. Fr. 26.-
54. Ecological conditions limiting the distribution of *Fagus sylvatica* L. and *Abies alba* Mill. near Schwarzenbach (Lucerne), Switzerland
By H. GADEKAR. 1975. Fr. 25.-
55. Zur Vegetation und Flora von Griechenland (Ergebnisse der 15. internationalen pflanzengeographischen Exkursion IPE durch Griechenland). Band 1: Einführung; allgemeine Vegetationskunde; Kreta und Aegäische Inseln
Redigiert von Sp. DAFIS und E. LANDOLT. 1975. Fr. 42.-
56. Zur Vegetation und Flora von Griechenland (Ergebnisse der 15. internationalen pflanzengeographischen Exkursion IPE durch Griechenland). Band 2: Mazedonien, Thessalien-Epirus,

- Attika, Peloponnes; einzelne Pflanzengesellschaften; botanische Literatur über Griechenland.
Redigiert von Sp. DAFIS und E. LANDOLT. 1976. Fr. 44.-
57. Pflanzensozioökologische und ökologische Untersuchungen an insubrischen Trockenwiesen karbonathaltiger Standorte
Von M. MEYER. 1976. Fr. 33.-
58. Fragmenta Phytosociologica Raetica III, IV, VII (Peucedano-Cirsium, Wegbordgesellschaften, Halbtrocken- und Trockenrasen; Verzeichnis der geobotanischen Arbeiten über Graubünden)
Von J. BRAUN-BLANQUET. 1976. (vergr.)
59. Quercus pubescens-Wälder und ihre ökologischen Grenzen im Wallis (Zentralalpen)
Von J. BURRAND. 1976. Fr. 33.-
60. Beschreibung exzentrischer Hochmoore mit Hilfe quantitativer Methoden
Von O. WILDT. 1977. (vergr.)
61. Verbreitungsbiologie der Blütenpflanzen
Von P. MÜLLER. 3., durchgesehene Auflage. 1983. Fr. 36.-
62. Die Epiphytenvegetation im Raume Zürich als Indikator der Umweltbelastung
Von S. ZÜST. 1977. Fr. 30.-
63. Stickstoff-Düngungseinflüsse von Intensiv-Grünland auf Streu- und Moorbiesen
Von K. Ch. BOLLER-ELMER. 1977. Fr. 30.-
64. Ökologische Zeigerwerte zur Schweizer Flora
Von E. LANDOLT. 1977. Fr. 18.-
65. Comparaison d'écosystèmes de prairies permanentes exploitées de manière conventionnelle et biodynamique
Par M. PORRET. 1978. Fr. 33.-
66. Soziologisch-ökologische Untersuchungen im Grenzbereich Fagus silvatica L./Pinus sylvestris L. in der nördlichen Schweiz
Von Ch. ROTH. 1979. Fr. 27.-
67. Die ökologische und pflanzensoziologische Stellung der Eibe (Taxus baccata) in der Schweiz
Von Ch. LEUTHOLD. 1980. Fr. 42.-
68. Contributions to the knowledge of flora and vegetation in the Carolinas (Beiträge zur Kenntnis der Flora und Vegetation der Carolinas) - Ergebnisse der 16. internationalen pflanzengeographischen Exkursion (IPE) 1978 durch die südöstlichen Vereinigten Staaten. Band 1.
Edited by H. LIETH and E. LANDOLT. 1979. Fr. 44.-
69. Contributions to the knowledge of flora and vegetation in the Carolinas (Beiträge zur Kenntnis der Flora und Vegetation der Carolinas) - Ergebnisse der 16. internationalen pflanzengeographischen Exkursion (IPE) 1978 durch die südöstlichen Vereinigten Staaten. Band 2.
Edited by H. LIETH and E. LANDOLT. 1980. Fr. 36.-
70. Biosystematic investigations in the family of duckweeds (Lemnaceae). (Lemnaceae). Band 1. Key to determination. Cytological variation. Amino acid composition and sugar content. Effects of nitrogen and phosphorus. Bibliography. List of studied material.
Edited by E. LANDOLT. 1980. Fr. 50.-
71. Biosystematic investigations in the family of duckweeds (Lemnaceae). 2. The family of Lemnaceae - A monographic study. Volume 1.
Von E. LANDOLT. 1986. Fr. 69.-
72. Die Vegetation der Nationalstrassenböschungen der Nordschweiz und ihre Eignung für den Naturschutz
Von A. KLEIN. 1980 (vergr.)
73. Keimverhalten und frühe Entwicklungsphasen einiger Alpenpflanzen
Von A. FOSSATI. 1980. Fr. 38.-
74. Zum Wasserhaushalt verschiedener Strassenbaumarten unter dem Einfluss der winterlichen Streusalzanwendung
Von A. SPIRG. 1981. Fr. 25.-
75. Phenological methods in permanent plot research. The indicator value of phenological phenomena - A study in limestone grassland in northern Switzerland
Von B. KRÜSI. 1981. Fr. 24.-
76. Alpine Rasengesellschaften auf Silikat bei Davos (mit farbiger Vegetationskarte 1:2500)
Von L. VETTERLI. 1982. Fr. 38.-
77. Contributions to the knowledge of flora and vegetation in the Carolinas (Beiträge zur Kenntnis der Flora und Vegetation der Carolinas) Ergebnisse der 16. internationalen pflanzengeographischen Exkursion (IPE) 1978 durch die südöstlichen Vereinigten Staaten. Band 3.
Edited by H. LIETH, E. LANDOLT and R.K. PEET. 1981. Fr. 36.-
78. Genetisch-ökologische Untersuchungen an Ranunculus montanus Willd. s.l. aus der alpinen Stufe von Davos (Graubünden)
Von R. DICKENMANN. 1982. Fr. 24.-
79. Einfluss der Feldmaus, Microtus arvalis (Pall.), auf die floristische Zusammensetzung von Wiesenökosystemen
Von A. LEUPERT. 1983. Fr. 33.-
80. Biosystematic investigations in the family of duckweeds (Lemnaceae). Band 3. Das Wachstum von Wasserlinsen (Lemnaceae) in Abhängigkeit des Nährstoffangebots, insbesondere Phosphor und Stickstoff
Von A. LÜND. 1983. Fr. 33.-
81. Pflanzensozioökologische Untersuchungen von gemähten Magerrasen bei Davos.
Von G. ZUMBÜHL. 1983. Fr. 33.-
82. Schaffung artenreicher Magerwiesen auf Strassenböschungen. Eignung von verschiedenem Saatgut für die Neuschaffung Mesobrometum-artiger Bestände. Eine Untersuchung in der Nordschweiz.
Von Th. WEGELIN. 1984. Fr. 33.-
83. Bedeutung des Pflanzenwasserhaushaltes für Koexistenz und Artenreichtum von Trespen-Halbtrockenrasen (Mesobromion).
Von U. KUHN. 1984. Fr. 36.-
84. Ertragskundliche Untersuchungen von gedüngten Mähwiesen der subalpinen Stufe bei Davos.
Von R. PETERER. 1985. Fr. 36.-
85. Verbreitungsbiologie der Blütenpflanzen Graubündens.
Von P. Müller-Schneider. 1986. Fr. 45.-
86. Genetic-ecological investigations in Biscutella levigata L.
Von M. GASSER. 1986. Fr. 36.-
87. Aspekte der Geobotanik. Festschrift Elias Landolt.
Redigiert von K.M. URBANSKA. 1986. Fr. 55.-
- 88a. Vegetationskartierung und Untersuchungen zum landwirtschaftlichen Ertrag im MAB6-Gebiet Davos
b. Kartenband zu Heft 88a.
Redigiert von E. LANDOLT, B.O. KRUESTI, G. ZUMBÜHL. 1986. Fr. 45.-

89. Auswirkungen und Beseitigung von Düngungseinflüssen auf Streuwiesen. Eutrophierungssimulation und Regenerationsexperimente im nördlichen Schweizer Mittelland.
Von Th.B. EGLOFF. 1986. Fr. 42.-
90. Analyse vegetationskundlicher Daten. Theorie und Einsatz statistischer Methoden.
Von O. WILD. 1986. Fr. 36.-
91. Contributions to the knowledge of flora and vegetation of northern Argentina. (Beiträge zur Kenntnis der Flora und Vegetation Nordargentinens). Ergebnisse der 17. Internationalen Pflanzengeographischen Exkursion (IFE) 1984 durch Nordargentinien.
Edited by U. ESKUCHE and E. LANDOLT. 1986. Fr. 51.-
92. Pilzökologische Untersuchungen in Wiesen und Brachland in der Nordschweiz (Schaffhauser Jura).
Von I. BRUNNER. 1987. Fr. 48.-
93. Vegetation of stabilizing and eroding slopes in eastern Nepal.
By Ruth SCHAFFNER. 1987. Fr. 45.-
94. The impact of draining, burning and fertilizer treatments on the nutrient status of *Typha glauca* floating mats in water-level stabilized marshes.
By B.O. KRUESI (in press)
95. Biosystematic investigations in the family of duckweeds (Lemnaceae)
4. The family of Lemnaceae - A monographic study. Volume 2.
By E. LANDOLT and R. KANDELER. 1987. Fr. 64.-

Die Hefte sind zu beziehen beim Geobotanischen Institut ETH, Stiftung Rübli, Zürichbergstrasse 38, CH-8044 Zürich

ca induced by two successive blue light pulses. In contrast to L. trisulca, the O_2 uptake in L. aequinoctialis shows no stimulation in blue light (WITZTUM et al. 1979). Further details on light-dependent chloroplast movements are given in chapter 2.4.1.6.

Far-red light and to a lesser extent blue light elongate fronds and stalk of L. trisulca at low light intensities. This effect may partly be attributed to the phytochrome system. Red, yellow, and green light does not have any other effect than white light. Full darkness resulted in a slight elongation. High intensities of red and far-red but not of blue light also lead to elongation of the frond (ZURZYCKI 1957b). BATA and NESKOVIC (1982b) observed greatly shortened stalks and fronds of L. trisulca at high intensities of blue light compared to the effect of white and red light. This effect was not changed by the addition of kinetin, IAA, or GA.

TSUDZUKI and KONDO (1979) observed, in L. gibba, a potassium uptake rhythm which only showed up at relatively high light intensities (below 700 lux, the rhythm was rapidly damped). The amplitude of the rhythm was independent of the growth rate and of the potassium content of the medium (for details see chapter 2.5.9).

The influence of light on biosynthesis of carotenoids was studied by MONEGER (1968d) in S. polyrrhiza. The production of carotenoids (carotene, lutein, epoxylutein, violaxanthin, neoxanthin) is not fully prevented in darkness. However, light has a very stimulating effect. It seems probable that beside chlorophyll other photoreceptors are involved in the effect (compare chapter 2.5.8.2). The flavonoids vitexin and orientin are formed in S. intermedia in darkness as well as in light. However, cyanidin-3-monoglucoside is produced only in white or blue but not in red light. The vitexin content reaches the same level at light intensities of 4000 and 9000 lux. In contrast, the content of orientin and cyanidin-3-monoglucoside is lower at the lower intensity. Kaempferol behaves similarly as vitexin, quercetin similarly as orientin (McCLURE 1968). Anthocyanin in S. punctata has a high peak of synthesis at 705 nm and at 300 nm. Blue and green light are relatively ineffective. There is no evidence that the formation of anthocyanin is controlled by phytochrome (NG et al. 1964). However, MANCINELLI (1984) and MANCINELLI and RABINO (1984) assume that phytochrome as well as photosynthetic pigments are involved in biosynthesis of anthocyanin in S. polyrrhiza. In S. intermedia, a further photoreceptor (called cryptochrome) seems to be

active in addition. For further details on light-dependent anthocyanin biosynthesis see chapter 2.5.8.5.

The Hill reaction activity of isolated W. arthiza chloroplasts is lower under red illumination compared with white light. Under blue light, it is stimulated at high growth rates and reduced at low growth rates in comparison with white light. The effect of blue light is strongly connected with an increase in the endogenous cytokinin level in fast growing cultures (EICHORN and AUGSTEN 1983a).

The synthesis of the 5×10^{-5} dalton plastid mRNA does not seem to be dependent on phytochrome, but is stimulated by irradiation in the blue and red regions (GRESSEL 1978).

Pigments which absorb blue light and might be considered as active in some of the mentioned processes are flavins and carotenoids (VOSKRESEN-SKAYA 1979).

UV rays stimulate the growth of S. polyrrhiza, L. gibba, and L. minor at a daily exposure of 2.5 to 10 minutes. They promote the activity of L-phenylalanine ammonia lyase and raise the anthocyanin content of S. polyrrhiza (GORDON 1977). LUKINA (1983b) also observed a stimulation of growth rates in S. polyrrhiza, L. gibba, and L. minor by UV irradiation (2-10 minutes with BUV 30 lamps). Different strains responded differently. However, BIGGS (1983) recorded a reduction of growth rate in L. minor (probably L. obscura) and a repression of flowering after UV-B irradiation (280-320 nm). Photosynthetically active radiation is required for photoreversal of these inhibitions. UV irradiation changes the colorless flavanes in S. punctata into reddish-brown (phlobaphene-like) pigments (WITZTUM 1974a). Further effects of UV irradiation in S. punctata are: lower IAA level, and abscission of daughter fronds. Exogenous supply of IAA neutralizes the effect of UV irradiation (WITZTUM 1974b, WITZTUM and KEREN 1978a, WITZTUM et al. 1978). DIJKMAN et al. (1964) observed a 20% life-span extension in L. minor and L. aequinoctialis by short exposure to UV irradiation. A longer exposure reduces the life-span. It is supposed that the UV irradiation may selectively cause the destruction of certain protoplasmic factors that shorten life-span. HICKS (1932a) noted necrotic tissue in different Lemnaceae species after UV irradiation of at least 5 minutes at an intensity of $0.1 \text{ J m}^{-2} \text{ s}^{-1}$. He also reported flower induction with the same treatment in L. minor, L. turionifera (named as L. minor var. purpureus), L. trisulca, L. valdiviana, and L. minuscula, but not in S. polyrrhiza, W. gladiata, and W. bo-

realis. WANGERMANN and LACEY (1952, 1953) noted damaged fronds of L. minor after irradiation with UV rays of 250 nm. The effect is smaller in aerated cultures. The authors as well as HILLMAN (1961a) suppose that the injuries might have been caused not by the UV irradiation, but by the simultaneously developing ozone.

2.3.5.1.5. Heterotrophic growth

GORHAM (1945) was the first to demonstrate that Lemnaceae are able to grow in darkness if sucrose is added to the solution. Further cultures in the dark have been carried out by GORHAM (1950), THIMANN and EDMONDSON (1949), HUTNER (1953), LANDOLT (1957), HILLMAN (1957), and many others.

The fronds of Lemnaceae grown in darkness are yellowish white, smaller than in the light, and have reduced air spaces. Spirodela and Lemna species develop only rudimentary roots (S. punctata, L. gibba, L. minor) or very small roots (S. polyrrhiza, L. aquinoctialis, L. minuscula) (LANDOLT 1957, EFRAT et al. 1975, LEUCHTMANN 1979, LIEBERT 1980a). The protein content in S. punctata is lower in fronds grown in the dark (2.5% of the dry weight) than in those grown in the light (7.8%), the amount of starch greater (50% versus 27%) (DE HEIJ et al. 1984).

The bleaching of the fronds in the dark is reversible if the plants are again transferred to the light. Young chloroplasts of S. punctata develop in the dark to amyloplasts; old chloroplasts remain unchanged for up to three months (PORATH 1979). Chlorophyll-free clones could not be induced experimentally in darkness or in light by treatments with streptomycin or 3-amino-1,2,4-triazole (SCHER and AARONSON 1958). FRICK (1972, unpubl. results) reports that cytidine desoxyriboside (C_{dr}), a natural metabolite, quantitatively induces bleaching of the fronds of photoheterotrophically growing L. minor. The bleaching consists of a specific inhibition of plastid replication in the fronds.

Some Lemnaceae species or clones are not able to grow in the dark with the sole addition of sucrose. It has not been worked out extensively which organic substances are needed besides sucrose by different species. As HILLMAN (1957, 1961a) pointed out, the light intensity needed for sufficient production of certain vitamins or growth factors is very low. Unfortunately, most experiments dealing with growth in the dark of Lemnaceae were not made in complete darkness because the cultures had to

be controlled or the fronds counted. This resulted in short exposition to light. GORHAM (1945) found that green light had a very small effect on production of growth factors.

S. polyrrhiza (GORHAM 1945, LANDOLT 1957, HILLMAN 1961a, EFRAT et al. 1975, LEUCHTMANN 1979) and S. punctata (GORHAM 1950, HUTNER 1953, LANDOLT 1957, SCHER and AARONSON 1958, EFRAT et al. 1975, LEUCHTMANN 1979) are reported to grow with sugar alone in complete darkness.

A special effect in S. punctata was observed by MCCOMBS and RALPH (1972a,b). In culture with sucrose as the only organic source (beside EDTA), growth stopped after 2-3 days following the transfer of the fronds to darkness. The fronds began to grow again 3 to 4 weeks later. Using different special treatments (cold shock, heat shock, addition of distilled water, of chloramphenicol, of actidione, or of kinetin), but also with the addition of yeast extract or with intermittent red light irradiation, the interruption of growth could be overcome. Adding kintin to dormant S. punctata initiated increased synthesis of DNA, RNA, and protein within one hour. The authors suppose that S. punctata normally contains a growth inhibitor which is blocked by cytokinins. Since cytokinins are only formed in the light, the inhibitor still works in the dark until it is used up or inactivated after 3 to 4 weeks. Some amino acids (aspartic acid, asparagine, glutamic acid, glutamine, alanine), α -ketoglutaric acid, and pyruvate also neutralize the inhibitory effect of darkness on the growth of S. punctata (BUCKLE and RALPH 1977). Differently from Spirodela, L. minor was found to grow continuously in the dark only if it is supplied additionally with casamino acids and yeast extract. With sucrose alone, the species was not able to utilize inorganic nitrate (GORHAM 1950). According to the same author, vitamin B_1 (thiamine) turned out to be the active substance in yeast extract. The optimum concentration amounted to 0.005 mM. The thiazole component of vitamin B_1 is active whereas the pyrimidine component is inefficient. BORNKAMM (1970) was able to grow L. minor with the addition of yeast extract only (casamino acids proved not to be necessary). However, the protein content was higher in solutions with supplementary casamino acids, the carbohydrate/protein ratio being 8 instead of 16. Also ROM-BACH (1974a,b, 1976) and LEUCHTMANN (1979) succeeded in growing L. minor in solutions without casamino acids but with yeast extract. In contrast to these results, GORHAM (1950) had to supply casamino acids in addition to yeast extract for growth of his L. minor in darkness. It is not sure

if this difference in need of casamino acids is due to a different clone used (GORHAM possibly had a clone of *L. turionifera* instead of *L. minor*) or 'if other factors, e.g. culture conditions, are responsible (GORHAM used a modified Hoagland solution without ammonium). The addition of casamino acids without vitamin B₁ was able to delay the death of *L. minor* compared with a medium without casamino acids; with casamino acids, the fronds died after 25 times multiplication, without casamino acids, death was observed after 6 times multiplication of the fronds (ROMBACH 1974a). HILLMAN (1957) cultivated *L. minor* successfully in the dark if he added 15 or 45x10⁻⁴ mM kinetin or related compounds to the solution containing sucrose. However, his experiments lasted for only relatively few days. ROMBACH (1974a,b, 1976) showed that the cytokinins BA and kinetin (10⁻³ mM) stimulate the growth of *L. minor* in the dark for the first 20 days, but did not enable long-lasting growth without the addition of vitamin B₁. If vitamin B₁ was present, the stimulating effect of BA lasted for at least 80 days. Up to 0.1 J m⁻² s⁻¹ the stimulation of kinetin is the same, at higher light intensities it decreases (ROMBACH 1974b). Also TRASSERON-DE JONG and VELDSIPA (1971b) observed, in *L. minor*, a stimulation of BA or kinetin only for 7 days without addition of vitamin B₁ under dark conditions. After 3 weeks, the cultures died. Short illumination with red light (10 minutes every 3 to 4 days) made the addition of yeast extract superfluous (GORHAM 1950, HILLMAN 1957). The effect of red light is accomplished by the phytochrome system. The vitamin B₁ synthesis is stimulated by phytochrome (P_{fr}) (ROMBACH 1974a,b, 1976).

According to LEUCHTMANN (1979), *L. aequinoctialis* needs not only vitamin B₁ for continuous growth in the dark, but still another component out of the yeast extract.

LANDOLT (1957 and unpubl. results) cultivated different species under two different nutrient conditions (table 2.22). Since he did not keep absolute darkness, his results are only indicative. He grouped the investigated clones in four categories according to their different behaviour in the three solutions (table 2.23). The negative effect of amino acids and yeast extracts on *S. polyrrhiza* can also be observed under light conditions. Therefore, it must be concluded, that either the amino acids are not in a balanced composition, or some factors of the yeast extract are present in too high concentration for good growth in species of the group A). In group B) the species multiply with the same growth

Table 2.22. Grouping of different species and clones of Lemnaceae according to their behaviour towards yeast extracts and casamino acids in darkness (after LANDOLT 1957 and unpubl. results). Conditions and reactions are presented in table 2.23.

Clones with an asterisk (*) were also investigated by LEUCHTMANN (1979) with essentially the same results

Group	species	clone No.
A)	<i>Spirodela polyrrhiza</i>	6581, 6593, 6613, 6627, 6731, 7003
	<i>Wolffiella neotropica</i>	7010, 7344*
	<i>Wolffiella linguata</i>	7225 6748, 7289
B)	<i>Spirodela punctata</i>	6631, 6725
	<i>Lemna gibba</i>	6566, 6583, 6729, 6745, 6861*, 7007, 7135, 8428*
	<i>Lemna minuscula</i>	6584, 6589, 6597*, 6600*, 6726, 6747
	<i>Wolffia angusta</i>	7476
C)	<i>Lemna disperma</i>	7223, 7782
	<i>Lemna obscura</i>	7856
	<i>Wolffiella gladiata</i>	7000
	<i>Wolffia microscopica</i>	8359
	<i>Wolffia arrhiza</i>	6862, 7014
	<i>Wolffia columbiana</i>	7222
D)	<i>Wolffia globosa</i>	6592
	<i>Lemna minor</i>	6568, 6570, 6578*, 6579, 6580, 6591*, 6625, 6728, 7008, 7011
	<i>Lemna turionifera</i>	6573, 6601, 6619, 6727, 6736, 6853
	<i>Lemna trisulca</i>	6601, 6624, 6722, 7013
	<i>Lemna aequinoctialis</i>	6609, 6612, 6746, 6748, 7001*, 7006, 7806*
	<i>Lemna valdiviana</i>	7002, 7005
	<i>Lemna minuscula</i>	6752
	<i>Wolffia australiana</i>	7211

Table 2.23. Meaning of the categories A), B), C), and D) in table 2.22.

++ good growth, + slow growth, 0 dieing

Substances added to Hutner 1/5	A	B	C	D
1% sucrose	++	++	+	0
1% sucrose + 0.004% yeast extract + 0.08% casamino acids	+	++	++	++

rate in both solutions whereas in group C) casamino acids and yeast extract have a stimulating effect. It is possible that some or all clones of group C) might in effect need a very low amount of a growth factor in absolute darkness. The group D) with obligatory need of yeast extract and/or casamino acids can possibly be divided into two or more groups if vitamin B₁ is not the only factor needed for some clones. LEUCHTMANN (1979) noted a slightly inhibiting effect of casamino acids and yeast extract on his two clones of L. minuscula (6597, 6600) which are here attributed to group B). It is possible that all or some clones of L. minuscula should be changed to group A).

2.3.5.2. Radiations outside the visible and UV range

2.3.5.2.1. X-rays (γ-radiation)

FELDMANN (1968, 1969, 1971, 1975) succeeded in stimulating the growth rate of L. minor and in increasing the size of the frond by treatments with low intensities of X-rays. This was especially the case when light and temperature conditions were kept suboptimal. POSNER and HILLMAN (1960) did not observe any effect on L. aequinoctialis with radiation of 300 r. At higher intensities, the growth rate was inhibited, and at 5000 r no growth occurred any longer. The preference of the side on which the first daughter frond appears was often changed after irradiation. JOHNSON (1941) noted deformed or damaged fronds in L. minor as a result of short-term radiation of 1000 and 2500 r. According to LEINERTE (1969), the accumulation of Sr, Cs, and Ce is lowered with increasing irradiation doses of X-rays.

Table 2.24. Radiosensitivity of different Lemnaceae species

DL 100 = doses needed for 100% lethality

Species	DL 100 in mC kg ⁻¹	Authors
<u>W. arrhiza</u>	800	SAROSIEK and WOZAKOWSKA-NATKANIEC (1984)
<u>L. minor</u>	1806	KASINOV (1966)
<u>S. polyrhiza</u>	2580	WOZAKOWSKA-NATKANIEC (1977a)

WOZAKOWSKA-NATKANIEC (1977a,b) investigated the tolerance of two populations of S. polyrhiza and 14 populations of L. minor to γ-irradiation (with Co). The populations showed different behaviour. S. polyrhiza appeared to be less sensitive than L. minor. The 50% lethal dose for L. minor was achieved between 4000 and 10000 r depending on the population. The corresponding dose for S. polyrhiza was reached between 6000 and 8000 r. Some defects could be detected already at 1000 r. L. gibba shows abnormalities at 800 r (10-60% of the fronds). This proportion was held for a long time under the same conditions. Chromosome changes within the meristem cells are assumed. The sexual descendants proved to be mostly normal (KASINOV and KASINOVA 1971a,b). The different radio sensitivities of 3 Lemnaceae species are summarized in table 2.24).

Mg and especially Ca increase the resistance of W. arrhiza to acute radiation. Therefore, high concentrations of these minerals have a protective effect on Lemnaceae which are exposed to γ-radiation in nature (SAROSIEK and WOZAKOWSKA-NATKANIEC 1984). LYSENKO (1981) observed a similar higher resistance of S. polyrhiza to γ-radiation after pretreatments with alkylating mutagens (N-nitroso-N-methylurea, N-nitroso-N-ethylurea, ethyleneamine, diethyl sulfate). Postradiation treatment with 10 mM caffeine inhibited the protective effect of the mutagens.

2.3.5.2.2. Radionucleides and other radioactive substances

Radionucleides which serve for disinfection of water are taken up by Lemnaceae to 50-60% (GUSKOVA et al. 1970) (see also chapter 3.5.3). The

uptake of radionucleides by L. gibba generally decreases with increase of the pH from 5 to 9 (EL-SHINAWY and ABDEL-MALIK 1980). S. polyrrhiza can be used as an indicator for radioactive contamination of water by measuring the content of radionucleides. However, submerged plants such as Potamogeton are more accumulative (VINTSUKOVICH and TOMILIN 1984). Lemnaceae are able to accumulate more ^{106}Ru than small animals in an aquarium ecosystem (HONDA et al. 1971).

2.3.5.2.3. Laser radiation

COUCH and GANGSTAD (1974) investigated the response of L. minor to $\text{CO}_2\text{-N}_2\text{-He}$ laser radiation ($10.6\ \mu$). Plants irradiated with $100\ \text{J cm}^{-2}$ laser energy died within three weeks. Ten J cm^{-2} stopped the growth rate completely without destroying the fronds. Growth was not affected by $1\ \text{J cm}^{-2}$ treatment.

2.4. DEVELOPMENT

2.4.1. VEGETATIVE DEVELOPMENT

2.4.1.1. Morphogenesis of fronds

Environmental and endogenous factors regulate not only growth but also morphogenesis of duckweed fronds. In L. trisulca relative frond length (length/width ratio) and length of the stipe are increased, when plants were cultivated under low intensity far-red light (far red applied solely to cultures containing sugar or as an extension of the daily high-intensity white light period). Under high-intensity light conditions red as well as red plus far red have an elongating effect on frond and stipe (ZURZYCKI 1957b). Possibly, gibberellins may participate in the mediation of the cited light effects. GUERN (1965) found a stipe lengthening after application of gibberellin A_3 in L. trisulca and S. polyrrhiza. Another possibility to lengthen the stipe of a certain frond in L. trisulca is removal of the two daughter fronds (GUERN 1963a, 1965). In L. gibba a strong enhancing effect on stipe length is achieved, if benzyladenine is added to an ammonium-containing nutrient medium (DEKOCK et al. 1974). Salicylic acid, acetylsalicylic acid and salicylhydroxamic acid, but not benzoic acid and several other related substances have the same effect as benzyladenine (inducing in addition, however, chlorosis and growth inhibition in daughter fronds). Salicylic acid is effective only when iron and/or copper as well as ammonium are present in the nutrient medium. Addition of either iron or copper caused the salicylic acid effect to appear and the effect was more pronounced when iron or copper were increased. In S. polyrrhiza, however, benzyladenine ($2 \cdot 10^{-6}\text{M}$) causes abnormal stipe growth also in the absence of ammonia (LE PABIC 1976a,b). In L. minor, the size of the frond and the length of the stipe are greatly enlarged after supply of 10^{-3}M arginine (FELLER and ERISMANN 1976). The fact that arginine is one of the precursors of polyamines (see fig. 2.50) should be considered in this context.

Papillae, which can be developed on the upper frond surface above the point of root insertion and along the middle nerve in S. punctata and L. aequinoctialis, are very distinctly visible when the ethylene-releasing agent ethephon is added to the culture medium (BUECHELE and KANDELER,

SCHARFETTER and KANDELER, unpublished results). In *S. punctata* the effect of ethylene can be achieved also by supply of ACC (SCHARFETTER et al. 1986), EDDHA or salicylic acid (SCHARFETTER et al. 1978). Naphthaleneacetic acid reverses the effect of EDDHA.

Another characteristic, which is promoted under the influence of ethylene, ethylene, EDDHA, and salicylic acid, is the development of air spaces on the lower side of the frond (CLELAND 1974c, CLELAND et al. 1982, ELZENGA et al. 1980, PIETERSE 1976, PIETERSE et al. 1970b, 1970c). Especially in *L. gibba* this development gives rise to the gibbosity of the fronds. The effect of EDDHA can be reversed by benzyladenine (PIETERSE and MUELLER 1977), decomposition products of gibberellin A_3 , i.e. allogibberic acid, epiallogibberic acid and gibberic acid (PIETERSE 1976), β -naphthol (PIETERSE 1978b), and free (unchelated) copper ions (PIETERSE 1975a), but not by iron deficiency (PIETERSE 1975a). EDDHA changes the uptake of certain heavy metals into the plant. In *L. gibba* G3 iron uptake is increased and copper uptake decreased by EDDHA (PIETERSE 1975a). Presumably, chelation of copper is only one of the effects of EDDHA, which lead to gibbosity, because EDDHA is effective also in a medium without copper (SCHARFETTER and KANDELER, unpublished results). Manganese as an agent, which activates enzymatic ethylene release from ACC, should be investigated in this context. Preliminary investigations of ELZENGA et al. (1980) have shown that EDDHA increases the ethylene evolution in *L. gibba* G3. In *S. punctata* the formation of visible air chambers is promoted by EDDHA and SA (SCHARFETTER et al. 1978), by the anti-auxin PCIB, by kinetin, but not by ACC (SCHARFETTER et al. 1986). The auxin NAA and the blockers of ethylene synthesis AVG and AOA abolish or diminish the effect of EDDHA. Thus, EDDHA may act in at least two-fold way: by a decrease of the endogenous auxin level and by an increase of endogenous ethylene formation. In *L. minor* development of intercellular air spaces needs low-intensity red light (ROMBACH 1976). In fronds grown in darkness (with or without kinetin) intercellular cavities are absent. The effect of red light is strengthened by addition of 3×10^{-6} M kinetin to the nutrient solution (fig. 2.28).

Cell and tissue differentiation in fronds of Lemnaceae has been studied in connection with stomata and the vascular system. FERNANDEZ and MUJICA (1973) determined the stomatal index (stomata-epidermal cell ratio) in *S. intermedia* grown under different regimes of light and temperature. Whereas lowering of light intensity from 7000 to 4000 lux decreases the

stomatal index, lowering of night temperature from 25 to 4°C increases this value.

In the genus *Lemna* stomata are inflexible after differentiation and open all the time (see chapter 2.5.1.1). The width of the stomatal fissure is fixed during development. This process can be modified by ABA supply to the nutrient medium. The mean aperture of stomata in *L. minor* is 73% of control after 8 days growth in the presence of 10^{-6} M ABA, and 44% of control after 14 days (MCLAREN and SMITH 1976).




Fig. 2.28. The effect of red light and kinetin on the development of intercellular air spaces in *Lemna minor* (ROMBACH 1976). Fronds were grown in darkness or low-intensity red light (0.6 or 8.5 μ W cm^{-2}) without (-K) or with kinetin (+K) for 21 days. Diagrams of fronds show the intercellular air spaces in black.




Fig. 2.29. The vascular system in fronds of *Lemna minor* as influenced by four concentrations (0-15 mg/l) of triiodobenzoic acid (SARGENT and WANGERMANN 1959).

The elaboration of the vascular system in fronds of *L. minor* is regulated by auxins and ethylene. Indoleacetic acid (IAA), naphthaleneacetic acid (NAA), as well as ethylene cause the production of 5 veins per frond instead of 3 or 4 in the controls. The application of trilobobenzoic acid (TIBA), an inhibitor of auxin transport, has a by far more dramatic effect (fig. 2.29). 15 mg/l of this substance increases the number of veins per frond to 10-12 (by differentiation of longer or shorter branches from the 3 veins, which arise from the "node", i.e. the insertion point of the root). 10 mg/l TIBA in combination with 0.05 mg/l NAA lead to a further increase of vein elaboration. Several other halogen-substituted benzoic acids have the same, but a less pronounced, effect as TIBA (SARGENT and WANGERMANN 1959, see also FLETCHER 1965).

The branching pattern of *Spirodela* and *Lemna* fronds is homodromous in most cases: One of the two daughter fronds, produced in one of the two pouches of the mother frond, dominates over the other during development and this dominance is preserved on the same side of the frond in all successive, vegetatively produced frond generations. In this way right-handed and left-handed clones occur, which normally are stable over decades. Nevertheless, the extent of dominance can vary and even a change of dominance can appear.

In *L. trisulca* the asymmetry of frond branching is expressed very weakly under high intensity light, but comes out in the following frond generations after lowering of the light intensity (GOEBEL 1921). Such colonies are spiral in shape. Supply of a sugar (sucrose, glucose or fructose), of kinetin, or an increase of medium pH (to 6-8) markedly reduces the dominance phenomenon in *L. trisulca* under low intensity light (BATA 1973, GUERN 1965). Gibberellin A₃ has a similar (GUERN 1965) or no effect (BATA 1973) in comparison to kinetin.

Experiments with excision of frond primordia have shown that there is a correlative inhibition between the different meristematic parts of a frond colony. Removal of the dominant daughter leads to development of the dominated daughter in the opposite pouch (GOEBEL 1921, GUERN 1965). In a similar way removal of grandchildren from the dominant daughter promotes development of grandchildren in the dominated daughter and vice versa. Furthermore, withdrawal of the first (= dominating) and second (= dominated) daughter or removal of grandchildren from the first and second daughter induces outgrowth of the third daughter (= second organ in the "dominating" pouch). The fourth and fifth daughter are

The hypothesis of WITZTUM (1979) that the "axillary frond" induces dominance in the grandchild positioned near by may be questioned in the light of Guern's results. Both developmental events, position of the "axillary frond" and of the dominant grandchild, could be under control of the same - hitherto unknown - hormonal gradients.

Another mode of determination of dominance takes place after sexual propagation. When the embryo grows out from the zygote, the handedness of the first frond (and in consequence all following frond generations) is determined at random. KASINOV (1968c), POSNER and HILLMAN (1960) and KANDELER (unpublished results) have shown that an 1:1 ratio of left-handed and right-handed progeny occur after germination of a lot of seeds.

Concluding this section, two examples of anomalous frond development may be referred to. Bell-shaped fronds, which are curved on the upper side and dip into the water with the border have been observed in *S. punctata* after application of IAA (1 ppm) and/or zeatin (0.025-0.25 ppm) (KERN and NAEF-ROTH 1975). In *L. gibba* some clones derived from X-ray (800 R) treated material show an irreversible tendency to produce monstrous individuals (between 10 and 60%) (KASINOV and KASINOVA 1971a). The following varieties occurred in all such clones: 1. Deformed individuals (proximal part of frond shortened and thickened, inflorescence often suppressed); 2. Funnel-shaped individuals (the two pouches join up to form a single pouch which is open on the upper side); 3. Scoop-shaped individuals (complete lack of stomatal epidermis; the frond consists of a single pouch with daughter fronds and inflorescence); 4. Individuals with gaps (1 or 2 gaps on one or both sides of the frond). This tendency to develop monstrous individuals persists only during vegetative reproduction and is lost after sexual propagation (self-fertilization).

2.4.1.2. Morphogenesis of roots

The number of root primordia developed at the "node" on the lower side of the frond primordium can vary in *S. punctata*. Addition of EDDHA or salicylic acid to the nutrient medium reduces the number of roots per frond from 2-4 to 1 (SCHARFETTER et al. 1978). A partial reversion of the EDDHA effect is achieved by blockers of ethylene synthesis AVG (aminoethoxyvinyl glycine) and AOA (aminooxyacetic acid) as well as by the ethylene-releasing agent ethephon and the ethylene precursor 1-aminocyclopropane.

developed when the first three are removed (GUERN 1965). Correspondingly, kinetin or gibberellin A₃ (GUERN 1965) resp. the combination of sucrose with kinetin or pH increase (BATA 1973) stimulates the simultaneous production of two daughter fronds in one pouch.

Cancelling of meristem competition by optimal growth conditions may lead to a further consequence. WITZTUM (1979) has shown that in *L. aequinoctialis* 6746 8-10 percent of frond colonies are symmetrical under such conditions. The first two daughter fronds are both well developed and bear the dominating grandchild toward the proximal end (stipe end) of the mother frond. In this way on one side of the colony the homodromous branching pattern is interrupted by a heterodromous link and the progeny of this link shows an inversion of handedness. The same mode of branching has been found by BOECKER (1936) in a strain of *L. valdiviana* (collected from aquarium plants). In the southern part of Italy GIUGA (1973, and pers. comm.) collected many specimens of *L. gibba* colonies with symmetrical, proximally oriented grandchildren from the natural environment. In the latter case not only eutrophy of water and high intensity light may be responsible for the altered branching, but also the occurrence of certain herbicides. In addition GIUGA found another type of symmetrical frond colonies, at least in one case together with the described type. The second type has symmetrical grandchildren, which are distally oriented in relation to the mother frond. As WITZTUM (1979) pointed out, the two types of symmetrical frond colonies behave differently. The third and fourth daughter ("axillary fronds"), which are well developed in the first type of branching, are lacking in the second type. In the second type a change of handedness occurs too, but if the branching pattern is perpetuated, a regular heterodromous branching results, i.e. a change of handedness in each generation. An example of partial heterodromous branching in *L. trisulca* (cultured under high intensity green light) can be taken from figure 2 in ZURZYCKI (1957b). The change of handedness, which can be induced by X-rays (KASINOV 1973, POSNER 1961, POSNER and HILLMAN 1960), tritiodobenzoic acid (POSNER 1961, WANGERMANN and LACEY 1953, WITZTUM 1979), 2,4-D (KASINOV 1973, KASINOV and KASINOVA 1974, KASINOV and PAVLOVA 1977) and - in *L. aequinoctialis* - by long-lasting short day (leading to double flowering fronds: DOSS 1978, WITZTUM 1979), seems to be based on a transitory appearance of the second type of symmetry. This has been shown especially in the investigations of KASINOV and co-workers and WITZTUM.

clopropane-1-carboxylic acid (ACC) (SCHARFETTER et al. 1986). Naphthaleneacetic acid reduces the EDDHA effect to a very small extent only (SCHARFETTER et al. 1978).

Root growth seems to depend on the ratio of endogenous sugars to nitrogen compounds (WHIRE 1937b, 1938). High intensity light and nitrate deficiency increase the length of the root in L. minor. Potassium deficiency which inhibits the net assimilation rate and enhances starch accumulation, decreases the length of the root. In accordance with these results SCHUSTER (1968) found a promotion of root growth under long day (in comparison to short day) and an inhibition of root growth by the photosynthesis blocker DCMU in L. aequinoctialis 6746. Exogenous sucrose (1-4%), however, also causes a distinct shortening of roots. Experiments with L. gibba G1 and G3 (BUECHELE and KANDELER, unpublished results) revealed that long day more than doubles the length of the root and that exogenous sucrose greatly diminishes this effect. Also under short day, with end-of-day far red, sucrose has a growth-inhibiting effect on roots of L. gibba. In S. punctata, on the other hand, not addition but removal of sucrose from the medium leads to a marked reduction of the length of the root (EFRAT et al. 1975). A complete lack of root growth occurs in S. punctata and L. gibba, when plants are cultivated heterotrophically, i.e., in darkness with a medium containing sugar (EFRAT et al. 1975, LIEBERT 1980a, 1986a). In S. polyrrhiza root growth is more or less independent from light and sucrose application (EFRAT et al. 1975). The inhibition of root length by adequate nitrogen supply has been observed in S. polyrrhiza only with increasing concentrations of NH_4^+ but not of KNO_3 (MENSCHICK 1970). Later, the different behaviour of certain Lemnaceae against varying NH_4^+ and K_2HPO_4 -concentrations was investigated by LUEBEND (1983). In L. gibba and L. minuscula root growth is optimal, when nitrogen as well as phosphorus are present in very low concentrations. L. minor needs low phosphorus combined with moderate nitrogen and S. polyrrhiza moderate nitrogen and phosphorus concentrations for optimal root lengthening.

The effect of phytochrome on root growth has been investigated in L. gibba (G1 and G3) and L. minor (CHAMURIS and NIELSEN 1983, KANDELER 1963). A short far red irradiation given after the daily light period reduces the growth rate of roots and their attainable length. Red light given after far red reverses the far-red effect.

A treatment of plants with the auxins 2,4-dichlorophenoxyacetic acid

(ONO 1952b) or naphthaleneacetic acid (SCHARFETTER et al. 1978), with gibberellin A_3 (LOOS 1962), higher concentrations of benzyladenine (>0.1 ppm) (LIEBERT 1980a) or with abscisic acid (NEWTON 1972a, 1974a, 1977) reduces the length of the root in some Lemnaceae. In the case of abscisic acid not only root growth but also frond growth and frond multiplication are inhibited. The other hormones may rather work in an indirect way. In the same experiments naphthaleneacetic acid promotes the size of the frond and gibberellin A_3 enhances multiplication of them.

Low concentrations of benzyladenine (0.01 ppm) slightly promote root growth in L. gibba (LIEBERT 1980a). Other root-lengthening agents are EDDHA, salicylic acid, ethephon, ACC, AVG, and AOA (SCHARFETTER et al. 1978, SCHARFETTER et al. 1986). Cobaltous ions inhibit root growth in S. punctata. Some other heavy metals (Cd , Cu , Zn , Mn , Mo) inhibit light-induced root lengthening in heterotrophic cultures of L. gibba (LIEBERT 1986a).

2.4.1.3. Abscission

In a series of papers Witztum and co-workers analysed the separation of daughter fronds from the mother in S. punctata (WITZTUM 1974b, WITZTUM and KEREN 1978a,b, WITZTUM et al. 1978). Two abscission zones are present in fronds of this species, one at the proximal end and one at the distal end of the stipe. In both cases the abscission layer consists of small dividing cells. Subsequent cells of mother and daughter frond, respectively, contain red-brown phlobaphene-like compounds after dying and seem to form a protective layer (for two abscission zones in S. polyrrhiza see NEWTON et al. 1978).

UV irradiation is one of the agents which activates the processes leading to abscission (preferably in the proximal abscission zone). UV lowers the IAA level drastically and in this way seems to remove the block against ethylene induction of abscission. Exogenous supply of IAA abolishes the effect of UV irradiation. The addition of 0.015-0.1 M sucrose (ca. 0.5-3.5%) to the medium has a similar effect as exogenous IAA. In this case the effect of UV on abscission is outplayed by a serious enhancement of endogenous IAA production.

A high concentration of sucrose (0.4 M, ca. 13.5%), which causes an osmotic stress situation, induces abscission in normal frond colonies and

increases the effectivity of UV. The effect of 0.4 M sucrose may come about by a change in the levels of ABA, ethylene, and IAA. The ABA content of *S. punctata* rises six-fold within 24 h, when plants are floated on 0.4 M sucrose (in comparison to distilled water). Application of 10 ppm ABA, on the other hand, induces abscission to a great extent.

Ethylene release is greatly promoted during the first 24 h of 0.4 M sucrose treatment, but falls off to a value below the control during the second day. The ethylene outburst may be effective in frond separation, if one assumes that the osmotic stress simultaneously reduces the IAA content (compare the results of HARTUNG and WITT 1968, for *Anastatica*).

The results of Witztum and co-workers were confirmed and extended by other authors. In *S. punctata* naphthaleneacetic acid, EDDHA, and salicylic acid increase the size of the colonies of fronds, i.e., inhibit frond separation (SCHARFETTER et al. 1978). In *L. minor*, which develops only one abscission zone, exogenous ABA accelerates abscission even at 0.2 ppm. Cellulase but not pectinase activity is increased by the ABA treatment. Benzyladenine inhibits abscission in this way enhancing the size of the colonies (OSTROW-SCHWEBEL 1973, 1979). In *S. polyrrhiza* separation of sub-colonies is facilitated by 1 ppm ethylene. Under 10 ppm ethylene, which simultaneously inhibits growth, only single fronds and groups with two fronds occur (control colonies consist of 2-4 fronds) (NEGBI et al. 1972). WCISLO (1963) mentioned that bacterial contamination accelerates frond separation in *L. trisulca*. Possibly such infections give rise to an enhanced ethylene production. LOOS (1962) stated that gibberellin A_3 promotes frond separation more than frond multiplication in *L. minor*. Most plants consist of a single frond after cultivation with 50 mg/l gibberellin A_3 .

2.4.1.4. Senescence and rejuvenation

Senescence is a complex of physiological processes which limits morphogenetic potentialities and life-span of a certain organ. During senescence protein degradation overcomes protein synthesis and - in interrelation with this - the hormonal state of the organ changes. The extent of senescence depends on environmental stress factors as well as on the presence of other organs which compete for nutritional and hormonal factors. On the other hand, the senescing organ may influence other parts of the organism. Especially the morphogenetic potentialities of meri-




Fig. 2.30. The length of life of *Lemna minor* fronds (see +), the number of daughter fronds produced during the life-span of one mother frond (see the number of bars), and the length of daughter fronds (•) as influenced by the N- and P-content of nutrient medium (BOESZÖRMENYI and BOESZÖRMENYI 1957).

stems can be changed (vegetative vs. generative development) or restricted (number and size of organs produced). In principle, organ senescence itself as well as the senescence-induced meristem state are reversible, thus, a true rejuvenation occurs.

In *Lemna*, the life-span of fronds is reduced by nitrogen- or phosphorus-deficiency (fig. 2.30 BOESZORMENYI and BOESZORMENYI 1957), high temperature (30°C) (WANGERMANN and ASHBY 1951), UV-irradiation (2 min or more) (DIJKMAN et al. 1964), or ABA (KANG and CLELAND 1985). In the cited cases production of daughter fronds is diminished simultaneously or - at least - not enhanced. Therefore the life-shortening agents may be effective directly as stress factors. If, on the other hand, nitrogen deficiency prolongs the life-span (WANGERMANN and LACEY 1955), an indirect effect mediated by an inhibition of the development of daughter fronds may be responsible. The same can be said of short UV-irradiation (60-90 sec) which prolongs the length of life of the fronds, but diminishes their multiplication' (DIJKMAN et al. 1964). Long day is another factor, which can increase or reduce life-span depending on the species or on the culture conditions (OSTROW and DIJKMAN 1969, WANGERMANN 1952). The life-shortening effect of short day in *L. aequinoctialis* (SCHUSTER 1968) is caused without doubt by the fact that flowers are produced under these conditions. If flowering is prevented by a short night break, life-span exceeds that under long day. The connection between flowering and frond senescence can be inferred also from results with *L. perpusilla* p146 (HUEGEL et al. 1979). Treatment of short-day cultivated plants with end-of-day far red inhibits flowering and frond senescence (see number of sister fronds produced by one mother) only to a certain degree. A night break with red light, on the other hand, inhibits flowering completely and retards frond senescence more distinctly.

The effect of stress factors on protein turnover rates was investigated in *L. minor* by TREWAVAS (1972b). If the plants are placed on pure water, there is a reduction in the rate of synthesis of protein and an increase (3- to 6-fold) in the rate of degradation. Similar results were obtained when only one of the following minerals were omitted from the medium: nitrate, phosphate, sulfate, magnesium, or calcium. The cited stress effects could be mediated by an increase of the ABA and decrease of cytokinin content within the plant. Supply of 5 µM ABA to plants grown in sucrose-mineral medium has the same effects on protein turnover as the

stress factors themselves. Benzyladenine has antagonistic effects to ABA, but - surprisingly - alters in plants grown on sucrose-mineral salts only the rate of synthesis and in plants grown in pure water only the rate of degradation. An increase of endogenous ABA level and a decrease of endogenous cytokinin level 24-48 hrs after withdrawal of nitrogen from the medium has been found in *L. aequinoctialis* (MERTEN, HARTUNG, GRUNTZEL, LOEPFERT and KANDELER, unpublished results).

Turnover of ribosomal RNA behaves like that of proteins after transfer of plants to water or to a medium without nitrate, phosphate, calcium, or magnesium. The rate of rRNA degradation is increased and the rate of rRNA synthesis is decreased (TREWAVAS 1970). ABA (5x10⁻⁶M) markedly reduces the rate of rRNA synthesis but leaves the degradative rate unaltered. In plants grown on water, benzyladenine increases both the synthetic and degradative rates of nucleic acid metabolism (TREWAVAS 1970). The inhibiting effect of ABA on RNA synthesis has been found also in roots of *L. minor* (NEWTON 1974a).

The investigations of Trewavas were continued by Davies and co-workers in a series of papers. In spite of the increased protein degradation under nitrogen starvation the concentration of free amino acids decreases in *L. minor* (HUMPHREY et al. 1977). Whereas protein half life is reduced from 143 to 73 hrs under these stress conditions, the total amount of amino acids recycling seems to be increased (DAVIES and HUMPHREY 1978). Four days after set-up of nitrogen deficiency the lowering of soluble protein content of fronds ceases (COOKE et al. 1979b). The same holds true for mineral deficiency (water culture) or osmotic stress (0.5M mannitol). The rates for protein synthesis and degradation tend to return to values comparable with unstressed plants. Obviously, an adaptation to the special stress condition takes place, which includes a replacement of several proteins by others. During nitrogen starvation activities of glutamate dehydrogenase and malate dehydrogenase sustain a reduction which is stronger than under osmotic stress (mannitol). Activities of glucose-6-phosphate dehydrogenase, 6-phosphogluconate dehydrogenase, and PEP carboxylase, however, are more reduced during mannitol treatment than during nitrogen deficiency (COOKE et al. 1979b). In the case of aldolase the decrease in activity after withdrawal of nitrogen is shown to be due, at least partly, to formation of a specific inhibitor, which appears to be a protein with a molecular weight of ≤ 15.000 (DAVIES 1978).

Another factor which alters protein turnover is deuterium oxide ($^2\text{H}_2\text{O}$). When plants are transferred to medium containing 50% $^2\text{H}_2\text{O}$, protein synthesis is inhibited for 4 h and then proceeds at a slower rate than in non-deuterated medium. Protein degradation is increased 9-10-fold initially and after 6 days in $^2\text{H}_2\text{O}$ is still 2-3 times higher than normal. All tested enzyme activities are reduced to a similar extent by the $^2\text{H}_2\text{O}$ treatment (COOKE et al. 1979a).

The effects of $^2\text{H}_2\text{O}$, nitrogen starvation, and 0.5M mannitol do not come about by an activity increase of proteinases but rather by an alteration of tonoplast permeability (COOKE et al. 1980a, b). Utilizing the different $^3\text{H}/\text{H}$ exchange on amino acids in the cytosol and vacuole (s. transaminases in the cytosol only) the authors pointed out that the stress conditions mentioned above increase the permeability of the tonoplast for amino acids. ABA has the same effect on amino acid "leakage" from vacuole to cytosol as all the stress factors. Benzyladenine, on the other hand, reduces the effects of nitrogen deficiency and mannitol on amino acid permeation through the tonoplast. The results with amino acids have led to the hypothesis that primarily the tonoplast passage of certain substances - protons for an activation of proteinases in the cytosol or proteinases themselves - is increased, when stress factors - possibly mediated by hormonal changes - alter protein metabolism.

The effect of ABA on plasma membrane permeability seems not to be restricted to the tonoplast. ANDRES and SMITH (1976) ascertained a substantial increase of amino nitrogen in the medium within 3 h of transfer of *S. polyrrhiza* to 10^{-6} M ABA. In *L. aequinoctialis* three days after ABA supply, the amino nitrogen in the medium amounts to 61% of the total in comparison to 21.9% in the control. In *L. minor* the values are 36.2 and 8.5%, respectively.

The refined kinetics for cytokinin action on nucleic acid and protein metabolism have been investigated on *L. minor*. The RNA content shows a transitory increase with a peak after 30 minutes, when plants are treated with 10^{-5} M kinetin (ERISMANN and FANKHAUSER 1967, FANKHAUSER and ERISMANN 1969b). Within the same time the protein content increases too, and remains at the higher level for a long time (3 days investigated). Similar results have been found with darkened non-growing plants of *S. punctata* (MCCOMBS and RALPH 1972a). Within 1 hour DNA-, RNA-, and protein synthesis (measured by ^3H thymidine-, ^3H uridine-, and ^{35}S incorporation, respectively) are enhanced by about 10^{-6} M kinetin in this

plant material. The rate constant for RNA synthesis is doubled by kinetin and RNA degradation is diminished by 60%. Uptake of ^{35}S , uridine, and ^{14}C glucose, however, are not affected by kinetin in darkened *S. punctata*.

Not only cytokinins but also polyamines could be agents for inhibition of senescence and promotion of rejuvenation in *Lemma*. FLORES and GALSTON (1982) have shown great amounts of agmatine and spermidine to be present in vegetative and flowering *L. gibba* G3. GIOVANELLI et al. (1981) found spermidine in *L. aequinoctialis* as one of the main by-products of the methionine - 5'-methylthioadenosine cycle. Ethylene, another possible by-product of this cycle, could not be detected in these experiments. The effect of sugars on senescence may be rather complex. *Lemma minor* cultivated without sucrose for 4 weeks shows a reduced rate of rRNA synthesis and an increased rate of rRNA degradation in comparison to plants grown on sucrose-containing medium (TREWAVAS 1970). This result may be considered in connection with the fact that sucrose application causes a drastic increase of the endogenous IAA level in *S. punctata* (WITZTRUM et al. 1978). But the growth promotion rendered possible by supply of sugars in consequence can lead to an exhaustion of endogenous and exogenous nitrogen reserves and in this way to an acceleration of senescence. Yellowing - as an indication of senescence - in fronds of *S. punctata* is retarded by 1% glucose only when the nutrient medium is renewed frequently. When, however, the glucose containing medium is not changed, a great acceleration of thylakoid and chlorophyll degradation proceeds (GROB and EICHENBERGER 1973). Transfer of yellowing fronds to fresh (glucose-free or glucose-containing) medium induces a regeneration of thylakoids and regreening.

Sugars which enhance senescence directly are galactose (DEKOCK et al. 1979) and sorbose (STROTHER 1981). Mature and young *Lemma* fronds become progressively chlorotic and growth is arrested in the presence of one of these sugars. In the case of galactose the changes in mineral content of fronds has been followed up. The potassium content is lowered and levels of sodium, calcium, magnesium, and iron are raised. Largely the same changes in mineral composition of *L. gibba* have been found after application of 1 ppm ABA (DEKOCK et al. 1978). Benzyladenine (1 ppm) can partially counteract the ABA effect. LIEBERT (1977), investigating the ABA dependence of mineral content in the same species (*L. gibba*) but using another nutrient medium, has stated a drastic decrease of K^+ level and

only a slight decrease of Ca^{++} level after 13 days of plant cultivation with 1 ppm ABA. A decrease of K^+ and increase of Ca^{++} content has been ascertained in plants grown for 26 days in an unchanged, pure inorganic medium (LIEBERT 1980b).

Stress factors such as (high) temperature, intense photosynthetically active radiation, UV-B radiation, and high osmoticum, induce the production of several "stress proteins" in *L. minor* (BIGGS and KOSSUTH 1985). The cell wall protein extensin is one of the stress-induced proteins. High concentrations of CuSO_4 (1 mM) enhance breakdown, among others, of the large and small subunits of ribulose diphosphate carboxylase, 26 kDa-light harvesting chlorophyll a/b apoprotein, and the O_2 thylakoid protein (MATTOO and CONLON 1985). These effects appear to be independent of ethylene production that is markedly stimulated in the presence of 1 mM CuSO_4 (see chapter 2.5.7.2).

The effect of frond senescence on daughter frond development was first investigated by ASHBY and WANGERMANN (1949). Daughter fronds are produced within the whole life-span of a given mother, but fronds of such a series successively produced from one mother frond behave differently. The later during the lifetime of the mother frond the daughter frond is developed, the smaller the adult frond remains (compare fig. 2.30) and the shorter is its own life span. This gradual reduction in the size of successively produced daughter fronds holds true in *L. minor* (BOESZÖERMENYI and BOESZÖERMENYI 1957, BOSS et al. 1963a, WANGERMANN 1952, WANGERMANN and ASHBY 1951, WANGERMANN and LACEY 1953), in long-day cultivated *L. aequinoctialis* 6746 (SCHUSTER 1968), and in *L. aequinoctialis* 6746, held at 30°C and 12 h light per day (CLAUS 1972). Another mode of daughter frond development is seen in *L. aequinoctialis* 6746, when grown under short day, or short day with night break (SCHUSTER 1968), and in *L. aequinoctialis* "clone 1", held at 25°C and 12 h light per day (BOSS et al. 1963a). In this case, some or several daughters - first produced by the mother - reach about the same size of frond. After that, further daughters with smaller fronds are produced by the mother. This "one-step reduction" of daughter frond development may be an interesting hint to a special type of frond senescence (fig. 2.31).

The restriction of morphogenetic potential induced by the senescent frond in its daughter fronds is reversed during the development of granddaughter and great-granddaughter fronds. A rejuvenation takes place especially in the firstborn fronds. These fronds become larger than

those of their mother. In this way small lateborn fronds produce a progeny of maximum size within a line of about four first-daughter generations (WANGERMANN and ASHBY 1951, compare vol. 1, fig. 2.11).

A special case of rejuvenation combined with a change in the morphogenetic program is the proliferation of disordered cell material. When intact mature *Lemna* fronds are grown on agar medium with a certain combination of growth substances, outgrowth of calli takes place. Whereas in *L. gibba* G3 10 mg/l 2,4-D combined with 1 mg/l N^6 -s(Δ^2 -isopentenyl)-adenine (2iP) induces callus formation (CHANG and CHIU 1976, 1978), only 0.1 mg/l 2,4 D together with 10 mg/l 2iP is effective in *L. aequinoctialis* (CHANG and HSING 1978). In *L. gibba* large friable undifferentiated




Fig. 2.31. Senescence and frond development in *Lemna aequinoctialis* 6746 (after SCHUSTER 1968). Frond lengths of a mother frond (first bar) and its daughter fronds (following bars) are plotted at the time of frond appearance. The cultures of single mother fronds were held under long day (LD), short day with night break (SD+NB), or short day (SD). The end of the life-span of the mother frond is marked by +.

calli were obtained by the end of 8 weeks. The callus could be subcultured and proliferated in fresh medium under illumination or in the dark. In *L. aequinoctialis* 3 months were needed for callus induction. Up to now a complete regeneration of fronds from callus tissue has been achieved only in *L. gibba*. A basal medium supplemented with indoleacetic acid (4 mg/l) and kinetin (1 mg/l) was able to induce frond regeneration. A regenerated strain, named *L. gibba* strain Sinica No. 1, exhibited a slower growth rate but no distinct morphological features.

2.4.1.5. Chloroplast development

Ultrastructure of chloroplasts at various developmental stages has been investigated using electron microscope techniques. Meristematic cells of *S. polyrrhiza* and *L. minor* contain proplastids with only a few thylakoids (RAO 1968, FRICK and JONES 1976, LE PABIC 1972, WROBLEWSKI 1973). By contrast, the primordial cells of *W. arrhiza* show no proplastids, but well-developed chloroplasts containing grana (ANDERSON et al. 1973). In mature frond tissues of *S. polyrrhiza* grana stacking is more marked in the palisade parenchyma than in the spongy parenchyma (LE PABIC 1972). Chloroplasts of root cap, epidermis, cortex, and endoderms in *L. minor* exhibit only few grana thylakoids (WROBLEWSKI 1973). When frond colonies of *S. punctata* are darkened for a longer time, the chloroplast ultrastructure in mature fronds remains unchanged, but in young green fronds a retrogressive development occurs. Chloroplasts are transformed to etioplasts with a prolamellar body. Both crystalline and vesicular forms of prolamellar body can be observed (PORATH 1979). The reverse process has been investigated in *L. minor*. As in other plants, a radiation of thylakoids from the prolamellar body can be seen at an early stage of de-etiolation. Plastid division occurs only in primordial cells of green and etiolated plants (FRICK and JONES 1976).

In the mutant 1073 of *L. aequinoctialis* no. 6746, which is incapable of photosynthesis and ammonium ion uptake, the grana of chloroplasts contain many more thylakoid membranes than the wild-type chloroplasts. In addition, tubular clusters are present in mutant chloroplasts, which differ from prolamellar bodies found in etioplasts (MONSELISE et al. 1984). Supply of L-cystine normalizes the ultrastructural features of the mutant, but not its physiological behaviour (MONSELISE et al. 1986). Greening of etiolated plants under continuous white or red light pro-

ceeds after a lag phase of 20-24 h (*S. polyrrhiza* and *S. punctata*) (PORATH and BEN-SHAUL 1973) or 10-12 h (*L. minor*) (FRICK 1975). In all cases the lag phase can be eliminated by a pre-illumination with a short red light given about the length of lag period before the onset of continuous light. Phytochrome seems to mediate the effect of short red light, because far red given simultaneously with or after red reduces the red light effect. When *L. minor* is grown under light of different spectral qualities, grana stacking is highest under blue light. The appearance of appressed thylakoid membranes also increases after blue irradiation (MAENPAE and ARO 1986).

Mature Lemna chloroplasts (as well as whole plants) possess an outstanding resistance against extremely high centrifugal forces (350.000 x g for 15-60 min). Thylakoids are not so greatly stretched as in spinach leaf chloroplasts. This may be due to the presence of a denser stroma in the duckweed chloroplast (BEAMS et al. 1979).

Several changes of chloroplast ultrastructure take place in *L. minor* when plants are cultivated in a medium containing atrazine (0.25 ppm), a specific inhibitor of Hill reaction (BEAUMONT et al. 1980). The number of grana per chloroplast and the length of grana lamellae increases, whereas the number of lamellae per granum decreases. Intralamellar vesicles disappear and outgrowths are formed at the ends of the chloroplast. The stroma is reduced and accumulates osmophilic globules.

Many agents - metabolic and growth inhibitors, metabolic intermediates, as well as high temperature - can cause a severe inhibition of chloroplast development, visible as production of bleached daughter fronds. A first group of bleaching substances are inhibitors of protein synthesis (chloramphenicol, cycloheximide, tetracycline, streptomycin, and kanamycin) and inhibitors of nucleic acid synthesis (fluorodeoxyuridine, nalidixic acid, and aminotriazole). Azaguanine and azauracil are effective only when given together with certain ribonucleosides. The effects of most of the cited substances are reversible after removal. This means that mature cells of bleached fronds regreen in the absence of the inhibitor. Only effects of fluorodeoxyuridine and aminotriazole are irreversible. Bleaching of growing fronds by aminotriazole (59 µM) and kanamycin (5 µM) is specific in a sense that frond growth is not inhibited by these agents (FRICK 1972, FRICK and JONES 1975).

Some intermediates of nucleic acid metabolism are another group of bleaching substances. Cytidine deoxyriboside (Cdr), uridine deoxyribo-

promoting agents (see chapters 2.4.1.4, 2.5.7.3 and 2.5.8.3). The degradation of chloroplast ultrastructure under the influence of nitrogen or mineral deficiency was demonstrated by GROB and RUFENER (1969) and HUBALD and AUGSTEN (1977b). Destruction of the chloroplast becomes irreversible latest when the envelope bursts on account of excessive starch deposition. Such a process has been found in *L. minor* after applying 10^{-4} M kinetin (ERISMANN and WEGNER 1967) and in *L. gibba* senesced artificially by 0.1% galactose (DEKOCK et al. 1979). Another monosaccharide which causes bleaching in mature and young fronds of *L. minor* is sorbose (STROTHER 1981). Toxicity of galactose is counteracted by glucose added simultaneously and the effect of sorbose can be diminished by fructose.

2.4.1.6. Movement of chloroplasts

After reviewing the developmental processes of chloroplasts the movements of these organelles within the cell may also be reported. Under dark conditions, chloroplasts in the mesophyll layer of *L. trisulca* are distributed more or less at random (apostrophe position). In unilateral, low-intensity light, the chloroplasts migrate to the periclinal cell walls lying approximately perpendicular to the direction of illumination (epistrophe position). Under high-intensity light, the chloroplasts move to the anticlinal cell walls (parastrophe position). Other environmental factors as well as endogenous rhythms have no definite influence on the movements of chloroplasts in *L. trisulca* (ZURZYCKA and ZURZYCKI 1953). In experiments with white light the transition to epistrophe was accomplished with a light intensity as low as 9 lux. In the range from about 800 to 12000 lux epistrophe is replaced gradually by parastrophe. The velocity of movement increases with the "jump" of light intensity, i.e. the difference between the initial and the final light intensity. The migration from parastrophe (at 13300 lux) to epistrophe (at 40 lux), for example, is completed within about 15 min. The movement from parastrophe (5500-13300 lux) to apostrophe proceeds through an epistrophe transition stage occurring 30 min after the beginning of darkness. Apostrophe then is reached after a further 30-60 min (ZURZYCKA and ZURZYCKI 1953). The path of single chloroplasts from epistrophe to parastrophe has been followed by a cinematographic method. The path is very complicated; direction as well as rapidity of the movement change constantly (ZURZYCKI and ZURZYCKA 1953). During the parastrophe - epistrophe reaction, on the

side, and guanosine deoxyriboside, all induce quantitative bleaching in *L. minor*, when grown under continuous light in a sucrose-containing medium (FRICK 1978). Cdr-induced bleaching is not accompanied by a reduction in frond multiplication rate and is fully prevented by an additional supply of thymidine, cytidine, or uridine. These facts are interpretable by the assumption that Cdr interferes with biosynthesis of pyrimidine nucleotides and acts in a compartment which provides DNA precursors for plastids only. Further analysis of Cdr effect has shown that plastid replication is affected in the first place (PIZZOLATO and FRICK 1979). The number of plastids in cells of comparable size is reduced during bleaching by more than 50%, their volume by 10-30%.

A third group of substances capable of inhibiting chloroplast development are certain nitrophenols (PRICE and WAIN 1976). White fronds are produced by *L. minor*, when 4-hydroxy-3-nitrobenzoic acid, 4-hydroxy-3-nitrobenzaldehyde, or 4-hydroxy-3-nitrobenzyl alcohol is added to the medium. 10^{-5} M of these substances are effective to 69, 72, and 66%, respectively.

Growth-inhibiting concentrations of benzyladenine (10^{-6} - 10^{-4} M) are a further possibility to inhibit elaboration of chloroplast ultrastructure primarily in young frond primordia. Bleached fronds of *S. polyrrhiza* grown in the presence of 10^{-6} M benzyladenine contain plastids with only a few thylakoids (LE PABIC 1976b). Mature fronds, however, can also show a partial loss of pigments and thylakoids after treatment with high benzyladenine concentrations. Accumulation of starch is greatly enhanced under these conditions. The content of phospholipids increases and the surfaces of the thylakoid membranes are more appressed (LE PABIC 1976b, 1980).

S. punctata, grown in light at 37°C for a period of 3-6 days, loses the ability to synthesize chlorophyll, to propagate, and to evolve O_2 . Grana thylakoids are destroyed and the number of droplets in the chloroplast increased. This "heat bleaching" appears first in younger fronds. Plants, kept in the dark at 37°C for at least 5 days, are not affected by the thermal treatment (PORATH and BEN-SHAUL 1971). The bleaching by exposure to raised temperatures is reversible in *Spirodela* (SCHER and AARONSON 1958).

Not only proplastids and developing chloroplasts are sensitive to inhibiting factors. Ultrastructure, protein, and pigment content of mature chloroplasts are the primary sites, which are attacked by senescence-

other hand, the paths are rather straight, and the velocity of movement is fairly constant, ranging from 0.6 to 1.0 $\mu\text{m}/\text{min}$ (ZURZYCKI and ZURZYCKI 1957). Under dark conditions, transient rearrangements of chloroplasts can be induced by short light pulses. An 0.8 s pulse of strong blue light (30 W m^{-2}) causes a transitory migration to the weak-light position (epistrophe). With longer pulse duration (8 s), a biphasic response takes place: The initial movement to a partial strong-light position is followed by a wave of translocation to a weak-light arrangement (GABRYŚ et al. 1981).

The blue component of light is responsible for induction of the oriented chloroplast movements in *L. trisulca*. The action spectra for weak-light and strong-light responses are identical and have a major peak at 450 nm and two lower ones at 485 and 380 nm (ZURZYCKI 1962). Also in *S. polyrrhiza* and many other plants only blue light is effective (INOUE and SHIBATA 1973). Consequently, the photosynthetic mutant, strain 1073, of *L. aquinoctialis* which has a block in the electron transport chain between plastoquinone and cytochrome *f* is capable of light-induced chloroplast movements (WITZTUM et al. 1979). A flavin has been proposed to be the main photoreceptor for chloroplast orientation (HAUPT 1982). When the fluence rate dependence of chloroplast translocations is tested with blue light, some rearrangement occurs even under an intensity near 0.15 mW m^{-2} . At temperatures of 20 to 25°C maximum weak-light arrangement is achieved approximately 150 mW m^{-2} and complete strong-light position is obtained in the range $3\text{--}10 \text{ W m}^{-2}$ (ZURZYCKI et al. 1983). At a temperature of 5°C , however, the optimum intensities of blue light for the weak-light and strong-light positions are shifted to lower values (HAUPT and WEISENSEE 1966, WEISENSEE 1968a).

In healthy, growing material of *L. trisulca* the light-controlled displacement of chloroplasts depends not only on the intensity and wavelength of radiation but also on the direction of polarization. Under high-intensity light, chloroplasts avoid the side walls situated parallel or nearly parallel to the E-vector of light (fig. 2.32). From this reaction pattern it can be concluded that the responsible photoreceptor molecules are oriented, their vector of absorption being parallel to the cell surface (ZURZYCKI 1969, ZURZYCKI and LELATKO 1969). In some *L. trisulca* material (especially from old cultures in the autumn) the action dichroism cannot always be observed. The appearance of undulations in the cytoplasmic membrane has been considered to be the cause for this

behaviour. EDTA, ATP, and phosphate buffer of pH 4.7 are agents which can restore the action dichroism in otherwise not reacting material (HAUPT and WEISENSEE 1967, ZURZYCKI 1969).

The speed of chloroplast displacements can be influenced by strong red light. The action spectrum of this photokinetic activity reveals a peak at 680 nm. Thus, photosynthetic pigments seem to mediate this light effect (ZURZYCKI 1962, 1964). The source of energy for all types of chloroplast rearrangements is ATP as revealed by studies with metabolic inhibitors and by ATP-feeding experiments (ZURZYCKI 1965, ZURZYCKI and ZURZYCKA 1955). Respiration as well as photosynthesis can contribute to the ATP delivery. Thus, the effect of strong red light on the photokinetic activity of chloroplasts obviously comes about by photosynthetic ATP formation. The effects of temperature on the velocity of chloroplast movements (WEISENSEE 1968b, ZURZYCKA and ZURZYCKI 1950) may also be explained by the dependence on temperature of energy generation.




Fig. 2.32. Arrangement of chloroplasts in the same cells induced by polarized light. Arrows indicate the plane of the E-vector of light. Cells with chloroplasts in low light intensity - position were irradiated 1 hour with an intensity of 38 W m^{-2} polarized blue light (ZURZYCKI 1969).

The principal mechanism of chloroplast displacement in Lemna seems to be the passive transport with the creeping or streaming cytoplasm (ZURZYCKI and ZURZYCKI 1957, ZURZYCKI and ZURZYCKA 1953). The light regime decides, in which particular regions of the ectoplasm chloroplasts are anchored. An indicator for the ability of ectoplasm to attach chloroplasts should be the low transferability of chloroplasts by centrifugation. In fact, weak light and to some degree also strong light diminish the passive displacement of chloroplasts by centrifugation (ZURZYCKI 1960, 1962). On the other hand, the transition from epistrophe to parastrophe or apostrophe needs a local weakening of cytoplasm stickiness. Ions which diminish the viscosity of cytoplasm, K^+ and Li^+ , accelerate especially the transitions from epistrophe to parastrophe and apostrophe, whereas Mg^{2+} and Ca^{2+} retard the same reactions (ZURZYCKA and ZURZYCKI 1951). As an early link in the transduction chain from light signal to response ZURZYCKI (1972) envisaged glyoxylate to be an "attractant". This compound is assumed to be formed from glycine by the membrane-bound deaminating enzyme glycine oxidase, which is activated by blue light via FMN. In correspondence with the hypothesis, an enhancement of O_2 uptake by very low blue light intensities has been demonstrated in DCMU-treated plants (ZURZYCKI 1970, 1972; see chapter 2.5.1.2). Chloroplasts should accumulate at those regions in the cytoplasm which are enriched in glyoxylate. In Funaria leaves, a glyoxylate solution applied unilaterally causes a preference of chloroplasts to the cell wall facing the test solution (GODZIEMBA-CZYŻ 1973).

The influence of chloroplast displacements on the optical properties of leaves has been elaborated in detail (ZURZYCKI 1961). The work of ZURZYCKI (1953, 1955a, 1955c) with regard to the influence of chloroplast rearrangements on photosynthesis has been reported in chapter 2.5.1.1.1.

2.4.2. Dormancy

True dormancy can be characterized as an developmental intermission, in which growth and other developmental processes are arrested by certain internal conditions. In Lemnaceae dormancy lasts for some time in turions, turion-like fronds, and seeds. Turions are small rootless fronds, which sink to the bottom of the water because of reduced or absent air spaces and a high starch content. They have been found in S. polyrrhiza (HEGELMAIER 1868, VAN HOREN 1869), L. turionifera (LANDOLT 1957, 1975), N-2 type strains of L. aequinoctialis (BEPPU and TAKIMOTO 1981c), W. brasiliensis (LANDOLT 1957), W. arhiza (HEGELMAIER 1868), and W. columnata (LANDOLT 1957). Also W. borealis, W. australiana, W. angusta, and W. globosa are able to form turions (LANDOLT 1986). Small, thick, fleshy, non-growing fronds, which contain much starch but float, are functionally comparable with turions and therefore can be called turion-like or resting fronds. They occur in S. punctata (HARRISON 1964, SCHARFETTER and KANDELER, unpublished results), L. perpusilla (KANDELER and HUGEL 1974a), L. gibba (VAN HOREN 1869), L. minor (SCHULZ 1962), and W. gladiata (PIETERSE et al. 1970d).

Recently, SMART and TREMAVAS (1983b) compared ultrastructural features of turions and vegetative fronds in S. polyrrhiza. Turions show smaller air spaces, smaller cell vacuoles, thicker cell walls, and accumulation of numerous starch grains. No distinct quantitative differences were found for ground cytoplasm, endoplasmic reticulum, mitochondria, chloroplasts (envelope, stroma, and thylakoids) and lipid droplets.

2.4.2.1. Induction of turions and turion-like fronds

Most factors inducing development of turions or turion-like fronds in Lemnaceae are agents which simultaneously promote senescence in the mature parts of the plant. This holds true for several forms of mineral deficiency or imbalance, as well as for the effects of short day, red light, and abscisic acid.

Turion induction in S. polyrrhiza has been found after exhaustion of minerals (CZOEK 1963a, HENSSEN 1954), withdrawal or lowering of nitrogen (JACOBS 1947, MALEK and COSSINS 1983a, MALEK and ODA 1979, NEWTON and DUFFEY 1975, NEWTON et al. 1978, PERRY 1968), sulfate deficiency (MALEK and COSSINS 1983a), and phosphate deficiency (AUGSTEN and JUNGNIKKEL

1983, JUNGnickel 1978, 1986a). Different clones, however, behave differently and, in addition, an endogenous seasonal factor can influence the extent of turion production. Nitrogen deficiency was effective in only two of four clones investigated by PERRY (1968). In clone P115 (ETH No. 7505), used by MERTEN and KANDELER (unpublished results), turion induction by nitrogen deficiency proceeded from January to March, but not during May. HENSEN (1954) obtained turions by mineral deficiency during the whole year, but turion formation in full nutrient medium occurred in the "Gr" clone during the winter only. In a similar way turion formation in *L. turionifera* can be observed after exhaustion of minerals, especially phosphate deficiency. Under short day conditions, turions of some clones are formed under otherwise optimal conditions (LANDOLT, not publ.). BEYER (pers. comm.) observed turion formation during winter in a clone from Southern California even under long day and otherwise optimal conditions. It is not investigated whether turion formation is induced by an endogenous rhythmic factor or by the air pollution which is greater during the winter.

Short day hastens senescence in mature fronds and induces development of turion-like fronds in *L. perpusilla* Pl46 (ETH No. 7507) (HUEGEL et al. 1979). The short-day effect is truly photoperiodic, because a short night break with red light cancels both responses. Another light program, which abolishes the short-day effects on senescence and production of turion-like fronds in *L. perpusilla*, is short irradiation with far red at the end of the daily light period. The connection between senescence and formation of dormant storage organs seems to be very close in *L. perpusilla*, as can be seen by the fact that first-daughter fronds are the last which reach the dormant state (HUEGEL et al. 1979).

Far red at the end of short day inhibits turion development also in *S. polyrrhiza* (MERTEN and KANDELER, unpublished results). Furthermore, when the infrared content (beyond 750 nm) of irradiation is relatively high during long day, production of turions in *S. polyrrhiza* is reduced (DEUTCH and RASMUSSEN 1974). Far red as well as infrared may be effective by lowering the level of active phytochrome. Continuous blue light diminishes turion formation in *S. polyrrhiza* in comparison to red light (MALEK and ODA 1979). The light effect may act through an influence on the degree of senescence, because blue light in comparison to red shifts the ratio of protein to carbohydrate synthesis to the former (VOSKRESSENSKAYA 1979). In accordance with this assumption are the facts that w.

arrhiza cultivated under continuous blue light contains distinctly more cytokinins than under red light (EICHORN and AUGSTEN 1980), and that turion formation in *S. polyrrhiza* (induced by nitrate or sulfate deficiency) is prevented by addition of isopentenyl adenine or benzyladenine (MALEK and COSSTIN 1983a).




Fig. 2.33. Effect of abscisic acid on growth and turion formation (STEM-ART 1969). (●) growth rate; (▲) turion ratio. (All points are the mean of three determinations.)

Abscissic acid (ABA) as a senescence-promoting hormone induces turions in S. polyrrhiza (fig. 2.33) (PERRY and BYRNE 1969, SAKS et al. 1980, SMART and TRENAVAS 1983a, STEWART 1969) and in L. turionifera (VAN OVERBEEK et al. 1968), and turion-like fronds in S. punctata and L. gibba (SCHARPETER and KANDELER, unpublished results). No effect of ABA has been found in one of four clones of S. polyrrhiza (PERRY and BYRNE 1969) and in W. gladiata (PIETERSE et al. 1970d). Presumably a high cytokinin content of plants hinders the realization of the ABA effect. Kinetin applied simultaneously with ABA prevents turion formation in S. polyrrhiza (STEWART 1969). In old cultures of S. polyrrhiza ABA is released from the plants to the medium, in this way acting not only within the ABA-synthesizing individuals but also on their neighbours. The time of appearance of released ABA in the solution was correlated with the beginning of turion formation in the old culture (SAKS et al. 1980). SMART and TRENAVAS (1983a) stated that only frond primordia <0.7 mm long, i.e., before the stage of rapid cell expansion, could be induced by ABA to change morphology to turion formation. During about 3 days the effect of ABA is reversible. Plants transferred to ABA-free medium after this time did not produce turions. Cytokinins and ABA are not the only hormones which influence senescence and turion formation in Lemnaceae. Gibberellin A₃ retards senescence as well as induction of turion-like fronds in L. perpusilla (HUEGEL et al. 1979).

As in senescence sugars can have an inhibiting or promoting effect on turion production. In L. perpusilla short-day induced development of turion-like fronds is prevented by addition of sucrose to the medium. Simultaneously, senescence of mature fronds is retarded (HUEGEL et al. 1979). A promotion of turion production by exogenous sucrose or glucose, on the other hand, has been shown in S. polyrrhiza (CZOPEK 1963a, HENSSSEN 1954, NEWTON and DUFFEY 1975, NEWTON et al. 1978), W. gladiata (PIETERSE et al. 1970d), and W. arhriza (GODZIEWBA-CZYZ 1969). Furthermore, factors which increase the endogenous production of photosynthates, as high intensity light, long daily light period and CO₂ enrichment, promote turion formation in S. polyrrhiza (JACOBS 1947). The effects of exogenous and endogenous carbohydrates on turion formation may come about in part indirectly by increasing growth and, therefore, exhaustion of minerals in the medium. Sugars and photosynthesis-promoting agents, however, are turion inducing also under growth-inhibiting conditions. They increase the turion-inducing effect of low night temperature

(10-15°C) (JACOBS 1947), and mineral deficiency (NEWTON and DUFFEY 1975, NEWTON et al. 1978).

2.4.2.2. Dormancy and germination of turions

Turions in comparison to vegetative fronds of S. polyrrhiza are characterized by a lack of many proteins as revealed by polyacrylamide gel electrophoresis (PERRY 1963, SMART and TRENAVAS 1984a, see chapter 2.5.7.3.4). The capacity for photosynthesis is lowered drastically (CZOPEK 1967. See, however, BEER 1985). Also in turions of W. arhriza rates of gas metabolism are reduced. The water content of these turions amounts to 78.9% in comparison to 97% of vegetative fronds (GODZIEWBA-CZYZ 1970). In turion-like fronds of W. gladiata the content of gibberellins (similar to GA₁, GA₂, GA₃ and GA₈ in the chromatograms) is lowered distinctly and another fraction with gibberellin-like activity is raised (PIETERSE 1972, PIETERSE et al. 1971a).

Turions of S. polyrrhiza are very tolerant to anaerobic conditions, but not resistant to air drying. They are tolerant to temperature extremes of -8° and 50°C for at least 24 hrs. Two days at -12° or one day at 60°C, however, are lethal (JACOBS 1947). Interestingly, there are some observations that turions are able to survive in India in ponds that dry out provided that they are completely covered by mud (B. GOPAL, pers. comm.). In W. arhriza floating fronds are more resistant to desiccation than turions and resistance to low temperatures is similar in turions and vegetative (floating and immersed) fronds. Only darkness is endured better by turions (GODZIEWBA-CZYZ 1970).

The dormant state of turions lasts to a different degree depending on the kind of turion-inducing factors, culture conditions during turion development, and genetic properties of the clone. Turions of S. polyrrhiza strain 0-381 induced by low temperature possess a dormancy, which is broken either by storage in N-deficient medium and darkness (>12 days) or by storage under continuous light (>20 days). Turions of the same strain, however, which are induced by long-lasting nitrogen deficiency, need no further treatment, before germination can proceed in N-containing medium (SIBASAKI and ODA 1979). Supply of sucrose or another sugar (NEWTON and DUFFEY 1975) or gibberellin A₃ (LACOR 1969) during induction of turions removes the requirement of chilling to break dormancy. Long day can replace cold treatment to break dormancy in strain 16-

62 of *S. polyrhiza*, but not in strain 8-64 (PERRY 1968). In principle, the following factors break dormancy in turions of *S. polyrhiza*: certain metabolic inhibitors, osmotic stress, nitrogen deficiency in combination with sucrose, chilling, long day, gibberellins and cytokinins. HENSSEN (1954) has shown that a treatment of turions with 0.1% potassium cyanide or 0.2% 2,4-dinitrophenol (40-48 hours) breaks dormancy at any time. Similarly, dormancy can be broken by treatment (1-2 hours) with 3% sodium sulfocyanate, 95% ethanol or 0.1% IAA (YOSHIMURA 1950). Germination of turions amounts to 90-100% within 2 days, if treated with the osmoticum polyethylene glycol 6000 (5-20%) for 4 days prior to the germination test. Mild drying of turions in air has a similar, but less pronounced effect (MALEK 1981). Turions of strain 0-381, which were stored in the turion-inducing medium (1/20 nitrate Hoagland's solution with 1% sucrose) or in 1% sucrose solution, reached full germination capacity about 20 days after onset of turion formation. In 1/20 nitrate medium lacking sucrose, however, turions need 51 days to reach partial germination ability (MALEK and ODA 1980). For chilling the following temperature programmes have been found to be effective in different strains of *S. polyrhiza*: 14 days at 10°C or below (JACOBS 1947), 20-30 days at 4°C (NEWTON et al. 1978, PERRY 1968), 28 days at 5.5°C (HENSSEN 1954), and 1 month at 0-3°C, turions in water (CZOPEK 1962, 1964a). When turions are stored at 25°C, dormancy can last 6 months or longer (JACOBS 1947). Turions collected by HENSSEN (1954) during autumn from the natural location remained in the dormant state for 10 months, when stored at 20°C in the dark. Some germination capacity, however, began to appear already after 4 months, when turions were given light during the germination test. Application of long day or continuous light during turion storage is a further agent for shortening of the dormant state. Unchilled turions of strain 16-62 germinate within 21 days only when held under long day, but not under short day. Supply of 10 ppm gibberellin A₃ has about the same effect as long day. Gibberellin and long day, given together, cause an immediate germination as after chilling (PERRY 1968). Dormant turions, strain 0-381, stored under continuous light germinate after a lag period of about 24 days. When, however, the turions are darkened during the first 6, 12, or 18 days of storage period, then the onset of germination is delayed to a corresponding extent (SIBASKI and ODA 1979). A positive effect of gibberellin A₃ as well as of kinetin on breaking dormancy has been demonstrated by LACOR

(1969). The highest germination rate in light is obtained when both gibberellin and kinetin are added to the medium. When the dormancy of young turions has been varied by a different phosphorus supply during turion development, benzyladenine strongly increases germination independent of the dormant state (JUNGNIKEL 1986a).

For light and some other factors it is important to distinguish between effects on breaking dormancy as a precondition of germination and on germination itself. Light promotes germination also in turions which have been chilled for an adequate time (HENSSEN 1954, NEWTON et al. 1978). Ten minutes of white or red light per day are sufficient to stimulate turion germination. Phytochrome seems to mediate the light effect, because far red irradiation after red abolishes the effect of red light, at least partially (CZOPEK 1962). Kinetin (10⁻⁶M) is another agent which promotes germination of chilled turions. The substance is effective in the dark as well as in combination with a short red light. Gibberellin A₃ (10⁻⁵-10⁻⁷M) inhibited dark or red light-stimulated germination in these experiments (CZOPEK 1964a). Benzyladenine and isopen-tenyladenine render germination of osmotically pretreated turions possible even in pure water (MALEK and COSSINS 1983a). Without added cytokinins these turions need at least supply of the anions which were omitted from the medium during turion induction for germination. Turions formed in nitrate-deficient medium germinate in 1 mM Ca(NO₃)₂, but not in 1 mM MgSO₄. The reverse holds true for turions which were produced in a sulfate-deficient medium. DUDLEY (1983) observed germination of turions of *L. turionifera* in old cultures if phosphorus was added to the solution.

The floating up of turions to the surface of the water is made possible by gas bubble formation in the light (JACOBS 1947). The bubbles seem to consist of photosynthetically evolved O₂, as photosynthetic rates of submerged turions correlate positively with their surfacing rate. Enriched CO₂ concentrations increase photosynthesis as well as surfacing (BEER 1985).

2.4.2.3. Dormancy and germination of seeds

In contrast to turions, seeds are more or less resistant to desiccation. In Israel, WITZTUM (1977) observed fruiting *L. gibba* in a shallow pool which were desiccated completely by August. Seeds taken from this loca-

tion germinated within a week in the laboratory under continuous light at 25°C. In the natural environment the seeds survived until the next rainy season and germinated during December and January. Seeds of L. gibba, however, collected in August from a Czechoslovakian fishpond and kept under dry conditions at room temperature for more than one year showed only a very small percentage of germination (REJMANKOVA 1976). Control seeds stored in diluted medium in the dark exhibited 70% germination under long day at the same time, and lost their viability only during the course of the following ten months.

Seeds of L. gibba are able to germinate soon after ripening. Germination does not occur in the dark, but a short dark pre-treatment seems to be necessary for rapid germination. Increasing water temperature (17° to 33°C) promotes the rate of germination (REJMANKOVA 1976).

Similar results have been found with seeds of L. aquinoctialis 6746 (GLICKMAN and POSNER 1966, POSNER and HILLMAN 1962). No germination proceeds without light, also when the medium has been complemented with yeast extract plus casein hydrolysate, kinetin, or gibberellin A₃. On the other hand, germination takes place only after a lag period of 6-7 days, when short day, long day, or continuous light is applied without preceding 1-4 days darkness. Gibberellin A₃ enhances the germination rate under such conditions. The most rapid germination is achieved when 4 days darkness (at >10°C) are followed by continuous high-intensity light (800 ft-c). Kinetin or - especially - casein hydrolysate in combination with lower light intensity (200 ft-c) can replace the requirement of high-intensity light. In S. intermedia germination has been observed in darkness in a medium complemented with casein hydrolysate, yeast extract, kinetin, and gibberellin A₃ (ROSSI 1969).

In L. perpusilla PL46 germination proceeds only after chilling of seeds, for example, by keeping the culture in a refrigerator (4-6°C) for one month (KANDELER 1975). Releasing the embryo from its seed coat, however, makes germ-bud development possible without chilling on a sucrose- and kinetin-containing medium (KANDELER and HUBGEL 1974a). A germination-inhibiting substance, presumably ABA, seems to be located chiefly in the seed coat (and possibly the pericarp).

2.4.3. Flowering

2.4.3.1. Flower induction

Using the advantages of Lemnaceae as axenically cultivable small water plants, several workers examined the effects of diverse organic and inorganic substances on flower induction. From the summary of these investigations (table 2.25) one can make a first survey on the factors which have an influence on flower initiation. As in other plants flowering depends on all the known phytohormones, their metabolism and interactions. Nitrogen and micro nutrient supply, light intensity and colour, as well as day length and temperature are the main environmental conditions, which are tested by the plant to come to the "decision" to flower or not. In table 2.25 the rough direction (positive or negative) of substance effectiveness is entered. It should be taken into consideration, however, that many agents work at a certain concentration range, under certain experimental conditions, or in certain species or strain only. In dependence of concentration and experimental conditions one and the same substance may even have a flower-promoting or flower-inhibiting effect. In some cases the classification of agents is rather hypothetical (EDDHA and salicylic acid, for instance). These and other details are discussed in the following chapters. A more circumstantial review on physiology of duckweed flowering has been given recently (KANDELER 1985).

2.4.3.1.1. The hormonal state

Phytohormones seem to play a central role in flower induction. Presumably they are important for mediation, integration, and processing of environmental stimuli as well as for internal coordination of flowering with other developmental processes.

All the known phytohormones modify the extent of flowering if added to the nutrient medium of duckweed cultures (table 2.25). In L. gibba G3 (long-day plant = LDP) flowering is promoted by gibberellin A₃ (10⁻⁶-10⁻⁵ M) under long-day conditions (CLELAND and BRIGGS 1969, CLELAND et al. 1982, OOTA 1965, OOTA and TSUDZUKI 1971). Higher concentrations of gibberellin A₃ (10⁻⁴ M), however, as well as the inhibitor of gibberellin synthesis, CCC, are flower inhibiting (CLELAND and BRIGGS 1969,

Tab. 2.25. Substances influencing flower initiation in *Lemnaea*

+ promoting
- inhibiting
0 not or only slightly promoting

LDP long-day plants
SDP short-day plants

Experiments with amino acids of low concentration (10^{-6} M) are not included in the table (see text).

	LDP	SDP	References	
			LDP	SDP
A. Hormones. Inhibitors and promoters of hormone synthesis				
Gibberellin A ₁	+/-	-	9, 48, 58, 89, 90	15, 25, 26
CCC	-	+	9	37
Allogibberic acid	-	-	59	69
Abscisic acid	+/-	+/-	70, 89, 90, 99	15, 37
Zeatin (and other cytokinins)	+/-	+/-	48, 58, 61, 89, 90, 106	1, 15, 16, 18, 45, 48, 86, 95
Indoleacetic acid	+/-	-	48, 58, 90	15, 18
Ethephon	-	-	59, 70	
ACC	+/-	+/-	70, 103	103
AVG, AOA	-	-	70	
Co ²⁺	-	-	70	17
Polyamines	-	-	104	
MGBG	+	+	104	
B. Some heavy metals, chelating agents, and phenolic substances				
Mn ³⁺	+	+	42	17, 22, 43, 74, 101
Fe	-	-	49	
o-Phenanthroline, α, α'-dipyridyl, azide (iron reagents)	+	+	3, 60, 71	17, 43, 74
EDDHA (compare also EDTA, group C)	+	+	2, 6, 7, 8, 10, 27, 54, 55, 61, 71, 77, 79, 90, 106	2, 10, 15, 27, 41, 78, 94
Salicylic acid, benzoic acid, acetylsalicylic acid	+	+		96, 98
8-Hydroxyquinoline	+	+		97
Tannic acid	+	+		93
Dicoumarol	+	+		93
4-Hydroxycoumarin	+	+		105
2-Hydroxy-1-phenyl-1, 4-pentadione	+	+		105
Phenylglyoxal	+	+		105
Chlorogenic acid	-	-	84	
β-Naphthol	+	+	102	
Benzo (a)anthracene	-	-	4	

Table 2.25 (continued)

	LDP	SDP	References	
			LDP	SDP
C. Substances involved in nitrogen metabolism				
NO ₃ ⁻	+/-	+/-	14	14
MoO ₄ ²⁻	+	-	5	76, 83
WO ₄ ²⁻	-	+	87	76, 83
Cu ²⁺ (and other sulfhydryl blockers:	+/-	+	23, 42, 87	23, 72, 75
Ag ⁺ , Hg ²⁺ , iodoacetamide)				
Salicylaldehyde, diethyldithiocarbamate, hydroxyquinoline	+		49	
(copper chelators)				
EDTA	+	+/-	21, 22, 28, 51, 53	20, 43, 44, 75
NH ₃	-	-	33	24, 80
Urea	-	-		80
Casamino acids	-	-		80
Cysteine, cystine, arginine, phenylalanine, tryptophane	-	0/-	46, 47	65, 80, 81
Aspartate, glutamate	0	+ / 0	46, 47	64, 65, 80, 81, 83
Alanine, glycine, serine	-	+/-	46, 47	64, 65, 80, 81, 83
Threonine, proline, iso-leucine	-	0/-	46, 47	65, 80, 81
Lysine	+/-	0/-	46	65, 80, 81
Leucine, tyrosine, valine, methionine, hydroxyproline, histidine	-	0	46, 47	65, 81
Asparagine, glutamine		+ / 0/-		65, 80, 81, 83
D. CO ₂ , carbohydrates, and energy budget				
CO ₂	+/-	-	29	65
Sucrose	+/-	+/-	21, 30, 31, 38, 52	13, 26, 38, 62, 63, 64, 65, 66, 67, 72, 73
Glucose, fructose	0/-	0/-	38, 50	38, 62, 63, 64
NADH, NADPH	+	-	35	35
Ascorbic acid	-	+	35	35
DCMU	+	+/-	32	68, 72, 73
ATP, ADP	+	+/-	32, 33, 38	34, 40, 65, 67

Table 2.25 (continued)

	LDP	SDP	References	
			LDP	SDP
Arsenate (and other respiratory inhibitors)	-		30,49	
2,4-Dinitrophenol	-	-	49	40
Glucose-6-P (and other respiratory intermediates)	-	+		65
E. Effectors of membrane transport				
Acetylcholine, eserine	-	+	36,55	36
Valinomycin, gramicidin	+		56	
Li ⁺	-	+	34	34
Ca ²⁺	+	+/-	100	19,23,63
PO ₄ ³⁻	+	+/-	100	17,19,23,63,67,101
F. Some further substances				
Ferricyanide, ferro-cyanide, CN	+	+	77,87	75,76,82,83
Nicotinic acid	+	+	91	91
cyclic AMP	+	+	39,50,55,57	67
Isoproterenol (and other catecholamines)	+		52,55	
Vitamin K ₅	-	+		92
5-Fluorodeoxyuridine	-	-	85	
5-Fluorouracil	-	-	85	12
2-Thiouracil	-	+/-	85	11,12
Cycloheximide, ethionine, chloramphenicol	-	-		12
Agar	-		88	

References:

- 1 BENNINK AND DE VRIES 1975)
- 2 BEPPU AND TAKIMOTO 1981b)
- 3 BHALLA AND SABHARWAL 1972a)
- 4 BHALLA AND SABHARWAL 1974)
- 5 BUECHELE AND KANDELER, unpublished results
- 6 CLELAND 1974c
- 7 CLELAND 1979
- 8 CLELAND AND AJAMI 1974
- 9 CLELAND AND BRIGGS 1969
- 10 CLELAND AND TANAKA 1979
- 11 DOSS 1975c

Table 2.25 (continued)

12 DOSS 1975d	59 PIETERSE 1976
13 ESASHI and ODA 1964	60 PIETERSE et al. 1970b
14 ESASHI et al. 1972	61 PIETERSE and MUELLER 1977
15 FUJIOKA et al. 1983b	62 POSNER 1967b
16 GUPTA and MAHESHWARI 1969	63 POSNER 1969a
17 GUPTA and MAHESHWARI 1970a	64 POSNER 1970
18 GUPTA and MAHESHWARI 1970b	65 POSNER 1971
19 HALABAN and HILLMAN 1970a	66 POSNER 1973b
20 HILLMAN 1959b	67 POSNER 1973a
21 HILLMAN 1961c	68 POSNER et al. 1977
22 HILLMAN 1961b	69 PRYCE 1973b
23 HILLMAN 1962	70 SCHARFETTER et al. 1986
24 HILLMAN and POSNER 1971	71 SCHARFETTER et al. 1978
25 HODSON and HAMNER 1971	72 SCHUSTER 1968
26 HUEGEL et al. 1979	73 SCHUSTER and KANDELER 1970
27 KAIHARA et al. 1981	74 SETH et al. 1970
28 KANDELER 1962	75 TAKIMOTO and TANAKA 1973
29 KANDELER 1964a	76 TAKIMOTO and TANAKA 1976
30 KANDELER 1967	77 TANAKA and CLELAND 1980
31 KANDELER 1968	78 TANAKA and CLELAND 1981
32 KANDELER 1969b	79 TANAKA et al. 1979a
33 KANDELER 1969a	80 TANAKA and TAKIMOTO 1975
34 KANDELER 1970	81 TANAKA and TAKIMOTO 1977
35 KANDELER 1971a	82 TANAKA and TAKIMOTO 1978
36 KANDELER 1972	83 TANAKA et al. 1979b
37 KANDELER and HUEGEL 1973	84 UMEMOTO 1971
38 KANDELER et al. 1975	85 UMEMURA and OOTA 1965a
39 KESSLER and STEINBERG 1973	86 VENKATARAMAN et al. 1970
40 KHUHAIRI and MAENG 1973	87 WATANABE and TAKIMOTO 1977
41 KHURANA and MAHESHWARI 1978	88 CLELAND 1985
42 KRAJNCIC 1976a	89 CLELAND et al. 1982
43 MAHESHWARI and GUPTA 1967	90 FUJIOKA et al. 1985
44 MAHESHWARI and SETH 1966b	91 FUJIOKA et al. 1986b
45 MAHESHWARI and VENKATARAMAN 1966	92 KAIHARA and TAKIMOTO 1985a
46 NAKASHIMA 1964	93 KAIHARA and TAKIMOTO 1985b
47 NAKASHIMA 1965	94 KHURANA and MAHESHWARI 1983a
48 OOTA 1965	95 KHURANA and MAHESHWARI 1983b
49 OOTA 1969	96 KHURANA and MAHESHWARI 1984
50 OOTA 1972a	97 KHURANA and MAHESHWARI 1986a
51 OOTA 1972b	98 KHURANA et al. 1986
52 OOTA 1974	99 KRAJNCIC 1985
53 OOTA 1975b	100 MAENG and BAE 1984
54 OOTA 1975c	101 MORI 1984b
55 OOTA 1977a	102 PIETERSE 1978b
56 OOTA 1977b	103 SCHARFETTER et al. 1987
57 OOTA and KONDO 1974	104 TSAO and JIN 1985
58 OOTA and TSUDZUKI 1971	105 SUZUKI et al. 1985
	106 WATANABE et al. 1983

frond senescence has the same effect on flowering as in *L. aequinoctialis* (LACOR 1970). A low concentration of ABA (0.1 mg/l) has a flower-promoting effect in this species, when given under long day and in combination with EDDHA (KRAJNCIC 1985). In contrast to the above cited results FUJIOKA et al. (1983b) found a flower-inhibiting effect of low ABA concentrations in the SDPs *L. aequinoctialis* 151 and 381, when flowering under continuous light was forced by addition of benzoic acid to the nutrient medium. KRAJNCIC (1985) demonstrated some promotion of flowering by 0.1-1 µg/l ABA in the LDP *L. minor*. 10 µg/l ABA, however, were inhibitory in all investigated photoperiodic types: the LDPs *L. minor* and *L. gibba*, the long-short-day plant *W. arhiza* and the day-neutral plant *S. polyrrhiza*.




Fig. 2.34. The contrary influence of mother frond senescence on flower induction in daughter fronds of *Lemna gibba* G1 and *L. aequinoctialis* 6746 (after data from KANDELER et al. 1974). Single mother fronds (four in both species) were cultivated under long-day conditions, and the resulting daughter fronds (with ordinal numbers 1 to 9) were examined for flowering. Senescence of the mother frond is responsible for the lack of flowers in late-produced daughter fronds of *L. gibba*, as the series of first-daughter frond generations (derived from the same mothers and cultivated in parallel under the same conditions) forms flowers all the time. In *L. aequinoctialis*, first-daughter fronds do not flower under long day, but daughter fronds with a higher ordinal number (i.e., produced by a senescing mother frond) can produce a flower in spite of the non-inductive day length.

CLELAND et al. 1982, OOTA and TSUDZUKI 1971). Likewise the decomposition products of gibberellin A_3 after autoclaving, allogibberic acid, gibberic acid and epiallogibberic acid, suppress flowering in *L. gibba* G3 (PIETERSE 1976). In the short-day plant (SDP) *L. perpusilla* p146 short-day induced flowering shows a distinct reduction with 5×10^{-7} M gibberellin A_3 (HUEGEL et al. 1979). The same results were found in the SDP *L. aequinoctialis* 381 with 1 ppm gibberellin A_3 (FUJIOKA et al. 1983b). CCC on the other hand, can force flowering in the SDP *L. aequinoctialis* 6746 under long day (KANDELER and HUEGEL 1973, TSAO et al. 1985). High concentrations of gibberellin A_3 (10^{-4} M) and gibberellin decomposition products are both flower inhibiting in *L. aequinoctialis* 6746 (HODSON and HAMNER 1971, PRYCE 1973b). The opposite behaviour of the two photoperiodic reaction types against low concentrations of gibberellin A_3 and against CCC may be explained by the hypothesis that the endogenous accumulation of gibberellins is different in the two groups of plants and must be corrected by day length in a different way to reach a flower-inducing level. In fact, uptake of [3 H]-gibberellin A_1 reaches a higher saturation level in *L. aequinoctialis* 6746 than in *L. gibba* G1 (AL-SHA-LAN and KANDELER 1979, HARTUNG and KANDELER 1976). Moreover, preliminary determinations of the endogenous gibberellin content in both species cultivated under the same conditions showed higher values in strain 6746 than in strain G1 (LADENBURGER, BAUER and KANDELER, unpublished results). Recently, TSAO et al. (1985) revealed that the level of gibberellins in *L. aequinoctialis* 6746 decreases after transfer of plants to short day.

Abscissic acid (ABA) is another hormone which has an opposite effect in the LDP *L. gibba* G3 and the SDP *L. aequinoctialis* 6746. 10^{-8} - 10^{-7} M ABA inhibit long-day induction of flowering in *L. gibba* G3 (CLELAND et al. 1982, SCHARFETTER and KANDELER, unpublished results). In *L. aequinoctialis* 6746 2×10^{-9} - 2×10^{-8} M ABA strengthen the flower-inducing effect of CCC under long-day conditions (KANDELER and HUEGEL 1973). Correspondingly, the level of ABA in strain 6746 increases transiently about threefold, when plants are transferred to short day (TSAO et al. 1985). In both species frond senescence has the same effect as ABA. Daughter fronds, which are produced under long day during the second part of the life-span of a mother frond, failed to flower in *L. gibba* G1, but came into flower in *L. aequinoctialis* 6746 in spite of the non-inductive day length (fig. 2.34) (KANDELER et al. 1974). In the day-neutral plant *S. polyrrhiza*

Cytokinins promote flowering in the LDs L. gibba (CLELAND et al. 1982, OOTA 1965, OOTA and TSUDZUKI 1971, PIETERSE et al. 1970b) and L. minor (WATANABE et al. 1983) as well as in the SDs L. aequinoctialis (FUJIOKA et al. 1983b, GUPTA and MAHESHWARI 1969, 1970b, KHURANA and MAHESHWARI 1983b), L. perpusilla (BENNINK and DE VRIES 1975, and W. microscopica (MAHESHWARI and VENKATARAMAN 1966, VENKATARAMAN et al. 1970) and in the long-short-day plant W. arthiza (KRAJNCIC 1983). Benzyladenine has the highest flower-promoting effectiveness of 5 different cytokinins in W. microscopica (VENKATARAMAN et al. 1970). In L. gibba 10^{-6} M kinetin enhances flowering only under short-day conditions (in combination with gibberellin, OOTA 1965), but inhibits flower initiation under long day (OOTA and TSUDZUKI 1971).

Indoleacetic acid (IAA) causes inhibition of flowering in L. gibba and L. aequinoctialis (FUJIOKA et al. 1983b, GUPTA and MAHESHWARI 1970b, OOTA 1965, OOTA and TSUDZUKI 1971), but a weak flower-promoting effect of 10^{-9} M IAA has been found in L. gibba (OOTA and TSUDZUKI 1971). Ethephon (Ethrel) as an ethylene releasing agent is flower inhibiting in L. gibba G3 (PIETERSE 1976). The ethylene precursor 1-aminocyclopropane-1-carboxylic acid (ACC), however, promotes flowering in L. gibba G1 as well as in L. aequinoctialis 6746, when plants are induced to low or moderate flowering by other agents (SCHARFETTER et al. 1987). The optimum concentration is lower in L. gibba (10^{-8} M) than in L. aequinoctialis (10^{-7} - 10^{-6} M). Higher ACC concentrations are inhibitory in both species. In S. punctata O5, ethylene-releasing substances (Ethrel, ACC) and inhibitors of ethylene formation (AVG, AOA, Co^{2+}) are effective only in the case that flowering has been induced by EDDHA (SCHARFETTER et al. 1986). Then both groups of substances greatly inhibit flower induction. It is suggested that flower initiation depends on an intermediate ethylene level, but only after accomplishment of certain other flower-inducing processes. EDDHA may reduce the level of free auxin (see next chapter) and, simultaneously, may adjust ethylene formation to a flower-promoting level. A flower-inhibiting action of Co^{2+} were found also in L. aequinoctialis (GUPTA and MAHESHWARI 1970a).

Summarizing the results hitherto, it can be stated that levels of gibberellins and ABA have to be changed in an opposite way for flower initiation in LDs and SDs and seem to be under photoperiodic control. Cytokinins and auxins, on the other hand, exert their effects on flowering more or less independently from the photoperiodic type of plant. Possi-

bly, certain polyamines belong to the first of the two groups of growth regulators. In L. aequinoctialis, the endogenous spermidine content decreases during a 4 day dark induction (putrescine increases concomitantly). Correspondingly, exogenous spermidine and spermine inhibit the short-day induction of flowering more strongly than putrescine. The inhibitor of spermidine and spermine synthesis, MGBG, induces flowering under continuous light (TSAO and YIN 1985). Generally, polyamines are known to delay senescence and, therefore, counteract the effects of ABA and ethylene. Polyamines and ethylene can inhibit their formation mutually (see PALAVAN et al. 1984). Thus, the higher demand of ethylene formation for flowering in the SD L. aequinoctialis (see above) may be related to the required reduction of polyamine synthesis in this plant.

2.4.3.1.2. Some heavy metals, chelating agents, and phenolic substances

Manganese and iron seem to be of special importance for flowering in Lemnaceae. Omission of manganese or iron from the nutrient medium lowers or prevents flowering in L. aequinoctialis (GUPTA and MAHESHWARI 1970a). In strain 6746 of L. aequinoctialis a temporary lowering of the iron level during short-day induction inhibits flowering without an effect on vegetative frond multiplication (HILLMAN 1961b). Especially under continuous darkness, the flower-retarding effect of medium dilution is abolished by supply of iron or phosphate ions. No other component of Huter's medium is effective in such a way in strain 6746 (MORI 1984a, 1984b). Furthermore, excessive doses of iron can force flowering in the Sohna strain of L. aequinoctialis and in W. microscopica under otherwise non-flowering conditions (GUPTA and MAHESHWARI 1970a, MAHESHWARI and GUPTA 1967, SETH et al. 1970). In L. gibba G3 PIETERSE et al. (1970b) found no inhibition of flower initiation by deletion of iron from the medium, but some iron reagents (o-phenanthroline, α , α' -dipyridyl, azide) inhibit flowering in this plant (OOTA 1969).

Subsequently the strong effect of some chelating agents on flowering may be reported. In LDs L. gibba (PIETERSE et al. 1970b), L. minor (BHALLA and SABHARWAL 1972), and S. punctata (SCHARFETTER et al. 1978), as well as in SDs L. aequinoctialis (GUPTA and MAHESHWARI 1970a, MAHESHWARI and GUPTA 1967) and W. microscopica (SETH et al. 1970) flower initiation is rendered possible (under certain experimental conditions) only when EDTA (ethylenediaminetetraacetic acid) or EDDHA (ethylenediamine-di-o-hydro-

xypenylacetic acid) is added to the nutrient medium. This effect can be made understandable by the well-known fact that such chelating agents increase the availability of iron and manganese for plants (for copper decontamination compare chapter 2.4.3.1.3). In L. gibba G3 addition of EDDHA to the medium doubles the iron content of the plants (PIETERSE 1975a), and this is also the case in L. aequinoctialis 6746, when the plants are cultivated under short day conditions (KHURANA and MAHESHWARI 1984). Under long day, EDTA is more effective than EDDHA to increase the endogenous iron level of L. aequinoctialis. In flower induction, EDDHA is distinctly more effective than EDTA in most cases, but in strain 6746 of L. aequinoctialis long-day flowering is achieved by EDTA but not by EDDHA (YUKAWA and TAKIMOTO 1976). Interestingly, ELZENGA et al. (1980) found an enhancement of ethylene production in L. gibba G3 after addition of EDDHA to the medium (preliminary result).

A group of substances, whose effectivity is very similar to that of EDDHA, are salicylic acid (SA), benzoic acid (BEA), and acetylsalicylic acid (ASA). The flower-promoting activity of SA was found by CLELAND (1974c; CLELAND and AJAMI 1974), when he tested with axenic L. gibba cultures the phloem sap of Xanthium plants for flower-inducing agents. In continuation of this work several authors confirmed the flower-inducing or flower-promoting effect of SA, BEA, and ASA in LDPs L. gibba (CLELAND and BEN-TAL 1982, CLELAND and TANAKA 1979), S. punctata (SCHARFETTER et al. 1978), L. minor (KAIHARA et al. 1981, WATANABE et al. 1983), and S type strains of L. aequinoctialis (BEPPU and TAKIMOTO 1981b, KAIHARA et al. 1981); moreover in many short-day-dependent strains of L. aequinoctialis (BEPPU and TAKIMOTO 1981b, CLELAND and TANAKA 1979, FUJIOKA et al. 1983b, KAIHARA et al. 1981, KHURANA and MAHESHWARI 1978), in the SDP W. microscopica (MAHESHWARI and KHURANA 1978, KHURANA and MAHESHWARI 1983a), and in the day-neutral plants S. polyrrhiza (KHURANA and MAHESHWARI 1980), L. obscura (CLELAND et al. 1982), and L. aequinoctialis K type strain (BEPPU and TAKIMOTO 1981b). With only one exception, SA generally is more effective than BEA in LDPs (CLELAND 1974c, KAIHARA et al. 1981), whereas BEA is more effective than SA in SDPs (KAIHARA et al. 1981, WATANABE et al. 1981, WATANABE and TAKIMOTO 1979).

Analysing the effect of SA in L. gibba G3 CLELAND and BEN-TAL (1982) found an immediate decrease of flowering after removal of SA from the medium. Thus, for maximal effectiveness SA must be present in the medium

for the entire experiment. An explanation for this behaviour is given by the observation that [¹⁴C] SA after uptake is partially converted to one or more bound forms and then the free and bound SA are stored within the cells (presumably the vacuoles) without further transport or metabolism (BEN-TAL and CLELAND 1982). Correspondingly, FUJIOKA et al. (1983a, 1985) found no positive correlation between the endogenously produced level of BEA and flowering in L. aequinoctialis strains 151, 381, and 321, and in L. gibba G3.

Besides SA and BEA, certain other phenols (respectively derivatives of phenols) have been found to induce or promote flowering in Lemnaceae. 8-Hydroxyquinoline increases the critical day-length in L. aequinoctialis (KHURANA and MAHESHWARI 1984) and makes long-day flowering possible in W. microscopica (KHURANA et al. 1986). With 10⁻⁵M tannic acid L. aequinoctialis 6746 flowers to a high degree even under continuous light (KHURANA and MAHESHWARI 1986a). Dicummarol has a strong flower-promoting effect especially under near critical photoperiods in strains 6746 and 441 of L. aequinoctialis, but is highly effective also under continuous light in strain 151 of the same species (KAIHARA and TAKIMOTO 1985b). 4-Hydroxycoumarin, a component of dicoumarol, is active like dicoumarol in strain 151. A flower-inducing substance, which has been isolated from Pharbitis purpurea, 2-hydroxy-1-phenyl-1,4-pentadione, as well as phenylglyoxal, a component of this substance, show flower-inducing activity as high as benzoic acid in strain 151 (SUZUKI et al. 1985). In contrast, chlorogenic acid and β-naphthol are phenols, which abolish flowering in the LDP L. gibba (UMEMOTO 1971, PIETERSE 1978b). The flower-inhibiting effect of benzo(a)anthracene in L. gibba may be mentioned as an appendage in this context (BHALLA and SABHARVAL 1974).

At least in the case of 8-hydroxyquinoline and EDDHA there are arguments that the flower-inducing activity may be based on chelation of heavy metals. 8-Hydroxyquinoline increases the iron content of L. aequinoctialis 6746 more strongly than EDTA and EDDHA, increases the copper content instead of a decrease caused by EDTA and EDDHA (see chapter 2.4.3.1.3), and has a higher flower-inducing activity than EDTA and EDDHA in the same experiments (KHURANA and MAHESHWARI 1984). Long ago, MAHESHWARI and GUPTA (1967) demonstrated that in another strain of L. aequinoctialis the flower-inducing effect of EDTA and EDDHA can be mimicked by supply of high doses of ferric citrate. In W. microscopica, iron uptake is higher with Fe-EDDHA than with EDTA + Fe-citrate, and, correspondingly,

long-day flowering is achieved with Fe-EDDHA, but not with EDTA + Fe-citrate (SETH et al. 1970).

Presumably, chelation of heavy metals is not the only way, in which phenols act on flower induction. Activation or inhibition of IAA-oxidases, formation of conjugates with IAA or with polyamines, and free radical scavenging (leading to a lower ethylene formation) are further processes, which - singly or in combination - may participate in the action mechanism. A further working point for SA and related substances may be the signalling action of oligosaccharides. DOHERTY et al. (1987) have shown that ASA abolishes the effect of oligosaccharides on the activity of serine proteinases in tomato leaves. Agar (CLELAND 1985) as well as oligosaccharides (ALBERSHEIM et al. 1987) inhibit flowering in L. gibba G3, and the effect of agar is nullified by SA (see chapter 2.4.3.1.8). Interestingly, TAKIMOTO (1987) has reported that BEA is bound to apiofuranosyl-1,6-glucopyranoside, aspartate, isocitrate, and malate only, when supplied to L. aequinoctialis 151.

In any case, lowering of the endogenous free auxin level seems to be one of the major consequences of application of flower-promoting phenols. When in L. gibba G3 flowering is induced by supply of SA, a distinct part (40%) of endogenous IAA is converted to a bound form (EPSTEIN and BENTAL 1985). SA itself may react with IAA, as SA is converted in part to one or more bound forms after uptake (BEN-TAL and CLELAND 1982). From p-hydroxybenzoic acid, which has virtually no effect on flowering in L. gibba, a higher percentage remains as the free acid after uptake (CLELAND and KANG 1985). In S. punctata, the effects of EDDHA on flowering and several other developmental processes are counteracted by the auxin NAA (SCHARFETTER et al. 1978). The anti-auxin PCIB while not acting on flowering mimics the EDDHA effects on seven vegetative attributes in S. punctata (SCHARFETTER et al. 1986). The lack of flower-inducing activity of PCIB may be caused by the fact that EDDHA changes not only the auxin but also the ethylene metabolism. The similarity of SA and EDDHA effects has been explained by the assumption that EDDHA is degraded by the plant to SA-like fragments (CLELAND et al. 1982).

2.4.3.1.3. Significance of nitrogen metabolism

As in many other plants, nitrogen supply has a distinct influence on flower initiation in Lemnaceae. ESASHI et al. (1972) investigated the

effect of different nitrate concentrations and found strong effects especially under conditions of high-intensity light (7000 lux). In the SDP L. aequinoctialis 6746 flowering under short day is optimum at medium or high nitrate concentrations, whereas optimal vegetative growth is achieved at lower NO_3 levels under the same conditions. In continuous light the same strain needs relatively low or medium concentrations of nitrate for good flowering (medium or high concentrations for vegetative growth). In the LDP L. gibba G3 the nitrate effect on flowering depends on the pH of nutrient medium. At pH 5.5 optimal long-day flowering occurs with medium nitrate supply (optimal vegetative growth with high nitrate supply). At pH 4.2 a high NO_3 level is needed for optimal flowering and a medium NO_3 level for vegetative growth. In summary, the claims of vegetative and generative development for nitrate supply are different in both photoperiodic reaction types and under diverse experimental conditions.

Several observations have led to the hypothesis that activity of nitrate reductase is of great importance for flowering in Lemnaceae (TAKIMOTO and co-workers). Deletion of molybdate, an activator of nitrate reductase, from the medium makes long-day flowering possible in the SDP L. aequinoctialis 6746 (TAKIMOTO and TANAKA 1976, TANAKA et al. 1979b). In the LDP L. gibba G1, on the other hand, addition of molybdate to the nutrient medium enhances long-day flowering (BUECHELE and KANDELER, unpublished results). Correspondingly, tungstate, a specific inhibitor of nitrate reductase, promotes long-day flowering in L. aequinoctialis, but suppresses flowering in L. gibba (TAKIMOTO and TANAKA 1976, TANAKA et al. 1979b, WATANABE and TAKIMOTO 1977).

More or less identical with the effect of tungstate is that of copper and other sulphhydryl blockers (Ag^+ , Hg^{++} , iodoacetamide), which may also work through inhibition of nitrate reductase. All these agents induce long-day flowering in L. aequinoctialis 6746 (HILLMAN 1962, SCHUSTER 1968, TAKIMOTO and TANAKA 1973, NASU et al. 1984). The effect can be cancelled by supplemental addition of cysteine (TAKIMOTO and TANAKA 1974). In L. gibba G3 the same SH inhibitors inhibit flowering under continuous illumination (HILLMAN 1962, WATANABE and TAKIMOTO 1977). Copper chelators as salicylaldehyde, diethyldithiocarbamate and hydroxyquinoline, on the other hand, enhance flowering in L. gibba G3 (OOTA 1969). Certain effects of EDTA on flowering - inhibition of long-day flowering in L. aequinoctialis 6746 and promotion of flowering in L.

gibba G3 - are also caused by sequestration of free copper from the medium (HILLMAN 1961d, 1962). In both species, Cu^{2+} uptake is prevented almost completely by EDTA (TANAKA et al. 1982a, NASU et al. 1983). Compare also KHURANA and MAHESHWARI 1984).

In their later publications TAKIMOTO and co-workers rejected the original hypothesis that SH inhibitors act on flowering by blocking nitrate reductase (TAKIMOTO and TANAKA 1976, TANAKA and TAKIMOTO 1978, WATANABE and TAKIMOTO 1977), because under their experimental conditions lowering of nitrate concentration in the medium does not act like inhibitors of nitrate reductase, but rather like molybdate (that is flower inhibiting in L. aequinoctialis 6746). This discrepancy, however, can be explained by the assumption that nitrate has a twofold function for flower-inducing processes. On the one hand, nitrate may be effective by being a substrate of nitrate reductase and in consequence a material for protein synthesis, in this way leading to a retardation of senescence and the connected hormonal changes (lowering of gibberellin/ABA balance). On the other hand, nitrate may have a more direct influence on the hormonal state of the plant. In L. aequinoctialis 6746 the presence of nitrate is needed for an adequate cytokinin synthesis. Twenty-four - 48 hours after omission of NO_3^- from the medium the level of cytokinins is lowered in this plant (JOEPFERT, GRUNTZEL and KANDELER, unpublished results). Recently, TANAKA et al. (1986a) confirmed that the dependence on concentration of copper-, silver-, tungstate-, and molybdate-deficiency-effects on nitrate reductase activity (inhibition) is about the same as in the corresponding effects on long-day flowering (promotion) in L. aequinoctialis 6746. They now conclude again that one of the flower-inducing actions of copper, silver, tungstate, or the deletion of molybdate is to suppress the nitrate assimilation. There is also a relation between effects of ferricyanide on flowering and nitrate reductase activity as with the other above-named agents (TANAKA et al. 1986a). In hybrids of strain 6746 with strain 371 of L. aequinoctialis, however, only the ferricyanide effectivity, but not the copper and silver effectivity is maintained (BEPPU and TAKIMOTO 1983). Furthermore, ferricyanide acts on flowering in both photoperiodic reaction types in the same way, i.e., not in an opposite way as the other nitrate reductase inhibitors. Therefore, the principal action mechanism of ferricyanide in connection with flowering seems to differ from that of nitrate reductase inhibitors (for the details of ferricyanide effects see chapter 2.4.3.1.8).

Other nitrogen sources, which have been tested for their effect on flower initiation, are nitrite, ammonia, urea, casamino acids, glutamine and asparagine. All these substances abolished long-day flowering in L. aequinoctialis 6746 induced by Cu^{2+} , Ag^+ or ferricyanide (TANAKA and TAKIMOTO 1975). In connection with the effect of ammonia TANAKA et al. (1982a) have stated that this substance inhibits the uptake of copper from the medium (see also NASU et al., 1983). Ammonia is flower inhibiting in strain 6746 also under short-day conditions when the nutrient medium is diluted (1/10 conc.) and complemented with 1% sucrose (HILLMAN and POSNER 1971). In the LDP L. gibba, strains G1 and G3, ammonia suppresses flowering without a simultaneous effect on the vegetative multiplication of fronds (KANDELER 1969a).

The effectivity of single amino acids was investigated by several authors. As can be seen from tab. 2.25 most of the amino acids inhibit flower production in the LDP L. gibba, if given in a concentration range of 10^{-5} - 10^{-3} M (NAKASHIMA 1964, 1965). Aspartate and glutamate have a small effect only and lysine, which is flower inhibiting in itself, can partially or wholly reverse the floral inhibition caused by some other amino acids. In L. aequinoctialis 6746 a group of amino acids, aspartate, glutamate, alanine, glycine, and serine, has a clear promotive effect on flowering in reverting the sucrose-induced inhibition (POSNER 1971) or supporting the inductive effect of a single 96 h dark period (TANAKA and TAKIMOTO 1977). All the other amino acids are not, or only slightly, promotive under these conditions. When the above named amino acids are tested with the same strain under long-day conditions, they cause an enhancement or an inhibition of flower initiation, depending on culture conditions (suboptimal concentration of ferricyanide or Mo-deficient medium on the one hand or slightly higher concentration of ferricyanide or tungstate addition on the other hand) (TANAKA et al. 1979b, TANAKA and TAKIMOTO 1975). MAENG and KHUDAIRI (1973), working with a lower concentration of amino acids (10^{-6} M), found, for strain 6746, a flower promoting action of serine, threonine, and tryptophane and a flower-inhibiting effect of cysteine, if amino acids were added to the nutrient medium during the last dark period of three inductive short-day cycles. KRAJNCIC and DEVIDE (1982b) studied the effect of amino acids in the LDP L. minor, strain Barje, and the day-neutral plant S. polyrrhiza, strain Petanjci. Using the same low concentration as MAENG and KHUDAIRI, they observed flower-promoting effects with glutamate, alanine, glycine,

serine, threonine, and some other amino acids under long day in both species. Under short day, however, preferably other amino acids are flower enhancing in S. polyrrhiza: valine, proline, iso-leucine, and histidine.

Aspartate, glutamate, alanine, glycine, threonine, and some other amino acids are repressors of nitrate reductase in tobacco (FILNER 1966). Assuming the validity of this statement for Lemnaceae, one can speculate that flower inhibition in the LDP and flower promotion in the SDP by such amino acids comes about in the same way as the effect of tungstate, Mo-deficiency, and sulphydryl blockers, namely by lowering of nitrate reductase activity. The observation that one and the same amino acid can exert a positive or negative effect depending on experimental conditions, may be explained by the fact that the molecule is substrate for protein synthesis and effector for enzyme repression at the same time.

2.4.3.1.4. Significance of carbohydrates and energy-generating processes

Under long-day conditions, addition of 1% sucrose to the nutrient medium causes an inhibition of flowering in LDPs L. gibba G1 and G3 (HILLMAN 1961c, KANDELER 1968, OOTA 1974), but - especially in combination with a copper supply - a flower promotion in the SDP L. aequinoctialis 6746 (ESASHI and ODA 1964, SCHUSTER 1968, SCHUSTER and KANDELER 1970). Under short day, however, supplemented with a short far-red irradiation at the end of the daily light phase, sucrose enhances flowering considerably in L. gibba G1 (KANDELER 1968). L. aequinoctialis 6746, precultured under long day with sucrose and grown in diluted ammonium-containing Hutner medium, shows a suppression of flowering with sucrose under short day (POSNER 1967b, 1969a, 1970).

Glucose and fructose seem to be effective like sucrose only when the plants were precultured with sucrose: in L. gibba G3 under long day (compare OOTA 1972a and KANDELER et al. 1975), as well as in L. aequinoctialis 6746 under short day or long day respectively (compare POSNER 1970 and KANDELER et al. 1975).

CO₂ enrichment of the air or high-intensity light results in the same effect as sucrose in most of the above-named situations. CO₂ (3.5%) inhibits long-day flowering, but enhances flowering under short day with end-of-day far red in L. gibba G1 (KANDELER 1964a). In L. aequinoctialis 6746 1% CO₂ inhibits flowering like sucrose under short day (POSNER

1971). High-intensity light inhibits flowering under long day in L. gibba G1 (KANDELER et al. 1975). In L. aequinoctialis 6746 high-intensity light in combination with copper makes flowering possible under long day, if a Hoagland-type medium is used (ESASHI and ODA 1964, SCHUSTER 1968). On the contrary, with a 1/2 strength Hutner-sucrose-medium (ammonium during preculture only) high-intensity light inhibits flower initiation in strain 6746 under long day (POSNER et al. 1977).

The correspondence between certain effects of sucrose, CO₂ enrichment, and light intensity suggests that photosynthesis is the physiological process which mediates CO₂- and light-intensity effects and works by delivering of assimilates. In fact, the effects of high-intensity light, cited above for L. aequinoctialis 6746, can be cancelled by application of the photosynthesis inhibitor DCMU (SCHUSTER 1968, SCHUSTER and KANDELER 1970, POSNER et al. 1977). There are, however, some experimental results which cannot be explained by the assumption that photosynthetically active light acts on flowering by production of assimilates. Shortly as well as long-day flowering can be achieved by DCMU in L. aequinoctialis 6746 although sucrose is present in the nutrient medium (POSNER 1973b, POSNER et al. 1977). (Compare experiments under long day with another nutrient medium where DCMU is not effective in the presence of sucrose; SCHUSTER and KANDELER 1970). The DCMU-mediated lowering of photosynthetically produced reductants may be responsible for this effect (POSNER et al. 1977). Addition of NADPH or NADH to the medium suppresses long-day flowering in strain 6746 whereas ascorbic acid promotes it (KANDELER 1971a). In connection with these statements it should be remembered that in L. aequinoctialis a low nitrate reduction is a prerequisite for long-day flowering (chapter 2.4.3.1.3).

During the light phase DCMU lowers the size of the ATP pool in L. aequinoctialis 6746 (GOWER and POSNER 1979). The DCMU effect on flowering in this strain under long day - reversion of high-intensity-light inhibition of flowering (see above) - is, however, not mediated by this reduction of the ATP pool, because the DCMU action proceeds also after addition of ATP to the medium (POSNER et al. 1977). Nevertheless, ATP or ADP supply has a distinct effect on flowering in strain 6746 under certain other experimental conditions. The percentage of flowering increases, when ATP is applied during the last dark period of three inductive short-day cycles (KHUHAIRI and MAENG 1973). Correspondingly, the uncoupler of oxidative phosphorylation 2,4-dinitrophenol diminishes flow-

ering, when applied during the same time. Long-day flowering of strain 6746, on the other hand, when induced in Pirson-Seidel medium with copper, is drastically reduced by addition of ADP to the medium (KANDELER 1970). In reverting the sucrose inhibition of flowering under short day not only ATP, but also other adenylates, inorganic phosphate, glucose-6-phosphate, and other respiratory intermediates are effective in strain 6746 (POSNER 1971, 1973a).

In L. gibba G1 the flower-promoting effect of DCMU (which blocks noncyclic electron transport, but permits cyclic photophosphorylation) in 1/10 Pirson-Seidel medium can be imitated by application of ADP (KANDELER 1969b). Several uncouplers of oxidative and/or photophosphorylation as arsenate, atebtrin, and 2,4-dinitrophenol inhibit flower initiation with or without promoting effect on frond multiplication in this species (KANDELER 1967, OOTA 1969). The effects of arsenate and atebtrin as well as the flower-inhibiting effects of sucrose and ammonia can be cancelled by addition of ATP or ADP (KANDELER 1969a, KANDELER et al. 1975). ADP is more effective than ATP in most cases, possibly because of the action of adenylate kinases. ATP and ADP were effective in compensating atebtrin inhibition, but not so AMP, adenosine, or adenine. Taking all the results together it seems likely that not only assimilates and photosynthetic reductants but also ATP from photophosphorylation or oxidative phosphorylation can contribute to the physiological processes which induce or inhibit flower initiation in Lemnaceae.

2.4.3.1.5. Photoperiodism and effects of coloured light

Depending on the species, clone, or environmental conditions, the effect of daylength on flowering can be strong, moderate, or lacking in Lemnaceae. A qualitative response is seen in the LDPs L. gibba (KANDELER 1955), and L. turionifera (LANDOLT 1957; BENNINK et al. 1970, named as L. minor), in the SDPs L. aequinoctialis (most strains, YUKAWA and TAKIMOTO 1976) and L. perpusilla (HUBGEL et al. 1979), and in the long-short-day plant W. arhriza (KRAJNCIC and DEVIDE 1980). A quantitative daylength effect was found in the LDPs S. polyrrhiza (one strain only, KRAJNCIC and DEVIDE 1982a), S. punctata (SCHARFETTER et al. 1978) and S type strains of L. aequinoctialis (BEPPU and TAKIMOTO 1981b), and in the SDPs L. aequinoctialis (strain 151, WATANABE and TAKIMOTO 1979) and Wolffia brasiliensis (MAHESHWARI and SETH 1966b). Day-neutral plants are

S. polyrrhiza (most strains, KRAJNCIC and DEVIDE 1980) and L. aequinoctialis (K type strain 351, YUKAWA and TAKIMOTO 1976). Strain 6746 of L. aequinoctialis behaves as a qualitative SDP under some experimental conditions, but as a day-neutral plant under others (HILLMAN 1959b), and therefore can be denoted as conditional SDP. Strain Sohna of W. microscopica shows a qualitative short-day response, when cultivated in Hoagland medium, but only a quantitative response in Bonner-Devirian medium (VENKATARAMAN et al. 1970). Critical daylengths and the minimal number of inductive light-dark cycles vary from strain to strain and are listed in table 1 in KANDELER (1985).

Most of the effects of coloured light, which have been pointed out in Lemnaceae, are summarized in table 2.26. Blue, red, and far red are the spectral regions, which are preponderantly active in light regulation of flowering, and their effect depends on the timing within the daily light-dark cycle. At least in part the effects of red and far red are phytochrome-mediated: The effect of short far-red irradiation given at the end of the daily light phase (short day), which is flower-inducing in L. gibba G1 and flower-inhibiting in L. aequinoctialis, can be reverted by a following short red light (KANDELER 1962, PURVES 1961, SAIJ et al. 1982). Also the effect of a short night break with red light during long night, which induces flowering in L. gibba and suppresses flowering in L. aequinoctialis, is cancelled by subsequent short far red under certain experimental conditions (KATO 1979a, HILLMAN 1966, ISHIGURI and ODA 1974). The distinctness of these phytochrome effects varies in dependence of strain properties and environmental conditions. The end-of-day far-red effect is lacking in strain G3 of L. gibba and in blue-light cultivated L. aequinoctialis 6746 (KANDELER 1968, ISHIGURI and ODA 1976). Further modifying factors of the end-of-day far-red effect in strain 6746 are light intensity and duration of the main light period as well as the copper content of the nutrient medium (SCHUSTER 1968). Night break effectivity is demonstrable in strain G3 only under threshold conditions for flowering (CLELAND and BRIGGS 1968, KANDELER 1968, KATO 1979a, NAKASHIMA 1968).

If far red is applied not at the beginning but during the course of a long dark period, then the effectivity of far-red irradiation decreases gradually with time in L. gibba G1 (KANDELER 1956). Therefore, an hour-glass mechanism may be responsible for the change in sensitivity to this phytochrome effect. Night break effectivity of red light, however, de-

Table 2.26. Effects of blue, red, and far-red irradiation on flowering in *Lemna gibba* (LDP) and *Lemna aequinoctialis* (SDP)

+ promotion
- inhibition
0 no effect in comparison to control
in the same paper (compare reference number)

	Main light period				End-of-day treatment		Night break treatment
	white	white with far red	blue	red	far red	red after far red	
<i>L. gibba</i> G1							
control (LD)							
9, 11							
control (SD)							
9, 10, 11							
<i>L. gibba</i> G3							
control (LD)							
11							
control (SD)							
11, 12							
<i>L. aequinoctialis</i> 6746							
control (SD)							
15, 17							
control (LD)							
13, 17							
Low Int. +							
<i>L. aequinoctialis</i> 335, 421, 391							
control (LD)							
18							
Low Int. : 0							
<i>L. aequinoctialis</i> T-101							
control (SD)							
14							
control (LD)							
14							
<i>L. aequinoctialis</i> 441							
control (SD)							
16							

References:

- 1 CLELAND and BRIGGS 1968
- 2 ESASHI and ODA 1966
- 3 HILLMAN 1959a
- 4 HILLMAN 1965
- 5 HILLMAN 1966
- 6 HILLMAN 1967
- 7 ISHIGURI and ODA 1976
- 8 ISHIGURI et al. 1975
- 9 KANDELER 1956
- 10 KANDELER 1962
- 11 KANDELER 1968
- 12 KATO 1979a
- 13 ODA 1962
- 14 OHTANI and ISHIGURI 1979
- 15 PURVES 1961
- 16 SAJI et al. 1982
- 17 TAKIMOTO 1973
- 18 YUKAWA and TAKIMOTO 1976

pend on a circadian clock. A brief interruption of a long dark period with red irradiation has a maximal flower-inhibiting effect in *L. aequinoctialis* 6746, when applied 7 or 9 hours after the beginning of darkness (HILLMAN 1959a, PURVES 1961). In *L. gibba* G3 a 30 min light interruption, given at various times during a 51 h dark period after a 9 h short day, shows two maxima of effectivity: 12 and 36 hours after the beginning of a dark period (NAKASHIMA 1968). The significance of endogenous rhythms for light regulation of flowering, which is demonstrated by such results, is discussed further in chapter 2.5.9.

Spectral composition of radiation during the main light period also has a strong influence on flower initiation. Supplementation of white light with far red enhances long-day induction of flowering in *L. gibba* (KANDELER 1956, 1968). Far red or red as the sole source of light during long day, however, cannot induce flowering in strain G3 of *L. gibba*. Only blue light (ESASHI and ODA 1966) or violet and green (ISHIGURI et al. 1975) have similar effects as white under long day in this plant. When far red and red are applied together during continuous light, a ratio of one to one is optimal for flowering in strain G3. With mixed blue and red light the optimal ratio for flowering amounts to 4.9 (ESASHI and ODA 1966). All these results can be interpreted by the assumption that an intermediate level of active phytochrome is needed during the main light period for optimal flowering in *L. gibba*.

In the SDP *L. aequinoctialis* 6746 long-day flowering can be achieved when blue or far red is used instead of white or red (HILLMAN 1965, ISHIGURI et al. 1975, ODA 1962). Under continuous light with two combined spectral regions optimal flowering occurs at values far red/red = 14.8 and blue/red = 13.5 respectively (ESASHI and ODA 1966). Therefore, a relatively low level of active phytochrome seems to allow long-day flowering in strain 6746. Another possibility, to achieve long-day flowering in strain 6746 is reduction of white or red light intensity (TAKIMOTO 1973). Both cited effects, long-day flowering by use of blue or far-red irradiation or by lowering of light intensity, are restricted to strain 6746. Other strains of *L. aequinoctialis*, 335, 391, 421, and T-101, cannot be brought to flower under long day with these special light programmes (OHTANI and ISHIGURI 1979, YUKAWA and TAKIMOTO 1976). In strain T-101 short-day flowering is even suppressed when blue or far-red irradiation is used instead of white or red (OHTANI and ISHIGURI 1979).

With a special light program OOTA (1977a) tested the effect of acetylcholine in strain G3. He found that this substance eliminates the effect of photoperiodic "L1-phase", which works as the dawn signal for entrainment of the circadian clock (see chapter 2.5.9.). Other membrane effectors, the K⁺-ionophores valinomycin and gramicidin, can substitute for the light requirement of "L1-phase" (OOTA 1977b). A further argument that effectors of membrane transport exert their effects on flowering through an influence on the circadian rhythm, comes from experiments with lithium chloride. On the one hand, lithium acts like acetylcholine on flowering in L. gibba and L. aequinoctialis (KANDELER 1970). On the other hand, this ion lengthens the period of the free-running K⁺-uptake rhythm in L. gibba G3 (KONDO and TSUDZUKI 1980c). Lowering of calcium or phosphate concentration in the nutrient medium allows or enhances long-day flowering in L. aequinoctialis 6746 (HILLMAN 1962), but is flower-inhibiting under short day in combination with sucrose and ammonium (HILLMAN and POSNER 1971, POSNER 1969a). In the same plant material calcium and phosphate ions inhibit the leakage of one or more flower-promoting substances, when plants are transferred to distilled water for 4 hours during long night (HALABAN and HILLMAN 1970a, 1971). Thus, calcium and phosphate may influence flowering by changing the conditions for membrane transport. In L. gibba G3, lowering of calcium or phosphate concentration diminishes the flower-promoting effect of suboptimal salicylic acid concentrations (MAENG and BAE 1984). DMSO, rendering the cell membrane more permeable to small molecules, has a similar effect as low Ca²⁺ or phosphate in NH₄⁺-free 1/2 Hutner medium.

2.4.3.1.8. Some further factors

Ferricyanide has a flower-promoting effect in both L. gibba G3 and L. aequinoctialis 6746. In L. gibba some short-day flowering (TANAKA and CLELAND 1980, WATANABE and TAKIMOTO 1977) and in L. aequinoctialis flowering under long day (TAKIMOTO and TANAKA 1973, 1976, TANAKA and TAKIMOTO 1978, TAKIMOTO et al. 1979b) is made possible. Ferrocyanide and cyanide have the same effect as ferricyanide in strains G3 and 6746 (TAKIMOTO and TANAKA 1976, TANAKA and CLELAND 1980, TANAKA and TAKIMOTO 1978, TANAKA et al. 1983), indicating that released cyanide may be the active component within the plant. In L. gibba ferricyanide and salicylic acid show a synergistic interaction. Applied together, they induce some flow-

2.4.3.1.6. Effects of temperature

Flower induction proceeds in L. gibba G3 with optimal rate, if culture temperatures between 26 and 31°C are used (OOTA 1973). Especially night temperature seems to be critical. Lowering of the temperature to 21°C has a flower-inhibiting effect only when applied during the second half of the daily light-dark cycle. Under continuous light, the same diurnal change in sensitivity to temperature proceeds, if acetylcholine, eserine, sodium lauryl sulfate, ouabain, or indoleacetic acid is added to the culture medium (HOSHINO 1979, OOTA 1973, OOTA and HOSHINO 1974). In comparison to L. gibba L. aequinoctialis needs lower night temperatures for flowering. A night temperature of 23°C instead of 25°C can compensate the flower-inhibiting effect of sub-critical dark periods in many strains of this species (BEPPE and TAKIMOTO 1981b). In strain 6746 of L. aequinoctialis flower initiation by inductive skeleton photoperiods is optimal at 23°C. Temperatures above and below 23°C markedly diminish flowering (DOSS 1975). Under conditions of full short day a 4 hour period of 33°C (instead of 28°C) is flower-inhibiting when given during the dark period but not during the light period (HILLMAN 1959b). Long-day flowering in a copper-containing medium can be achieved in strain 6746 with a culture temperature of 27°C but not 29.5°C (HILLMAN 1959b). In W. microscopica, a quantitative SDP, a temperature of 22°C has to be used for optimal flower induction (RIMON and GALUN 1968b).

2.4.3.1.7. Effectors of membrane transport

Feeding experiments with axenically cultivated Lemnaceae have revealed many further substances, which influence flowering in a specific way (this means without, or only relatively small, simultaneous effects of vegetative multiplication). Most interesting are some effectors of membrane transport (tab. 2.25E). Acetylcholine, a transmitter substance which depolarizes the postsynaptic membrane in many animal nerves, suppresses flowering in L. gibba G1, but promotes long-day flowering in L. aequinoctialis 6746 (KANDELER 1972). The acetylcholine action seems to be not simply a drug effect, because eserine, an inhibitor of acetylcholine esterase, acts like acetylcholine, presumably by increasing the endogenously produced level of acetylcholine. The occurrence of acetylcholine in L. gibba G3 has been demonstrated by HOSHINO and OOTA (1978).

ering even under light-dark cycles with only 3 hours light (TANAKA and CLELAND 1980).

Nicotinic acid induces flowering in L. gibba G3 and L. aequinoctialis 151 and 381 when they are grown in one tenth-strength M medium under continuous light (FUJIOKA et al. 1986b). The effect is modified by supply of hormones. GA₃ and ABA are inhibiting in both species, GA₃ being more active in L. aequinoctialis and ABA being more active in L. gibba. Zeatin increases the effect of nicotinic acid in L. aequinoctialis, but decreases this effect in L. gibba. IAA has a relatively small inhibiting effect in both species. Preliminary studies indicate that nicotinic acid is present in Lemna in significantly higher levels than in benzoic acid (FUJIOKA et al. 1986a). Little or no difference in the endogenous levels of nicotinic acid between vegetative and flowering Lemna has been found. Another flower-promoting agent in L. gibba and L. aequinoctialis is cyclic adenosine-3'-5'-monophosphate (cyclic AMP) a "second messenger" in animal cells. Addition of cyclic AMP to the nutrient medium enhances flowering in L. gibba G3 under a variety of experimental conditions (KESSLER and STEINBERG 1973, OOTA 1972a, 1977a, OOTA and KONDO 1974). In L. aequinoctialis 6746 cyclic AMP acts like other adenine derivatives in reversing the sucrose inhibition of flowering under short day (POSNER 1973a). Endogenous cyclic AMP may be involved in flower initiation, because adenyl cyclase-activating catecholamines such as isoproterenol and epinephrine have a similar effect as exogenous cyclic AMP in strains G3 and 6746 (IVES and POSNER 1982, OOTA 1974, 1977a). In L. aequinoctialis, strains 151 and 6746, vitamin K₅ is a further flower-inducing agent, when plants are held in one tenth-strength M medium under continuous light (KAHARA and TAKIMOTO 1985a). The substance acts synergistically with benzyladenine (strain 151), benzoic acid, ferri cyanide (both strains), copper, and high-intensity light (strain 6746).

Inhibition of flowering has been found in L. gibba G3 by some antimetabolites of nucleic acid synthesis. At certain concentrations 5-fluorodeoxyuridine (blocker of DNA replication), 5-fluorouracil (blocker of DNA replication and transcription), and 2-thiouracil (blocker of DNA transcription) reduce flowering percentage without effect on frond multiplication (UMEMURA and OOTA 1965a).

Short-day flowering in L. aequinoctialis 6746 is inhibited to some extent by 5-fluorouracil and 2-thiouracil as well as by some antimetabo-

lites of protein synthesis (cycloheximide, ethionine, and chloramphenicol), when applied for only 2 hours during the first four inductive dark periods (DOSS 1975c). The strong flower-inhibiting effect of a short night break, however, is diminished by a two hour treatment with 2-thiouracil or actinomycin D (DOSS 1975b).

Concluding this chapter, a flower-inhibiting effect of agar may be mentioned. In L. gibba G3 various agars and even the most highly purified are very inhibitory to flowering (CLELAND 1985). The effect of 0.5% agar can be abolished by supply of 3.2 µM salicylic acid.

2.4.3.2. Flower and fruit development

In many plants photoperiodic requirements for flower induction and further development of flower primordia are different, but Lemnaceae belong to a group of plants, which need the same daylength for initiation as well as further growth of the flower (compare type 3 in: VINCE-PRUE, Photoperiodism in Plants, McGraw-Hill, 1975). In the LDP L. gibba G3 at least 6 long days are required for reaching mature flowers, whereas a minimum of two long days is sufficient to induce the formation of flower primordia (CLELAND and BRIGGS 1967). When plants of strain G1 were transferred from continuous light to short day, then in the generation, which is the last but one at the beginning of short-day treatment, shrunken primordia of stage II (stamens and pistil fully meristematic) dominate. In the following generation only dried stage I primordia (without organ differentiation) or no flower residues can be seen (KANDELER, unpublished results). Obviously, flower primordia dye and shrink immediately after transfer of plants to non-inductive photoperiodic conditions. This holds true also in SDPs L. aequinoctialis and L. perpusilla. A regression or abortion of flower primordia after transfer to long day or night-break treatment has been observed in these species by several authors (BENNINK and DE VRIES 1975, DOSS 1975a, HILLMAN 1959a, 1969c, KIRKLAND and POSNER 1974, SATO and ODA 1977, SHIBATA and TAKIMOTO 1975). The inhibition of flower differentiation in SDPs by long day can be overcome by some treatments: low temperature SATO and ODA 1978), low light intensity (HILLMAN 1969c, KIRKLAND and POSNER 1974), blue light, DCMU (KIRKLAND and POSNER 1974), or benzyladenine (BENNINK and DE VRIES 1975). In L. aequinoctialis 6746 at each temperature there is a certain threshold level of light intensity, which is without any inhib-

iting or promoting activity for flower-bud development (SHIBATA and TAKIMOTO 1975).

The statement that flower initiation and further flower development are influenced in the same manner seems to be valid not only for the effect of day-length. This can be drawn from the general rule that conditions, which are flower-inducing to only a small extent (s. low flowering percentage), simultaneously lead to an increase in dried up flower primordia. In L. gibba G1 and G3 this has been found time and again in experiments, where flowering was modified by very different agents (for example far red irradiation, medium composition, CO₂, sucrose, ATP and ADP, DCMU, arsenate, and atebirin) (BUCHELE and KANDELER, unpublished results). In L. aequinoctialis 6746 long-day flowering has been induced only to a certain degree by supply of CCC plus ABA (KANDELER and HUEGEL 1973) and also in this case appearance of aborted flower primordia was enhanced.

The equilibrated development of male and female flower organs depends on a certain balance of gibberellins and ethylene. Hormones are produced by the flower primordium only in part and must be completed by hormonal imports from the leaf. When flower primordia of L. gibba G1 are explanted to an agar medium which contains minerals, sucrose, and kinetin, then the development of the pistil preponderates. The two stamens remain relatively small. A normalization of in vitro flower development, due to a promotion of stamen growth, is achieved by addition of gibberellin A₃ to the agar medium, whereas CCC (blocker of gibberellin synthesis), ethephon (ethylene-releasing agent), or IAA enhance feminization by a further promotion of pistil growth (HUEGEL 1976a, 1976b). Explants of L. aequinoctialis 6746 cultured on the above cited minimal medium preferably develop the stamens. This masculinization can be corrected by supply of CCC or ethephon which allow better pistil growth also in strain 6746 (HUEGEL 1976a,b). Low concentrations of kinetin as well as ABA enhance the one-sidedness of in vitro sex expression in both species, presumably by a more general promoting resp. inhibiting effect on organ development (HUEGEL 1976c). The opposite behaviour of L. gibba and L. aequinoctialis may be caused by a different endogenous production or accumulation capacity of gibberellins (compare the statements in connection with photoperiodic flower induction, chapter 2.4.3.1.1), but ethylene metabolism may also be different in the two species. The tendency to feminization of L. gibba has been observed in several strains also in

sterile cultures, i.e. in vivo (LANDOLT, unpublished results). DNA duplication in connection with microsporogenesis has been investigated in W. microscopica using autoradiographic methods (RIMON and GALUN 1968b). Unexpectedly, optimal incorporation of tritiated thymidine into the nuclei of future pollen cells was obtained when labelling was done at a stage at which the whole floral bud was not yet anatomically recognizable. The authors conclude that thymidine-³H incorporation into the nuclei of microspores occurs in all probability during one of the premeiotic mitotic interphases.

The physiology of pollination and fruit development has not yet been investigated in Lemnaceae. HUEGEL (1974) mentioned a special development of pistil and parthenocarpic development of seeds in L. gibba, when advanced flower primordia were explanted on to an agar medium with 16% sucrose. POSNER (1962b) induced some aberrant strains of L. aequinoctialis 6746 by X-irradiation which produced very few seeds and these always lacked embryos. These strains may be a good material for further investigations.

cit. in HUBER and SANKHLA 1979). For example, 10^{-5} M ABA added to the medium for 24 h has no effect on the degree of opening in *L. gibba* G1 (HARTUNG, personal comm.). In *L. minor* CMU, FCCP, valinomycin, and nigericin are not effective as in *S. punctata* (fig. 2.35). Nevertheless, young stomata of *L. minor*, which are still viable and occur especially in the basal part of fronds, exhibit a weak movableness (HABERLANDT 1987, REUTER 1948).

As in other plants light intensities considerably above those experienced during growth inhibit photosynthesis reversibly. This so-called photoinhibition has been found in *L. trisulca* (ZURZYCKI 1957a), *L. minor* (LINDEMAN 1979), and *L. gibba* (NILSEN and DANIELSEN 1984, OEGREN et al.




Fig. 2.35. Movability of stomata in Lemnaceae (SEVERI and BARONI FORNASTIERO 1983b). In *Spirodela punctata* a 6 hour treatment with CMU, FCCP, valinomycin, or nigericin results in a strong closing of stomata (b) in comparison to untreated controls (a). In *Lemna minor*, on the other hand, treated fronds (d) and control fronds (c) show about the same degree of aperture (SEM pictures; bar = 10 μ m). (With permission of Dr. A. Severi)

2.5. METABOLISM

2.5.1. Generation of energy

2.5.1.1. Photosynthesis

2.5.1.1.1. Factors limiting or restricting photosynthesis

In older work photosynthesis has been investigated by measuring dry-weight increase or growth rate (see chapter 2.3.5.1.1.). ZURZYCKI (1953, 1955a,c) used a microtechnique based on a capillary tube respirometer for photosynthesis measurements in *L. trisulca*. He showed that during the first 15-30 minutes of low-intensity illumination photosynthesis depends on the rearrangement of chloroplasts. O_2 production increases with the displacement of chloroplasts from a random position (apostrophe in darkness) to a flat position (epistrophe in weak light). Under high light intensities gas exchange proceeds with a constant rate independent of simultaneous changes in chloroplast arrangement (transition to profile position, parastrophe in strong light). Epistrophe lasts between 10 and 800 lux, full parastrophe is reached approximately in those light intensities (12000 lux) in which a further increase of intensity has no influence on photosynthesis (for analysis of chloroplast movements see chapter 2.4.1.6).

A farther reaching adaptation to different light intensities seems to exist on the basis of developmental processes. Plants of *S. polyrrhiza* precultivated under 1400 lux show a saturation of photosynthesis at about 4400 lux. Exposure to strong light (10000-50000 lux for 12 h) leads to a substantial increase (to 60-85%) in the rate of photosynthesis, which is not eliminated by a subsequent exposure to 1400 lux (GAPO-NENKO and STAZHETSKII 1969).

A further limiting factor for photosynthesis, effective in floating plants only, is the behavior of stomata on the upper side of the fronds. In *S. punctata* stomata close during water deficit and - to a various extent - after application of ABA, metabolic inhibitors (CMU, FCCP), or ionophores (valinomycin, nigericin), but not after darkening (SEVERI and BARONI FORNASTIERO 1982, 1983b). In *Lemna*, however, stomata are not movable and remain open under all conditions (HABERLANDT 1987, WAGNER 1973,

1984, OEGREN and OEQUIST 1984a, b). Factors which restrict photosynthesis such as chilling (2-3°C, LINDEMAN 1979, OEGREN et al. 1984), or N₂ atmosphere (NILSEN and DANIELSEN 1984) lower the light intensity necessary for the inhibitory effect of bright light. The papers dealing with *L. gibba* come to the conclusion that inactivation of photosystem II-mediated electron transport occurs during photoinhibition. Polarization of light (both linear and circular) has no effect on photosynthesis in *L. trisulca* (ZURZYCKI 1955b).

In *S. punctata* and *L. minor* maximum rates of O₂ production and ¹⁴CO₂ uptake are reached at 35°C (WEDGE and BURRIS 1979). The range of favorable temperatures is broader in *L. minor* than in *S. punctata*, but at 10°C and 45°C there is no O₂ evolution in either species. At 35°C *S. punctata* evolves 1.6 µl O₂/h/mg fresh wt. (*L. minor*: 1.1 µl O₂/h/mg fresh wt.): 12-hour experiments with *L. minor* grown in continuous light showed that a diurnal fluctuation in photosynthesis does exist, with a peak in photosynthesis appearing at midday.

Availability of CO₂ is another factor which determines the rate of photosynthesis in many cases. The great differences in accumulation of photosynthetic starch appearing in *L. gibba* after 3 h cultivation in open dishes, open or cotton-plug closed vessels, may be caused mainly by differences in CO₂ supply (KOENIGSHOFER, personal comm.). When the CO₂ concentration of the atmosphere is raised from 300 to 1000 ppm, under high-intensity light conditions (17000 lux, equivalent to 240 µE m⁻² s⁻¹ PAR; temperature 27±1°C) the photosynthetic rate increases from about 107 to 165 mg CO₂/h/g dry wt. in *S. punctata*, from 160 to 250 mg CO₂/h/g dry wt. in *L. minor*, and from 135 to 300 mg CO₂/h/g dry wt. in *L. valdiviana* (LOATS et al. 1981). Correspondingly FUHRER and ERISMANN (1983) found a highly significant correlation between CO₂ input (100-700 ppm) and dry weight increase in *L. minor*. In *L. gibba* the net assimilation rate increases with CO₂ enrichment of the air till 1500 ppm CO₂ (ANDERSEN et al. 1985, BJØERNDAL and NILSEN 1985). A dramatic deterioration of CO₂ supply for photosynthesis results from vacuum infiltration of fronds. O₂ evolution in *L. minor* (211 µmoles/h/g/ fresh wt. under 18000 lux of incandescent light, at 25°C) is eliminated completely by vacuum infiltration, whereas O₂ absorption in the dark is not lowered by such treatment (McDONALD 1975). FILBIN and HOUGH (1985) concluded from in situ investigations that in field populations of *L. minor* an average of 86% of fixed carbon was from aqueous inorganic carbon. ESHEL and BEER (1986),

however, demonstrated in laboratory experiments with *S. polyrrhiza* that carbon assimilation from the aqueous phase amounts only up to 5% of that from the air.

Restriction of photosynthesis can be achieved by many other factors: limitation of nitrogen supply (INGEMARSSON et al. 1984), inactivation of glutamine synthetase (JOHANSSON and LARSSON 1986a, b), phosphate deficiency (LINDEMAN 1951), feeding with sodium acetate (BRUNOLD and ERISMANN 1968), salt stress with sodium chloride (HUBER and SANKHLA 1979), supply of ABA (BAUER et al. 1976, McLAREN and SMITH 1976, TILLBERG et al. 1981), fumigation with SO₂ (LOATS et al. 1981), "synthetic smog", that is a mixture of ozone and hexene (ERICKSON and WEDDING 1956), application of certain herbicides (BIELECKI and SKRABKA 1976), and preculture with the morphactin "chloroflurenol" (TREICHEL 1974b).

The effect of reduced nitrogen supply on photosynthesis in *L. gibba* is apparent only when calculated on a dry weight basis, but not on a chlorophyll basis. Thus reduced nitrogen works in a rather indirect way (presumably restricting the differentiation of chloroplasts). The light compensation point was unaffected by the nitrogen regime (INGEMARSSON et al. 1984). Nevertheless, blocking of glutamine synthetase by MSO (L-methionine-D,L-sulphoximine) decreases CO₂ fixation in *L. gibba* to 50% within 100-150 min (JOHANSSON and LARSSON 1986a, b). The inhibiting effect of P-deficiency on photosynthesis in *L. minor* can be abolished within a few hours by supply of phosphate. At rate limiting light intensities reactivation is greater at 650 nm than at 700, 711, and 717 nm (LINDEMAN 1951, 1972). Sodium acetate ($\approx 10^{-3}$ M) taken up from the medium may consume photosynthetic ATP and NADPH for assimilation and therefore compete with CO₂ fixation. In addition, fatty acids are produced from assimilated acetate and may inhibit some photosynthetic reactions as ATP synthesis, carboxylation of RuDP, and dephosphorylation of FDP (BRUNOLD and ERISMANN 1968).

High concentrations of NaCl (85 mM) slow down the photosynthetic electron transport (HILL reaction) in *L. minor*, but have no effect on the pattern of ¹⁴CO₂ fixation (HUBER and SANKHLA 1979). Exogenous ABA acts in a similar way (BAUER et al. 1976, McLAREN and SMITH 1977). As NaCl increases endogenous ABA level up to 1.5 times and exogenous ABA lead to an endogenous ABA concentration about twice as high as in the control plants, salt stress may act on photosynthesis through the "stress hormone" ABA (HUBER and SANKHLA 1979, SANKHLA and HUBER 1979). Preliminary

experiments of McLAREN (cited in McLAREN and SMITH 1977) indicate that ABA decreases the chloroplast membrane permeability to 3-phosphoglyceric acid. This could be the explanation for the fact that starch is accumulated in chloroplasts of ABA-treated plants (McLAREN and SMITH 1976). Fumigation with 0.75 ppm SO₂ during preculture for 8 h each day leads to an inhibition of net photosynthesis in S. punctata and L. valdiviana, but not in L. minor (LOATS et al. 1981). L. minor seems to oxidize SO₂ to SO₄²⁻ (BRUNOLD and ERISMANN 1976) more efficiently than the other species. The inhibition of photosynthesis after fumigation with 1 ppm ozone can be explained by chlorophyll degradation. In the case of an ozone-hexene mixture, however, reduction of photosynthesis exceeds that of chlorophyll loss (ERICKSON and WEDDING 1956). Herbicides which inhibit photosynthesis are Prometryne, Ametryne, Diuron (DCMU), Fluometuron (CEDENO-MALDONADO and LIU 1976), and Aflon (50% Linuron), Gramoxone (20% paraquat), Simazine and Krezamon (50% DNOC) (BIELECKI and SKRABKA 1976). The morphactin, 2-chloro-9-hydroxyfluorene-9-carboxylate-(9) (chl-roflurenol) influences photosynthesis and related processes only if the plants are treated with the substance for at least 9 days. Morphogenetic changes seem to be involved during this time. Photosynthetic O₂ production, CO₂ consumption, and ¹⁴C incorporation all are reduced after prolonged morphactin treatment, whereas starch, ATP, and chlorophyll content increase (TREICHEL 1974b).

2.5.1.1.2. Steps of the photosynthetic process

In several cases Lemnaceae have been used to investigate certain steps of the photosynthetic process: action of the two photosystems, the chain of electron transport, generation of reduction energy, production of ATP, assimilation of CO₂, and light respiration.

The presence of the Emerson enhancement effect in L. minor and L. gibba shows that photosystems I and II work as in other plants (LINDEMAN 1973, CHATURVEDI et al. 1982). Capacity for photosynthetic O₂ evolution depends on the mode of preillumination. Lemna illuminated for 5 to 7 days every 15 min by a 1 ms strong polychromatic flash has no capacity for O₂ evolution, and stroma thylakoids of these plants are unfused. Three to 6 minutes of illumination with photosynthetic actinic light are enough to induce photosystem II-activity and a fused state of stroma thylakoids (STRASSER 1974). The author hypothesizes that a manganese reducing sys-

tem may be created during the induction period.

The DCMU-sensitive step in photosynthetic electron transport has been investigated with S. punctata (WARTOO et al. 1981, 1984a). DCMU seems to bind reversibly to a 32 kDa polypeptide, which is the main thylakoid protein. Plants depleted of this protein by incubation with low levels of chloramphenicol (WEINBAUM et al. 1979a) show a selective inhibition of e⁻ transport at the reducing side of photosystem II, especially between the primary quencher and plastoquinone pool. The 32 kDa protein may be a "proteinaceous shield" covering the primary electron acceptor of photosystem II and acting as a regulator of electron flow between photosystem II and photosystem I. The protein is not rate limiting for CO₂ fixation. Photoassimilation of CO₂ remains at 86% of controls in chloramphenicol-treated plants containing less than 20% of 32 kDa protein (WEINBAUM et al. 1979a). Biosynthesis, turnover, and light induction of the 32 kDa protein have been investigated extensively (chapter 2.5.7.3.2). The binding site of triazine herbicides in photosystem II seems not to be identical with the 32 kDa protein but located in close connection (GRSSEL 1982). In L. minor 1.0 ppm atrazine reduces photosynthetic O₂ evolution to 50%. Lower concentrations (0.02-0.10 ppm) however, distinctly stimulate photosynthesis (BEAUMONT et al. 1976). A block between plastoquinone and cytochrome f in the photosynthetic electron transport system has been found in a mutant strain 1073, which was isolated from an x-irradiated culture of wild type strain 6746 of L. aequinoctialis (SHAHAK et al. 1976). Isolated chloroplasts of the mutant are able to reduce diaminodurene (DAD) or phenylenediamine (lipophilic electron acceptors at the level of plastoquinone) but not its own cytochrome f. Electron paramagnetic resonance analysis of mutant chloroplasts reveals a lack of the "Rieske" iron-sulfur center signal (WALKIN and POSNER 1978). This Rieske center may be either absent or defective in the mutant.

Moderately high Hill activity of crude chloroplasts was demonstrated in L. minor and L. borealis as early as 1950 (CLENENNING and GORHAM 1950). DYAR (1953) then used blue tetrazolium as an in vivo indicator of the Hill reaction and showed photochemical reduction of the tetrazolium salt to be located especially in grana-like regions of chloroplasts in fronds and roots of L. minor. Manganese requirement during culture of L. minor for optimum Hill reaction of chloroplasts has been found to be 5x10⁻² ppm (EYSTER et al. 1958). Hill activity of isolated chloroplasts

from *W. arhiza* was determined by EICHORN and AUGSTEN (1983a). Plants were cultivated in chemostat steady state populations under continuous white, blue, or red light. Deviations in Hill activity caused by blue or red light preculture do not exceed 10-15% of white light control. Supply of boron (0.5-1 ppm) to the nutrient medium, however, reduces Hill activity to a greater extent (to about 50% with material grown under red light and high growth rate conditions). Isolated chloroplasts from *L. gibba* have been used for measurement of Hill activity in connection with photoinhibition of photosynthesis (see chapter 2.5.1.1.1). Pretreatment of plant material with high photosynthetic photon flux density reduces the capacity of whole chain electron transport, as well as efficiency of electron transport to the primary electron acceptor Q of photosystem II, measured as variable chlorophyll fluorescence at 20°C (OEGREN and OSQUIST 1984a). ABA supply during preculture of *L. minor* has an inhibiting effect on the Hill reaction of isolated chloroplasts (BAUER et al. 1976). A method to prepare isolated chloroplasts from *S. polyrrhiza* is given by BAHL (1971).

A method for determination of minute amounts of reduced and oxidized NADPH and NAD from *S. polyrrhiza* was worked out by MONEGER et al. (1977) and VERMEERSCH et al. (1977b). Oxidized forms of pyridine nucleotides are largely predominant and NADPH represents a high proportion of reduced forms (plants cultivated with or without addition of sucrose). Sucrose feeding provokes a decrease in content of all pyridine nucleotides (measurements from the whole tissue as well as from isolated chloroplasts (LECHEVALLIER et al. 1977, VERMEERSCH et al. 1977a). Plants cultivated in a medium enriched with calcium (20-40 mg/l calcium instead of 4) have an increased total pyridine nucleotide content and a decreased NADPH/NADP ratio (LECHEVALLIER 1977a).

A small part of photosynthetic reduction energy is used for driving the xanthophyll cycle. Within the chloroplast violaxanthin is reduced to antheraxanthin and zeaxanthin, and these substances are re-transformed to violaxanthin by oxygen (fig. 2.36). In *L. gibba* G3 zeaxanthin is accumulated only during the first 10-20 minutes of illumination (transition phase) and - under steady state conditions - in the case of high light intensities (> 10000 lux) (STIEFERMANN 1971, 1972). Under the conditions used in these experiments CO₂ consumption is saturated at 35000 lux.

Energy charge of the adenylate system, ATP + 1/2 ADP/ATP+ADP+AMP, is not affected by light intensity (500-4400 lux) and remains nearly constant throughout a 1 day period when plants of *L. gibba* G3 in mixotrophic culture are exposed to short day, continuous light, or continuous darkness. Depletion of sucrose from the medium, at least under continuous light and continuous darkness, causes no change in energy charge (KONDO and NAKASHIMA 1979). Correspondingly, EICHORN and AUGSTEN (1977a) found only slight changes of energy charge after supply of diverse carbohydrates




Fig. 2.36. Scheme of the xanthophyll cycle (after STIEFERMANN 1972). E^{1,2} are pH-dependent enzymes of deepoxidation; E^{3,4} are enzymes of epoxidation.

to *W. arrhiza* cultures under light and dark conditions. In *L. aequinoctialis* pool sizes of ATP were determined by GOWER and POSNER (1979). Wild type fronds (strain 6746) show a higher ATP level than the photosynthetic mutant (strain 1073) under light and dark conditions. DCMU (1 μ M, not effective in the mutant) lowers rather than raises the ATP level in wild type plants during prolonged exposure to light. Supply of NH_4NO_3 instead of KNO_3 to tenth-strength Hutner's medium (containing 1% sucrose) increases the ATP pool in both strains. In steady-state populations of *W. arrhiza* adenosinephosphate pools and energy charge are higher in blue-light cultivated plants, as compared with plants cultivated in red light (EICHORN and AUGSTEN 1977b). Protein and chlorophyll content, however, are also elevated under blue light.

Photosynthetic CO_2 assimilation proceeds through the reactions typical for C_3 plants as indicated by the lack of Kranz leaf anatomy (RAGHAVENDRA and DAS 1976), the pattern of $^{14}\text{CO}_2$ fixation (BAUER et al. 1976) and a distinct photorespiration (ERISMANN 1972). In *L. minor* BAUER et al. (1976) found labelling to be highest in 3-phosphoglyceric acid and sulphate-esters after a $^{14}\text{CO}_2$ pulse of 10 sec. Relatively fast labelling occurs also in some amino acids: alanine, aspartate, serine, and glycine (BAUER et al. 1976, ERISMANN and KIRK 1967). Supply of ammonium instead of nitrate changes the steady-state levels of photosynthetically produced sugars and amino acids. The formation of sucrose is favoured at the cost of glucose and fructose. Ammonium increases aspartic acid and glutamic acid (ERISMANN and KIRK 1969). Carbonic anhydrase, catalyzing the reversible hydration of CO_2 , shows a very low activity in *L. trisulca*, but an intermediate activity in *L. minor* and *W. columbiana*. The results correspond with those of other submersed or floating macrophytes (WEAVER and WETZEL 1980).

Much work has been done by Ziegler and co-workers to investigate the light-induced increase in the activity of NADP-dependent glyceraldehyde-3-phosphate dehydrogenase (NADP-GPD) in *L. gibba*. Activation of the enzyme by light proceeds within some minutes and needs high light intensities (11-37 $\text{mW}\cdot\text{cm}^{-2}$) for an optimal effect. Simultaneously, activity of the NAD-dependent GPD decreases (ZIEGLER et al. 1968). The light induction seems to be mediated by chlorophyll (SCHMIDT-CLAUSEN and ZIEGLER 1969) and depends on an operating non-cyclic electron transport. DCMU and some other metabolic inhibitors block the light effect completely (ZIEGLER and ZIEGLER 1966). Actinomycin C has no influence, but chloram-

phenicol (2.5 mg/ml) suppresses the light activation of NADP-GPD (ZIEGLER and ZIEGLER 1965). On the basis of further kinetic studies for light activation and dark inactivation and of their earlier results the authors proposed a model in which activation of NADP-GPD is accomplished by association of four subunits of the enzyme to the activated tetrameric form. NADPH, created during the non-cyclic electron flow, could be the allosteric effector (SCHMIDT-CLAUSEN et al. 1969). Blue light (450-500 nm) not only has an activating effect on NADP-GPD through chlorophyll, but also an inhibiting effect as shown under optimum light-inducing conditions by elimination of particular bands of wavelengths from the complete spectrum (SCHMIDT-CLAUSEN and ZIEGLER 1969). Blue-light enhanced respiration with a concomitant decrease of photosynthetic O_2 evolution could be the cause for this phenomenon.

Photorespiration was substantiated in *L. minor* by CO_2 outburst phenomena occurring after darkening in cultures held under pure oxygen (fig. 2.37) or in light after short CO_2 fixation and a following change from N_2 to O_2 fumigation (ERISMANN 1972). Also several other criteria suggest the presence of photorespiration in *L. minor* (FILBIN and HOUGH 1985). The ratio α between oxygenation and carboxylation of ribulose-1,5-diphosphate was calculated by FUHRER and ERISMANN (1984) on the basis of CO_2 compensation concentration and the intercellular CO_2 concentration. This ratio α is lowered when plants are transferred from 25 to 15°C. Supply of ammonium instead of nitrate, however, has no influence on α . Under conditions allowing a low photorespiratory $^{14}\text{CO}_2$ release only (2% O_2 fumigation), DCMU increases photorespiration in *L. gibba* significantly (CHATURVEDI et al. 1982). The authors come to the conclusion that electrons from the oxidation of glycolate are donated to photosystem I when the electron supply from water is low. Glycolate oxidase in *L. minor* has a high specific activity (FREDERICK et al. 1973). The enzyme activity is doubled in plants grown with an ammonium nitrogen source as compared with plants grown on nitrate. In addition, the quaternary structure of glycolate oxidase is altered. Ammonium-consuming plants contain a high-molecular-weight form of enzyme (about 500000). After transition to nitrate nutrition molecular weight of glycolate oxidase amounts to about 250000 (EMES and ERISMANN 1982). As expected, NH_4^+ plants evolve more $^{14}\text{CO}_2$ from labeled glycolate than NO_3^- plants (MARQUES and ERISMANN 1984, MARQUES et al. 1985). Formate is labeled only from (2- ^{14}C) glycolate and in NH_4^+ plants it is markedly less labeled in the light than

in the dark. TILLBERG (1980) found a stimulation of growth in *L. gibba* G3 by supply of 0.5 mM glyoxylate or glycolate and interpreted the effect as mediated by an inhibition of photorespiration.




Fig. 2.37. Photorespiration in *Lemna minor* as shown by the course of CO_2 evolution during repeated light-dark changes (20:20 min) under pure oxygen (ERISMAN 1972). Curve sections from every fourth light-dark period are shown. The magnitude of CO_2 outburst at the beginning of the dark phase corresponds to the value of transient CO_2 evolution during the preceding light phase. Weakening of both phenomena during the course of the experiment may be caused by an exhaustion of substrate for photorespiration. (A short supply of glycolate to the plants can restore the CO_2 outburst phenomena.)

2.5.1.2. Respiration

In *L. aequinoctialis* respiration of intact plants has been compared with gas exchange of isolated mitochondria (LOEPPERT 1983). In both cases O_2 consumption shows the same dependence on dissolved oxygen concentration. Half-maximum respiration rate (K_m) for intact plants is reached at $1.15 \pm 0.08 \mu\text{M O}_2$ and for isolated mitochondria at $1.07 \pm 0.06 \mu\text{M O}_2$. From this fact it can be concluded that diffusion of oxygen within the tissue is obviously not rate-limiting for respiration in *Lemna*. At $2.5 \mu\text{M}$ dissolved O_2 (in equilibrium with $1\% \text{ O}_2$ in air) oxygen consumption amounts to about 70% of maximum rate. Plants of *Lemna spec.*, grown heterotrophically and fumigated with argon containing about $1\% \text{ O}_2$, show an adaptation to such conditions. After a lag-period of 10 days, growth rate is recovered up to approximately one-half of the aerobic rate (SIEGEL 1961).

The ATP level of intact *L. aequinoctialis* plants decreases roughly in




Fig. 2.38. ATP content of intact *Lemna aequinoctialis* plants as a function of respiration rate (LOEPPERT 1983). Values of respiration rate and ATP content were determined at various oxygen concentrations and then used for a plot expressing both parameters as percentage of the maximum value.

parallel with the respiration rate (fig. 2.38) when the oxygen concentration is reduced below $10 \mu\text{M O}_2$ (LOEPPERT 1983). Both, the CN^- sensitive and the CN^- insensitive pathway participate in respiration of Lemna, as an inhibition of respiratory gas exchange can be achieved by simultaneous supply of CN^- and SHAM (salicylhydroxamic acid) only (LOEPPERT, personal comm.; ULLRICH-EBERIUS et al. 1983). Cyanide, given alone, has no, or even an enhancing, effect on respiration in L. aequinoctialis (LOEPPERT, personal comm.). In L. gibba CN^- inhibits or stimulates respiration depending on its magnitude. In sucrose-fed plants with high respiration ($7-9 \mu\text{M O}_2 \text{ g}^{-1} \text{FW h}^{-1}$), O_2 uptake rate is inhibited more than 60% by CN^- ; in plants depleted of sucrose for about 7 days and exhibiting low respiration ($2 \mu\text{M O}_2 \text{ g}^{-1} \text{FW h}^{-1}$), O_2 uptake is stimulated by CN^- (ULLRICH-EBERIUS et al. 1983). In L. aequinoctialis the ATP level remains high in the presence of CN^- , presumably on the basis of site-I phosphorylation (LOEPPERT 1981). In sucrose-nourished L. gibba, however, cyanide reduces the ATP level in the dark by more than 50% (ULLRICH-EBERIUS et al. 1983). Corresponding to the stimulating effect of cyanide under some circumstances, DALY and BROWN (1954) found an enhancing effect of carbon monoxide on respiration in Lemna spec. (blocking of cytochrome oxidase). Other agents which lead to an increase in respiration, are mechanical damage (S. punctata: DVORAKOVA-HLADKA, 1964), certain aromatic sulphonates (anionic detergents: "Sulfapol", "Nekal"), and boron deficiency (EICHORN and AUGSTEN 1974). The effect of "Nekal" may be caused - at least partially - by an increase of monophenolase activity which has been found in homogenates from L. minor (GODZIK 1964). Also the effect of B-deficiency has been interpreted in a similar way (see strengthening of salicylaldehyde sensitivity at B deficiency).

In photosynthetically active tissue respiration (CO_2 evolution other than from photorespiration) is greatly lowered by light. Based on the oxygen-dependence of mesophyll resistance and carbon dioxide compensation concentration, FUHRER (1983) calculated the fraction of respiration not inhibited by light in L. minor. At light intensities above $150 \mu\text{E m}^{-2} \text{sec}^{-1}$ "day" respiration amounts to 23% of dark respiration, but increases rapidly when the light intensity is decreased.

A more or less close connection between nitrogen provision, growth, and respiration exists in L. minor (WHITE and TEMPLEMAN 1937; HUMPHREY et al. 1977). When nitrate concentration of the nutrient medium is lowered to $<1 \text{ mM}$, frond multiplication as well as respiration decrease greatly.

A number of enzymes, including aldolase show decreased activity. Further analysis of the effect of nitrogen starvation on aldolase activity reveals that not pyridoxal phosphate (SARAWEX and DAVIES 1977a) but rather a newly produced, specific inhibitor (low molecular weight protein) is responsible for the loss of enzyme activity (SARAWEX and DAVIES 1977b). Another agent which inhibits frond multiplication as well as respiration in Lemna, is the morphactin chloroflurenol (TREICHEL 1974a).

Obviously, respiratory gas exchange comes about not only by metabolism of mitochondria but also by other energy-generating and oxidative processes. Examples are the direct oxidation of glucose through the pentose-phosphate path in chloroplasts and the cytosol, energy-generating processes presumably located at the plasmalemma (compare chapter 2.5.2.1) and the action of phenolases (see above). The key enzyme of the oxidative pentose-phosphate path, glucose-6-phosphate dehydrogenase, was investigated by EICHORN and AUGSTEN (1977a,b). Activity of this enzyme in W. arrhiza is influenced by addition of certain carbohydrates to the nutrient medium. For example, glucose and ribose are stimulating under different growth conditions, whereas galactose, rhamnose, and especially lactose have an inhibiting effect.

A hint to respiratory gas exchange at the plasmalemma comes from the work of ZURZYCKI (1970, 1971). The author studied the effects of light on oxygen uptake in DCMU-treated leaves of L. trisulca. Blue light of very low intensity (saturation at $10 \text{ erg/cm}^2 \text{ sec}$) enhances O_2 uptake by 30%. With high-intensity light of short-wave range, however, the effect is diminished. When linear polarized light is used, the direction of the polarization plane has no effect in Lemna, but - to a different degree - in leaves of Elodea and two mosses. Discussing these results in connection with cell geometry of the investigated plants, ZURZYCKI comes to the conclusion that the photoreceptor for low-intensity blue-light effect on respiration may be dichroic and oriented parallel to the cell surface, that is, located in the plasmalemma. The effect of light on O_2 uptake shows a close correlation with the effect of light on the arrangement of chloroplasts (chapter 2.4.1.6). A blue light effect, which decreases photosynthetic O_2 evolution possibly through an enhancement of respiration, was already mentioned in chapter 2.5.1.1 (see SCHMIDT-CLAUSEN and ZIEGLER 1969).

2.5.2. Membrane transport

2.5.2.1. Membrane potential, proton extrusion, and proton co-transport processes

Electrophysiological methods have been adapted to measure the membrane potential in single subepidermal cells of *Lemna* fronds (LOEPPERT et al. 1978, NOVACKY et al. 1978b) as transmembrane potential changes can indicate ion-transport processes through cell membranes. Following this line much work has been done to substantiate the existence of proton pumps at the plasmalemma which extrude protons to the apparent free space, and in this way make proton co-transport processes possible for uptake of the anions nitrate, phosphate, and sulfate, as well as hexoses and amino acids. Furthermore, the energy provision for proton extrusion, the modulation of membrane potential (E_m) by phytochrome and the effect of ABA on E_m have been investigated.

L. aquinoctialis 6746 and *L. gibba* GL, pre-cultured in an anorganic nutrient medium, show E_m values of -180 to -220 mV (LOEPPERT et al. 1978, LOEPPERT 1979, 1981, KANDELER et al. 1980). *L. gibba* GL, pre-cultured in medium with 29 mM sucrose, exhibits E_m values which are more variable (-200 to -280 mV) (NOVACKY et al. 1978a, b; ULLRICH-EBERIUS et al. 1981, 1983). After elimination of sucrose from the nutrient medium, high E_m values are maintained under continuous darkness (at least for 7 days), but not, when 7 long days are followed by 6 to 72 hours of darkness. E_m then decreases to -90 mV.

Without doubt, the high E_m values cited cannot be caused by passive diffusion of ions only. The diffusion potential, calculated for currently used potassium concentrations (as K^+ is the most important ion in this connection), would amount to -80 to -125 mV. Active transport processes must contribute to the high charge difference between the cell interior (vacuole) and the surrounding medium. In fact, membrane potential can be reduced to diffusion potential when respiration is blocked by concurrent supply of cyanide and salicylhydroxamic acid (ULLRICH-EBERIUS et al. 1983). Evidence for the fact that proton extrusion plays a dominant role in maintaining high membrane potentials in *Lemna* was obtained in two ways. LOEPPERT (1979) manipulated the pH at both sides of the plasmalemma and monitored the accompanying changes of E_m . Reversible depolarisation of E_m occurs after an increase of external or a decrease of inter-

nal proton concentration. These results are in agreement with the postulated E_m -generating proton pump. BOECHER et al. (1980) achieved a hyperpolarization of E_m in *L. gibba* by supply of fusicoccin, which was reported to stimulate a H^+ -pumping ATPase at the plasmalemma. Membrane potential of *L. gibba* depolarizes transiently when certain inorganic or organic nutrients are added to the bathing solution. This holds true for nitrate (NOVACKY et al. 1978a; ULLRICH and NOVACKY 1981), phosphate (ULLRICH-EBERIUS et al. 1981, 1984), sulfate (LASS and ULLRICH-EBERIUS 1984a), glucose and fructose (NOVACKY et al. 1978b), glycine and other neutral or acidic amino acids (NOVACKY et al. 1978a, FISCHER and LUETTGE 1980, JUNG and LUETTGE 1980, 1982a, JUNG et al. 1982). These E_m changes can be interpreted by the action of carriers at the plasmalemma, which facilitate uptake of certain specific substrates especially in the case that protons are co-transported. A transient lowering of the proton gradient at the plasmalemma and thus of the membrane potential should occur when an inward proton co-transport is set in motion by supply of a carrier-accepted substrate. Some lines of evidence can support this view. Firstly, NOVACKY et al. (1980) and ULLRICH-EBERIUS et al. (1980) measured extracellular pH changes on the lower surface of single *L. gibba* fronds with a microelectrode and found the transient E_m depolarization, occurring after supply of glucose, to be correlated with an extracellular alkalization. Then, LUETTGE et al. (1981) pointed out the enhancement of alanine uptake by fusicoccin to be correlated with the extent of fusicoccin-triggered E_m increase. Finally, JUNG et al. (1982) calculated proton electrochemical gradients (from E_m , medium pH and assumed cytoplasmic pH) and showed a linear correlation between this proton gradient and amino acid uptake rates (at amino acid concentration of 10^{-4} M). A selective inhibition of the amino acid co-transport carrier can be achieved by pretreatment of plants with the amino acid analogue p-fluorophenylalanine (JUNG and LUETTGE 1982b). Alanine uptake and alanine-induced transient E_m depolarization are reduced, but 3-O-methylglucose uptake is only slightly, and proton pump-dependent resting potential not affected. A poison which acts on the amino acid carrier as well as the glucose carrier is 10^{-5} M Hg Cl_2 (GOLLE and LUETTGE 1983). At higher concentrations of Hg Cl_2 (5×10^{-5} M) membrane potential and passive Na^+/K^+ -permeabilities are also affected (LOOS and LUETTGE 1984).

Investigating the energy provision of the proton pump ULLRICH-EBERIUS et

al. (1983) and LOEPPERT (1981, 1983) reached different conclusions. ULLRICH-EBERIUS et al. stated a similar reduction of E_m and ATP level during supply of cyanide, the ATP-level decay being more rapid than the E_m decay. In analogy of what is known from other higher and lower plants (compare NOVACKY et al. 1978b), they postulated the proton pump to be an ATPase. LOEPPERT, on the other hand, found a cancellation of the active component of E_m after supply of cyanide or after reduction of oxygen concentration below a certain value, despite a sufficient availability of ATP. Furthermore, DCCD, an inhibitor of membrane-bound ATPase, reduced the ATP level to 11%, but leaves cell potential uninfluenced. The author concluded that ATP cannot be the fuel for the electrogenic proton extrusion. A redox pump may be envisaged as an alternative. Obviously, the different results of LOEPPERT and ULLRICH-EBERIUS et al. are caused not only by the use of different plants (LOEPPERT: *L. aequinoctialis*; ULLRICH-EBERIUS et al.: *L. gibba*), but also by different pre-culture conditions. *L. aequinoctialis* was held in anorganic nutrient medium under long day, whereas *L. gibba* was fed with 1% sucrose and cultured under short day. The differing behaviour of both experimental plants in respiration (see CN⁻ effect: chapter 2.5.1.2) and membrane potential per se (see above) has already been mentioned.




Fig. 2.39. Transmembrane potential changes in single subepidermal cells of *Lemna aequinoctialis* fronds induced by short red (r) and far-red (fr) light pulses (LOEPPERT et al. 1978). Light treatments were carried out after DCMU poisoning (for suspension of photosynthetic electron transport) and after a dark pre-treatment of 1 hour. The horizontal bars indicate the onset and duration of light treatment. The number at the beginning of each plot denotes the cell potential (inside negative); downward movement of the trace indicates depolarization.

A high concentration of ABA (10^{-5} M), which stops vegetative growth of *L. minor* almost completely (VAN OVERBEEK et al. 1968), exerts similar effects on membrane potential in *L. gibba* as fusicoccin (HARTUNG et al. 1980). E_m was increased by 85 mV in especially pre-treated (stressed) plants, which possess E_m values near the diffusion potential. Glucose and glycine uptake was stimulated, and glucose- and glycine-induced changes of E_m were diminished. It seems to be obvious to the writer that the results are brought about by an extensive ABA blocking of metabolism and energy consumption within the plant. Thus, more energy can be offered to transport processes at the plasmalemma.

Phytochrome-mediated changes of E_m have been pointed out in *L. aequinoctialis* (LOEPPERT et al. 1978) and *L. gibba* (KANDELER et al. 1980). After inhibition of photosynthesis by DCMU membrane potential is modulated by short far red and red light irradiations (fig. 2.39). Far red hyperpolarizes E_m by 5 to 9 mV within one minute. The hyperpolarization lasts after the end of irradiation and is abolished by a following red light pulse. Phytochrome seems to act on the active component of E_m as NH_4^+ -treated plants show no light-mediated changes of diffusion potential.

2.5.2.2. Uptake and release of electrolytes

2.5.2.2.1. Alkaline cations

RATHJE (1952) demonstrated that in *L. minor* potassium is taken up from the medium during light and released during darkness, whereas protons show an opposite behaviour. When a high sodium concentration is present in the nutrient medium, sodium is taken up during darkness and released during light, and that to a much higher degree than potassium carries out the reverse movements (SCHEFFER et al. 1952). Potassium uptake under conditions of continuous light was investigated by YOUNG et al. (1970) and YOUNG and SIMS (1972, 1973) with exponentially growing plants of *L. minor*. Influx of $^{42}K^+$ at different external K concentrations closely matches the demands of expanding tissue. At low external concentrations (i.e., below 2×10^{-4} M) movement is effectively unidirectional, and passive fluxes in either direction are negligible. Potassium uptake seems to be mediated by an ATPase, because ATP stimulates, but CCCP (a phosphorylation uncoupler) blocks K^+ uptake. Correspondingly, LOEPPERT (1981) found a close correlation between Rb^+ influx and ATP level after

treatment of *L. aequinoctialis* with arsenate or DCCD. AYADI et al. (1974; see also THELLIER and AYADI 1971) studied K^+ uptake in *L. minor* after keeping plants on distilled water for 24 h. Presence of calcium inhibits K^+ absorption at moderate K^+ concentrations, but activates the process for the lowest K^+ concentrations. The inhibition is interpreted by structural perturbation of the catalytical K^+ uptake site and the activation by an influence on coupled (probably metabolic) processes. Addition of $MgCl_2$ instead of $CaCl_2$ to the KCl solution exhibits an inhibitory effect on K^+ uptake, but no activating effect. When the velocity of K^+ absorption is plotted against K^+ concentration, the classical dual phase behaviour appears (fig. 2.40). Leaving the molecular interpretation of this behaviour open, the data are treated with an 'electrokinetic' formulation of absorption processes (THELLIER 1970, 1973), which employs linear 'flows and forces' relations when conditions are close to equilibrium, and proposes non-linear formulations through analogies with classical electrokinetics when far from equilibrium (high concentrations in the external medium).




Fig. 2.40. Velocity (V) of K^+ absorption plotted against K^+ concentration. Data from experiments with *Lemna minor* (AYADI et al. 1974)

AYADI et al. (1974) probing K^+ uptake rate during different times of the day, found some variations but no diurnal oscillation. In contrast, KONDO and TSUNZUKI (1978) pointed out a circadian rhythm of potassium uptake in *L. gibba*, when plants were held in a weak and constantly flowing nutrient medium under continuous light (see chapter 2.5.9).

The competitive interaction between potassium and rubidium uptake was studied by AYADI et al. (1971) and LASSALLES et al. (1973). Uptake of rubidium at moderate concentrations is inhibited by calcium in a similar way as K^+ uptake (AYADI and THELLIER 1970). Also caesium-137 is absorbed actively by *L. minor* like potassium. The concentration factor (Cs internal : Cs external) ranges from about 84 to 184 (BERGAMINI et al. 1979).

Lithium transport has been investigated in *L. gibba* with the help of stable isotopes, 6Li and 7Li , as tracers (BIELENSKI et al. 1984a, b). Kinetic parameters taken from influx and efflux data showed lithium to be distributed to at least three cell compartments: cell wall, cytoplasm, and vacuole. The passive permeabilities of the plasmalemma and of the tonoplast are rather low (order of 10^{-12} to 10^{-13} m. s.⁻¹). Li^+ is actively pumped out of the cytoplasm, both towards the exterior and towards the vacuole (like Na^+ in other plant cells).

2.5.2.2.2. Ammonium

When the nutrient medium contains ammonium as well as nitrate, ammonium is taken up preferentially (FERGUSON 1969a, FERGUSON and BOLLARD 1969, FELLER and ERISMANN 1971, GROB et al. 1973, TATKOWSKA and BUCZEK 1983, INGEMARSSON et al. 1984, ULLRICH et al. 1984b). In cultures of *S. punctata* without supply of glucose 80% of NH_4^+ and 20% of NO_3^- are consumed from the medium within 10 days. Uptake of NO_3^- increases after practical exhaustion of NH_4^+ ($NH_4^+ = 5\%$). In cultures with glucose-containing medium 40% of NH_4^+ and 20% of NO_3^- are used up within 3 days (GROB et al. 1973). When in sucrose-grown cultures of *S. polyrrhiza* one fifth of the NO_3^- is replaced by NH_4^+ (1.6 mM), uptake and assimilation of NO_3^- are reduced to one half after 18 days of growth (TATKOWSKA and BUCZEK 1983). In cultures of *L. gibba*, which were held in a sucrose-containing medium and were adapted to limited nitrogen supply, uptake of NO_3^- was depressed as long as NH_4^+ remained in the medium (INGEMARSSON et al. 1984). Similar results were obtained with *L. minor*

in diluted anorganic medium (FELLER and ERISMANN 1971). Even a lag-phase of about 10 hours was stated before NO_3^- uptake began after exhaustion of NH_4^+ . NH_4^+ , added to NO_3^- -absorbing plants of L. gibba (N-starved before the experiment), reduced the NO_3^- uptake rates to about one third until the NH_4^+ was consumed. In plants provided with NO_3^- , however, the slow NO_3^- uptake was not inhibited by addition of NH_4^+ (ULLRICH et al. 1984b). In all cases, NH_4^+ uptake can follow the electrochemical gradient. The concentration-dependence curve of NH_4^+ uptake shows a saturating as well as a linear component. Thus carrier-mediated NH_4^+ uniport as well as diffusion of NH_3 may contribute to the uptake of NH_4^+ (ULLRICH et al. 1984b).

2.5.2.2.3. Alkaline-earth cations

Absorption of calcium into L. gibba was studied by DEKOCK et al. (1973). An increase of Ca concentration in the medium from 0.05 to 2.5 mM causes a raising of internal Ca content from 1.6 to 3.5 meq/100 g dry weight. At the same time internal Mg content is lowered from 3.6 to 2.1 meq/100 g dry weight. Addition of oxalate (0.1-2.5 mM), to the medium increases Ca absorption of plants. EDTPA (0.1-0.5 mM), when supplied in the presence of 2.5 mM oxalate, reduces ⁴⁵Ca absorption as well as ¹⁴C-oxalate uptake. BA impedes the absorption of Ca, ABA stimulates it (DEKOCK and HALL 1981).

2.5.2.2.4. Anions

At fairly low medium concentrations nitrate, phosphate, and sulfate, are all taken up by an anion/ H^+ co-transport (see chapter 2.5.2.1). In the case of nitrate it was demonstrated by LOEPPERT et al. (1977) that during steady-state conditions of growth the NO_3^- uptake rate parallels alkalization of the medium with a molecular ratio of 1:1 (experiments with L. aequinoctialis). Furthermore, when the proton gradient across the plasmalemma is discharged with a permeant acid, NO_3^- uptake is inhibited completely (LOEPPERT and KRONBERGER 1979). Under such conditions even in plants well provided with NO_3^- a passive efflux of NO_3^- occurs.

The existence of a NO_3^- -carrier in Lemnaceae can be deduced from a diagram plotting NO_3^- absorption against NO_3^- concentration, which shows

a biphasic pattern for W. arrhiza (SWADER et al. 1975). In Wolffia the carrier is present also after NO_3^- starvation, but its quantity or effectivity is raised 4 to 5 hours after NO_3^- supply. In L. gibba an induction effect of low NO_3^- concentrations (0.1 mM KNO_3) on NO_3^- -uptake rate after N-starvation has been found (ULLRICH et al. 1981). The effect appears after 1 hour, is saturated after 6 hours, and then remains constant for a fairly long time (10 to 20 hours). Also in this case formation of the nitrate carrier may be involved.

In Wolffia an increase of nitrate uptake during illumination occurs in plants with very low nitrate reduction (SWADER et al. 1975). Light via photosynthesis seems to provide energy for the uptake process. In L. aequinoctialis, on the other hand, light increases net nitrate uptake via an enhancement of metabolic NO_3^- consumption, which lowers passive nitrate efflux (LOEPPERT and KRONBERGER 1979).

Biphasic uptake kinetics have been worked out also for phosphate (L. minor: THELLIER and AYADI 1968, MONNIER and THELLIER 1970. L. gibba: NOVACKY and ULLRICH-EBERIUS 1982, ULLRICH-EBERIUS et al. 1984). NOVACKY and ULLRICH-EBERIUS (1982) suggested that the transport systems are both located at the plasmalemma and function with a single, conformational changeable phosphate/ H^+ carrier. Arsenate causes a competitive inhibition of phosphate uptake (MONNIER and THELLIER 1970).

Dependence on concentration of sulfate uptake was investigated in L. minor (THOIRON et al. 1969, THELLIER et al. 1971b), and in L. aequinoctialis (DATKO and MUDD 1984a). In the first two cited publications the biphasic character of kinetic curves was interpreted as being a consequence of non-equilibrium thermodynamics, and it was concluded that postulation of a double system of catalysts would not be necessary. DATKO and MUDD (1984a) obtained SO_4^{2-} uptake kinetics in L. aequinoctialis, which include a saturating and a linear component. As in other plants, molybdate competes with SO_4^{2-} for uptake by the saturating transport system. The effectivity of this transport system is regulated by the internal sulfate and cystine concentration. Between 'low- SO_4^{2-} ' provided and 'high- SO_4^{2-} ' provided plants a 500-fold change in effectivity can be observed. The linear, 'nonsaturating' transport system is not regulated in a similar way. It is much less affected by pH and temperature than the saturating system and may represent a diffusion-mediated uptake.

Fumigation of L. minor with 18 ppm H_2S (BRUNOLD and ERISMANN 1970) or

with 0.5 ppm SO_2 (SCHAEFER et al. 1975) inhibits sulfate uptake very rapidly. Both gases can be used in Lemna as an S-source (see chapter 2.5.6.6). Furthermore, the SO_4^{2-} level within the plant increases gradually during fumigation with H_2S or SO_2 (BRUNOLD and ERISMANN 1970, SCHAEFER et al. 1975). Thus, the inhibiting effect of H_2S and SO_2 on SO_4^{2-} uptake may be caused - at least in part - by the above cited regulation of SO_4^{2-} uptake by internal cystine and sulfate concentration.

Under the conditions used by DATKO and MUDD (1984a), $^{35}\text{SO}_4^{2-}$ efflux from equilibrium-labelled L. aquinoctialis is very low (1.1% of the total ^{35}S within 22.4 hours). By contrast, radioactive SO_4^{2-} efflux from equilibrium-labelled L. minor amounts to 30% of total ^{35}S within the same time (calculated by DATKO and MUDD from the data of THOIRON et al. 1981). Presumably, the rate of SO_4^{2-} metabolism (species-specific and dependent on experimental conditions) may be crucial in this respect.

Uptake of chloride and iodide has been investigated with L. minor (THELLIER et al. 1967). Within the concentration range studied (0.055 - 1 mM), chloride uptake rates reach saturation at about 0.3 mM. Iodide uptake rates, on the contrary, show a linear dependence upon external concentration (0.002 - 4 mM).

When analyzing borate uptake it must be taken into consideration that borate forms very stable borodiester complexes with cell wall constituents possessing pairs of adjacent cis-hydroxyl groups. Using the stable isotope ^{10}B , which can be detected by a (n, α) nuclear reaction, THELLIER et al. (1979) and DUVAL et al. (1980) carried out a compartmental analysis of borate with plants of L. minor. Evaluation of efflux and influx kinetics allowed determination of the borate capacities of the various compartments and of the borate unidirectional fluxes between the different compartments, at borate flux equilibrium. Phosphate uptake (THELLIER and AYADI 1967) and chloride uptake (THELLIER et al. 1967) are inhibited by borate, but it has no effect on sulfate uptake (THELLIER and TROMEUR 1968) and iodide uptake (THELLIER et al. 1967).

2.5.2.2.5. Leakage

Release of electrolytes into the medium has been investigated in connection with the effect of several herbicides (O'BRIEN and PRENDEVILLE

1979). Dinoseb, sodium azide, linuron, prometryne, and simazine were the most active in increasing leakage in L. minor, followed by oxyfluorfen, amitrole, and 2,4-D. Glyphosate and dalapon were the least active.

2.5.2.3. Uptake, transport, and release of organic substances

2.5.2.3.1. Sugars and amino acids

As indicated by changes of membrane potential during uptake experiments (see chapter 2.5.2.1), hexoses as well as neutral and acidic amino acids are presumably taken up from the medium by proton co-transporters (carriers) at the plasmalemma. This view is strengthened by the fact that uptake of glucose and 3-O-methylglucose (ULLRICH-EBERIUS et al. 1978), leucine (NEWTON 1974b), and other neutral and acidic amino acids (BORST-LAP 1977a, JUNG and LUETTGE 1980, 1982a), shows saturation kinetics. Biphasic saturation was found for DL-leucine absorption by intact root tips of L. minor (NEWTON 1974b), and also for uptake of glycine, L-alanine, L-leucine, L-isoleucine, and L-valine by intact fronds of S. polyrrhiza (BORST-LAP 1977a). A common carrier of all neutral amino acids seems to exist in the low as well as in the high concentration range, because competitive uptake inhibition between these amino acids can be demonstrated (BORST-LAP 1974, 1977a, 1978). Deviating results were obtained with L. gibba by JUNG and LUETTGE (1982a). Dependence on concentration of amino acid uptake showed three-phase kinetics in the range from 10^{-6} to 10^{-2} M. Uptake in the lowest concentration range (<0.02 mM) exhibits competitive inhibition between different neutral amino acids but only slight inhibition of L-alanine uptake by aspartate. In the concentration range of 0.02 to 0.5 mM competition between neutral amino acids and L-aspartate has been shown. This second uptake system possesses kinetic data which correspond about to the 'low-concentration' system in S. polyrrhiza. Proton co-transport works especially in this system. The third uptake system in L. gibba in the concentration range >0.5 mM is assumed to come about by diffusion.

Efflux of $\text{L-}^{14}\text{C}$ leucine has been determined from preloaded fronds of S. polyrrhiza. At a concentration of about 2.5 mM (both external and internal) the efflux was 1% of the influx (BORST-LAP 1977a). Boron deficiency has no influence on leucine uptake in L. minor (THELLIER and LE GUEL 1967b).

Uptake of the basic amino acid arginine in L. gibba was investigated by JUNG and LUETTGE (1982a). The authors conclude from their results that arginine moves as a cation into the cells via a uniporter, driven by the negative cell potential. Recently, discrete transport systems for (a) neutral and acidic amino acids, and (b) basic amino acids have been shown to occur also in S. polyrrhiza (BORSTLAP et al. 1986) and L. aequinoctialis 6746 (DATKO and MUDD 1985). In both species and for both systems saturable uptake conforms to the sum of two Michaelis-Menten terms. In addition to the dual saturable uptake the presence of a non-saturable component has been indicated in S. polyrrhiza.

Besides the two uptake systems for amino acids (a and b) six further transport systems have been identified in L. aequinoctialis (DATKO and MUDD 1985, 1986). Reciprocal inhibition studies defined the preferred substrates for these systems as follows: (c) purine bases, (d) choline, (e) ethanolamine, (f) tyramine, (g) urea, and (h) aldohexoses. Sucrose is taken up probably only after cleavage into its hexose moieties. ULLRICH-EBERIUS et al. (1978) pointed out invertase activity to be present in the cell walls of L. gibba. Light stimulates hexose uptake in L. gibba (ULLRICH-EBERIUS et al. 1978) and W. arhiza (BIANCHETTI 1963). Photosynthetic energy seems to contribute to preservation of the transmembrane proton gradient which drives proton co-transport processes.

2.5.2.3.2. Phytohormones and other growth substances

Auxins, gibberellins, and abscisic acid are weak acids which can penetrate cell membranes passively as undissociated molecules. Therefore, short-term uptake of such substances depends on the pH of the medium. Lowering of solution pH (from 5.1 to 4.1, for example) markedly increases the uptake of 2,4-dichlorophenoxyacetic acid (2,4-D) in L. minor (BLACKMAN et al. 1959) and S. intermedia (FERNANDEZ et al. 1972). Similar results have been obtained with 2,3,5-triiodobenzoic acid (TIBA) (BLACKMAN and SARGENT 1959). When several chlorinated derivatives are compared, uptake rate increases with progressive chlorination of phenoxyacetic acid and decreases with progressive chlorination of benzoic acid (KENNEY-WALLACE and BLACKMAN 1972). This behaviour corresponds roughly to the graduated lipophilicity of these substances (see transfer rates through a layer of n-octanol).

Capacity for uptake of weak acids depends on the pH within the various

cell compartments (s. ion trapping by dissociation). This may give an at least partial explanation for the fact that roots of L. minor accumulate much more TIBA than fronds (BLACKMAN and SARGENT 1959). Correspondingly, 2,4-D is more rapidly, and to a higher degree, taken up by roots than by fronds in S. intermedia (FERNANDEZ et al. 1972). Accumulation of gibberellin A₁ proceeds in young daughter fronds of L. aequinoctialis to a higher level than in mother fronds (AL-SHALAN and KANDELER 1979). Daughter fronds of L. gibba show no such dominance over the mother fronds for GA₁ uptake (HARTUNG and KANDELER 1976). In L. gibba the capacity for GA₁ uptake is relatively low in comparison to L. aequinoctialis. Only in L. gibba the GA₁ accumulation can be increased by a pre-treatment with far-red irradiation (10 min far red at the beginning of the daily dark period for four days).

Uptake investigations with 1-¹⁴C-indole-3-acetic acid (IAA) are made more difficult by photolytic decarboxylation of IAA in the medium and subsequent assimilation of ¹⁴CO₂ (BHATNAGAR and TILLBERG 1982). With the α-pyrone fluorescence method for IAA determination the authors revealed a conjugation of IAA immediately after uptake in L. gibba. Also in the case of abscisic acid (ABA) transformation or destruction occurs after uptake (TILLBERG et al. 1980). Compounds containing ¹⁴C other than ABA appear in the solution during long-term experiments with L. gibba.

2.5.2.3.3. Further organic substances

Passive permeability of several small organic molecules has been investigated on root epidermis cells of L. minor with the deplasmolysis-time method (MARKLUND 1936). Generally, the permeability in Lemna is relatively high (but not so high as in Chara) and in accord with lipoid solubility of the respective substances. Urea and thiourea permeate faster, but methylurea and dimethylurea more slowly than in Chara (comparison on the basis of glycerol permeability).

Uptake of uridine in L. gibba was studied by NAKASHIMA and TSUDZUKI (1976, 1977). Saturation of the uptake process at a low concentration of substrate as well as competitive inhibition by compounds structurally analogous to uridine shows uridine uptake to be a carrier-mediated process. Calcium seems to be necessary for the molecular connection of transport protein(s) to the cell membrane. A treatment of plants with sodium citrate reduces the rate of uridine uptake and CaCl₂ is the most

efficient compound in offsetting the citrate reduction. Ascorbic acid is a further substance taken up by *L. gibba* with a saturable transport system (BARBER and CALDWELL 1976).

2.5.2.3.4. Leakage

Already NICKELL and FINLAY (1954; cited after HILLMAN 1961c) stated that *Lemna* cultures excrete organic substances into the medium. FILBIN and




Fig. 2.41. Accumulation of released ABA in the medium of a *Spirodela polyrrhiza* culture. The time scale is related to the day of appearance of detached turions, which is denoted as zero. The curves represent three separate but similar experiments (SAKS et al. 1980).

HOUGH (1979) found a release of organic ^{14}C into the solution after pre-labeling of *L. minor* plants with $^{14}\text{CO}_2$ under photosynthetic conditions. The loss of organic material increases from 1.19 to 8.88% of recently fixed $^{14}\text{CO}_2 \text{ h}^{-1}$, when the concentration of CuSO_4 is elevated from 2.5 to 3.94 mg l^{-1} in standard Hutner's medium (with high EDTA content). Leakage of flower-promoting material into distilled water from fronds of *L. aequinoctialis* was detected by HALABAN and HILLMAN (1970a, 1971). ABA is one of the compounds which are released even into a complete nutrient medium (fig. 2.41). This has been demonstrated for *S. polyrrhiza* (SAKS et al. 1980), *L. gibba*, and *L. aequinoctialis* (MERTEN 1979). Also WOLEK (1979) concluded from his investigations on allelopathy between *S. polyrrhiza* and *W. arrhiza* that ABA may be released into the medium by *W. arrhiza*. JURD et al. (1957) found a caffeic acid ester (similar to, but not identical with chlorogenic acid) to be excreted into the medium by *S. punctata*. Similarly, WALLACE (cited in MCCLURE 1970) stated that cinnamic acids and - after several days of culture - flavonoids can be detected in the medium of axenically cultivated *Lemnaceae*. For the release of amino nitrogen to the medium see chapter 2.4.1.4.

2.5.3. Location and accumulation of minerals

2.5.3.1. Location in organs and tissues

Fronds and roots of *Lemna* spec. contain very different amounts of nitrate (LEHMAN et al. 1981). Already in the youngest adult frond generation the root has 6 to 8 times more nitrate per dry weight than the frond. When the fronds are two generations older, the root contains 11 to 16 times more nitrate per dry weight than the frond.

The distribution of several mineral elements in roots of *L. minor* was investigated by ECHLIN et al. (1982a). X-ray microanalysis on fracture faces of bulk-frozen tissue has shown that K^+ , P , S , Mg^{++} , Cl^- , and Ca^{++} all are present in distinctly higher concentrations in the apical meristem than in the somewhat older tissues which have begun to differentiate. In a region approximately 90 μm from the root tip the potassium values are more or less the same in the xylem and phloem parenchyma and in the endodermis and inner cortex, although the mean value for this element is somewhat higher in the tissues surrounding the stele than in the stele itself. In approximately 300 μm from the root tip there appears to be a higher relative concentration of potassium in the two sectors radiating outwards from the differentiated sieve element and companion cell compared to the region between these two tissues.

2.5.3.2. Location within the cell

Concentrations of Ca^{2+} , K^+ , and Cl^- in cytoplasm and vacuole of vegetative fronds and ABA-induced turions of *S. polyrrhiza* have been determined by SMART and TREWAVAS (1984b). Data from efflux experiments and stereological cell compartment analysis (SMART and TREWAVAS 1983b) have been used for this purpose.

From the results given in table 2.27 it can be drawn that distribution of calcium and potassium within the cells changes, when turions are produced instead of vegetative fronds. In turions the calcium content of the cytoplasm increases 2.5 times and the potassium content decreases to one-third. The differences are caused by changes in developmental processes. Turion-inducing ABA ($10^{-7} M$) has no rapid effect on Ca^{2+} , K^+ , or Cl^- transport.

To some degree calcium is fixed in the Donnan free space. The cell wall

Table 2.27. Concentration (in $mol\ m^{-3}$) of some ions in cell compartments of *Spirodela polyrrhiza* (after SMART and TREWAVAS 1984b)

	Cytoplasm				Vacuole		
		Ca^{2+}	K^+	Cl^-	Ca^{2+}	K^+	Cl^-
Vegetative fronds		16.18	61.09	15.38	3.29	188.14	41.31
Turions		39.94	20.87	3.27	0.52	246.81	3.43

contains compounds with negative charges (polyuronic acids, in the first place) and, therefore, behaves like an ionic exchanger (DEMARTY et al. 1977, 1978). MORVAN et al. (1979, 1980) titrated isolated cell walls of *L. minor* and showed that cell walls have at least two kinds of binding sites. One site is titratable within one hour and can be attributed to polyuronic acids. Both sites have a much greater affinity for Ca^{2+} than for Na^+ and K^+ . In comparison to other plants *Lemna* has a medium capacity for Ca^{2+} fixation (STELZ et al. 1975).

A compartmental analysis for boric acid has been made with the help of ^{10}B , ^{11}B , $^{10}B_3$, detectable by a (n,q) nuclear reaction (THELLIER et al. 1979). *L. minor* plants were grown in nutrient solution with 0.16 mM ^{10}B -boric acid and then the efflux kinetics into an ^{11}B -boric acid solution were determined. From the data borate capacities of the various compartments were calculated. The capacity of free space (including easily dissociable borate monoesters) amounts to 7.2. $\mu mol\ g^{-1}$ dry weight. The values for cell wall (very stable borate diesters), cytoplasm, and vacuole are 19.4, 34.6, and 32.1 $\mu mol\ g^{-1}$ dry weight, respectively. The values obtained from influx experiments are lower (about 1/3) but show the same result with regard to the relative distribution of borate within the cell (DUVAL et al. 1980). In comparison to other aquatic macrophytes *L. minor* is a strong accumulator of boron. Plants grown in a pond together with *Ceratophyllum demersum* show boron concentrations 10-45 times greater than *Ceratophyllum* throughout the growing season (GLANDON and MCNABB 1978). In most determinations boron concentration in *L. minor* amounts to about 2-3 mg B g^{-1} ash-free dry weight.

The significance of chemically, rather than spatially separated compart-

ments has been shown for the compartmentation of cadmium in *W. gladiata* (SCHREINEMAKERS 1986). Three compartments can be divided within the cytoplasm: Cytoplasmic soluble and easily exchangeable Cd, cytoplasmic weakly complexed Cd (or vacuolar Cd), and cytoplasmic strongly complexed Cd.

2.5.3.3. Accumulation

Accumulation of alkaline cations in the vacuole proceeds as long as the permeability of the tonoplast is low. The drastic lowering of potassium content in *L. gibba* after application of a high concentration (1-10 ppm) of ABA (LIEBERT 1977) seems to be the result of a change in permeability of plasma membranes. Also the strong reduction of ^{137}Cs accumulation in *L. minor* after a 32 krad γ -irradiation (LEINERTE 1969) may be explained in the same way.

As mentioned in chapter 1.1 and 1.2.4. Lemnaceae accumulate phosphate in the form of condensed anorganic phosphates and/or inositol phosphates. Without doubt, these poly-anions bind not only alkaline-earth cations, but also several heavy metals. In a number of investigations concentration factors have been determined, which are given by the ratio: content per g dry weight plant / content per ml solution (EL-SHINAWY and ABDEL-MALIK 1980; KOVACS et al. 1984; SILVEY 1967; SKLAR 1980; VERMAK et al. 1976). Interestingly enough, especially the physiologically active micro-nutrients Fe, Mn, Zn, Cu, and Co are accumulated to a very high degree (concentration factors between 10^4 and 4.6×10^5). Phosphate also reaches a factor of about 10^4 . Other elements with concentration factors in this range are Al, Ti, Sn, and Sb. Concentration factors of 10^2 to 10^3 were found for Pb, Pr, Ce, La, Cs, Mo, Nb, Zr, Y, Ni, Cr, and V. Absolute values for Mn are 1500 (*L. gibba*), 2800 (*L. minor*), 3900 (*L. tri-sulca*), and 2000 (*S. polyrrhiza*) $\mu\text{g/g}$ dry weight, each (BORESCH 1935/1936).

OLSEN (1934a,b) found 500 μg Mn/g dry weight only after culture of *S. polyrrhiza* in 2.5 mg $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$ l^{-1} (The culture solution was of pH 7.0. The Mn uptake is optimum at pH 6-7). The Fe content of *S. polyrrhiza* was 2370 $\mu\text{g/g}$ dry weight after supply of 7 mg Fe l^{-1} as ferric citrate at pH 4, which is optimum for Fe uptake.

Another factor which contributes to accumulation of heavy metals, is the production of several chelating organic compounds (organic acids, amino

acids, peptides, nucleotides, RNA, DNA, and others). In *L. minor* nitrate-grown plants contain 3-4 times more iron and malate than ammonium-grown plants (FELLER and ERISMANN 1973). Correspondingly, supply of ammonium to the nutrient medium reduces the copper content in *L. aequinoctialis* (NASU et al. 1983).

Heavy metal accumulation in *S. polyrrhiza* has been investigated varying the supply of iron, magnesium, zinc, and copper (SCHREINEMAKERS 1984, SCHREINEMAKERS and DORHOUT 1985). Under most conditions, the accumulation changes more or less in parallel for pairs of elements, namely calcium and zinc, magnesium and manganese, and copper and iron.

2.5.4. Water relations

Stomata are movable in Spirodela (at least S. punctata), but not in the genus Lemna (s. chapter 2.5.1.1). Therefore, transpiration proceeds in Lemna in parallel to evaporation. DEBUSK (1980) compared evapotranspiration of L. minor with evaporation of the free water surface and found a ratio evapotranspiration/evaporation = 0.9. The average water loss of plants amounted to $4.4 \text{ l m}^{-2} \text{ d}^{-1}$.

Duckweeds can grow on liquid media as well as on wet substrates (loamy soil or agar), but the under side of fronds is very sensitive to desiccation. Plants of W. arrhiza can survive without water (at 65% relative humidity) for 20-30 min. Plants of S. polyrrhiza, L. minor, L. gibba, and L. trisulca endure desiccation for about 2.5 hours by subsequent outgrowth of daughter fronds (WOLEK 1981).

The osmotic pressure in young, still viable stomata of L. minor fronds is stated to be equivalent to 0.25-0.3 M CaCl_2 (REUTER 1948). The surrounding epidermis cells show values which are only slightly lower. In subepidermal cells of roots, however, the osmotic pressure lies in the range of 0.2-0.3 M glucose (PIRSON and SEIDEL 1950). From the tip to the basis of the root there is an osmotic gradient with a minimum at the end of the cell elongation zone. Replacement of Ca^{2+} by K^+ or Na^+ in the surrounding medium for 7-12 days causes a slight increase of osmotic values. A similar effect comes about by culturing plants in medium containing 0.5% glucose. Replacement of K^+ by Na^+ , on the other hand, decreases osmotic values somewhat, as also 4 hours of darkness. PIRSON and GOELLNER (1953) stated that reduction of nitrate supply to 1/100 or phosphate supply to 1/300 (replacement of NO_3^- and PO_4^{3-} by Cl^-) increases the osmotic pressure especially in the cell elongation zone.

The relation of osmotic quantities (osmotic pressure, turgor pressure, and suction force) to the relative cell volume of young daughter fronds (about 3 mm in length) of L. aequinoctialis was determined by TAKAKI (1969). The increment of volume is 27%, when a frond is transferred to water from sucrose solution with a concentration identical to the incipient plasmolysis concentration (0.24 M).

Plasmolysis of L. trisulca with diverse hypertonic solutions for 30-40 min destroys all the mature cells, but meristems can grow out (KARZEL 1926). During the following 8 months (i.e., many frond generations) growth is decelerated and fronds and stalks remain smaller. A less ex-

treme form of osmotic stress comes about by application of subplasmolytic solutions. 0.2 M polyethyleneoxide, glucose or mannitol, supplied for 4 hours to L. minor plants, stops cell lengthening in the elongation zone of roots irreversibly (SCHAEFER 1958). The treatment reduces the adhesion between cytoplasm and cell wall as can be seen by subsequent plasmolysis. The plasmolysis time is shortened and the plasmolysis form is more convex.

Investigations on plasmolysis time and plasmolysis forms in root cells of L. minor have been made by several authors. STRUGGER (1934) demonstrated concave plasmolysis to occur in the elongation zone but convex plasmolysis in mature cells. Correspondingly, plasmolysis time is maximum at the end of cell elongation zone (PIRSON and SEIDEL 1950, SCHAEFER 1955). Darkening, CO_2 , elevated temperature (26 versus 20°C), and Ca^{2+} deficiency are factors which shorten the plasmolysis time (PIRSON and SEIDEL 1950, PIRSON and SCHAEFER 1957, SCHAEFER 1956, 1958). Moderate N- or P-deficiency, however, extend this parameter (PIRSON and GOELLNER 1953). Investigations with the upper epidermis of L. minor and S. polyrrhiza fronds have shown that the plasmolysis form changes from the concave to the convex type with increasing senescence of the fronds (PI-SCHER 1949).

2.5.5. Carbon metabolism

2.5.5.1. Starch

As stated in chapter 2.5.1.1.2 *Lemnaceae* belong to the C_3 type of photosynthetic CO_2 assimilation. Products of photosynthesis which are not directly used up by cell metabolism, are stored within the chloroplast in the form of starch. Accumulation of starch depends primarily on provision of the plants with sufficient CO_2 . Under continuous light, the starch content of *L. minor* increases with increasing CO_2 supply (to the five-fold level from 100 to 9000 ppm CO_2 ; MUELLER et al. 1977). At 330 ppm CO_2 (normal air) starch production is adequate only when the plants are kept in the open air or are provided with a continuous air stream. Under such conditions, *L. gibba* shows an accumulation rate of 16 μg starch mg^{-1} dry weight h^{-1} (light intensity 9000 lux; temperature 26°C; medium pH 4.8. KOENIGSHOFER, unpublished results). During the daily light phase starch content increases continuously and decreases during the night phase. Short-period oscillations of starch level, which were found in plants cultivated in vessels with cotton stoppers (ROTTENBURG et al. 1981), cannot be ascertained in open air. Also the effect of end-of-day far red on starch content (KANDELER et al. 1980) cannot be reproduced when gas exchange is improved.

In nitrate- as well as in ammonium-grown *L. minor* plants sugar and starch content increases, when the pH of the nutrient medium is lowered (KOPP et al. 1974a). SATAKE and SHIMURA (1983) showed that *S. polyrrhiza* takes more CO_2 from water at low pH than at high pH, because the dissolved CO_2 content in the water is much higher. Other factors which enhance starch accumulation, are P deficiency (REID and BIELESKI 1970a), K^+ deficiency (WHITE 1939), supply of branched-chain amino acids (VAN MAZIJK 1975), and B deficiency (SCHOLZ 1962). In the first three of these cases an inhibition of growth may lead to a restricted starch utilization. B-deficiency, however, even enhances frond multiplication without a reduction of dry weight per frond. The author presumed an effect of B on metabolism of carbohydrates. Fumigation of *L. minor* with sublethal concentrations of SO_2 (0.6 ppm) diminishes the starch content immediately by 20% and also the growth rate is reduced during the first two weeks by 25% (FANKHAUSER et al. 1974, 1976).

2.5.5.2. Organic acids

In *L. aequinoctialis*, malate is produced as an end product of photosynthesis. As can be seen in fig. 2.42, accumulation of malate proceeds only during the light phase and is blocked by 10^{-6} M DCMU (KANDELER and HELDWEIN 1979). The extent of malate synthesis seems to depend on a control mechanism, which regulates the cytoplasmic pH. When protons are consumed (especially during nitrate reduction) and cannot be delivered subsequently from outside the cell (i.e., at relatively high medium pH), then malic acid production is enhanced. This view is stressed by the finding that the inhibitor of nitrate reduction, tungstate, lowers the




Fig. 2.42. The effects of continuous light, DCMU, and sucrose on malate accumulation in *Lemna aequinoctialis* 6746 (KANDELER and HELDWEIN 1979). Plants were held in Pirson-Seidel medium without nitrate at pH 6. Malate content of plants at the beginning of the 24-hour experiment was 0.11 μmoles/g FW.

malate accumulation in L. aequinoctialis and L. gibba (HELDWEIN and KANDELER 1981). Transfer of L. minor from NH_4^+ to NO_3^- -containing solution raises malate content of the plants after a lag period of about 4 hours (KOPP et al. 1974b). Adaptation of plants to end-of-day irradiation with far red, which enhances malate accumulation, also seems to act through an improvement of nitrate assimilation (HELDWEIN and KANDELER 1981). Furthermore, a rise of medium pH from 4.8 to 6.0 increases malate production in L. aequinoctialis (KANDELER and HELDWEIN 1979).

Supply of ABA (10^{-6} M) is another treatment which promotes malate accumulation in L. aequinoctialis and L. gibba. This finding may explain the fact that nitrogen deficiency leads to a higher malate content in L. aequinoctialis (KANDELER and HELDWEIN 1979). Preliminary data of MERTEN (1979) have shown that endogenous ABA production is increased by N deficiency in this plant. ABA (3.8×10^{-6} M) enhances the activities of PEP-carboxylase and NADP-malate dehydrogenase in L. minor (BAUER et al. 1976). NaCl as another stress factor, increasing ABA level up to 1.5 times, has no increasing effect on malate synthesis in L. minor (HUBER and SANKHIA 1979). In this case, however, the malate level of control plants lies one order of magnitude higher than that found in L. aequinoctialis and L. gibba.

Oxalate synthesis has also been considered to be a component of the biochemical "pH-stat". In correspondence with this hypothesis, BORNKAMM (1965) found a close connection between oxalate accumulation and protein content in L. minor. Plants grown with various concentrations of NO_3^- , or H_2PO_4^- , or chloramphenicol, show a parallelism between oxalate and protein content in every case. Water-soluble as well as water-insoluble oxalates are affected in the same way.

In L. minor and L. trisulca there are both forms of glycolate oxidase, the FMN-activated form as well as the glycolate-activated form (BORNKAMM 1969a). This enzyme catalyzes not only the oxidation of glycolate to glyoxylyate, but also the oxidation of glyoxylyate to oxalate. Thus, oxalate production in Lemna could proceed within peroxisomes in connection with photorespiration. In cell-free extracts of L. minor glyoxylyate- ^{14}C is transformed to oxalate. Furthermore, in vitro oxidation of glyoxylyate is inhibited by addition of 25 μM oxalate to the cell-free system (BORNKAMM 1964, 1965). Plants grown in nutrient medium with 1 mM glyoxylyate, reach a higher content of water-insoluble oxalate. Glycine, on the other hand, inhibits oxalate accumulation. NUSS and LOEWUS (1978)

stated that L-ascorbic acid can serve as another precursor of oxalic acid, but - in contrast to what is seen in several other plants - the incorporation of ^{14}C from [$1\text{-}^{14}\text{C}$] ascorbic acid into oxalate in intact plants of L. gibba is relatively low. Incorporation of ^{14}C -labeled glycolate, glyoxylyate, and ascorbate into calcium oxalate crystals has been found microautoradiographically in L. minor (FRANCESCHI 1985). The glycolate oxidase pathway of oxalate formation is operational as shown by the inhibiting effect of αHPMS (2-pyridylhydroxymethane sulfonic acid) and MHA (methylhydroxybutanoic acid). Both inhibitors are effective also during incorporation of ^{14}C from ascorbate, indicating that glycolate and/or glyoxylyate are intermediates of ascorbate catabolism.

Factors which inhibit oxalate synthesis in vivo to some extent, are high temperature (30°C ; BORNKAMM 1964, 1965, 1970c), higher light intensity (4000 versus 600 lux), 1.8% CO_2 , 0.5% glucose, and manganese deficiency (BORNKAMM 1964, 1965). The water-insoluble portion of oxalate is deposited in idioblasts within special membranes as the monohydrate of calcium oxalate (ARNOTT and PAUTARD 1965, AL-RAIS et al. 1971). BORNKAMM (1965) could not detect any degradation of calcium oxalate crystals (raphides) even under calcium deficiency. On the contrary, under such conditions the oxalate accumulation is not reduced, so that the portion of water-soluble oxalate increases from 7.7 to 85.4%.

2.5.5.3. Myo-inositol

This compound plays a role in many metabolic processes. Two of them have been investigated with duckweeds: the production of phytic acid as a phosphate-storing substance (s. chapter 2.5.6.2), and biosynthesis of apiose, a component of pectic polysaccharides in the cell wall of Lemna-ceae (see below).

Myo-inositol is derived from glucose-6-phosphate in two steps (fig. 2.43). Firstly glucose-6-phosphate is transformed to myo-inositol-1-phosphate. The enzyme myo-inositol-1-phosphate synthase is present in L. aequinoctialis (LOEWUS and LOEWUS 1971) and has been isolated carefully from L. gibba (SCHWARCZ et al. 1974, OGUNYEMI et al. 1978). The purified enzyme has a molecular weight of 135000 ± 5000 and consists of 3 subunits with identical electrophoretic behaviour. It is inhibited by EDTA and can be reactivated by Mn^{2+} . The specific activity of the enzyme is about 500 times higher than that of the animal enzymes so far investigated.

The second step of myo-inositol formation, the cleavage of myo-inositol-1-phosphate by a phosphatase, should also occur in Lemnaceae, as free myo-inositol can be handled in phytic acid metabolism (ROBERTS and LOEWUS 1968, BOLLMANN et al. 1980) as well as in biosynthesis of apiose and xylose (ROBERTS et al. 1967).

2.5.5.4. Apiose and apiogalacturonans

A very characteristic feature of Lemnaceae are apiogalacturonans in the cell wall. Besides Lemnaceae apiose-containing polysaccharides have been found only in some species of Hydrocharitaceae, Potamogetonaceae, and Zannichelliaceae (see GRISEBACH 1980). Such polysaccharides with high D-apiose content are not degraded by pectinase (HART 1969, HART and KINDEL 1970b) and thus, make plants resistant to the attack of several microorganisms. When isolation of Lemna protoplasts is required during physiological investigations, an enzyme mixture must be used containing not only cellulase "Onozuka R-10" and Mazerozyme R-10, but also Driselase (LOEPPERT and KOENIGSHOFER, unpublished results).

Occurrence of apiose, a branched-chain sugar, in Lemnaceae has been ascertained several times (s. chapter 1.2.3). Biosynthesis proceeds via UDP-bound intermediates from D-glucuronic acid (fig. 2.43). UDP-glucuronic acid is oxidized with NAD to a 4-keto-intermediate. Then decarboxylation leads to a second intermediate, which undergoes a ring contraction reaction. The resulting aldehyde is reduced by NADH to UDP-D-apiose. Extensive work has been done by several working groups to elucidate this biosynthetic reaction chain in Lemna. Reviews have been given by WATSON and ORENSTEIN (1975), GRISEBACH (1980) and BECK (1982). The original papers on this topic are the following: MENDICINO and PICKEN (1965), BECK and KANDLER (1966), PICKEN and MENDICINO (1967), SANDERMANN et al. (1968), GUSTINE and KINDEL (1969), MENDICINO and HANNA (1970), SANDERMANN and GRISEBACH (1970), KELLEHER and GRISEBACH (1971), KINDEL and WATSON (1973).

The enzyme managing the reaction chain from UDP-glucuronic acid to UDP-apiose also catalyzes the formation of UDP-D-xylose from the second above-named intermediate and, therefore, has been named UDP-apiose/UDP-xylose synthase (WELLMANN and GRISEBACH 1971, GUSTINE et al. 1975). The enzyme, which has been purified more than 1000-fold, possesses a molecular weight of 110000 and a pH optimum of 7-8.5 (WELLMANN and GRISEBACH




Fig. 2.43. Biosynthesis of myo-inositol, apiose, and xylose in Lemna.

1971, MENDICINO and ABOU-ISSA 1974). In addition, another enzyme (UDP-glucuronate carboxylase) exists, which catalyzes the transformation of UDP-glucuronic acid to UDP-xylose solely (GUSTINE et al. 1975).

The glucuronate units for biosynthesis of apiose are derived from glucose-6-phosphate via glucose-1-phosphate and UDP-glucose or via myo-inositol-1-phosphate and myo-inositol. ROBERTS et al. (1967) showed that myo-inositol-2-¹⁴C is incorporated into apiosyl and xylosyl units of cell wall polysaccharides in *L. gibba* G3.

Apiogalacturonans have been isolated from *L. minor* (BECK 1967). One fraction consisted solely of D-apiose and polygalacturonic acid, another one contained xylose in addition. HART and KINDEL (1970b) separated at least five polysaccharides containing D-apiose from *L. minor*. The apiosyl side-chains are attached to the galacturonan backbone in form of apioside, a disaccharide consisting of two apiose moieties (HART 1969, HART and KINDEL 1970a, MASCARO and KINDEL 1977). Transferase activity which attaches D-apiosyl- and D-xylosyl-units from UDP-D-apiose and UDP-D-xylose to unknown endogenous acceptors (apiogalacturonans), has been demonstrated in a particulate preparation of *L. minor* (PAN and KINDEL 1977). Also a particulate enzyme incorporating UDP-galacturonic acid into pre-existing galacturonans has been isolated (LEINBACH 1975). This D-galacturonosyl transferase is stimulated by Mn²⁺ and possesses optimum activity at pH 6.0-6.2.

In connection with the development of intercellular air spaces *Lemna* should have the ability to degrade apiogalacturonans. Even an oxidation of apiose to CO₂ occurs when D-[U-¹⁴C] apiose is supplied to sterile *L. minor* plants (HANNA et al. 1973).

2.5.6. Assimilation of phosphorus, nitrogen, and sulfur

2.5.6.1. Assimilation of phosphate

The use of phosphate within cell metabolism of *S. punctata* has been investigated by BIELESKI (1968a). After 9 days of growth in a solution containing phosphate-³²P the various radioactive compounds were extracted and determined quantitatively. In the principal fractions, inorganic phosphate, RNA, phospholipids, and soluble phosphate esters, the proportion of phosphorus amounts to 71.1, 15.1, 7.5, and 4.9% of total P, respectively. Within phospholipids phosphatidyl choline (46.3%), phosphatidyl ethanolamine (22.5%), phosphatidyl glycerol (13.5%), and phosphatidyl inositol (6.3%) are the main components. In the fraction of acid-soluble phosphate esters glucose-6-phosphate (22.4%) and phosphoglyceric acid (10.2%) predominate.

In well-nourished plants inorganic phosphate is present in two pools: a metabolic (12%) and a non-metabolic pool (88%), which may be located in cytoplasm and vacuole, respectively (BIELESKI 1968b). Under conditions of phosphate deprivation, however, the non-metabolic pool is exhausted and more than 90% of the remaining inorganic phosphate is traceable in the metabolic pool. Phosphate ester content then falls from 3.5 to 0.6 µmoles/g, phospholipid content from 3.5 to 1.2 µmoles/g, and residual phosphate (mainly RNA) content from 7.5 to 2.0 µmoles/g. Relative proportions of the various phosphate esters and relative proportions of the various phospholipids, are not markedly affected by phosphate deficiency.

2.5.6.2. Storage of phosphate

Within minerals phosphorus is the factor which limits occurrence of *Lemna* in the field (LUEBEND 1983). Obviously, the ability of duckweeds to store P excessively is one of the evolutionary adaptations to the habitat of these floating water plants. Besides the accumulation of orthophosphate within the vacuole, phosphate can be stored in the forms of condensed inorganic phosphates and of phytic acid. In vegetatively growing fronds of *L. aquinoctialis* and *L. gibba* linear oligophosphates with 2 to 7 residues, cyclic tri-, tetra-, penta- and hexametaphosphates, and high-molecular condensed phosphates are produced rapidly, when phos-

phate-³²P is applied to the plants (NIEMEYER 1975, INHUELSEN and NIEMEYER 1975). Subsequently the condensed phosphates are degraded in connection with the synthesis of nucleic acids. Condensed phosphates, therefore, seem to function as a short-term reserve pool of inorganic phosphate. Phytic acid, on the other hand, is produced preferentially under conditions, which favour the development of turions or turion-like fronds (a. chapter 2.4.2), and can serve as a long-term P-storage compound. Nitrogen deficiency, sucrose supply, and short day in combination with low temperature (8-10°), are all factors inducing phytic acid synthesis in *L. gibba*. Moreover, turions of *S. polyrrhiza* have a very high content of phytic acid in comparison to vegetative fronds (SCHEINER et al. 1978).

Besides the inositol hexaphosphate phytic acid, also the other inositol phosphates have been found in *L. minor* (INHUELSEN and NIEMEYER 1978), *L. gibba* and *Wolffiella gladiata* (ROBERTS and LOEWUS 1968). A series of three enzymes is needed to convert myo-inositol to phytic acid with ATP as the phosphate donor (BOLLMANN et al. 1980). Myo-inositol kinase phosphorylates myo-inositol to myo-inositol-1-phosphate. Then myo-inositol-1-phosphate kinase catalyzes the formation of myo-inositol trisphosphate which is transformed to phytic acid by myo-inositol trisphosphate kinase.

2.5.6.3. Acquisition of phosphate at P deficiency

Shortage of phosphate in the environment not only leads to consumption of endogenously stored phosphate, but also to attempts for acquisition of exogenous phosphate by enzymatic cleavage of P-containing organic compounds. At phosphate deficiency phosphatases and ribonucleases are synthesized, which either are fixed on the outer side of the plasmalemma, or are released to the medium. Induction of phosphatase synthesis by P deficiency has been demonstrated in *S. polyrrhiza* (JUNGnickel 1981), *S. punctata* (REID and BIELESKI 1970b, BIELESKI 1974, KNYPL and KABZINSKA 1977, KNYPL 1978), *L. minor* (LIEDKE and OHMANN 1969), and *W. arhriza* (JUNGnickel and VOELKSCHE 1979, JUNGnickel 1981). BIELESKI (1974) and KNYPL (1979a) found two alkaline phosphatases to be induced in P-starved plants, which are located (fig. 2.44) at the epidermal and subepidermal cell walls on roots and the lower side of fronds (BIELESKI and JOHNSON 1972). In addition, KNYPL (1979a; see also KNYPL and SOBOLEWSKA

1978, KNYPL 1979b) pointed out one acid phosphatase and two further alkaline phosphatases which are released to the medium at P deficiency. The two ribonucleases produced by *S. punctata* during P starvation, remain on the cell surface (KNYPL and KABZINSKA 1977, KNYPL 1978, 1979a). BIELESKI and JOHNSON (1972) and KNYPL (1978) have shown that inorganic phosphate appears in the medium of a P-deficient *S. punctata* culture,



Fig. 2.44. Location of phosphatase in control (left) and phosphorus-deficient (right) *Spirodela punctata* plants (BIELESKI and JOHNSON 1972). Plants were fixed, stained by the procedure of Gomori, embedded, and sectioned. Note that control and P-deficient plants were stained for different lengths of time, to compensate for their different phosphatase activities.

A. Cross section of control frond (x70)

B. Cross section of P-deficient frond (x75)

C. Cross section of control root (x300)

D. Cross section of P-deficient root (x250)

Note occurrence of phosphatase (black stain) in the region of vascular tissues in control plants (left), and in the epidermis of P-deficient plants (right).

fic NR is induced after supply of sucrose to the culture medium (YOUNG 1967, cit. in SIMS et al. 1968). Iron-deficiency lowers NR activity considerably (SPILLER et al. 1973) and tungstate prevents appearance of NR activity completely (STEWART 1972b). ABA (2×10^{-5} M) has no effect on NO_3^- induction of NR in *S. polyrrhiza* (STEWART and SMITH 1972). The ionic surfactant DBST (trimethylamine alkylbenzenesulfonate) distinctly lowers the activities of NR and NiR in *S. polyrrhiza* (BUCZEK 1984b). The non-ionic surfactant ENF (alkylphenoxypolyethoxy ethanol) inhibits only the NiR activity.

A wealth of papers has been published to answer the question of repression and/or inactivation of NR by ammonium (SIMS et al. 1968, FERGUSON 1969a, FERGUSON and BOLLARD 1969, JOY 1969a, STEWART 1972a,b, STEWART et al. 1974, OREBAMJO and STUART 1974, 1975a, 1975b, MELZER and EXLER 1982, VIJAYARAGHAVAN et al. 1982, TATKOWSKA and BUCZEK 1983). A difficulty not considered in the older works is the fact that an inhibitor of NR, tentatively characterised as a protein, seems to occur in duckweeds (EISENMEIER and SPILLER 1973, OREBAMJO 1973, cited in STEWART et al. 1974, BUCZEK 1984a) and that this inhibitor has its own dynamics (OREBAMJO and STEWART 1975a). The mode of interrelation between ammonium ions and the inactivating system is not yet known. A promotive NH_4^+ effect on formation or an activation of the inhibitor protein has been discussed (OREBAMJO and STEWART 1975b).

The different results obtained for the effect of ammonium on NR induction may be caused by certain pre-culture conditions and - in consequence - by different contents of NO_3^- , NH_4^+ , and the supposed inactivator. When plants were transferred to the induction (NO_3^-) medium directly from NH_4^+ - or medium containing amino acid, presence of ammonium in the induction medium strongly inhibits or prevents, the raise of NR activity (JOY 1969a, STEWART 1972a,b). A more detailed investigation of kinetics reveals, however, that under such conditions NH_4^+ inhibition of NR activity originates not before 3-4 hours after the beginning of substrate induction (OREBAMJO and STEWART 1975a). This holds true also for plants which have been pre-cultured in an N-free medium for 3 days (TATKOWSKA and BUCZEK 1983). Even a promotion of the NR induction process by ammonium can be seen during the first hours (FERGUSON 1969a) or one day (VIJAYARAGHAVAN et al. 1982), when plants were N-starved for two days before the experiment.

Taking all the data together, it seems that ammonium ions inactivate NR

when glucose-1-phosphate or certain adenosine monophosphates (5'-AMP, 2'-AMP) are supplied. Also 32P -RNA extracted from *S. punctata* is hydrolyzed externally by P-starved *S. punctata* plants (KNYPL 1980). Both oligo- and mononucleotides appear in the solution and the label is incorporated to nucleic acids and low molecular weight P compounds of the P-starved plants. Obviously, compounds containing P released from senescing plants can be re-utilized by younger neighbours. RNA-like material has been detected in the medium of senescing plants (KNYPL 1978).

Acquisition of phosphate or P-containing compounds even occurs in P-provided cultures with the help of constitutive enzymes. This fact may explain, why symptoms of P deficiency as well as induction of phosphatases and ribonucleases do not proceed, when inorganic phosphate in the solution is replaced by yeast RNA (KNYPL 1978). Inorganic P sources other than orthophosphate, such as pyrophosphate, tripolyphosphate, and shark teeth phosphorus, are accessible to *L. minor* (FITZGERALD 1971).

2.5.6.4. Nitrate reduction

Under natural conditions, nitrate is the main N-source for duckweeds in most cases. After uptake, nitrate must be reduced to ammonia before synthesis of N-containing organic compounds can proceed. The two enzymes responsible for this reduction are NADH-specific nitrate reductase (NR) and red. ferredoxin-dependent NADH-specific nitrite reductase (NiR). Both enzymes are produced adaptively, i.e., only in the presence of their substrates. Nitrate induces synthesis of NR as well as of NiR (*S. punctata*: FERGUSON 1969a; *L. minor*: JOY 1969a, STEWART 1972b). Nitrite is less effective than nitrate for induction of NiR in both species and very little effective for induction of NR (for nitrate induction of NR in *L. trisulca* see MELZER and EXLER 1982, and in *S. polyrrhiza* see SWÄDER and STÖCKING 1971). That really new synthesis of the enzymes proceeds during substrate induction, has been confirmed by experiments with inhibitors of nucleic acid and protein synthesis (JOY 1969a, STEWART 1968, 1972b, VIJAYARAGHAVAN et al. 1982). An enhancing effect of sulfate on NR induction in *L. minor* indicates some coordination of N- and S-metabolism (BRUNOLD and SUTER 1984a). Light and sucrose, alone as well as in combination, increase substrate induction of NADH-specific NR in *S. punctata* (VIJAYARAGHAVAN et al. 1982). The light effect is mediated by the photosynthetic apparatus as it is blocked by DCMU. In *L. minor* a NADPH-specific

but do not repress this enzyme (TATKOWSKA and BUCZEK 1983). Reversibility of NR inactivation in vivo and in vitro has been shown by OREBAMJO and STEWART (1975b).

Certain amino acids (alanine and arginine, for example), asparagine and urea exert similar effects on NR induction as ammonium ions. Depending on experimental conditions a strong or moderate inhibition (SIMS et al. 1968, STEWART 1972a, 1972b, VIJAYARAGHAVAN et al. 1982) or no effect (FERGUSON 1970) has been found.

Nitrite reductase is a more stable enzyme than NR (BUCZEK 1984a) and less accessible to inhibiting effects of ammonia and amino acids (SIMS et al. 1968). Nevertheless, STEWART (1972b) obtained strong inhibitions of NIR induction by ammonia, glycine, asparagine, and urea in *S. punctata*. SWADER and STOCKING (1971) pointed out that NO_2^- reduction but not NO_3^- reduction is located in the chloroplasts of *W. arizhiza*. The pro-viding of protons for nitrite reduction by formation of malic and oxalic acid has been reported in chapter 2.5.5.2.

2.5.6.5. Assimilation of ammonia, urea, and other exogenous organic

N compounds

Ammonia is taken up and utilized preferentially (s. chapter 2.5.2.2.2), when NH_4^+ and NO_3^- are both present in the nutrient medium (s. above: endproduct inhibition of nitrate reductase). However, in the case of ammonia and nitrite (FERGUSON and BOLLARD 1969) or ammonia and amino acids (FERGUSON 1970) both compounds are used up simultaneously.

BAUER et al. (1973) traced the $^{15}\text{NH}_4^+$ incorporation in the active turnover pools of glutamic acid, alanine, and glutamine in *L. minor*. They concluded that glutamic acid is the most important primary assimilation product of ammonium. In a more advanced analysis RHODES et al. (1980) revealed that *L. minor* assimilates $^{15}\text{NH}_4^+$ solely through the glutamine synthetase - glutamate synthase cycle when ammonium is provided at a relatively low concentration (2 mM). When activity of glutamine synthetase has fallen to zero after addition of azaserine and methionine sulphoximine, no ^{15}N labelling of glutamic acid occurs. This means that in spite of the presence of active NAD-glutamate dehydrogenase NH_4^+ cannot be incorporated directly into 2-oxoglutarate.

The glutamine pathway of ammonium assimilation (fig. 2.45) consists of two steps: the transformation of glutamate to glutamine with NH_3 by glu-

tamine synthetase (GS), and transfer of the amide N from glutamine to 2-oxoglutarate by glutamate synthase (GOGAT). One of the two molecules of glutamate released by the second reaction is recycled to the first reaction. The GS-GOGAT cycle is located in chloroplasts (for Lemna see STEWART, cited in RHODES et al. 1979) and, therefore, connected intimately with nitrite reduction. ATP (for the first step) and reduction energy (for the second step) are provided by photosynthesis.

The glutamine pathway is the principal way of NH_4^+ incorporation at least in plants grown on nitrate or relatively low concentrations of ammonium. STEWART and RHODES (1976) have shown that blocking of GS by methionine sulphoximine (MSO) causes an accumulation of ammonia and 2-oxoglutarate, and a decrease of glutamine level in nitrate-grown *L. minor*. Also in *L. gibba* MSO enhances the intracellular ammonium level (JOHANSSON and LARSSON 1986a). Furthermore, azaserine (inhibiting glutamine amide transfer of GOGAT) increases $^{14}\text{CO}_2$ incorporation into 2-oxoglutarate associated with a decrease of incorporation into glutamate in plants grown on 5 mM ammonia (RHODES 1976, cited in RHODES et al. 1979).




Fig. 2.45. The glutamine pathway of ammonium assimilation (from MIFLIN and LEA 1980). Enzymes: 1, glutamine synthetase; 2, glutamate synthase.

At relatively high NH_4^+ concentrations, however, a partial switching over to the glutamate pathway of ammonium assimilation seems to occur. With increasing NH_4^+ content of the medium (2-15 mM) the activities of GS and GOGAT decrease (RHODES et al. 1975, 1976) and the activity of glutamate dehydrogenase (GDH) increases (JOY 1969a [compare however JOY 1971], SHEPARD and THURMAN 1973, RHODES et al. 1976, EHMKE and HARTMANN 1976 [tab. 2]). The NH_4^+ effect on GS activity is provable especially when ammonium treatment is combined with darkness (RHODES 1976, cited in RHODES et al. 1979).

RHODES (1976, cited in RHODES et al. 1979) and RHODES et al. (1976) pointed out that glutamine - rather than ammonia itself - may be the substance exerting negative control on GS and GOGAT and positive control on GDH. Comparison of tissue glutamine levels with enzyme activities from plants cultivated with various N-sources reveals a negative correlation between glutamine and GS, but a positive correlation between glutamine and GDH (STEWART and RHODES 1977b, cited in RHODES et al. 1979). Glutamine synthetase from *L. minor* has a mol. wt of about 360000 and is probably composed of eight identical subunits. The physiologically active state needs Mg^{2+} (RHODES et al. 1979). The rapid loss of GS activity after transfer of plants to ammonium solution and darkness involves a reversible deactivation of the enzyme. A low molecular weight, heat-stable fraction has been isolated, which can promote GS deactivation and inhibit reactivation in vitro (RHODES et al. 1979). A feed-back control of GS, on the other hand, is exerted by some amino acids (alanine, glycine, serine, aspartate) and certain nucleotides. Both ADP and 5'-AMP competitively inhibit the enzyme with respect to ATP (STEWART and RHODES 1977a). The authors concluded that control of the enzyme by energy charge may be important under some conditions.

The enzyme for reductive amination of 2-oxoglutarate, glutamate dehydrogenase (GDH), has been investigated several times. The most advanced analysis was made by HARTMANN and co-workers with *L. minor* plants (EHMKE and HARTMANN 1976, 1978, SCHEID et al. 1980, EHMKE et al. 1980). The authors isolated a single protein (mol. wt 230000) consisting of four identical subunits. The enzyme exhibits catalytic activity preferably with NADH, but also with NADPH to some degree (ratio of activities NADH/NADPH = ca 12/1). The GDH activity was found to be located mainly in mitochondria. The complete loss of NADH-dependent activity of GDH by EDTA pretreatment as well as the moderately inhibiting action of various ade-

nylates can be fully reversed by adding Ca^{2+} . At low Ca^{2+} levels a pronounced substrate inhibition occurs, particularly with NH_4^+ and NADH. Hydride transfer from NADH to the substrate takes place from the 4B position. STEWART and RHODES (1977a) confirmed that only a single form of GDH exists in *L. minor* plants grown in nitrate and ammonium. The 5-fold increase of GDH activity within 3 days after transfer of *L. gibba* plants to a high ammonium concentration (15 mM) is independent of the presence of EDTA and is due to de novo synthesis on 80 S ribosomes as shown by experiments with inhibitors of protein synthesis and with deuterium oxide-labelling technique (SHEPARD and THURMAN 1973). In *L. aquinoctialis* DUKE and KOUKKARI (1977) found GDH activity in the chloroplast-rich fraction to be higher than in the mitochondria-rich fraction (in control plants and - to a higher degree - in plants grown in glucose). In plants grown in sucrose GDH activity was lower especially in the chloroplast-rich fraction.

Changing over from the glutamine to the glutamate pathway of ammonium assimilation may also be induced by certain stress factors. Addition of NaCl (20-80 mM) to cultures of *L. minor* decreases the activities of GS, GOGAT, and nitrate reductase, but slightly increases the activity of GDH (HUBER 1982). Finally, it should be mentioned that - in principle - GDH can display NAD-dependent activity, i.e., catalysis of glutamate deamination. TEIXEIRA and DAVIES (1974) suggested that at N-starvation glycolysis is inhibited (by high pyridoxal phosphate) thereby restricting the supply of carbohydrates to the Krebs cycle. Under such conditions GDH should be blocked to prevent the loss of organic nitrogen by deamination of glutamate. In fact, the level of pyridoxal phosphate (inhibiting GDH in pea mitochondria) increases, and the level of pyridoxamine phosphate decreases in *L. minor*, when the nitrogen supply is restricted.

Assimilation of urea and the ureid allantoin has been investigated with *S. punctata* (BOLLARD 1966, BOLLARD et al. 1968, BOLLARD and COOK 1968, COOK 1968). Under the conditions used, plants can grow with urea as the sole source of nitrogen provided that the pH of the medium is adjusted to <4.3. Correspondingly, urease activity appears after supply of urea or even at N-deficiency only in the case that the pH reaches <4.3. Urease activity has been confirmed by liberation of $^{14}\text{CO}_2$ from ^{14}C urea. The extent of $^{14}\text{CO}_2$ evolution was proportional to the level of urease as assayed by formation of ammonia in extracts. The results obtained suffer from the fact that nickel was added to the nutrient medium not directly

Organic S compounds which can be utilised by S. punctata as sole sources of sulfur in millimolar concentrations, are cysteine, cystine, and cysteic acid (FRASER 1974). At 0.1 mM L. gibba grows on cystine, cysteine, methionine, glutathione, and cysteic acid (ELLIS 1969).

DATKO et al. (1978b) determined quantitatively all the S-containing compounds which can be identified in L. aequinoctialis after labeling to radioisotopic equilibrium with $^{35}\text{SO}_4^{2-}$. These are inorganic sulfate, glutathione, homocyst(e)ine, cyst(e)ine, methionine, S-methylmethionine, sulfonium, S-adenosylmethionine, S-adenosylhomocysteine, cystathionine, chloroform-soluble (presumed to be sulfolipid), protein cyst(e)ine, and protein methionine. The amount of each component present in plants grown over a 3000-fold range of medium sulfate (0.32-1000 μM) was relatively constant except for inorganic sulfate.

For assimilatory sulfate reduction possibly the "free pathway" and the "bound pathway", are both used (fig. 2.46). In each case, firstly adenosine S-phosphosulfate (APS) is formed from SO_4^{2-} and ATP, catalyzed by ATP-sulfurylase. The enzyme has been extracted from L. gibba, L. aequinoctialis, and L. minor (ELLIS 1969, WILSON and KANG 1972, 1973, REUVENY 1974, BRUNOLD and SCHMIDT 1976, SCHMUTZ and BRUNOLD 1982). In the bound pathway the sulfonyl group of APS is transferred by APS sulfotransferase to a carrier and then reduced to a thiol group. Finally, this SH-group is incorporated into o-acetylserine (OAS) by cysteine synthase to form cysteine. The presence of APS sulfotransferase in L. minor has been demonstrated by SCHMIDT (1975). Activity of this enzyme is down-regulated by H_2S (BRUNOLD and SCHMIDT 1976, VON ARB and BRUNOLD 1980) and cysteine (BRUNOLD and SCHMIDT 1978). Cysteine synthase (= o-acetylserine sulphydrylase) has been found in L. minor (BRUNOLD and SCHMIDT 1976, VON ARB and BRUNOLD 1980) and L. aequinoctialis (THOMPSON et al. 1982a). The enzyme is not affected significantly by H_2S . Coordination of S- and N-assimilation is indicated by the fact that ammonium, arginine, asparagine, and glutamine cause a 50 to 100% increase in APS sulfotransferase activity (BRUNOLD and SUTER 1984b, VON ARB et al. 1985, SUTER et al. 1986).

APS sulfotransferase and cysteine synthase both also work in the "free" pathway of sulfate assimilation. APS sulfotransferase transfers the sulfonyl group of APS to reduced glutathione or another thiol and the product reacts nonenzymatically with a further thiol releasing free sulfite. Sulfite is reduced to sulfide and cysteine synthase then catalyzes

the reaction of sulfide and OAS to form cysteine and acetate. The existence of this pathway is indicated by direct incorporation of H_2S into cysteine without preceding oxidation of H_2S to SO_4^{2-} (BRUNOLD and ERISMANN 1975). Besides the enzyme reacting with OAS another cysteine synthase is present in L. minor which uses serine as substrate (BRUNOLD and ERISMANN 1972, 1975). Occurrence of sulfite reduction to H_2S has been demonstrated in L. valdiviana, S. punctata, L. gibba (TAKEMOTO and NOBLE 1984, TAKEMOTO et al. 1986), and L. minor (BRUNOLD and ERISMANN 1976).

H_2S and SO_4^{2-} are assimilated not only using the last steps of "free pathway", but also after oxidation to sulfate. Fumigation of L. minor with 6 ppm H_2S leads to a pronounced increase in internal sulfate concentration (BRUNOLD and ERISMANN 1974). Among other things sulfide may serve as an accessory electron donor in photosynthesis (KNOBLOCH 1966, ERISMANN et al. 1967). Atmospheric SO_2 is oxidized very effectively to SO_4^{2-} in L. minor as indicated by isotope competition experiments (BRUNOLD and ERISMANN 1976). Under the conditions used, all the SO_2 seems to be incorporated into proteins and sulfolipids following oxidation to SO_4^{2-} and subsequent reduction inspite of the presence of sulfite reductase. SCHAEFER et al. (1975) observed an increase of internal SO_4^{2-} level after fumigation with 0.3-0.6 ppm SO_2 . DATKO and MUDD (1984b) mentioned that even cystine is metabolized to sulfate in L. aequinoctialis.

Selenate and selenite are also assimilated by duckweeds. S. punctata absorbs and metabolizes selenite more readily than selenate (BUTLER and PETERSON 1967). After application of $^{75}\text{SeO}_3^{2-}$ or $^{75}\text{SeO}_4^{2-}$ selenocystine and oxidation products predominate in the soluble fractions, but selenomethionine and its oxide predominate in the protein hydrolysate.

2.5.7. Amino acids, proteins, and nucleic acids

2.5.7.1. Biosynthesis of amino acids

Principally, the amino groups of amino acids are provided by transaminase reactions. Two of these enzymes have been investigated in *W. arthimiza* (APPENROTH et al. 1986). Under continuous light, both alanine aminotransferase and aspartate aminotransferase exhibit a distinctly higher activity under blue, than under red or even white light. A short blue light pulse (10 min), given after darkness, immediately increases the activity of both enzymes.

Pathways and regulation of amino acid biosynthesis have been investigated mainly within the aspartate family. Thus, pathways are reviewed in the following order:

- (a) Aspartate - lysine
- (b) Aspartate - homocysteine - methionine
- (c) Homoserine - threonine - isoleucine
- (d) Pyruvate - valine - leucine
- (e) Other amino acids

2.5.7.1.1. Lysine

Experiments with ^{14}C -labeled aspartic acid, acetate, and alanine revealed that biosynthesis of lysine proceeds via aspartate (VOGEL 1959). The first step of this pathway, the phosphorylation of aspartate, is catalyzed by aspartate kinase. The enzyme exhibiting feedback inhibition by lysine and higher concentrations of threonine (but not homoserine, methionine, or isoleucine) has been isolated from *L. minor* (WONG and DENNIS 1973). Through the pathway from 3-aspartylphosphate to lysine meso- α,ϵ -diaminopimelic acid is the last intermediate. Decarboxylation of this compound is exerted by diaminopimelate decarboxylase, which has been extracted and characterized from *L. aquinoctialis* (SHIMURA and VOGEL 1966) and *S. punctata* (VOGEL and HIRVONEN 1971).

2.5.7.1.2. Homocysteine, methionine

The pathway from aspartate to methionine includes the following steps: aspartate \rightarrow 3-aspartylphosphate \rightarrow 3-aspartate semialdehyde \rightarrow homo-




Fig. 2.47. Methionine biosynthesis in *Lemna* (after THOMPSON et al. 1982a). The solid arrows represent the major pathway. The dashed arrows represent an alternative pathway which, however, seems not to be active in vivo (see text). Also indicated are the metabolic sites where lysine plus threonine (Lys+Thr), AVG, and propargylglycine block methionine formation.

serine \rightarrow o-phosphohomoserine (OPH) \rightarrow cystathionine \rightarrow homocysteine \rightarrow methionine. Corresponding to this line ^{14}C aspartate and ^{14}C homoserine are incorporated into methionine in S. polyrhiza (MALEK and COSSINS 1982). In L. aequinoctialis, inhibition of aspartate kinase by the combined supply of lysine and threonine prevents biosynthesis of OPH (fig. 2.47) and consequently the reaction of OPH with cysteine to cystathionine (DATKO and MUDD 1982). The step of OPH + cysteine to cystathionine is catalyzed by cystathionine- γ -synthase, an enzyme which is down-regulated in vivo by methionine (THOMPSON et al. 1982a, GIOVANELLI et al. 1985a) and inactivated irreversibly in vivo and in vitro by propargylglycine (PAG), vinylglycine, and cysteine (THOMPSON et al. 1982b). The inactivation is prevented (PAG, vinylglycine) or reduced (cysteine) by 40 mM OPH or o-succinyl-L-homoserine. Experiments with the inactivator PAG have shown that enzyme activity can be reduced to 16% without an effect on growth and methionine biosynthesis. When the cystathionine- γ -synthase activity is decreased to 9% by PAG treatment, the sublethal methionine deprivation which appears in the plants is abolished within two days on the grounds of an adaptation process (THOMPSON et al. 1983). Cystathionine- γ -synthase is located in the chloroplasts. The effect of inorganic phosphate in this cell organelle, which inhibits the enzyme in vitro, may be diminished by OPH (GIOVANELLI et al. 1986).

o-phosphohomoserine sulphydrylase, which combines OPH with sulfide directly to homocysteine (fig. 2.47), shows the same dependence on methionine and inhibitors as cystathionine- γ -synthase. Both activities are probably caused by one enzyme (THOMPSON et al. 1982a, 1982b). The affinity of OPH sulphydrylase for sulfide is relatively low in comparison to the S^{2-} -affinity of o-acetylserine sulphydrylase, i.e., cysteine synthase. Physiologically, the OPH sulphydrylase activity seems to be of no importance. Short term labeling experiments with $^{35}\text{SO}_4^{2-}$ revealed that radioactivity is incorporated predominantly in cystathionine (MACNICOL et al. 1981). The authors concluded that 85-100% of homocysteine biosynthesis proceeds through the transsulfuration pathway (correcting an earlier statement of DATKO et al. 1977).

The next step in methionine biosynthesis, the cleavage of cystathionine into homocysteine and pyruvate + NH_3 , is impaired by L-aminoethoxyvinylglycine (AVG). [^{35}S] cystathionine accumulates in L. aequinoctialis, when the medium contains $^{35}\text{SO}_4^{2-}$ and AVG (DATKO and MUDD 1982). During the final step of methionine biosynthesis methionine sulfoxide is

used for methylation of homocysteine. The precursor of methyl tetrahydrofolate, 5,10-methylenetetrahydrofolate, is provided by serine hydroxymethyltransferase. Activity of this enzyme is partially inhibited in L. minor, when lysine is added to the culture medium (WONG and COSSINS 1976).

2.5.7.1.3. Threonine, isoleucine

Biosynthesis of threonine from aspartate follows the methionine pathway till OPH. Then OPH reacts with H_2O catalyzed by threonine synthase. Five further steps are needed to transform threonine into isoleucine. Validity of this pathway in S. polyrhiza has been shown by ^{14}C aspartate- and ^{14}C homoserine-incorporation into threonine and isoleucine (MALEK and COSSINS 1982). Threonine synthase has been isolated from L. minor (SCHNYDER et al. 1975) and L. aequinoctialis (VELUTHAMBI et al. 1983, GIOVANELLI et al. 1984). The investigations with L. aequinoctialis revealed a reversible stimulation (over 30-fold) of threonine synthase by S-adenosylmethionine (SAM) and a down-regulation of the enzyme by methionine. As SAM is the first subsequent product of methionine in the methionine cycle (see chapter 2.5.7.2), it could be the signal of methionine overproduction which switches the pathway from methionine to threonine production. Methionine, on the other hand, may help to limit the overproduction of threonine. Threonine and also isoleucine have no feedback or repressing effect on threonine synthase. Inorganic phosphate and AMP are further inhibitors of threonine synthase, but OPH counteracts these effects (GIOVANELLI et al. 1986). If total OPH is restricted to the chloroplasts, in which threonine synthase is located, then the P_i concentration in this organelle does not limit the production of threonine.

The second step of the pathway from threonine to isoleucine, the trans-formation of 2-oxobutyrate into 2-acetohydroxybutyrate, depends on the activity of acetohydroxy acid synthase (fig. 2.48). This enzyme as well as the enzymes catalyzing the subsequent steps of isoleucine biosynthesis also work in valine-leucine biosynthesis. Therefore, it does not seem to be surprising that valine and leucine exert an inhibition of acetohydroxy acid synthase. The end-product inhibition has been deduced from the fact that 2-aminobutyrate, a side-chain product of 2-oxobutyrate, is accumulated in S. polyrhiza after addition of valine to the

nutrient medium (BORSTLAP 1972). The content of isoleucine decreases under these conditions as is the case after leucine supply.

2.5.7.1.4. Valine, leucine

Aceto-hydroxy acid synthase catalyzes, in addition to the above cited reaction, the transformation of pyruvate to 2-acetolactate, the first intermediate of valine-leucine biosynthesis (fig. 2.48). The effectivity of in vivo control of this enzyme has been investigated by isotope dilution experiments (BORSTLAP and VERNOOY-GERRITSEN 1985). An increase in the internal valine pool of *S. polyrrhiza* from 3.2 to 6.6 nmol mg⁻¹ dry weight reduced valine synthesis by 70%, but not isoleucine synthesis. From these and other results it was concluded that feedback inhibition of aceto-hydroxy acid synthase by valine and leucine can be exerted in such a way that synthesis of 2-acetolactate is appreciably reduced, whereas synthesis of 2-aceto-hydroxybutyrate (the isoleucine precursor) is not inhibited.

Another control point in the valine-leucine pathway is 2-isopropylmalate synthase, which switches the final intermediate of valine synthesis to the leucine pathway. BORSTLAP (1972, 1981) showed that activity of this enzyme is inhibited by isoleucine. The content of leucine in *S. polyrrhiza* decreases, but the content of valine increases, when isoleucine is added to the culture solution. Growth inhibitions caused by valine, leucine, isoleucine, valine + leucine, or leucine + isoleucine (BORSTLAP 1970) can be interpreted by deprivation of one or two of the afore-named branched-chain amino acids via inhibition of aceto-hydroxy acid synthase and/or isopropylmalate synthase (BORSTLAP 1981).

Experiments with ¹⁴C-labeled valine have shown that plants can convert valine to leucine (BORSTLAP 1975). Thus the last step of valine synthesis is reversible. Growth-inhibiting concentrations of valine increase the level of free leucine in the plant even thirty-fold (BORSTLAP 1972).

2.5.7.1.5. Other amino acids

As known for other plants, arginine metabolism is blocked by canavanine in *L. minor* (NICKELL 1956). The growth inhibition caused by 10 ppm canavanine can be overcome by 50 ppm arginine or 100 ppm citrulline. Biosynthesis of aromatic amino acids is inhibited by the herbicide glyphosate




Fig. 2.48. Biosynthetic pathway of branched-chain amino acids (after BORSTLAP and VERNOOY-GERRITSEN 1985).
TD = threonine deaminase; AHAS = aceto-hydroxy acid synthetase; IPMS = 2-isopropylmalate synthetase.

Gl, but strain 6746 exhibits only the first of the two peaks of ethylene evolution, and strain Gl only the second one.

Exogenous ACC is taken up by L. minor in the dark with no saturation (FUHRER 1985). ACC-dependent ethylene production, on the other hand, is saturated above 10 μ M ACC (equivalent to 32 μ M internal ACC). Only when 1 μ M or more ACC is supplied to light-grown plants, increasing CO₂ concentrations elevate the ethylene emanation. This effect seems to be based on a CO₂ inhibition of ethylene metabolism, as internal levels of ACC and MAACC are not influenced by the CO₂ treatment.

When toxic concentrations of copper (10^{-4} - 10^{-3} M) cause a damage of intracellular membranes in S. punctata, not ACC but other sources are responsible for the increase in ethylene evolution (MATTOO et al. 1984c, 1986a). The ¹O₂ quenchers and inhibitors of lipid peroxidation, viz. propyl gallate, 1,4-diazobicyclo [2,2,2] octane and selenomethionine, all markedly inhibit the Cu(II)-induced ethylene production. Selenomethionine also suppresses the membrane damage in Cu(II)-treated plants. The authors suggest that copper stimulates lipid oxidation by reacting with lipid peroxides to form new radical species. Both linoleic acid and methionine seem to act as precursors of ethylene after copper poisoning.

2.5.7.3. Biosynthesis and turnover of proteins

2.5.7.3.1. Dependence on light

The ability of several Lemnaceae to grow in complete darkness when provided with sucrose (see chapter 2.3.5.4) demonstrates that light is not needed for biosynthesis of at least most proteins. As early as 1896 and 1899 HANSTEEN found protein accumulation in darkness, when plants of L. minor obtained a certain combination of N source and carbohydrate. Asparagine, urea, or ammonium salt were effective as N source in combination with glucose, and urea or glycine were effective in combination with sucrose. A special case, however, are proteins of the photosynthetic apparatus. For economy, development of chloroplasts is completed only in the presence of light (see chapter 2.4.1.5). The bulk proteins of this cell organelle are especially affected. When L. gibba plants are transferred to darkness 19-21 hours, a dramatic decrease in biosynthesis of chlorophyll a/b-binding protein and the large and small subunits of ribulose-1,5-diphosphate carboxylase (RuDPCase) occurs (TOBIN 1978, 1981a, 1981c,

TOBIN and SUTTIE 1980). Re-lightening causes a strong increase in synthesis of RuDPCase subunits within 1-3 hours. In L. minor light stimulated amino acid incorporation into large subunit of Fraction I protein (=RuDPCase) and two plastid membrane proteins (BLACKWOOD and LEAVER 1977). In addition the assembly of large and small subunits of Fraction I protein is promoted by light. In S. punctata, ³⁵S methionine incorporation into soluble and membrane proteins (including a 32kDa protein) is enhanced especially 3 hours after dark-light transition (REISFELD et al. 1978a). Synthesis of a 33.5 kDa precursor of 32 kDa protein and its conversion are also light-dependent (EDELMAN and REISFELD 1978, 1980, MATTOO et al. 1980). Further proteins greatly increasing during greening of etiolated plants are the chlorophyll protein complex I, subunits of the ATPase complex, cytochrome f, and a putative cytochrome b₆ subunit (DE HEIJ et al. 1984).

In all cases investigated so far, the light effect is exerted primarily through increased amounts of translatable messenger RNAs. This holds true for the small subunit precursor of RuDPCase (TOBIN 1978, TOBIN and SUTTIE 1980), a 32kDa protein being the precursor of light-harvesting chlorophyll a/b-protein (TOBIN 1981a, 1981c), and another 32 kDa protein (REISFELD et al. 1978a) identified later on as shield protein for the electron acceptor II of photosynthesis (see chapter 2.7.1.1). In the case of small subunit of RuDPCase and the light-harvesting chlorophyll a/b-protein the light effect on mRNA content of plants is mediated by phytochrome (TOBIN 1981a, 1981b, STIEKEMA et al. 1983a). This result should be related to the fact that both proteins are encoded by nuclear genes and synthesized from polyadenylated RNA on cytoplasmic ribosomes as precursor polypeptides which are transported into the chloroplasts (TOBIN 1978, STIEKEMA et al. 1983a). The phytochrome effect on both messenger RNAs can be imitated or magnified by benzyladenine (FLORES and TOBIN 1986b). On ground of their experiments the authors suggest that BA exerts this effect at a post-transcriptional level, possibly by affecting the stability of the RNAs.

In the case of 32 kDa shield protein blue light is >5 times as effective as red light in stimulating the appearance of competent mRNA (GRESSEL 1978, MATTOO et al. 1980). Phytochrome is not involved in this light effect. It is interesting to note that the mRNA (5x10⁵ dalton) was isolated from plastids and does not contain poly(A) sequences (ROSNER et al. 1975a).

In addition to the light effects on mRNA content a light-dependence of translation has been found for the chlorophyll a/b-apoproteins (SLOVIN and TOBIN 1982). Under conditions of heterotrophic growth with intermittent red light, formation of mRNA and protein synthesis are uncoupled. Whereas the level of translatable mRNA is essentially the same as that found in plants grown in white light (TOBIN 1981b), the incorporation of [³⁵S] methionine into the apoproteins is very low. Possibly, the necessity of such an additional light effect on translation or processing also holds true for RuDPCase (TOBIN unpubl., cited in SLOVIN and TOBIN 1980).

2.5.7.3.2. Proteins of the chloroplast

Major protein components of thylakoid membranes are the two apoproteins of the light-harvesting chlorophyll a/b-protein complex. In *L. gibba* molecular weights of these apoproteins are 28000 and 28500, respectively (TOBIN 1981a, 1981c). The precursor of the apoproteins possesses a molecular weight of 32000 and is synthesized on cytoplasmic ribosomes with a poly(A) mRNA (TOBIN 1981a, 1981c, SLOVIN and TOBIN 1982). Chlorophyll a/b-protein is quite stable, as it is not degraded to an appreciable amount during a four-day dark treatment (TOBIN 1981c). The mRNA for the 32 kDa precursor, however, declines sharply during the first two days of darkness. Kinetin (3×10^{-6} M) can retard the degradation to some extent (TOBIN 1981a, TOBIN and TURKALY 1982). In the presence of glyphosate [N,N-bis(phosphonomethyl)glycine] synthesis of chlorophyll a/b apoprotein is extremely low compared to control plants, whereas the large subunit of RuDPCase is synthesized further (SLOVIN and TOBIN 1981).

A reversible phosphorylation of the light-harvesting pigment-protein complex of photosystem II (PS II) has been found in *L. aequinoctialis* (STEINBACK and WATSON 1981). Excitation of PS II by red light in vivo causes a phosphorylation of PS II and in this way an increased energy transfer to PS I at the expense of PS II. Excitation of PS I by far red results in a rapid dephosphorylation ($T_{1/2} = 4$ min) of PS II. Such a regulation of energy distribution between the photosystems via protein phosphorylation may be of ecological relevance in plants, shaded variably by leaves.

Another membrane protein of chloroplasts is the 32 kDa shield protein regulating electron transport (see chapter 2.5.1.1.2). Occurrence of

this protein has been demonstrated for *S. punctata*, several other angiosperms, and the alga *Chlamydomonas* (HOPFMANN-FALK et al. 1982). The shield protein contains no lysine (EDELMAN and REISFELD 1978, 1980, REISFELD et al. 1982) and is tightly bound to the thylakoids (EDELMAN and REISFELD 1978, 1980). Its half-life is relatively short. Decay proceeds at least one order of magnitude more rapidly than that of other major plastid polypeptides (EDELMAN and REISFELD 1980). Synthesis and processing take place within the chloroplast. Three hours after illumination with white light there is a burst in the synthesis of plastid mRNA coding for the 33.5 kDa precursor of shield protein (REISFELD et al. 1978a,b, EDELMAN and REISFELD 1980). The mRNA is characterized by a molecular weight of 50000 and the lack of poly(A). Translation on chloroplast ribosomes is confirmed by suppressing the process with low concentrations of D-threo-chloramphenicol (WEINBAUM et al. 1979a,b). Transcription of the 33.5 kDa precursor into the 32 kDa protein commences only after completion of the precursor polypeptide chain and insertion into the thylakoids (REISFELD et al. 1982). Processing of the 33.5 kDa protein, like synthesis, is dependent on light and takes place with a $t_{1/2}$ of ~3 min (EDELMAN and REISFELD 1980, MATTOO et al. 1980). Light is effective only in green plants, i.e., in the presence of differentiated thylakoids. In etiolated plants shield protein does not appear before 2 or 3 days of greening (MATTOO et al. 1980).

The photosystem I reaction center isolated from *S. punctata* contains seven different subunits (NECHUSHTAI et al. 1981). Inhibitor experiments with D-threo-chloramphenicol and cycloheximide revealed that subunit II is synthesized on cytoplasmic ribosomes, whilst subunits I, V, and VI are translated within the chloroplast. In the same way the origin of the 8 subunits of the proton-ATPase complex were investigated. Subunits γ and δ of CF₁ and subunit II of CF₀ are products of the cytoplasmic translation system. Subunits α , β , and ϵ of CF₁, as well as subunit III of CF₀ are synthesized on the chloroplast translation system. Protein synthesis at the two cellular locations is strictly coordinated, as the molar ratio of subunits produced (see CF₁- β and CF₁- δ) resembles the stoichiometrical amounts in the native protein complex (DE HEIJ et al. 1984).

The main polypeptides in the soluble part of chloroplasts are the large and small subunits (LS and SS) of ribulose-1,5-diphosphate carboxylase. Within Lemnaceae both subunits show some variation in the polypeptide

composition (CHEN and WILDMAN 1981). After electrofocussing four different types of the LS polypeptide clusters, and eight individual SS polypeptides have been found. Each of the 11 investigated species (representing the four genera) contains one type of LS and one, two, or four types of SS polypeptide. In *L. gibba* LS and SS are quite stable, as they do not show any decline during 4 days in the dark (TOBIN and SUTTIE 1980).

Messenger RNA for LS-RuDPCase is located in the chloroplast, contains no poly(A) sequences, and has a molecular weight range of $0.56-0.7 \times 10^6$ (ROSENER et al. 1976, REISFELD et al. 1978b,c). The poly(A) mRNA for the precursor of SS-RuDPCase has been isolated from cytoplasmic polysomes (TOBIN 1978). The rapid decline of this mRNA in *L. gibba* during four days in the dark can be diminished somewhat by supply of kinetin (TOBIN and TURKALY 1982).

2.5.7.3.3. Other proteins

Protein phosphorylation occurs not only in the photosystem II (see above), but also in ribosomes and chromatin. TREWAVAS (1973) labeled *L. minor* plants with $^{32}\text{P}_4$ and found radioactivity in the small subunit of ribosomes. After acid hydrolysis of the ribosomal protein serine phosphate appeared. The P-content per ribosome amounts to 0.75 atom in control plants, but only 0.36 atom of phosphorus in plants treated for 24 hours with $5 \mu\text{M}$ ABA. The enzyme catalyzing the phosphorylation of ribosomal proteins, protein kinase (ATP: protein phosphokinase), sediments with ribosomes during sucrose density gradient centrifugation. Seventy per cent of the enzymic activity dissociates from the ribosomes in 0.3M KCl, but the remaining protein kinase remains firmly bound in 0.7 KCl (KEATES and TREWAVAS 1974).

Phosphorylation of chromatin-associated proteins has been investigated in *L. aequinoctialis* (VAN LOON et al. 1975, CHAPMAN et al. 1975). After labeling of plants with inorganic ^{32}P , chromatin were prepared and then the associated proteins separated by gel electrophoresis. At least 15-20 proteins (with molecular weights ranging from 10000 to 100000) were found to be phosphorylated. Most of them are acidic proteins, but 25-30% may be histone material. Incubation of isolated nuclei with ^{32}P - ATP resulted also in labeling of chromatin proteins indicating the presence of an endogenous kinase. Partial acid hydrolysis of labeled proteins

yielded serine phosphate and traces of threonine phosphate. Pretreatment of plants with ABA (10 mg/l) causes substantial alterations in the labeling of three protein bands. There is an almost total loss of one protein band and an increase in labeling of two other protein bands.

Three soluble protein kinases (P I, P II, and P III) phosphorylating histones in vitro have been isolated from *L. aequinoctialis* by KATO et al. (1983, 1984). Activity of the kinases (with molecular weights of 165000, 85000, and 145000, respectively) is dependent on cyclic nucleotides in a different manner. P I is partially inhibited by cAMP, cGMP, or cIMP, while the P II enzyme is activated by these nucleotides. P III is not influenced by cAMP, but slightly inhibited by cCMP and cUMP. The three protein kinases are Mg-dependent. The P II enzyme, however, is stimulated more effectively by high concentrations of Co^{2+} , and the P III activity is maximum with Mn^{2+} . Cytokinins exert some inhibition in P III activity. At present, the in vivo functions of the three protein kinases are not known. The same holds true for a membrane-associated protein kinase, which has been isolated from endoplasmic reticulum-rich fractions (KATO and FUJII 1985). The enzyme (molecular weight 220000) shows optimum activity at pH 9.0-9.5 and is stimulated by Co^{2+} , Mg^{2+} , and Mn^{2+} . Cytokinins and cyclic nucleotides do not affect the activity.

Glycoproteins have been claimed to occur in *L. minor*, as the amino sugar D-[1- ^{14}C] glucosamine is incorporated into compounds of high molecular weight that resemble proteins in their physical properties (ROBERTS et al. 1971).

2.5.7.3.4. Protein metabolism in general

Growth and development are based mainly on synthesis and degradation of proteins. Thus it is not surprising that growth-regulating factors influence the overall protein metabolism. Addition of glucose to cultures of *S. punctata*, for example, accelerates the production of water-soluble and water-insoluble proteins (GROB et al. 1973), doubtlessly in parallel to the increased growth rate. Ammonium, more specifically, increases the protein content of *L. minor* plants per dry weight, reversibly. In translocation experiments from nitrate to ammonium the total protein rises by 25-30% within 40 hours. Re-transfer of plants to nitrate restores the original "steady state" value of proteins within 30-40 hours (DICHT et

al. 1976). The ammonium effect may come about through an enhancement of the endogenous cytokinin and auxin level. An increase in endogenous cytokinins after transition of plants to ammonium was found in *L. aquinoctialis* by GRUNTZEL (1982). Supply of kinetin or IAA, on the other hand, to cultures of *L. minor* increases the protein content per dry weight (KRZECZOWSKA and ZIMNA 1972, MACIEJEWSKA-POTAPCZYK et al. 1976). With kinetin especially the high molecular proteins are promoted whereas the level of mean and low molecular proteins decreases. IAA slightly decreases the level of high molecular proteins and increases the content of mean and low molecular proteins. A differential effect on protein synthesis has been demonstrated in detail for ABA (SMART and TREWAVAS 1984a). In connection with the change in developmental processes (turon formation) ABA changes the pattern of proteins seriously. Two-dimensional polyacrylamide gel electrophoresis of in vivo ³⁵S-methionine-labeled soluble proteins revealed that the majority of proteins decreases or disappears during the development of the turion, whilst some show increased labeling and others appear de novo. Also in the turion-producing mother frond certain changes of protein pattern are induced. ABA seems to exert its effects on the transcriptional as well as on the translational level. The rapid general inhibition of protein synthesis during the first two days of ABA action should be initiated at the translational level, as no decrease in translatable RNA occurs during this time. Induction of de novo synthesis of new proteins, however, could come about by changes in gene expression.

Evaluation of the rates of overall protein synthesis and degradation with labeled amino acids are complicated by the fact that at least two pools of amino acids exist in the plant, only one of which contributes to the synthesis of protein. TREWAVAS (1972a) used the trna-bound ³H-methyl-methionine for calculation of the flux rates of methionine into and out of protein and reached the following results: plants of *L. minor* cultivated under certain light and temperature conditions exhibit rate constants for protein synthesis and degradation of 0.48 and 0.09 days⁻¹, respectively, when grown in complete nutrient medium. Reduction of the MgSO₄ content of medium to 1/10 results in a decrease of protein synthesis and an increase in protein degradation. The rate constants for both processes then are 0.32 and 0.13 days⁻¹, respectively. Rate constants for fresh weight increase in these experiments (0.37 days⁻¹ for full-strength medium, and 0.18 days⁻¹ for 1/10 MgSO₄ medium) are in close re-

lation to the net production of proteins. Similar investigations with *S. polyrrhiza* have been carried out by MALEK and COSSINS (1983b). In this case, lowering of nitrate concentration to 1/20 does not affect the rate of protein synthesis, but increases the rate constant of protein degradation. Further work on protein turnover has already been reviewed in connection with the physiology of senescence and rejuvenation (see chapter 2.4.1.4.). Especially the effects of several stress factors and of plant growth regulators have been represented in some detail. The question which properties of proteins may be of significance for a selective protein degradation have been investigated in *L. minor* (COATES and DAVIES 1983). The authors found only a weak correlation between SH content of proteins and degradation, and little or no correlation between molecular weight or charge of proteins and degradation. They suggested that half-life of proteins is determined by the sum of many physical properties rather than by a single "signal" property.

2.5.7.4. RNA metabolism

2.5.7.4.1. Ribosomal RNA

Nucleic acid fractionations from duckweed material were done by HEMLEBEN (1972). After short labeling of *L. aquinoctialis* plants with [³H] uridine, nucleic acids were separated using MAX-column chromatography or polyacrylamide-gel electrophoresis. Fraction I (4 S rRNA) and fraction II (5 and 5.8 S) contained low-molecular RNA components, fraction III was DNA, fractions IV and V were high-molecular ribosomal (r) RNAs, and fraction VI contained components being presumably precursors of rRNA. Fraction IV was the 16 S rRNA of chloroplast and the 18 S rRNA of cytoplasm (with a molecular weight of 0.7x10⁶). Fraction V held the 23 S rRNA of chloroplasts and the 25 S rRNA of cytoplasm (MW of 1.3x10⁶). The components of fraction VI were separated further and had sedimentation constants (molecular weights) of 33-34 S (2.4 M), 29 S (1.7 M), and 27 S (1.4 M). Rates of [³H] uridine incorporation into single RNA fractions could be altered by certain environmental factors (photo-period, blue light, supply of sucrose, or phosphate deficiency). Most interestingly, transfer of plants from continuous light to short day caused - besides other reactions - an increase in the ratio of fractions VI/IV+V especially in the first half of the dark phase. Interruption of the dark phase

by light (30 min, 8 hours after the beginning of darkness) prevented this increase.

TREWAVAS (1970) determined rate constants of synthesis and degradation of cytoplasmic rRNA (25 S and 18 S) and chloroplast rRNA (23 + 13.7 S and 16 S) in mixotrophic cultures of *L. minor*. Half-life of both cytoplasmic rRNA components amounts to 4.3 days. Half-life of the chloroplast rRNA components is considerably longer and amounts to 15.5 days. The degradation rate for total rRNA increases after omission of either nitrate, phosphate, calcium, or magnesium. Transfer of plants to water causes an increase of rRNA degradation and a decrease of rRNA synthesis, but ABA reduces only rRNA synthesis.

Processing of presumptive precursors into mature rRNA has been followed in *S. punctata* by the time-course of [^3H] uridine incorporation into different RNA fractions and by inhibition of certain steps with darkness, transfer of plants to distilled water, or chloramphenicol (ROSNER et al. 1973, POSNER and ROSNER 1975, ROSNER et al. 1977a). After short labeling of plants two cytoplasmic RNAs with high molecular weight (2.8 M and 2.3 M) appear. The 2.8 M species seems to be cleaved to form 2.3 M RNA and "excess" RNA with a molecular weight of 0.5 M. The "excess" RNA is not methylated and may be discarded during the further metabolism (POSNER et al. 1974). The 2.3 M RNA then is processed to a 1.4 M intermediate and finally the light rRNA (0.7 M) and the heavy rRNA (1.3 M) are formed. Darkening of plants or incubation of plants on distilled water lead to an accumulation of the 2.3 M precursor. During prolonged darkening, however, a reduction of transcription and therefore a re-adjustment of transcription and processing takes place in the newly-produced, colorless fronds.

In chloroplasts precursors with 2.7 M, 1.2 M, 0.7-0.75 M and 0.5 M are labeled more rapidly than the mature rRNAs. The mature species with 1.05-1.1 M and 0.56 M are detectable only after labeling for more than two hours. Chloramphenicol treatment of plants inhibits processing of chloroplast rRNA, as incorporation of radioactivity into the mature rRNAs is inhibited, but the labeling of the above named precursors is enhanced. Darkening of plants also seriously retards processing of chloroplast rRNA. In such plants especially the 1.2 M, 0.75 M, and 0.5 M species accumulate. Supply of sucrose to the cultures can diminish not prevent the effect of darkening (on chloroplast rRNAs as well as on cytoplasmic rRNAs). The ability to synthesize high-molecular-weight rRNA

is retained to some degree in isolated chloroplasts of *W. arrhiza* (EICHORN and AUGSTEN 1984).

In addition to the high-molecular-weight rRNA species described above, ribosomes contain low-molecular-weight rRNAs, which are a further part of the large ribosomal subunit. Cytoplasmic ribosomes possess rRNA species of 5 S and 5.8 S, chloroplast ribosomes have rRNAs of 5 S and 4.5 S. Using *L. minor* as plant material complete nucleotide sequences were elaborated for the two 5 S rRNA species (DYER and BOWMAN 1976, 1979). The cytoplasmic 5 S rRNA has 118 nucleotides and a mol. wt. of 37800, the chloroplast 5 S rRNA contains 121 nucleotides and has a mol. wt. of 39300. Only few deviations in nucleotide positions were found when the sequences were compared with the respective 5 S rRNAs from other flowering plants. Comparison one with another, however, revealed that about 50% of the residues are different at equivalent positions in the two. There is a remarkably greater similarity between the chloroplast 5 S rRNA of *Lemna* and that of the prokaryote *Anacystis nidulans*. A detailed discussion of the results yielded a strong support of the endosymbiont hypothesis of chloroplast origin. The possible secondary structure of chloroplast 5 S rRNA is shown in fig. 2.50. The cytosol 5 S rRNA, although dissimilar in sequence from that of the chloroplast, may also be folded in the same way. The shape of both molecules seems to be essential for their function.

The chloroplast 4.5 S rRNA from *L. minor* has been fractionated by gel




Fig. 2.50. Possible secondary structure of *Lemna minor* chloroplast 5 S rRNA (DYER and BOWMAN 1979).

electrophoresis into three variants (BOWMAN and DYER 1979). The major component (70-80% of the total yield) contains 103 nucleotides and has a molecular weight of 33600. The two smaller, less-prevalent, variants contain 96 and about 63 nucleotides respectively. All the 4.5 S rRNA species have the same A-A-C_{OH} sequence at the 3'-end of the molecule; the 5'-end is not phosphorylated and apparently none contains any modified nucleotides. The 4.5 S rRNA seems to be synthesized as a segment of the 2.7 M precursor. It has been hypothesized that this rRNA may represent transcribed spacer RNA with a possible function in assembling of ribosomal components.

Density of the distribution of chloroplast and cytoplasmic ribosomes has been evaluated by electron microscopy in *S. punctata* (ROSNER et al. 1974a). In principle, density is considerably higher in light-grown than in dark-grown plants, but chloroplast ribosomes disappeared from the stroma in adult fronds. As the relative amount of chloroplast rRNA increases in adult fronds, chloroplast ribosomes seem to move from the stroma to the thylakoid lamellae. At this stage the heavy chloroplast rRNA loses its integrity and shows thermal instability. Cytoplasmic polyribosomes have sizes up to octamers (free polysomes) or heptamers (membrane-bound polysomes) in light-grown plants of *S. punctata* (VAN EE and PLANTA 1982). In dark-grown plants pentamers (free polysomes) and trimers (membrane-bound polysomes) are the largest polyribosomes.

2.5.7.4.2. Poly(A) RNA

COVEY and GRIERSON (1976) detected poly(A) sequences in the RNA of *L. minor* by hybridisation with [³H] polyuridylic acid. The poly(A) content of total RNA determined in this way amounted to 50 ng/100 µg RNA. Isolation of polyadenylated RNA from total RNA was achieved by affinity chromatography on oligo(dT)-cellulose or poly(U)-sepharose 4 B (TOBIN and KLEIN 1975, EICHHORN and AUGSTEN 1983b). The messenger RNA capacity of the separated poly(A) RNA from *L. gibba* was confirmed by its ability to serve as template in a cell-free translation system derived from wheat germ (TOBIN and KLEIN 1975). Poly(A) RNA from *W. arthiza* amounts to 5.0% of total RNA, when plants were cultivated under mixotrophic conditions in a chemostat and illuminated with continuous white light. Reduction of growth rate by stronger P limitation has no influence on the poly(A) RNA portion. Under continuous blue or red light the plants contain only 3.5

and 1.5% poly(A) RNA, respectively. Darkening of plants for 24 hours diminishes the poly(A) RNA content to 1.5% after white or blue light, and 0% after red light. Two bulk proteins of chloroplasts, which are synthesized on cytoplasmic ribosomes with light-dependent poly(A) RNA, the light-harvesting chlorophyll a/b-protein and the small subunit of ribulose-1,5-diphosphate carboxylase, have been reviewed in the foregoing chapter (2.5.7.3).

2.5.7.4.3. Inhibitor studies

Addition of 8x10⁻⁵ M 2-thiouracil (TU) to the culture medium inhibits frond elongation and differentiation of aerenchyma and vascular bundles in *S. punctata* (RIMON and GALUN 1967). The effect of TU is completely counteracted by uridine or uracil. TU works through an incorporation into all RNA species, in this way preventing uracil incorporation (RIMON et al. 1969). RNA synthesis is not suppressed, even after 20 h of TU treatment, but synthesis of proteins and DNA are reduced after TU incorporation into RNA. Another inhibitor which is incorporated into all RNA species is 5-fluorouracil (FU). This substance, however, inhibits not only frond elongation and differentiation, but also cell division, and these effects cannot be counteracted by either uridine, uracil, thymidine, or simultaneous addition of uracil and thymidine. A certain concentration of FU to 5-fluorodeoxyuridine (a DNA synthesis blocker) therefore may explain the FU action on development only partially. Extremely toxic to the growth of *L. minor* are the non-protein amino acids L-carnitine, L-canaline, and o-ureido-L-homoserine (ROSENTHAL et al. 1975). They are structural analogues of the ornithine-urea cycle amino acids, and their effect on growth can be abolished to some degree by arginine, ornithine, and citrulline. The substances inhibit [³H] uridine incorporation into RNA more than the [³H] thymidine incorporation into DNA (GULATI et al. 1977). The possible involvement of polyamines, which are derived from the ornithine-urea cycle, may be considered in this context, as they inhibit RNase activity and induce DNA synthesis in other organisms.

2.5.7.4.4. Ribonucleases

RNA degrading enzymes have been isolated from *S. punctata*. From plants

grown under standard conditions three ribonucleases (RNases) were separated by DEAE-Sephadex A 25 column chromatography (LASSOCINSKI 1982). The pH optima of enzyme activity were found to be 7.9, 7.5, and 7.2, respectively. All three enzymes have an endonucleolytic mode of action and produce, during poly(A) digestion, cyclic adenosine-2':3'-phosphate, which then is hydrolysed to adenosine-3'-phosphate. One of the RNases has a preference for poly(C) compared with poly(A), the other two RNases show the opposite behaviour.

With polyacrylamide gel electrophoresis KNYPL (1977b) revealed the presence of 10 RNase bands in extracts of S. punctata. When plants were treated with growth-inhibiting doses of N,N-dimethylmorpholinium chloride (DMMC) or CCC for 3-6 days before analysis, three of the RNase bands were markedly intensified, and three new isoenzymes appeared. Additional supply of gibberellin A₃ or benzyladenine reduces the increase in RNase activity. The DMWC-induced RNase has a pH optimum of 6.2-6.3, is activated by citrate, and has a molecular weight of about 31500. The growth retardant alden is another agent which increases the activity of RNase in S. punctata (KNYPL 1977a). Fusicoccin increasing fresh weight of S. punctata by 50%, enhances activity of RNases by more than 100% (KNYPL and JANAS 1980). Ribonucleases which are produced at P-deficiency and are located at the cell surface for acquisition of phosphate, have been mentioned in chapter 2.5.6.3.

2.5.7.5. DNA structure and function

2.5.7.5.1. Chloroplast DNA

S. punctata is the organism which has been used within the Lemnaceae for characterization of chloroplast DNA (cpDNA), for physical mapping, and for sequencing of some regions. The cpDNA isolated by CsCl gradient centrifugation is circular in shape, has a buoyant density of 1.698 ± 0.002 g cm⁻³, a G+C content of 37%, and an at least bimodal melting behaviour, resembling features of many other cpDNA species. Length measurements on electron micrographs, however, revealed a contour length of 54.1 ± 1.8 μ m (corresponding to a molecular weight of 115×10^6), which is about 10 μ m longer than the length of most other cpDNA species (VAN EE et al. 1980a). Analysing the digestion products of cpDNA after treatment with restriction endonucleases the order of cpDNA fragments within the

whole molecule could be determined (VAN EE et al. 1980b, 1982). The cpDNA is composed of two invertedly repeated DNA segments of 18 Md, separated by a small single-copy region of 19 Md and a large single-copy region of 67 Md. The surplus segment, which is responsible for the greater length in comparison to other cpDNAs, lies in the large single-




Fig. 2.51. Restriction fragment map of *Spirodela punctata* chloroplast DNA, elaborated with restriction endonucleases Xho I and Sac I, and the positions of low-molecular-weight rRNA genes ($\phi = 5S$ RNA, $\phi = 4.5S$ RNA) and tRNA genes ($\phi = 4S$ RNA). From GROOT and VAN HARTEN-LOOSBROEK, 1981 (see text).

copy region. Using hybridization techniques the positions of the low-molecular-weight rRNA genes and the tRNA genes within the restriction fragment map of cpDNA were determined (GROOT and VAN HARTEN-LOOSBROEK 1981). As can be seen from figure 2.51, the genes for 5 S rRNA and 4.5 S rRNA are located in the two sequence repeats of 17 Md. Three clusters of 4 S rRNA genes are situated at 120°, 170°, and 210° within the large single-copy region, and further 4 S rRNA genes are placed around map position 270° and in the inverted repeat region. 35-40 4 S rRNA genes have been calculated to be present in the cpDNA of *Spirodela*. The overall distribution of these genes is very similar to the distribution of tRNAs on spinach and maize cpDNA. Further investigations on cpDNA gene mapping in *S. punctata* have been made by VAN EE et al. (1982). Each of the two repeat units contains one set of rRNA genes in the order: 16 S rRNA gene-spacer - 23 rRNA gene - 5 rRNA gene. Within the large single-copy region genes for a 32Kd protein, a 55Kd protein, and the large subunit of ribulose 1,5-diphosphate carboxylase have been localized. The genes for 4.5 S rRNA and 5 S rRNA, their flanking regions and the spacer between the two genes have been sequenced by KEUS et al. (1983b). The results for the 5 S rRNA gene are in agreement with the sequence analysis for 5 S rRNA in *L. minor* of DYER and BOWMAN (1979). At the 3' end of both rRNA coding regions a prokaryotic termination signal (containing a palindrome of usual length followed by an AT rich region) has been found. A possible promoter site at the 5' end of the 5 S rRNA gene has been localized, too. In a further paper KEUS et al. (1983a) have communicated the nucleotide sequences of flanking regions of the genes coding for high-molecular-weight chloroplast rRNAs (23 S and 16 S), which also are located in the 17 Md repeats of cpDNA. The region 5'-proximal to the 16 S rRNA gene contains three prokaryotic promoter motifs and - between them - a gene coding for valine tRNA. Oligonucleotide sequences in the 16 S rRNA have been determined for *Lemna* (see STACKEBRANDT and WOESE 1981) and compared with that of several other plants (SEEWALDT and STACKEBRANDT 1982). There is no specific relationship between the 16 S rRNAs of *Lemna* and *Prochloron*, a unicellular prokaryote containing chlorophylls a and b and lacking phycobiliproteins.

2.5.7.5.2. Mitochondrial DNA

The mitochondrial genome of *S. punctata* has a size of at least 250 kilo

basepairs as revealed by restriction enzyme analysis (DE HEIJ et al. 1985). Certain restriction fragments of the mitochondrial DNA (mtDNA) were cloned. Then inserts of these clones were isolated, labeled by nick-translation, and used for back-hybridization. Each insert hybridizes to nearly all restriction fragments. The authors concluded from their results that the mtDNA contains a sequence which is reiterated about 100-fold.

2.5.7.5.3. Nuclear genes

The nucleotide sequence encoding the small subunit of ribulose 1,5-diphosphate carboxylase has been elaborated from *L. gibba* (STIEKEMA et al. 1983b). A cDNA clone has been used which was isolated from a poly(A) RNA-derived cDNA library (WIMPEE and TOBIN 1982, STIEKEMA et al. 1983a). The clone encodes 53 amino acids of the precursor transit peptide, the entire 120 amino acids of the mature small subunit polypeptide, and 241 nucleotides representing the 3' non-coding region of the mRNA. The amino acid sequence of the mature small subunit RuDPCase shows 70-75% homology to the reported sequences of other species.

2.5.7.5.4. In vitro transcription, DNA repair, plasmid uptake

Isolated nuclei from *L. gibba* have been used to investigate nuclear RNA synthesis and RNA polymerase activity. Incorporation rate of [³H]-uridine into RNA as well as RNA polymerase I activity depend on the nucleus isolation time during the day. Highest values for both parameters are reached during the night phase and a diurnal rhythm is exhibited with nuclei from plants held under continuous light. By contrast, activity of RNA polymerase II remains constant throughout the day (NAKASHIMA 1978, 1979a). Magnesium has three different functions on maintenance of RNA synthetic activity in isolated nuclei. Magnesium is a co-factor of RNA polymerase, it is a stabilizer for the binding of RNA polymerase I to nuclei, and for the binding of stimulating factors to the nuclei (NAKASHIMA 1979b).

Genetic information of chloroplast DNA from *S. punctata* has been expressed in vitro with a coupled transcription-translation system extracted from *Escherichia coli* (DE HEIJ and GROOT 1981). Within the products the α - and β -subunits of ATPase were identified by immunoprecipitation.

tation with heterologous antisera raised against spinach α - and β -subunits.

UV-induced DNA damage and repair have been investigated with W. microscopica and S. polyrrhiza (DEGANI et al. 1980). UV irradiation (250 Jm^{-2}) of plants induces the formation of cytosine-thymine and thymine-thymine dimers which are incorporated into the DNA. The capacity of plants to excise the dimers from their DNA has been examined by analysing DNA hydrolysates at different times after irradiation with UV. The ratio of thymine dimers to thymine in purified DNA amounts to 0.2% directly after irradiation. About 60% of dimers are removed during the following 7 hours of darkness and a total of 80% is removed after two days. In the light the disappearance of dimers from DNA proceeds faster and is complete within 3 hours (see 'photoreactivation').

Uptake of plasmid DNA into intact L. aequinoctialis has been tried by FREY et al. (1980). The plants were incubated with Escherichia coli plasmids pMB9 or pBR325 for 20-22 hours, then washed and treated with DNase I and proteinase K. The DNA isolated subsequently was tested for transformants by a bioassay (E. coli transformation to tetracycline resistance). The authors concluded from their results that in 7 out of 15 cases stable transformants were produced and that about 100-900 intact plasmid molecules were taken up by one Lemna plant.

2.5.8. Further compounds

2.5.8.1. Phospholipids and galactolipids

Metabolism of phospholipids has been followed in S. punctata after supplying plants with a 15-min pulse of ^{32}P -phosphate (BIELESKI 1972). The first phospholipid to become fully labeled is phosphatidic acid. Then phosphatidyl glycerol and phosphatidyl inositol become labeled more rapidly than phosphatidyl choline and phosphatidyl ethanolamine. Phosphatidyl serine is the slowest to label and comprises only about 1% of the ^{32}P -lipid. Synthesis of phosphatidyl glycerol is more sensitive to phosphorus deficiency than the other phospholipids. As phosphatidyl glycerol is located mainly in the chloroplast, the composition of lipids in the chloroplast changes. The phosphorus-deficient chloroplast has a 60% lower phospholipid content. Zeatin reduces the effect of P-deficiency on phospholipid synthesis.

The high percentage of galactolipids and unsaturated fatty acids is typical for the lipids of normal chloroplasts. In cultures of S. polyrrhiza provided with 1% sucrose not only the chlorophyll content per mg protein but also the galactolipid (especially digalactosyl diglyceride) content increase drastically (BAHL et al. 1971, LECHEVALLIER et al. 1971). High concentrations of benzyladenine ($2\text{-}5 \text{ mg/l}$), which increase or slightly decrease the chlorophyll content in L. minor, lower the proportion of galactolipids as compared with phospholipids (BERUBE et al. 1982).

Sublethal doses of the herbicide atrazine increase the number of grana per chloroplast in L. minor (BEAUMONT et al. 1980) respectively the number and length of grana lamellae in S. punctata (MATTOO et al. 1983b). Correspondingly, the percentage of diacylgalactosyl glycerol (compared with total phospholipids and total neutral lipids) increases in L. minor on ground of the atrazine treatment (GRENIER et al. 1979). In the presence of atrazine, labeled acetate is incorporated more strongly in total lipids, diacylgalactosylglycerol, diacyldigalactosylglycerol, phosphatidylglycerol, and diacylglycerol (GRENIER and BEAUMONT 1983). Furthermore, atrazine stimulates the desaturation of fatty acids. The specific radioactivity after acetate incorporation increases under the influence of atrazine in linoleic and linolenic acid in nearly all investigated lipids (GRENIER and BEAUMONT 1983). The ratio of linolenic to linoleic acid in galactolipids and phospholipids is raised by atrazine

sterol content and increases the ratio of stigmasterol to campesterol (D'HARLINGUE et al. 1976).

Within carotenoids (see chapter 1.2.6) the colorless lycopersene (a C₄₀ analogue of squalene) occurs in L. minor (PREVITERA and MONACO 1984). The well-known effect of Sandoz 6706 [4-chloro-5-(dimethylamino)-2-(trifluoromethyl)-3(2H)pyridinone] to eliminate the synthesis of carotenoids, has been confirmed for S. punctata and L. minor (BIGGS and KOSUTH 1980). Experiments with plants treated with Sandoz 6706 revealed that carotenoids play an important role as protectant against ultraviolet-B radiation. Light effects on the carotenoid content in Lemnaceae have been investigated by MONEGER (1968d,f), MONEGER and JACQUES (1968), LECHEVALLIER et al. (1976), and ROMBACH (1976). Under prolonged illumination the action spectra for carotenoid production and photosynthetic activity (CO₂ fixation) are very similar. With short illumination, however, the action spectrum for accumulation of carotenoids in etiolated plants shows 3 maxima: in the blue, green, and red range (MONEGER 1968d, 1968f, MONEGER and JACQUES 1968).

Gibberellin A₃ (10⁻⁵ M) decreases the carotenoid content somewhat in light-grown S. polyrrhiza (D'HARLINGUE 1976). Kinetin modifies the acetate incorporation into carotenoids mainly during a prolonged dark phase. The radioactivity incorporated into violaxanthin, for example, is diminished by 10⁻⁷ M kinetin, but enhanced by 3.3x10⁻⁶ M kinetin, and strongly lowered by 3.3x10⁻⁵ M kinetin (ADABRA-MICHANOL et al. 1975). The last-named kinetin concentration inhibits the production of chlorophylls a and b as well as of β-carotene, lutein, violaxanthin, and neoxanthin. In etiolated fronds of S. polyrrhiza, puromycin stimulates carotenoid synthesis and accumulation in the dark (MONEGER 1968d).

2.5.8.3. Chlorophyll

Very different amounts of chlorophyll have been found in Lemnaceae depending on species and culture conditions (see chapter 1.2.6). The precursor of chlorophylls, chlorophyllide, is accumulated to a determinable degree in L. trisulca, but not in S. polyrrhiza and L. minor (ANTONIELLI and CAGIOTTI 1976).


Chlorophyll synthesis takes place in intimate correlation with chloroplast development. Several factors which inhibit the development of chloroplasts have a bleaching effect on developing duckweed fronds.

in L. minor (GRENIER et al. 1979). Also in S. punctata supplied with atrazine the thylakoid membranes contain a higher ratio of linolenic to linoleic acid (MATTIOO et al. 1983b).

In connection with frond senescence in glucose-stimulated S. punctata cultures the content of monogalactosyl diglyceride decreases, but digalactosyl diglyceride increases. At the same time a marked decrease of the relative amount of linolenic acid and an increase of that of linoleic and palmitic acids can be observed (GROB and EICHENBERGER 1969). Similarly, the monogalactosyl diglyceride proportion of galactolipids as well as the relative content of linolenic acid in nearly all lipids decreases in S. polyrrhiza, when the plants are stressed by addition of polyethylene glycol to the medium (LECHEVALLIER 1977b). An enzyme system producing C₆ aldehydes from C₁₈-unsaturated fatty acids has been indicated from S. polyrrhiza (HATANAKA et al. 1978).

2.5.8.2. Isoprenoids

The diterpenes phytol, trans-1,3-phytadiene, and a novel substance (4R)-4-hydroxyisophytol (fig. 2.52) were isolated from L. minor (PREVITERA and MONACO 1984). The cyclic triterpenes sitosterol, stigmasterol, and campesterol were found repeatedly in Lemnaceae (see chapter 1.2.3). They are located mainly in the endoplasmic reticulum and are glycosylated to some extent (16%) in S. punctata (EICHENBERGER 1975). Supply of gibberellin A₃ to cultures of S. polyrrhiza enhances the total sterol content of the plants, with a higher proportion of campesterol and β-sitosterol, and a lower one of stigmasterol (D'HARLINGUE 1976). Addition of sucrose (1-4%) to the nutrient medium, on the other hand, lowers the total



1a R = H

1b R = Ac

Fig. 2.52. (4R)-4-hydroxyisophytol, a novel diterpene isolated from Lemna minor by PREVITERA and MONACO (1984).

These investigations have been reported in detail in chapter 2.4.1.5. Destruction of chlorophyll occurs in *L. trisulca* at light intensities higher than 50000 lux, i.e., in light of an intensity 5 times higher than at the saturation point of photosynthesis (ZURZYCKI 1957a). Ultra-violet radiation, which leads to a rapid destruction of chlorophyll in *L. minor*, exerts its effect largely through the formation of ozone (WANGMANN and LACEY 1952). In these and other cases not only the chlorophyll itself but also the thylakoid structure of chloroplasts may be affected. The herbicide nitrofen (2,4-dichlorophenyl-4-nitrophenylether) reduces not only the content of chlorophyll and carotenoids in *L. gibba* (WEJNAR and TAI 1983), but also destabilizes the chloroplast structure (WEJNAR and MICHEL 1983). Stress factors such as polyethylene glycol (LECHEVALIER 1977b) and senescence-promoting factors such as exhaustion of minerals, especially in cultures supplied with sucrose or gibberellin A_3 (D'HARLINGUE 1976, D'HARLINGUE et al. 1976), are further agents which lead to degradation of pigments and structures in chloroplasts (compare chapters 1.2.6 and 2.4.1.5).

Sucrose increases the chlorophyll content in *S. polyrrhiza* very greatly, when minerals are not limiting growth (BAHL et al. 1971). The positive effect of gibberellin A_3 on chlorophyll content in *L. trisulca* comes about by inhibition of senescence processes (BATA and NESKOVIC 1974). Growth-inhibiting agents can increase or decrease chlorophyll accumulation. Supply of 10^{-6} M ABA (ALBANELL et al. 1985) or $>0.6 \times 10^{-6}$ M CCC (SUPNIEWSKA 1963) to *L. minor* cultures causes a rise in chlorophyll content of plants. Growth-inhibiting concentrations of BA (10^{-5} - 10^{-4} M), on the other hand, diminish the chlorophyll level per weight of protein nitrogen in *S. polyrrhiza* (LE PABIC 1976a).

2.5.8.4. Cinnamic acids

The aromatic amino acids phenylalanine and tyrosine are the source for biosynthesis of cinnamic acids and all their derivatives (phenylpropanoids). Phenylalanine-ammonia-lyase (PAL) is the key enzyme for this metabolic branch as it deaminates phenylalanine to cinnamic acid and - in several cases - deaminates tyrosine to p-coumaric acid (PAL activity). GORDON (1977) isolated the PAL from *S. polyrrhiza*, *L. gibba*, and *L. aequinoctialis* and showed the enzyme to possess both PAL and TAL activities. In *S. polyrrhiza* PAL activity is enhanced by Mn^{2+} . Very high lev-

els of PAL activity are induced by continuous UV radiation which is increasing anthocyanin accumulation. During continuous light (but not darkness) the PAL and TAL activities exhibit a circadian rhythmicity in *S. polyrrhiza* and *L. aequinoctialis* (GORDON and KOUKKARI 1978).

Up to now, hydroxycinnamic acids and their esters have been detected in 5 members of the Lemnoideae. In *S. polyrrhiza*, 5-p-coumaroylquinic acid, 5-caffeoylquinic acid, 1-feruloylglucose and 1-sinapoylglucose occur (KRAUSE 1978, KRAUSE and STRACK 1979a). Also in *S. punctata* (NG and THIMANN 1962) and *L. gibba* (UMEMOTO 1971) 5-caffeoylquinic acid, i.e., chlorogenic acid, has been indicated. *L. trisulca* and *L. minor* contain caffeic acid, and *L. trisulca*, in addition, ferulic acid (REZNIX and NEUHAEUSEL 1959). Interestingly, such substances are secreted from the plants to the medium, as has been demonstrated by JURD et al. (1957) and WALLACE (cited in McCURE 1970).


Certain phenylpropanes are intermediates on the way to prenylquinones, e.g., plastoquinone. 4-hydroxyphenylpyruvic acid, which is derived from tyrosine by an oxidase reaction, can be converted into homogentisic acid by a thylakoid preparation isolated from *L. gibba* (LOEFFELHARDT and KINDL 1979). The enzyme responsible for this reaction, 4-hydroxyphenylpyruvate dioxygenase, shows maximum activity, when Fe^{2+} , high concentrations of ascorbate and catalase are added to the incubation mixture.

The most important mechanism for formation of benzoic acids in plants is the side-chain degradation of cinnamic acids. The occurrence of benzoic acid in *L. aequinoctialis* and *L. gibba* was established by FUJIOKA et al. (1983a, 1985). The endogenous level of this compound is very much influenced by the composition and concentration of the culture medium, but the benzoic acid content in vegetative and in photoperiodically-induced flowering plants does not differ significantly. Salicylic acid, i.e., 2-hydroxybenzoic acid, could not be definitely identified in *L. aequinoctialis*. If it exists at all, the amount must be less than 1 ng/g fresh weight (FUJIOKA et al. 1983a).

Benzoic acids are known to be the source for production of phenols by oxidative decarboxylation. The ability to convert such phenols to the corresponding mono- β -D-glucopyranosides for detoxification is widespread in angiosperms, but in *L. minor* and two other water plants (*Elodea canadensis*, *Utricularia vulgaris*) the ability to glucosylate quinol and resorcinol is lacking (PRIDHAM 1964). One can speculate that phenols may be released to the medium by water plants against the attack of microorganisms.

Fig. 2.54. Metabolic pathways leading to several subgroups of flavonoids in Lemnaceae, as elaborated essentially by WALLACE (see text).

A=oxidation
B=C-glycosylation
C=3'-hydroxylation
D=isomerization
E=O-glycosylation
F=3'-methyletherformation



Aglycones	R	R'	R ^d	R ^u
apigenin	H	H	H	H
luteolin	OH	H	H	H
witexin	H	H	H	glucose
orientin	OH	H	H	glucose
isovitexin	H	H	glucose	H
isoorientin	OH	H	glucose	H
kaempferol	H	OH	H	H
quercetin	OH	OH	H	H
Aglycones	R	R'		
cyanidin	OH	H		
petunidin	OCH ₃	OH		

Fig. 2.53. Flavanoid nucleus, numbering system, and substitutions (R) of the flavone, glycoflavone, and flavonols (above) and of the anthocyanidins (below) detected in Lemnaceae (after MCCLURE and ALSTON 1966).

Figure 2.54 summarizes the principal pathways leading to several subgroups of flavonoids in Lemnaceae. The scheme reflects the conclusions which can be drawn from the work of WALLACE (1967, 1969, 1973, 1975; WALLACE and ALSTON 1966a, 1966b; WALLACE et al. 1969; WALLACE and MARRY 1970; WALLACE and GRISEBACH 1973). Using ¹⁴C-labeled flavonoid aglycones WALLACE showed that Lemnoideae (at least S. polyrrhiza, S. punctata, L. minor, and L. obscura) are able to O-glycosylate supplied flavones. When ¹⁴C-apigenin is added to the nutrient medium, plants incorporate this flavone into apigenin-7-O-glucoside. Similarly, luteolin can be transformed to luteolin-7-O-glucoside. From these and other results it can be concluded that O-glycosylation of ring A and O-methylation of ring B evidently mark final stages of flavonoid biosynthesis. C-glycosylation of ring B, on the other hand, is an early biogenetic process occurring at the flavanone level. Naringenin but not apigenin or luteolin are incorporated into vitexin and orientin. Therefore, the enzyme oxidizing flavanone to flavone is irreversible. Furthermore, no labeled flavone is incorporated into an anthocyanin. The B-ring oxidation pattern of flavone-O-glucosides was determined at the flavone level of biosynthesis and was also essentially irreversible. The B-ring oxidation pattern of C-glycosyl flavones was noted to be prior to C-glycosylation, probably at the flavanone stage. Isomerization of the C-glycosyl flavones proceeds only from the 8-isomer to the 6-isomer (vitexin to isovitexin and orientin to isoorientin), but not in the reverse direction. HOESEL et al. (1972) have shown that L. minor, which does not accumulate any flavonol, nevertheless can convert kaempferol and quercetin to the corresponding 2,3-dihydroxy flavanone. The authors argue such reaction could be the first step of a flavonol degradation, and conclude that a rapid turnover of flavonols may proceed even in L. minor. However, flavonols do not occur in the whole genus Lemna. Thus, the responsible enzyme in Lemna may use primarily another substrate instead of flavonols.

Especially the biosynthesis of anthocyanins depends on some environmental factors: light and the mineral content of the medium. The light requirement for anthocyanin formation is different in the three investigated species of Spirodela. In S. polyrrhiza the action of light is mediated by phytochrome and photosynthetic pigments. Five minutes of red light (far-red reversible) increase the anthocyanin content of dark-grown etiolated plants. When such plants are irradiated with white light

for 43 hours, the anthocyanin formation is inhibited by streptomycin (inhibition of chloroplast development), or by norflurazon (photooxidation of chlorophylls in the absence of carotenoids), or by CMU (inhibition of photosynthetic electron transport). Red and blue light, but not far red enhance anthocyanin accumulation during prolonged irradiation (MANCINELLI 1977, 1984; MANCINELLI and RABINO 1984). S. intermedia needs (in addition) high-energy blue light for anthocyanin formation. Under continuous red light anthocyanin production is completely lacking (McCLURE 1967b, 1968, 1975). S. punctata shows another type of light dependence, when plants have been pre-cultivated under light intensities of approximately 4000 lux. Under this condition the plants develop a low anthocyanin content. A terminating irradiation with single wave-lengths between 250 and 750 nm (before 40 hours of darkness) reveals that only far red, red, and ultraviolet can increase anthocyanin formation (maximum of the action spectrum at 705 nm and at 300 nm). Both blue and green light are relatively ineffective. Red and far-red exert additive effects only; no red/far-red antagonism can be obtained (NG et al. 1964).

The action of photosynthesis on anthocyanin formation in S. polyrrhiza can be replaced by adding sucrose to the cultures (MANCINELLI 1977, 1984, MANCINELLI and RABINO 1984). Promotive effects of sucrose on anthocyanin formation have been found also in S. punctata (THIMANN and EDMONDSON 1949, THIMANN et al. 1951, NG et al. 1964). THIMANN and RADNER then analysed the light requirement of anthocyanin formation in more detail. They found that RNA biosynthesis is needed for asserting the light effect on anthocyanin formation. Purine and pyrimidine analogs (THIMANN and RADNER 1955b, 1962) as well as ribonuclease (RADNER and THIMANN 1963) inhibit the production of anthocyanins in S. punctata. A further step in the reaction chain between light and anthocyanins seems to be the synthesis of riboflavin. Azaguanine inhibits not only anthocyanin but also riboflavin formation in the plants. Exogenous riboflavin can reverse the inhibiting effect of azaguanine (and some other inhibitors). Furthermore, the combined supply of riboflavin and sucrose can replace the light requirement completely (THIMANN and RADNER 1958). The inhibitory effect of methionine on anthocyanin formation (THIMANN and RADNER 1955a) also can be explained as an intervention in riboflavin synthesis. The conclusions which can be drawn from the results of THIMANN and co-workers are summarized in figure 2.55. The necessity for co-operation of photosynthates with riboflavin is stressed by the observation that




Fig. 2.55. Intermediate steps in the light action on anthocyanin biosynthesis in *Spirodela punctata*, as elaborated essentially by THIMANN and RADNER (see text).

absence of CO_2 during a preillumination period greatly reduces the effectiveness of riboflavin in the following dark period (THIMANN and RADNER 1958).

Light effects on other flavonoids have been investigated with *S. intermedia* by McCURE and co-workers (see McCURE 1968, 1970, 1975). Vitexin and kaempferol are synthesized to some degree even in complete darkness, but orientin and quercetin are stated to need low-energy blue or white light for accumulation.

Shortage of several mineral elements also influences anthocyanin formation. Low supply or deficiency of nitrogen (BALL et al. 1967, MENSCHIK 1970), phosphorus (THIMANN et al. 1951, BALL et al. 1967, JUNGNIKEL 1978), iron (YOSHIMURA 1950), boron, zinc, and molybdenum (THIMANN and EDMONDSON 1949) promotes anthocyanin synthesis. In most, if not all cases a growth inhibition induced by the mineral deficiency may be the primary cause for the enhanced pigmentation. The effect of nitrogen deficiency in *S. polyrrhiza* can, but must not be connected with the induction of turions. Not only anthocyanin production is affected by nitrogen deficiency in *S. polyrrhiza*, but also the formation of other flavonoids (REZNIK and MENSCHIK 1969, MENSCHIK 1970). The relationship between growth inhibition (i.e., inhibition of protein synthesis) and enhanced anthocyanin formation exists also in experiments, in which copper (THIMANN and EDMONDSON 1949, EDMONDSON and THIMANN 1950) or a high concentration (1 mM) of EDTA (ELLIOTT 1977) were added to the medium. From three copper-chelating agents tested only one (phenylthiocarbamide) reduces anthocyanin formation without affecting plant growth (EDMONDSON and THIMANN 1950).

FURUYA and THIMANN (1964) investigated the effects of gibberellin A_3 on anthocyanin formation in *S. polyrrhiza* and *S. punctata*. In both species, anthocyanin synthesis is promoted by low GA_3 concentrations (10^{-7} - 10^{-6} M) and inhibited by high GA_3 concentrations (10^{-4} M) during the first 7-10 days of culture. After a further 10 days, all GA_3 concentrations are inhibitory. Then the sensitivity of plants to GA_3 is higher at lower light intensities. Under 3000 lux white light a concentration of 3×10^{-6} M GA_3 is sufficient for 50% inhibition. At this level, GA_3 has no effect on growth.

A prerequisite for physiological investigations on flavonoids other than anthocyanins would be an effective and dependable method to separate and quantify these substances. STRACK and KRAUSE (1978) showed reversed-

2.5.9. Circadian rhythms in metabolism and photoperiodism

Plants are well adapted to the daily changes of light and darkness. Obviously, endogenous oscillators ('clocks') do exist, which adjust the metabolic activities in relation to the expected environmental changes. In this way, a 'temporal compartmentation' (HILLMAN 1979a) between alternate metabolic pathways is established. In Lemnaceae circadian rhythms of respiration, potassium and nitrate uptake, nitrate reductase, and RNA metabolism have been thoroughly investigated. The relation between metabolic rhythms and the sensitivity changes to photoperiodic light signals has also been taken into consideration.

In L. aequinoctialis 6746 diurnal oscillations of CO₂ output were demonstrated under heterotrophic conditions (HILLMAN 1970a). In darkness following either continuous dim red light or entrainment to a 12(12) light (dark) schedule, the rate of CO₂ output oscillates with a circadian periodicity, before damping after one, two, or three days (see below). In continuous red light, the rate is linear. In darkness the CO₂ output rhythm can be prolonged by entrainment with a daily short red light. This phytochrome-mediated light effect acts as the synchronizer for the endogenous rhythm (HILLMAN 1971c, 1975b). O₂ uptake in strain 7646 is essentially parallel to CO₂ output under all conditions tested (HILLMAN 1977c).

The course of CO₂ output rhythm depends on the kind and extent of nitrogen supply as well as on temperature. With low or lacking nitrogen or with aspartate as the sole nitrogen source one daily peak of CO₂ output appears. With higher concentrations of nitrate or ammonia or with glutamine as the sole nitrogen source two daily peaks of CO₂ output are recorded (HILLMAN 1971a, 1972a, 1975b, 1979a). Damping of the CO₂ output oscillation in continuous darkness proceeds in the presence of nitrate after one single circadian cycle, with medium without nitrogen after two or three cycles, and with ammonium-containing medium after three or more cycles (HILLMAN 1975b). Lowering of the temperature from 28°C to 21°C reduces the relative peak height of CO₂ output rhythm roughly by 70% when plants were cultivated in nitrate, ammonium, or aspartate medium, but only by 20-30% in nitrogen-deficient medium (HILLMAN 1979b).

Similar CO₂ output rhythms as in L. aequinoctialis 6746 have been demonstrated also for other Lemnaceae [several strains of L. aequinoctialis, strains G1 and G3 of L. gibba, and strain 6573 of L. turionifera (under

phase high-performance liquid chromatography (HPLC) to be such a method for flavones and C-glycosylflavones from S. polyrrhiza. HPLC can also be used for separation of hydroxycinnamic acid conjugates (KRAUSE and STRACK 1979b).

the name of *L. minor*). Not only the distinctness of rhythm varies from strain to strain, but also the dependence on the kind of nitrogen provision (HILLMAN 1972a, 1979a).

Using mixotrophic conditions, MIYATA and YAMAMOTO (1969) pointed out a diurnal rhythm of O_2 -uptake in *L. gibba* G3. After exposure to 2 short days plants were exposed to continuous white light. Under this condition a circadian oscillation of O_2 -uptake proceeds, damping after about three days. A similar damped oscillation was found for $^{14}CO_2$ output after supply of glucose-1- ^{14}C , but not with glucose-2- ^{14}C or glucose-6- ^{14}C . Thus, the rhythm comes about by variations of activity of the pentose-phosphate pathway. The activities of NADP-linked and NAD-linked glyceraldehyde-3-phosphate dehydrogenase show diurnal changes in opposite directions, and an oscillation of acid phosphatase activity runs undamped for at least 3 days. Some other enzymes are active without rhythmical variations: NADP glucose-6-phosphate dehydrogenase, 6-phosphogluconate dehydrogenase, pyruvate kinase, and NADP isocitric dehydrogenase.

Further investigations with *L. gibba* G3 were made under autotrophic conditions (MIYATA 1970, 1971a, 1971b). The light-on signal for the O_2 uptake rhythm is mediated by phytochrome. The rhythm is set in motion after a lag-phase of 6 hours, in which the phytochrome has to be in the P_{fr} state. After the end of the lag-phase the expression of rhythm needs the presence of relatively high light intensities. Photosynthetic inhibitors such as CMU and DCMU abolish the O_2 uptake rhythm after a lag of 1 day. Only a part of the respiration is subject to the circadian oscillation. This rhythmic component is sensitive to azide, but not influenced by exogenous substrate or 2,4-dinitrophenol.

The physiological basis for the mutually inverse rhythmic changes in activity of NADP-dependent and NAD-dependent glyceraldehyde 3-phosphate dehydrogenases (GPDs) has been worked out by GOTO (1978, 1979a,b, 1983, 1984). The NADP-specific enzyme is located almost completely in the chloroplast and the NAD-specific enzyme largely in the cytoplasm. The activity changes are caused by alteration of the apparent K_m values, not the V_{max} values, of the GPD reactions for their substrate and coenzymes. In the case of NADP-GPD there are two molecular species, the NADP-GPD I with 520 Kdalton and low affinity for NADP, and the NADP-GPD II with 170 Kdalton and high affinity for NADP. The decrease in K_m value of NADP-GPD comes about by the conversion of NADP-GPD I to NADP-GPD II, which can be manipulated in vitro by addition of NADP(H). This result induced GOTO to

examine the levels of NADP(H) and NAD(H) in *L. gibba* G3. In fact, in vivo levels of total NADP and total NAD display mirror-image circadian rhythms, with NADP(H) peaking at the same time as NADP-GPD (12 hrs after the beginning of continuous light), and NAD(H) peaking at the same time as NAD-GPD (24 hrs after the beginning of continuous light). The enzymes converting NADP and NAD into one another also oscillate diurnally. NAD kinase activity is maximum 6 hrs after the beginning of continuous light, and NADP phosphatase activity (as of other phosphatases, see above) 12 hours later. Moreover, in vitro treatments with EGTA (a Ca^{2+} -chelator), chlorpromazine and W7 (both inhibitors of calmodulin) are able to both inhibit NAD kinase from its highest level of activity to its minimal one and activate NADP phosphatase from its lowest level of activity to its maximal one. From these results GOTO concluded that the in vivo level of Ca^{2+} -calmodulin should oscillate in phase with the rhythm of NAD kinase activity, in this way inducing mirror-image circadian rhythms of activities of NAD kinase and of NADP phosphatase. If this hypothesis were right, then Ca^{2+}/Ca^{2+} -calmodulin, NAD kinase/NADP phosphatase, and NADP/NAD could possibly be envisaged as components of a circadian clock, which is self-sustained and light-entrainable, and which controls the activities of NADP-GPD, NAD-GPD, and possibly several other enzymes. Figure 2.56 summarizes all the data of GOTO and also his working hypothesis.

The diurnal NAD/NADP-rhythm and the concurrent rhythm of respiration may be the basis for several other metabolic oscillations. First of all, the investigations of KONDO on rhythmic changes of uptake and release of inorganic ions, especially potassium, should be mentioned. When *L. gibba* G3 after the start of continuous light is transferred to distilled water at different times, then the amount of electrolytes leaking out from the plants changes diurnally (KONDO 1978). Maximum leakage appears in the early subjective day and minimum leakage in the early subjective night. As in respiration, the rhythm is reset by a light-on signal and persists under relatively high light intensities (3000 lux) for at least 3-5 days. Potassium and nitrate are the major ions, which contribute to the change in electroconductivity of the ambient water (KONDO and TSUDZUKI 1978, KONDO 1982b). Magnesium and phosphate ions show circadian rhythmicity of uptake and release to a far lesser extent and calcium ions are consumed by plants without any rhythmical feature. The potassium and nitrate uptakes are not linked together by a common ion permeating system,

because the nitrate uptake rhythm is not affected by the lowering of potassium concentration, and the potassium uptake rhythm persists to some extent in the absence of nitrate (KONDO 1982c). The attributes of potassium uptake rhythm were communicated in many further publications. The rhythm appears regardless of the potassium concentration in the culture medium (from 10 μM to 2 mM) (TSUDZUKI and KONDO 1979). The amplitude of oscillation is dependent on light intensity (KONDO and TSUDZUKI 1978, TSUDZUKI and KONDO 1979). In total darkness,




Fig. 2.56. Hypothetical scheme for a self-sustaining and light-entrainable rhythm working in *Lemna gibba* G3 (after data of GOTO, see text).

the rhythm fades out in c. 48 h, and under low-intensity light (below 700 lux) the rhythm is damped rapidly. With 13000 lux, on the other hand, the rate of potassium uptake increases almost linearly, probably due to the accelerated plant growth under this strong light. Therefore, a good rhythmic behaviour can be observed between 1500 and 8000 lux light intensity. Exogenous sucrose can largely replace the light requirement. A daily light pulse (4200 lux for 15 min) is sufficient under such conditions to sustain the rhythm (KONDO 1982a). Respiratory inhibitors or uncouplers (NaCN, 2,4-dinitrophenol, CCCP) reduce the average rate of potassium uptake without remarkably decreasing the amplitude of the rhythm. NaN_3 , however, preferably diminishes the amplitude at a concentration as low as 10^{-6}M (KONDO and TSUDZUKI 1980b). Factors changing the period of the free-running potassium uptake rhythm are lithium, rubidium, and ethanol (KONDO and TSUDZUKI 1980c, KONDO 1984b). Whereas Li^+ ($\geq 0.3\text{ mM}$) and Rb^+ ($[\text{Rb}^+/\text{K}^+] = 0.9$) lengthen the oscillation period under the routinely used experimental conditions, ethanol (0.3-3%) has a shortening effect. The lengthening of the period of potassium uptake rhythm due to 0.5 mM Li^+ can be removed by addition of 20 μM sodium (KONDO 1984a).

Phase resetting of the potassium uptake rhythm can be managed by light, darkness, extreme temperatures, and azide (KONDO 1982d, 1983a,b, KONDO and TSUDZUKI 1980a). Under conditions of continuous light a pulse of high temperature (39°C) shifts the phase in the same way as a pulse of high-intensity light (15000 lux): both pulses cause a phase delay when given early in the subjective night, but cause a phase advance when given late in the subjective night, with a distinct discontinuity (break-point) in the phase response curve about the middle of the subjective night. A pulse of darkness, or of low temperature (5 or 10°C), however, causes a phase shift that is the inverse of that caused by high-intensity light or high temperature. Temperature steps from 20 to 30°C or from 30 to 20°C do not modify the phase of the rhythm. A 6-hour application of 1 mM azide to plants has the same effect as the 6-hour pulse of darkness or low temperature.

In one publication, KONDO and TSUDZUKI (1978) investigated the circadian rhythm of potassium uptake not only in *L. gibba* G3, but also in *L. aequinoctialis* 6746. In correspondence to what is known about the respiration rhythm (see above), *L. aequinoctialis* 6746 displays a K^+ -uptake rhythm only in the dark and not in the light. The rhythm does not persist beyond the second cycle.

Diurnal fluctuations of nitrate reductase activity with a peak in the forenoon were found in L. aequinoctialis (DEVI and MAHESHWARI 1979) and W. microscopica (BAKSHI et al. 1978, 1979). In both species, the rhythmicity persists under extended dark conditions as well as under continuous light, at least for one, two, or three cycles. In W. microscopica, the phase of the rhythm shifts under inverted light-dark cycles. As shown by inhibitor studies, the increase in enzyme activity seems to depend on the new synthesis of both RNA and protein. Despite the fact that nitrate reductase is induced by its substrate (see chapter 2.5.6.4), there is no correlation between enzyme activity and the level of tissue nitrate. The nitrate level in the tissue declines constantly during a 24-h period of analysis, whereas the nitrate reductase level shows normal fluctuations. In relation to the NAD/NADP-cycle, however, it is of interest that the nitrate reductase extracted from W. microscopica is protected by NADH (BAKSHI 1983).

Other enzymes which display a circadian rhythmicity are phenylalanine ammonia-lyase (=PAL, determined from S. polyrrhiza and L. aequinoctialis 6746) and tyrosine ammonia-lyase (determined from L. aequinoctialis). The oscillations of enzyme activity can be demonstrated under continuous light, but not - at least the PAL activity in L. aequinoctialis - in continuous darkness (GORDON and KOUKKARI 1978). Recently, investigations on PAL rhythmicity in S. punctata have been made by KNYPL et al. (1986). Diurnal as well as 2-3-day rhythms have been found, when plants were held under continuous light and were provided with glucose or sucrose. For interpretation of some of the results it should be taken into account that plants may release cinnamic acids and ABA into the medium (see chapter 2.5.2.3.4). Trans-cinnamic acid is an inhibitor of PAL, and ABA has been shown to shift the PAL oscillation.

A circadian rhythm of RNA synthesis (indicated by ³H-uridine incorporation into RNA) has been shown to occur in L. gibba G3 under continuous light conditions (NAKASHIMA 1976). Uridine incorporating activity is reaching its highest level at 18 h and its lowest at 6 h after the beginning of continuous light. The length of the period of the rhythm is not affected by temperature over a range of 16° to 30°C. The distinctness of amplitude of the rhythm depends on the quantity of light (duration and intensity) during the first half of the cycle, and on the P_{fr} state of phytochrome during the second half of the cycle (KATO and NAKASHIMA 1979). The diurnal oscillation comprises the synthesis of ribosom-

al RNAs in the nucleus as well as in the chloroplasts (NAKASHIMA 1975, 1978). As stated already in chapter 2.5.7.5.4, in nuclei only RNA polymerase I, but not RNA polymerase II activity changes diurnally (NAKASHIMA 1979a).

As in other plants, circadian clocks work in Lemnaceae not only to arrange the timing of certain metabolic activities, but also to permit day-length measurement in connection with photoperiodic control of flowering. In chapter 2.4.3.1.5 it was already mentioned that the effectivity of short light pulses to induce or to inhibit flowering, changes diurnally during prolonged dark periods. In a series of papers HILLMAN (1963, 1964, 1972b, 1976a, 1976b, 1976c) pointed out that in heterotrophically grown L. aequinoctialis 6746 the light sensitivity of flowering and the oscillation of respiration may be dependent on the same clock. In both cases, the rhythm is entrained in the same way by "filled skeleton photoperiods" (that is a series of 2,4, or 6 short light pulses equally distributed during the regular photoperiod). A 6 h increase in length of the photoperiod delays the time of maximal light sensitivity for flowering and the time of maximal CO₂ output by 3.6 h and about 3 h, respectively.

The mechanism of the clock which controls the effectivity of light signals for regulation of developmental processes is unknown, but several details of its attributes are known. During the last decade OOTA has examined thoroughly, whether PITTENDRIGH's (1966) model of the photoperiodic clock may be applicable to the oscillator(s) responsible for light regulation of flowering in L. gibba G3 (OTA 1970, 1971, 1975a,b, 1981, OOTA and HOSHINO 1979, OOTA and NAKASHIMA 1978, OOTA and TSUBUZUKI 1979) and L. aequinoctialis 6746 (OTA 1983a, 1983b, 1984, 1985). In both species, a great number of special light-dark schedules have been tried out. For example, in experiments with 24 h-light-dark cycles and night breaks given additionally not only the position of the night break (short light pulse) but also the duration of the main light period have been varied systematically. From a wealth of data the following conclusions have been drawn. In plants entrained by light-dark cycles a light on oscillator works. A light pulse starts the circadian cycle, resetting the clock to its "LI-phase". In the following phase the clock runs independently whether plants are exposed to light or not. This means that complete or skeleton photoperiods are equally effective. After several hours the "L2-phase" is reached, which is identical with the terminating

part of the critical light phase. The L2-phase is light-sensitive and decides on the developmental effect (flower induction). In the long-day plant L. gibba G3 the critical light phase amounts to about 12 hours. Lightening of the L1- and L2-phase induces flowering. In the short-day plant L. aequinoctialis 6746 the critical light phase amounts to 14 hours. Lightening of the L1- and L2-phase prevents flowering. Light pulses given between a main light period and the light-sensitive L2-phase are registered as "false dusk", in this way establishing an "asymmetric skeleton photoperiod". Light pulses, however, which are supplied after the L2-phase, act as "false dawn". They constitute an asymmetrical skeleton photoperiod together with the following main photoperiod and shift the clock to the L1-state. After one (L. aequinoctialis) or two (L. gibba) transient cycles the entrainment is accomplished. Under conditions of continuous light, the light-on oscillator described in L. aequinoctialis becomes inactive. Then another clock is set in motion, when plants are transferred to darkness (OOTA 1985). The light-off signal starts the L1-phase of a light-off oscillator. The terminating part of the critical dark phase (L2-phase) is reached 7 hours later. For flower induction in L. aequinoctialis especially this light-sensitive L2-phase has to be darkened. Flower inhibition during the L2-phase is effected by red as well as far red light.

3. APPLICATION AND ECONOMIC IMPORTANCE

3.1. GENERAL

Lemnaceae have a few characteristics which make a versatile application promising:

- simple culture conditions (aseptic conditions possible, small need of space)
- vegetative multiplication (genetically uniform)
- high multiplication rate
- high protein content
- easy harvest possibilities
- high need of nutrients
- pronounced ability to accumulate heavy metals
- only few diseases

Accordingly, the main applicability must be looked for in the following directions:

- test plant for phytophysiological experiments and for detection of toxicants
- source of protein (food for animals)
- removal of surplus nutrients and heavy metals from (waste) water
- production of energy and phytochemicals
- regulator in aquatic ecosystems

A comprehensive survey on the economic possibilities of Lemnaceae was done by RUSKIN and SHIPLEY (1976) in a wide-spread booklet on "Making aquatic weeds useful: some perspectives for developing countries" with an annex on duckweeds and their uses. Since then, many other compilations on the applicability of duckweeds have been presented: HILLMAN and CULLEY (1978a), STEPHENSON et al. (1980), EDWARDS (1980), CULLEY et al. (1981), AUGSTEN (1984a;b), BJOERNDAL (1984), HUBAC et al. (1984), RATH et al. (1986). Earlier reviews can be found in HILLMAN (1961a) and SCHULZ (1962).

Lemnaceae are not always beneficial to man. In some cases when they cov-

er a water with a carpet centimeters deep, they may become a nuisance for motor-boating and fishing or for bathing. A typical action against duckweed invasion is described by HARGROVE (1976). The "Operation Duckweed" was performed with success in Alabama, USA, 70 ha of a water system was sprayed with 1 kg/ha diquat and 0.5 kg/ha copper. This action must be repeated at least once a year in order to keep the water free of a duckweed cover. Even if this method is effective and relatively inexpensive, it is ecologically not appropriate. The same is true for the frequent use of herbicides in fish ponds to achieve a better aeration of the water. The mass development of Lemnaceae certainly is a consequence of a high nutrient content in the water. Very often this is the result of industrial, agricultural or domestic pollution (especially with phosphorus and nitrogen). The high nutrient content should be counteracted by purification of the inlet water or by harvesting the Lemnaceae to lower the nutrient concentration in the water (see chapter 3.5).

3.2. BIOMASS, PRODUCTIVITY, ENERGY CONTENT AND NUTRITIVE VALUE

3.2.1. Productivity

The maximum growth rate in Lemnaceae is species and clone specific (LANDOLT 1957, REJMANKOVA 1975a, PORATH et al. 1979). The highest growth rate of Lemnaceae corresponds to about a doubling time of 24 hours. This was noted in L. aequinoctialis (LANDOLT 1957, CHANG et al. 1977, DATKO et al. 1980a), and in W. microscopica (VENKATARAMAN et al. 1970). This highest growth rate results in an increase of one gram per gram dry weight and day, or 64 grams per gram dry weight and week. In comparison, fast growing corn does not produce more than 2.3 g/g and week, according to HILLMAN and CULLEY (1978a). For S. polyrrhiza, the growth rate at high temperatures is not much below that of L. aequinoctialis. In this species, the dry weight amounts to at least 2 mg/cm² under optimal conditions. If we assume a doubling time of 24 hours and a completely covered water surface, the theoretical increase would be 20 g/m² and day or 73 t/ha and year, provided that the conditions are optimal during the whole year. This theoretical yield is never reached in nature, because temperature and nutrient condition are nowhere equally good at all seasons. However for shorter times, these values have been matched at least to 1/3-2/3 (LANDOLT 1957, SAHAI and SINHA 1970, PORATH and KOTON 1977, SAID et al. 1979, CORRADI et al. 1981, REJMANKOVA et al. 1983, MESTAYER et al. 1984, ORON et al. 1985, 1987, HENDERSON et al. 1984, and many others, see also table 3.1). Values of 1 to 15 g/m² and day have been achieved. Growth rate of Lemnaceae in axenic cultures is maximum at low frond densities (e.g. 10 g/m² dry weight) (REDDY and DEBUSK 1985a,b). The production of fixed carbon was calculated by WARD et al. (1963). The authors noted 7 mg carbon/g dry weight and hour in S. polyrrhiza under optimal nutrient conditions and at 12000 lux. WOHLER (1966) measured in pond water at 10000 lux 1.6 mg g⁻¹ h⁻¹ carbon in S. polyrrhiza, 1.4 mg g⁻¹ h⁻¹ in L. minor, 1.9 mg g⁻¹ h⁻¹ in L. trisulca and 2.7 mg g⁻¹ h⁻¹ in W. gladiata. The differences in the carbon content per dry weight are probably due to differences in assimilation area per dry weight for different species. FILBIN and HOUGH (1985) measured 2.5 mg g⁻¹ h⁻¹ carbon in L. minor during the growing season.

SAID et al. (1979) in Louisiana during summer time found 44 t duckweed

Table 3.1. Productivity of Lemnaceae in different regions of the world

* 0.65 t ha⁻¹ in January and 3.4 t ha⁻¹ in August

Region	dry weight in t ha ⁻¹ yr	authors
southern states of the USA	19.2 14.5 - 27 23.3* 13.5 16.1	STANLEY and MADEWELL (1975) MYERS (1977) CULLEY and MYERS (1980) DEBUSK et al. (1981) REDDY and DEBUSK (1985b)
Israel	39	HEPHER and PRUGININ (1979 cited from EDWARDS 1980)
	10	PORATH et al. (1979)
Egypt	10	EL-DIN (1982)
India	22	RAO et al. (1982)
northern Thailand	10.5	BHANTHUMNAVIN and MCGARRY (1971)
Uzbekistan	7 - 15	TAUBAEV and ABDIEV (1973)
GDR	16 (calculated)	SCHULZ (1962)
CSSR	7.5 - 8	REJMANKOVA (1975b, 1979)

(dry weight) calculated per ha and year which is nearly 2/3 of the theoretical maximum. The actual productivity in nature during a whole year is 8-30% of the maximal value (table 3.1). A mixture of different Lemnaceae species might reach higher values because then differently adapted clones are available for different seasons and different layers of the water can be utilized. For a high productivity in nature, the Lemnaceae should cover the water completely to avoid growth of algae (DEBUSK et al. 1981).

In the first three regions of table 3.1, the productivity is reduced and in the last three regions it is stopped during the winter. In India and Thailand the temperatures are mostly optimal but during the rainy season growth is slowed down due to a lack of nutrients.

KOLES (1986) developed a model to predict growth rates of duckweeds under various environmental conditions and to predict changes in water nutrient conditions accompanying the growth of these plants.

3.2.2. Biomass

The biomass of Lemnaceae in nature is relatively low compared with other water plants and with terrestrial plants. This is due to the special growth habit: Lemnaceae only form rather thin layers of fronds and are composed of relatively light assimilating and aerenchymatic tissue. Highest values of the dry weight of a single frond amounted to 0.6 mg in *S. polyrrhiza* (LANDOLT 1957); the lowest weight of a frond was found in *W. columbiana* (c. 0.008 mg, measured as 0.15 mg fresh weight by ARMSTRONG 1982).

The following values of biomass (in g dry weight per m²) during the summer have been reported: 44-56 (MOORE 1962, 1965), 50-150 (IKUSIMA 1963b), up to 192 (JERVIS 1969), 70-180 (REJMANKOVA 1978, 1982), up to 280 (EWEL and ODUM 1978), up to 250 (HEJNY et al. 1981), 114 (SASTROUTO-MO 1982), up to 180 (GHETTI et al. 1982), 20-184 (KUECHLER 1986), up to 220 (GEARHEART et al. 1986). In comparison, stands of *Phragmites* develop 1612 g/m² (KUECHLER 1986) or even up to 9860 g/m² (HEJNY et al. 1981) which is at least 40 times higher than the biomass of a duckweed cover. The biomass of Lemnaceae in waters which contain few nutrients is much lower. A *Lemno-Utricularietum* consists of 10 g/m² dry biomass (HEJNY et al. 1981), a "Lemnetum" 12 g/m² and 7 g/m² (KLOSE 1963 and VARFOLOMEEVA 1976, resp.). McLAY (1976) noted that *S. punctata* is able to develop a

Table 3.2. Number of fronds of Lemnaceae per m² water surface in nature

Species	number/m ²	author
<i>S. polyrrhiza</i>	10000-29000	KAUL and BAKAYA (1973)
<i>Lemna</i> sp.	40000 140000	KLOSE (1963) GHETTI et al. (1982)
<i>L. gibba</i>	16000-89000	KAUL and BAKAYA (1973)
<i>L. minor</i>	18000-84000	KAUL and BAKAYA (1973)
<i>L. trisulca</i>	10000-29000	KAUL and BAKAYA (1973)
<i>W. columbiana</i> , <i>W. borealis</i>	200000	HICKS (1937)

2.5 times higher biomass than L. minor and a 17 times higher biomass than W. arthiza.

Optimum stocking density for production was found to be 20 g (dry weight) /m² (DEBUSK et al. 1981) and 120 g/m² (PORATH et al. 1979), both for L. minor.

Numbers of fronds per m² were measured by different authors. The results have been put together in table 3.2. KLOSE (1963) counted fronds in Germany, GHETTI et al. (1982) in Italy, KAUL and BAKAYA (1973) in Kashmir (India) and HICKS (1937) in Indiana (USA).

According to RUSKIN and SHIPLEY (1976), a layer of L. minor may expand within 53 days from 6.4 cm² to 0.5 ha.

3.2.3. Energy content

The energy content of the Lemnaceae belong to the highest within water plants (STEUBING et al. 1980, see table 3.3). However, since the energy content is much dependent on the growth conditions, it is not clear if there are differences in the energy content between different species

Table 3.3. Energy content of different Lemnaceae

species	energy content in J/g	author
<u>S. polyrrhiza</u>	9660 15500-17090	SINGH and SHARMA (1975) SUTTON and ORNES (1977)
<u>S. punctata</u>	13880-17640 16840	SRIVASTAVA (1978) MESTAYER (1980)
<u>L. minor</u>	16920	MUZZAR et al. (1978a)
<u>L. aequinoctialis</u>	15360-16970	SRIVASTAVA (1978)
<u>L. minuscula</u>	15330	STEUBING et al. (1980)
<u>Egeria densa</u>	9310	STEUBING et al. (1980)
<u>Juncus procerus</u>	17380	STEUBING et al. (1980)

(table 3.3). SRIVASTAVA (1978) reports of somewhat higher energy content of dry weight during the cold season (average temperatures between 13°C and 19°C) in India than during the hot season. This is probably due to a higher starch content of the Lemnaceae frond at lower temperatures.

According to TRUAX et al. (1972), 4536 J/g of the energy content of the Lemnaceae are utilisable by animals. This corresponds to about 1/2-1/3 of the total energy content.

S. punctata has a mean solar energy conversion efficiency of 4.2% (highest value for a single sample up to 4.9%) of the photosynthetically active radiation (MESTAYER et al. 1984). Chapter 3.7.1 deals with the use of Lemnaceae as a source of energy.

3.2.4. Content of proteins and other nutritive substances

The high protein content of Lemnaceae has already been indicated in chapter 1.2.2. It reaches 15 to 45% of the dry weight. Starvation cultures sometimes contain only as little as 7% protein. After addition of an optimal amount of nutrients to the medium, the protein content is enhanced to 30% within two weeks (CULLEY et al. 1981). The lower limit of nitrogen content in the solution resulting in a high percentage of protein is 20-30 mg/l (FRYE and CULLEY 1980). The frond density in cultures of L. minor has no influence on the specific protein content nor on the cell wall components provided the nutrient supply is sufficient (TUCKER 1981).

Not only the quantity but also the quality of proteins makes the Lemnaceae economically interesting. With the exception of tryptophane and methionine, all essential amino acids used in human and animal food are satisfactorily present. Tryptophane is detected only in traces. Methionine content varies between 0.3% and 3% of the total protein, depending on investigated clone and author (PORATH et al. 1979, AMADO et al. 1980, RUSOFF et al. 1980 and further authors, see chapter 1.2.2). CHANG et al. (1977) analysed as much as 3.1-4.7% methionine, a value which would satisfy the requirements of the FAO. Possibly, the production of methionine can be improved by application of suitable nutrient solutions. Table 3.4 gives a survey of the protein content of Lemnaceae (according to AMADO et al. 1980) and several other plant and animal products.

The essential amino acid index (EAA) amounts to 76 for Lemnaceae (MEST-

Table 3.4. Content of essential amino acids (given in % of total protein content) of Lemnaceae (mean of 94 investigated clones calculated from AMADO et al. 1980), rice, soybean, *Chlorella*, and egg compared to FAO reference pattern. The need of amino acids (in % of total feed) for young pigs and chickens is also given.

amino acid content	in % of total protein	in % of total feed	amino acids
lysine	6.8	3.2	6.8
threonine	5.0	3.8	5.0
valine	6.6	6.2	6.6
methionine	1.0	3.4	1.0
leucine	9.6	8.2	9.6
isoleucine	4.8	5.2	4.8
phenylalanine	5.9	5.0	5.9
tryptophane	traces	1.3	traces
arginine	6.7		6.7
histidine	2.0		2.0
duckweed	1)		
rice	2)		
soybean	1)		
<i>Chlorella</i>	1)		
egg	3)		
FAO	2)		
pig	4)		
chicken	4)		

1 AMADO et al. 1980 2 RUSOFF et al. 1980 3 CHANG et al. 1977 4 MATSUMOTO 1981

LING and BOEHM 1980). In comparison, the EAA for eggs is reported as 97. The protein efficiency ratio for Lemnaceae feed to grass carp reached 2.36 (HAJRA and TRIPATHY 1985).

Considering the high productivity and the high protein content of Lemnaceae, the protein yield per area must be higher than that of any other crop plant. If we assume a maximum yield of 50 t dry weight per ha and year and a mean percentage of protein of 30% of the dry weight, we can expect a protein harvest of up to 15 t per ha and year. Accordingly, the results of ORON et al. (1987a) indicate that protein yield of duckweed grown in waste water systems may reach 12 t per ha and year. Soybean, belonging to crops richest in protein, yields maximum up to 0.7 t protein per ha and year which is about 1/20 of the potential yield of Lemnaceae. Even under suboptimal conditions, the Lemnaceae are able to produce a great amount of protein and also of carbohydrates. BHANTHUNAVIN and MCGARRY (1971) give examples of yearly yields of different crop plants in northern Thailand of protein, carbohydrate and fat (table 3.5), compared with *W. globosa* (named as *W. arifolia*). SAID et al. (1979) estimate that the production of the same amount of protein needs a ten times greater cultivation area for soybeans than for Lemnaceae and a 2.5-9 times greater area for alfalfa. MATSUMOTO (1981) calculated the total digestible crude protein in *S. polyrrhiza* as 23.5% of the dry weight and the total digestible nutrients as 44.2%.

Table 3.5. Content of protein, carbohydrate and fat in different crop plants from northern Thailand (from BHANTHUNAVIN and MCGARRY 1971)

	% dry weight	kg ha ⁻¹ yr ⁻¹				
		Wolffia	Wolffia	soya	nuts	rice
protein	19.8		2080	303	229	71
carbohydrate	43.6		4589	255	164	849
fat	5.0		533	158	397	4
fibres	13.3		1398	44	21	3
ash	18.3		1928	41	20	5

Vitamins A, B₁, B₂, B₆, C, E and PP are present in Lemnaceae (see chapter 1.2.10). Especially the content of vitamin E (0.02-0.04 mg per g fresh weight) and of vitamin PP (0.04-0.06 mg per g fresh weight) are remarkable (MUZAFFAROV et al. 1971).

3.3. CULTIVATION AND HARVEST

3.3.1. Cultivation

The cultivation methods are greatly dependent on the intended utilization of the Lemnaceae.

For physiological tests, aseptic cultures (as described in chapter 2.2.3) are necessary.

For cultivation in the field, the following guidelines may be useful:

- The cultivation area must be protected against wind in order to prevent an accumulation of the fronds in one corner; Lemnaceae should be able to cover the whole illuminated water surface evenly; larger areas should be divided into smaller ones; shores must preferably be steep to prevent the fronds from being washed ashore; water fluctuations should also be avoided.

- Since Lemnaceae are only able to utilize nutrients from the upper water layers, shallow waters with slow water circulation are best for growth; the optimal depth depends on the climatic conditions and on the intended utilization; in cool regions shallow water warm up faster but may not be favourable to survival in winter.

- Most Lemnaceae grow much slower at temperatures below 20°C; in regions where temperatures often remain under this point, species should be chosen which grow well in cooler waters: e.g. L. gibba, L. minor, L. trisulca, and S. punctata. CULLEY et al. (1978) recommend for Louisiana with a change of warm and fairly cool seasons a mixture of S. polyrrhiza, S. punctata, and L. gibba. Some species are not suited for warm regions (e.g. L. trisulca, L. minor), others do not grow in regions where temperatures often drop below 10°C (e.g. Wolffia Wolffii, W. neotropica, W. hyalina, W. arrhiza, L. aequinoctialis, S. polyrrhiza). Species which do well in warm regions are: S. polyrrhiza, L. aequinoctialis, W. microscopica, W. angusta, W. globosa.

- Lemnaceae only grow in waters relatively rich in nutrients; in waters where the conditions are not constantly optimal, a mixture of different duckweed species is preferable.

- The density of the Lemnaceae cover is important for maximum yield. MUZAFFAROV et al. (1971) received best yield with a density of 500 g fresh weight (corresponding to 25-30 g dry weight) per m². REJMANKOVA

(1978) and REJMANKOVA et al. (1983) noted 15-25 g dry weight per m² as optimal stocking density. DEBUSK et al. (1981), REDDY and DEBUSK (1985b) used the following operational plant densities in their experiments: 10-88 g m⁻² for *S. polyrrhiza* and 10-120 g m⁻² for *Lemna* sp. Growth rate was maximum at the lower plant densities used.

Cultivation plants are described by REJMANKOVA (1978), SAID et al. (1979), CULLEY and MYERS (1980), and CORRADI et al. (1981).

3.3.2. Harvest and processing

Pure cultures of *Lemnaceae* are relatively easy to harvest. The fronds can be skimmed off by some kind of net, or they can be collected at the outlet of the water by a grid. KOBAYASHI et al. (1977) developed an office type screen for harvesting *Lemna* in an irrigation channel. The highest yield is achieved if the close but not many-layered *Lemnaceae* cover is gathered at short intervals. According to SAID et al. (1979) and CULLEY and MYERS (1980), the daily harvest is more advantageous than the weekly removal (23.3 t dry weight per ha and year against 17.6 t). REJMANKOVA et al. (1983) developed best harvest strategy at 1 to 2 day intervals (800 g dry weight per m² in 90 days compared with 600 g if harvested every 14 days). Differently, RYTHIER et al. (1980) and DEBUSK et al. (1981) did not get a higher production if harvested every day compared with every 5 to 10 days. In northern Thailand, *W. globosa* is gathered every 3 to 4 days, an interval which proved to be favourable after many years of experience (BHANTHUNNAVIN and MCGARRY 1971). The different results of different authors are certainly due to different techniques applied. In general a shorter harvest interval which is much better suited for an evenly optimal utilization of sunlight must result in more productivity.

Drying is necessary to store the *Lemnaceae* yield. According to SCHULZ (1962), *Lemnaceae* become dry on a sunny day in Central Europe within 10 hours if turned over several times. Without the periodic turning over the drying takes much longer than for making hay, and it is only possible if no rain or strong wind occurs (CULLEY and EPPS 1973). The drying of a 5 cm thick layer of *S. punctata* at 130°C takes about 10 hours (LAWSON et al. 1974). At this temperature, some of the proteins are lost. Also PORATH and KOTON (1977) point out the fact that the content

of digestible proteins and amino acids is diminished during heat-drying. If the water is pressed out mechanically, a 66-71% loss of proteins is recorded by LAWSON et al. (1974).

BALDWIN and MYERS (1980) describe harvest methods of *Lemnaceae* for feeding cattle. From the skimmer, the *Lemnaceae* reach a drop box. After partial dehydrating and chlorinating they are transported to storage, drying facilities, or feed bins. The treatment with hypochlorite is recommended to lower the content of bacteria (AMBORSKI and LARKIN 1980).

Little is known about the possibilities of silage of *Lemnaceae*. According to a survey of EDWARDS (1980), the silage of water plants (e.g. *Eichhornia*) did not prove to be good because the water content is too high. Therefore it is necessary either to pre-dry the material or to add some concentrated organic substances (e.g. carbohydrates). EVERSULL et al. (1980) succeeded however to silage *Lemnaceae* together with a high dry matter corn crop.

3.4. UTILIZATION AS FOOD PLANT

3.4.1. General remarks

As pointed out earlier, Lemnaceae are easily harvested, have extended growing and harvesting periods, high protein and low fibre and lignin content and only very few and not severe pests (CULLEY and EPPS 1973). Therefore, they are generally suited as a food plant. The application for human nutrition and as an animal feed has been known for a long time. Lemnaceae ("duckweed") are favoured as a fodder plant for fish and birds in many countries. They are on offer at markets, e.g. in Mexico for fowl (according to a personal communication from M. Seidl, Greifensee) or in Taiwan where they are sold at 1 \$ for 10 kg wet weight as foodstuff for fish and duck (T.P. CHEN in EDWARDS 1980). In eastern Asia (Northern Thailand, Burma, Laos) W. globosa (named as W. arrhiza is cultivated under the name khai-nam ("eggs of the water") for many generations and sold at markets (BHANTHUMNAVIN and MCGARRY 1971). It is generally assumed that Lemnaceae will soon become more important as a crop plant (e.g. KRUGMANN-RANDOLF 1978). Extensive investigations are being made with Lemnaceae as protein suppliers in many regions of the world.

3.4.2. Human nutrition

The high content of protein, carbohydrate, and vitamins grant to Lemnaceae an outstanding nutritional value also for man. NAKAMURA (1960) mentioned the utilization of Wolffia as human food. In eastern Asia, W. globosa has been eaten by man for many generations (BHANTHUMNAVIN and MCGARRY 1971). The plants are cultivated in ponds of up to 100 m² area which are supplemented by rain water and shaded by bamboo. No artificial fertilizer is supplied. Every 3 to 4 days part of the Wolffia cover is harvested and eaten as a vegetable. The species flowers during the month soon between August and October and is then considered not wholesome. Possibly, the growth is very slow during that time, due to the very diluted water. Cultivating of W. globosa in this way yields 2 t protein, 4.5 t carbohydrate, and 0.5 t fat per ha and year. NAKAMURA (1960) re-

ports that the taste of Wolffia is excellent and sweet, resembling that of cabbage.

It is rather astonishing that Lemnaceae which are so wide-spread have not been used as human food in other regions. R.A. ALBERTS (Springfield, Virginia, USA, in lit. 1979) assumes that the Mayas in Guatemala used Lemnaceae as foodstuff under the name of Xim Ha ("water corn"). He thinks that in expansive irrigation systems the Mayas were able to nourish the numerous people living in a relatively small area with the productive Lemnaceae.

The explanation that the use of Lemnaceae as a human food is restricted might be the following:

- 1) Lemnaceae and especially Spirodela and Lemna contain a great amount of oxalic acid, partly in solution and partly crystallized. They are therefore not very agreeable in taste (rather harsh). It might be possible to prepare the plants by some treatment to make them tasty and desirable. SUTTON (1981a) describes an edible L. gibba salad, and NEY (1960) states that L. minor is far superior in taste to Chlorella. It is evident that Wolffia and Wolffiella which have the oxalate in the free form are more suited for food than Spirodela and Lemna.
 - 2) The harvested Lemnaceae are difficult to separate from other organisms such as snails, insects, worms, protozoa, algae and bacteria. There is also a certain danger of infection by pathogens when eating Lemnaceae that have been cultivated in waters polluted by waste water. AMBORSKI and LARKIN (1980) demonstrated that during the warm season in Louisiana one liter water of a pond covered with Lemnaceae and supplied with sewage of cattle contained 10⁸ coliform bacteria (of which 10⁶ were fecal), 10⁴ fecal Streptococcae and up to 300 Salmonella and Shigella germs. Also pathogenic viruses could be detected. A treatment of 20 minutes with hypochlorite resulted in a marked reduction of the bacteria content. NGUYEN (1978) detected trematoda in Lemnaceae covered waters which were in connection with a pig-breeding plant. However the toxic blue alga Anabaena never showed up in waters with a Lemnaceae cover (KELLY 1980). Further investigations on safe cultivation and harvest of Lemnaceae are needed.
- Another possibility to utilize Lemnaceae for human food is either via extraction of proteins (cf. RUSOFF et al. 1980) or in an indirect way via animal production. Certainly, this last possibility reduces the efficiency factor of the Lemnacean energy. CULLEY et al. (1981) cal-

culated a need of 50 m² pond area to nourish a family of 5 people via egg protein. To produce the 90 to 100 kg protein needed by the family each year, the egg producing chickens eat 40 kg of Lemnaceae protein and in addition 14 kg of animal protein (from worms and insects living in the soil of the surrounding area). The protein supply via fish still seems more favourable (see chapter 3.4.5.1).

3.4.3. Food for mammals

CULLEY et al. (1981) give a survey of all investigations made with Lemnaceae for use in farming.

3.4.3.1. Dairy heifers

RUSOFF et al. (1977, 1978) investigated the possibilities of using duckweed as a nutrient for Holstein cattle. The animals are able to take successfully more than 75% of fodder in form of Lemnaceae. The milk does not change in taste. Calves of 150 to 300 kg weight which have been fed with 67% duckweed (dry weight) and 33% silage of corn, showed an increase in weight of 0.95 kg per animal and day. In comparison, animals which received concentrated food and corn silage in addition to pasturing, only resulted in an increase of 0.5 kg per animal and day. According to CULLEY et al. (1981), 3.1 ha water area with Lemnaceae are sufficient to obtain enough protein for 100 dairy cattle. FRYE and CULLEY (1980) describe installations, functioning, and costs of a dairy farm in Louisiana based on a recycling principle and using Lemnaceae as main food for the animals.

3.4.3.2. Pigs

SCHULZ (1962) studied the nourishment of pigs with Lemnaceae. If fed with 75% grist of rye, wheat and barley, 5% of oat grist, 6% of wheat bran, 8% of fodder yeast, 5% of fish meal and 1% of minerals, the daily increase was still lower (455 g) than with 500 g fresh Lemnaceae fed in addition (546 g). In other experiments, the difference was still higher (up to 300 g per day). GALKINA et al. (1965) report a 14-20% weight increase of pigs in Russia if 1 kg fresh Lemnaceae was added to the normal

feed. HILLMAN and CULLEY (1978a) give some further examples.

3.4.3.3. Ram and sheep

TAUBAEV and ABDIEV (1973) noted 27% additional weight increase in ram and 14% in sheep if 0.5 kg Lemnaceae was added daily to the normal nutrient. PORATH et al. (1985) used duckweed as a substitute for animal protein rich feed in diets of young lambs and Awasi sheep.

3.4.3.4. Horses

STEWART (1972) mentions that horses feed on Lemnaceae in Dal Lake (Kashmir). However, no feeding under controlled and supervised conditions has been done so far.

3.4.3.5. Rabbits

MATSUMOTO (1981) made some investigations with rabbits. If he fed them with S. polyrrhiza alone, the increase was only 20-46% that with conventional fodder. Feeding a mixture of Spirodela with honey sugar resulted in an increase of 35-56%. Highest increase was reached if conventional fodder was supplemented with Spirodela.

3.4.3.6. Nutria and muskrat

Nutria (Myocastor coypus) and muskrat (Ondatra zibethicus) which are raised for fur, eat Lemnaceae in great amounts (WARKENTIN 1968, JACOBS 1947). For these animals, duckweeds are of great nutritional value and are recommended as a food in nutria farms. On account of the low content of fibre, indigestions may occasionally occur (SZUMAN and SKRZYDLEWSKI 1980).

3.4.3.7. Mice

WILKS (1962) demonstrated that mice can be sustained for indefinite periods on a strict diet of duckweeds without losing their normal physiological activity.

3.4.4. Bird food

It is well known that water fowl, especially ducks, feed on Lemnaceae (cf. JACOBS 1947, SCHULZ 1962, MACKENTHUN et al. 1964). A survey of the literature on experiments is given by CULLEY et al. (1981). In addition of small amounts of duckweed (2-5%) to the normal feed results in an additional daily weight increase of chickens by 10 to 32% (MUELLER and LAUTNER 1954, MUZAFFAROV et al. 1968, NAPHADE and MITHUJI 1969-1970, TRUAX et al. 1972, TAUBAEV and ABDIEV 1973). Other authors could not achieve an additional weight increase if the normal fodder was partly (2.5-25%) replaced by Lemnaceae. An addition of 50% Lemnaceae even gave negative results compared with controls (MUZTAR et al. 1976, 1977, JOHRI and SHARMA 1980). Positive results with layers were achieved if 25% of Lemnaceae was added (WILLIAMS 1978). SCHULZ (1962) reports on successful application of Lemnaceae in fattening of ducks. MUELLER and LAUTNER (1954) and NIKOLAEVA (1956) received additional weight gain (10-23%) in ducks if 2% or 37% of the foodstuff was replaced by duckweed. YAMANI et al. (1978) noted a favourable effect of Lemnaceae in protein nutrition of poultry. Caixina scutulata, the white winged wood duck, especially when young, prefers Lemnaceae (LUBBOCK 1975 from HUBAC et al. 1984). Successful effects of duckweed nutrition on ducks was also reported by GERGEL et al. (1985). Two phasianids (chukar and partridge) were fed effectively with duckweeds for 28 days (DØGEN 1987).

DYLIK et al. (1979) calculated for Anas platyrhynchos an assimilation index of Lemnaceae food of 0.64 which is, compared with other crops, very high.

An advantage of Lemnaceae for poultry food is the high content of carotenoids, especially carotene and xanthophyll, which favours the colouring of fat and skin of the birds (MUZTAR et al. 1979). Also the egg yolk is more intensely coloured when the birds are fed with Lemna minor (GRAF 1987).

One ha water area is sufficient to raise 4000 to 7000 chickens and ducks during a vegetation period (HARVEY and FOX 1973). REJMANKOVA (1981) calculated an area of 1 ha Lemnaceae cover as sufficient to produce protein for 480 ducks during the warm season.

3.4.5. Foodstuff for fish and other cold-blooded animals

3.4.5.1. Fish

Lemnaceae are a very valuable source of food for many fish. On the other hand, a closed cover of Lemnaceae may prevent the supply of oxygen to the water, thus resulting in unfavourable conditions for fish sensitive to low oxygen content. Trout, for instance, are not suited for raising in Lemnaceae ponds (WRIGHT 1973).

A distinct positive effect on prospering of the grass-carp or white amur (Ctenopharyngodon idella) is reported by many authors (VERIGIN 1962, GALKINA et al. 1965, NIKOLSKIJ and VERIGIN 1966, FISCHER 1968, OPUSZYN-SKI 1972, EDWARDS 1974, PRISHCHEPOV 1974, VARGHESE et al. 1976, SUTTON 1976, VOVK 1976, PORATH and KOTON 1977, ROTTMANN 1977, SHIREMAN et al. 1977, 1978, BAUR and BUCK 1980, MACEINA and SHIREMAN 1980, HAJRA and TRIPATHY 1985). In cultures with a mixture of L. gibba and L. minor, the weight of the fish was tripled (from 100 g to 300 g per fish) within 50 days (PORATH and KOTON 1977). According to VAN DYKE and SUTTON (1977), the grass-carp is able to use 65% of the dry weight of Lemnaceae for food: 61% of the gross energy content, 70% of the rough protein, 72% of the organic cell content, 30% of the organic parts of the cell wall. The food conversion rate on a dry weight basis of grass-carp feeding on Lemnaceae amounts to 1.6 for a fish of 3 g and 2.7 for a fish of 63 g. No other foodstuff (catfish chow, rye grass, or a mixture of both) was nearly as efficient (SHIREMAN et al. 1978). SUTTON (1976) measured a food conversion rate between 1.1 and 5.3, BAUR and BUCK (1980) between 1.55 and 4.07, and HAJRA and TRIPATHY (1985) between 3.10 and 3.15. Further food conversion rates, however on a wet weight basis, are listed in SINGH and SINGH (1967), MICHEWICZ et al. (1972), TAL and ZIV (1978), HEPHER and PRUGININ (1979) and EDWARDS (1980).

If we assume in a closed grass-carp - Lemnaceae ecosystem a mean food conversion rate of 3.0 and a mean yearly production of 50 t/ha, the yield of grass-carp per ha and year can amount to 25 t. If in a less favourable case the yearly duckweed production is only 10 t per ha and the food conversion rate 5, the fish production still achieves 2 t per ha and year. Hybrids of grass-carp with other species of the same genus Ctenopharyngodon show similar positive results as grass-carp if fed with Lemnaceae (DUTHU and KILGEN 1975, THERIOT and SANDERS 1975, CASSANI 1981, CASSANI et al. 1982).

Channel catfish (*Ictalurus punctatus*) was successfully fed with up to 20% Lemnaceae (dry weight) (ROBINETTE et al. 1980). Other fish which can be partly nourished with Lemnaceae are: e.g. common carp (Cyprinus carpio) (HICKS 1937, PORATH and KOTON 1977, PANICKER et al. 1985), common mullet (Mugil cephalis) (CHABRECK cited in CULLEY et al. 1981), goldfish (Carassius auratus) (CULLEY et al. 1981). Species of the genus Tilapia (or Oreochromis) partly eat Lemnaceae (T. rendalli: MANN 1967) and partly not (T. mossambica, T. aurea: BAUR and BUCK 1980). However, HENDERSON et al. (1984) got positive results by feeding T. aurea additionally with Lemnaceae. T. rendalli is able to use 42-55% of the protein of S. polyrrhiza and 52-68% of the crude fibre (MANN 1967). KIM and KHANG (1982) received a 12% weight increase in red Tilapia fingerlings by supplying them additionally with Lemnaceae. According to HENDERSON et al. (1984) and HECKMANN et al. (1984) Tilapia grows rapidly in tanks containing duckweed only. They used water from zero discharge power plants to cultivate the Lemnaceae. Tilapia hybrid (Oreochromis niloticus x O. aureus) was fed by GAIGHER et al. (1984) with L. gibba in a recirculating unit. Intake rate was low and food conversion rate good (1:1), relative growth rate poor (0.67% of body mass per day). 65% of the duckweed consumed was assimilated and 26% converted to fish. EHRLICH (1966) proposed a polyculture of duckweed and Daphnia with the duckweed providing shade for Daphnia. These two organisms have a similar turnover rate and could be harvested as fish food simultaneously with the same equipment.

Often, Lemnaceae are not used as direct food for fish (and prawns) but via small organisms feeding on duckweed. The relations between Lemnaceae and other organisms are surveyed in volume 1, chapter 5.5.3 (LANDOLT 1986). In Bengal, Lemnaceae are cultured to increase the zooplankton on which carp feed. First, a phytoplankton bloom is established which disappears later when shaded by Lemnaceae. The Lemnaceae cover is subsequently removed in order to let the zooplankton which is nourished by the dying algae develop. The carp feed mostly on the zooplankton (ALI-KUNHI et al. 1952). Also MALECHA et al. (1981) and HECKMANN et al. (1984) recommend the cultivation of Lemnaceae to increase the zooplankton for raising fish (e.g. carp or Tilapia) and prawns. In aquaculture of fish, L. gibba was shown to act as a biological ammonia stripper. A Lemnaceae mat was able to take up 80% of ammonia of a fish effluent in less than 48 hours (PORATH and POLLOCK 1982).

3.4.5.2. Crustaceae

L. minor is supposed to serve as a supplementary food source for the fresh water shrimp (Macrobrachium rosenbergii) (GODFRIAUX et al. 1975). HECKMANN et al. (1984) were able to raise the same prawn species in tanks containing duckweed only. Also MALECHA et al. (1981) report on Lemnaceae fed on by the shrimp. It is not investigated in detail if crayfish of the genera Oreonectes, Procambarus and Cambarus feed partly or predominantly on Lemnaceae. However, it is known that Procambarus clarkii consumes fresh duckweed if offered (CULLEY et al. 1981). In Louisiana and California, crayfish are released in irrigated rice fields to keep weed (e.g. Lemnaceae) under control. After drainage of the water, the crayfish are collected and used as human food.

The ostracod Cypris spec. which is eaten extensively by fish feeds predominantly on Lemnaceae. Up to 90% of duckweed mat of 50 g fresh weight per m² has been eaten by Cypris within 96 hours at temperatures of 21-24°C (MANISSERY et al. 1981).

3.4.5.3. Turtles

The wide-spread aquarium turtle Pseudomys scripta feeds predominantly on Lemnaceae and Eichhornia. Therefore, commercial turtle farms cultivate Lemnaceae extensively as a foodstuff for the turtles (CHABRECK cited by CULLEY et al. 1981).

3.5. UTILIZATION IN WASTE WATER

3.5.1. General remarks

LUDWIG (1909) already pointed out the fact that Lemnaceae are able to tolerate a relatively high degree of pollution and that they are sometimes the only vegetation growing vigorously in dirty village pools and ponds. In contrast to many other water plants, Lemnaceae species (especially L. gibba) colonize polluted waters and are not impeded severely by a relatively high content of detergents (e.g. Lemnaceae tolerate more than 15 ppm tetrapropylene benzolsulphonate) (AGAMI et al. 1976). Even in waste waters with up to 500 mg COD (chemical oxygen demand) per liter, they still survive (GHETTI et al. 1982).

Organic waste water (sewage from households and farms) is produced in the whole world in great quantities and becomes a great ecological problem. The biological purification and the recycling of nutrients (especially nitrogen and phosphorus) to plant proteins would be highly preferable to the present technical waste water treatments applied in the industrial states. Lemnaceae are well suited for biological treatment of waste water under certain conditions.

Lemnaceae have the following advantages for water purification:

- they are tolerant to high content of nutrients (S. polyrrhiza is still able to grow in solutions with 1 g nitrogen per liter and 1.5 g phosphorus per liter, according to EYSTER 1966)
- they are able to absorb and desintegrate toxic substances (cf. chapter 2.3.3.5.8) and to bring under control the content of some pathogens
- they are suited as food, energy producers and as manure
- they are able to accumulate heavy metals

Disadvantages of Lemnaceae are:

- they stop growth at low temperatures; outside the tropics and subtropics, the waste water plants have to be heated during the cold season
 - they live only in a thin layer on the surface of the water; large areas are needed to remove sufficient waste substances from the water
 - they accumulate heavy metals; it is not possible to use Lemnaceae for food or manure from waters containing heavy metals; waste water with heavy metals has to be purified separately from the normal waste water
- A survey of the possible utilization of different water plants to remove

nutrients and toxic substances is given by DINGES (1982).

3.5.2. Removal of nutrients

Numerous investigations have been made throughout the world in order to clean waste water and to recycle the nutrients. A bibliographic survey of macrophytes used in waste water treatment is given by BLAKE and DUBOIS (1979). Lemnaceae are shown to have the greatest capacity in assimilating the macroelements N, P, K, Ca, Na and Mg. Other studied macrophytes are less effective in this respect if the effluents are heavily loaded with nutrients (BLAKE and DUBOIS 1982).

In the CSSR according to KVET et al. (1979), Lemnaceae in basins of 0.5 m depth remove on an average 2 kg nitrogen per ha and day (calculated for the whole year). This corresponds to a 50% removal from water of 25 mg l⁻¹ nitrogen. If temperatures stay warm throughout the whole year, the effect is much higher. In waste water ponds of Louisiana covered with Lemnaceae CULLEY et al. (1978) observed a 20-40% lower nitrogen (predominantly ammonium) content than in ponds without Lemnaceae. Examples of the daily removal of elements are given in table 3.6.

CULLEY et al. (1981) calculated that on the average, a mixture of Lemnaceae could remove annually 1378 kg nitrogen, 347 kg phosphorus and 441 kg potassium from a ha water area in Louisiana. Further investigations with Lemnaceae alone or mixed with other water plants for purifi-

Tab. 3.6. Daily removal of N, P, and K by Lemnaceae during warm seasons

Region	species	daily removal in kg/ha			author
		N	P	K	
Louisiana	duckweeds	4.7	1.6	2.1	2
Florida	<u>S. polyrrhiza</u>		0.15		4
Italy	<u>L. gibba</u> and <u>L. minor</u>	4.15	0.97		1
CSSR	duckweeds	2.0			3

- 1 CORRADI et al. 1981
- 2 CULLEY et al. 1978, 1981, CULLEY and MYERS 1980
- 3 KVET et al. 1979
- 4 SUTTON and ORNES 1977

cation of sewage water were made by SUTHERLAND and BEVIS (1979) in Michigan, USA, WOLVERTON and McDONALD (1981) in Mississippi, USA, RYTHIER et al. (1980), KNIGHT et al. (1985) and REDDY (1984b) in Florida, USA, CONN and LANGWORTHY (1984) and GEARHEART et al. (1984) in California, USA, COPELLI et al. (1982) in Italy, ORON et al. (1985) in Israel, RAKHIMOV and RAKHIMOVA (1983) in USSR, SMITH et al. (1983) in Australia, and MATSUMOTO (1981) in Japan. EDWARDS (1980) reports on waste water systems with Lemnaceae from Southeastern Asia.

In mixtures with other water plants, the removal of nutrients might be even better. KUMAR et al. (1983) propose a mixture of S. polyrrhiza and Azolla pinnata. A combined mat of Lemnaceae with Eichhornia removes 6 times more nitrogen and twice as much phosphorus from the water than Lemnaceae alone (EDWARDS 1980). This can be explained by the much bigger biomass and the deeper rooting of Eichhornia. According to MATSUMOTO (1981), S. polyrrhiza, L. aequinoctialis, and Eichhornia absorb about the same amount of N (86-91%), P (35-85%), and K (12-15%) each out of sewage water which still contained (after first purification) 12 mg N, 8 mg P and 14 mg K per liter. The removal of N and P from water decreases considerably as soon as the content of these nutrients falls below 4 mg/l (REJMANKOVA 1982). At these relatively low concentrations, other water plants are required to complete purification. A reduction of the phosphorus level down to 0.1 mg per liter as is proposed for strict water quality standards in the USA cannot be achieved with Lemnaceae alone (CULLEY et al. 1978). The efficiency of nutrient uptake varied within a day: it was highest between 9 a.m. and 3 p.m. (highest light intensity) and lowest between 9 p.m. and 5 a.m. (1/3 to 1/2 of the highest value) (MATSUMOTO 1981). MATSUMOTO aerated the waste water basins to increase the efficiency of the purifying system. In this way, the nutrient became evenly distributed throughout the water. MARTIN et al. (1978) report on a small waste water plant in southwestern France for tertiary (biological) treatment of the municipal sewage using Lemnaceae and Nasturtium. The water basins had a depth of 12 cm and the water was renewed every 5 days; the biomass was harvested twice a week. At the outlet, the waste water contained 13-48% of the original content of BOD (biochemical oxygen demand; on the average 6.9 mg/l), 18-27% COD (chemical oxygen demand; on the average 0.9 mg/l), 3% MBAS (methylene blue active substances; on the average 28.3 mg/l), 1.5-4.5% nitrogen and 3-7% phosphorus. A need of 650 m² water area per 100 inhabitants was calculated for purifi-

cation. With more elaborate installations, the area might be reduced to about 200 to 300 m² per 100 inhabitants. Further studies with biological waste water purification using, at least partly, Lemnaceae have been made in France by CHASSANY-DE CASABIANCA (1982a,b), SAUZE (1982) and CHASSANY-DE CASABIANCA and SAUZE (1981), in Mississippi, USA, by WOLVERTON (1979), in California, USA, by GOLUEKE (1979), in Israel by ORON et al. (1985), and in Western Australia by SMITH et al. (1983). DEGHI and EWEL (1984) simulated the effect of waste water application on phosphorus distribution in cypress domes of Florida. An addition of waste water at the rate of 2.5 cm per week resulted in a 200fold increase of the phosphorus content within the duckweed. ELLIS and DAVIS (1984) noted that the passage of municipal effluent through floating duckweed communities generally increases diversity of algal species and decreases phytoplankton biomass. Also ROSETTI and PATIL (1986) measured reduced algal activity in systems with L. minor. However, if algae are used to remove ammonia and phosphorus from waste water an addition of duckweeds is not advisable (KOLES et al. 1986). According to BURTON et al. (1978), the regeneration of eutrophic lakes by Lemnaceae is only to be recommended if the additional input of nutrients is relatively low. Duckweeds in a rice paddy field irrigated by secondary treated sewage water play a beneficial role in reducing the excessive nutrient supply and in purifying the sewage effluents (TATSUKAWA 1986).

Closed or semiclosed systems of waste water purification and utilization of the harvested Lemnaceae for food have been investigated many times and for different purposes. CHEN (cited in EDWARDS 1980) reports on a simple system of ponds in Taiwan. Sewage from households drains directly into the water system covered with Lemnaceae. The duckweeds are regularly collected and used as foodstuff for ducks and fish. In India NASKAR et al. (1986) established a semiclosed system with W. globosa (named as W. arrhiza). The duckweeds cover a basin with sewage effluent medium and serve as food for carp. 10358 kg fish per ha and year are produced. The conversion ratio W. globosa (wet weight) to fish amounted to 6:1. Cattle ponds are wide-spread in the USA. TRUAX et al. (1972), CULLEY and EPSS (1973), RUSKIN and SHIPLEY (1976), and KELLY et al. (1978) investigated closed systems of nutrient flow from dairy cattle into waste water, into cattle pond, into Lemnaceae cover and back to dairy cattle. According to CULLEY et al. (1978), Lemnaceae covering one ha of waste water are able to purify the sewage of 15.5 cows for nitrogen, 34 cows for phosphorus

and 8.8 cows for potassium. It is recommended to first collect the methane gas from the sewage, by fermentation, before transporting the water to the pond. The feces of 100 cows develop enough energy via the methane production to supply the whole dairy farm with operation energy (except transport energy) (HUFFMAN 1980, FRYE and CULLEY 1980, CULLEY et al. 1981). Pathogens (coliforms, streptococci, Salmonella, Shigella) in the water must be destroyed by hypochlorite (AMBORSKI and LARKIN 1980). In a similar way, the recycling of pig sewage over Lemnaceae is possible (CULLEY and EPPS 1973, STANLEY and MADEWELL 1975, GHETTI et al. 1982). Lemnaceae are also used in closed systems for pisciculture. It is possible to reduce the ammonium content in the water by cultivating duckweeds (KIM and KANG 1982, PORATH and POLLOCK 1982). According to the latter authors, the circulation of fish effluent through water containing a duckweed mat succeeds in an ammonia removal of 80% within 48 hours.

3.5.3. Removal of heavy metals and other toxicants

3.5.3.1. Heavy metals

In chapter 1.1 the ability of Lemnaceae to accumulate certain heavy metals is described. The relatively high tolerances to heavy metals are mentioned in chapter 2.3.3.4.6. These two characteristics of Lemnaceae make it possible to use the plants for removal of heavy metals from polluted waters. A survey of the suitability of different water plants in this respect is given by DINGES (1981, 1982). The accumulation factor* of Lemnaceae for certain heavy metals depends greatly on the concentration of the metals and of other metals in the water as well as on the species. Table 3.7 summarizes the results of different authors. The relatively high accumulation factor of SILVEY (1967) is due to the long stay of the fronds in the slowly flowing water of very low metal content. It seems that the concentration factor is much higher at very low concentrations than at medium and high concentrations. In medium concentrations of Pb (1 mM to 10 mM) the content of Pb within the Lemnaceae

* The accumulation factor is always given on a dry weight basis in this chapter. Original values on a wet weight basis have been multiplied by 20.

raises in direct proportion to the concentration in the water (see fig. 1.2). The same is true for Cd between the concentrations of 0.1 mM and 1 mM. Within these concentrations, the accumulation factor is similar (c. 1000). In contrast, the accumulation factor for Cu and Zn is lower at high concentrations (5 mM and 10 mM) than at lower ones (0.5 mM and 1 mM) (VAN DER WERFF 1981), VAN DER WERFF and PRUIT (1982). VAN DER WERFF (1981) points out the mutual effect of Zn and Cu. A reciprocal uptake stimulation of Cu and Ni was observed by HUTCHINSON and CZYRSKA (1975). Further interrelationship between different metals are dealt with in chapter 2.3.3.4.7. Differences for species have been noted by VAN DER WERFF (1981) and HUTCHINSON and CZYRSKA (1975). According to HUTCHINSON and CZYRSKA, L. valdiviana accumulates much more Cu than L. minor (accumulation factor 500-54000 compared with 80 to 8000). The accumulation factor of Cu and Cd is much higher, that of Pb is slightly higher and that of Zn is slightly lower in L. gibba than in S. polyrrhiza (HUTCHINSON and CZYRSKA 1975; see table 3.7). The high accumulation factors for Al and Mn (up to 660'000 and 850'000, respectively) are remarkable. RUSKIN and SHIPLEY (1976) report that L. minor and L. trisulca absorb 10 times more boron from waters in Michigan than all other float-

Table 3.7 (p. 396 and 397). Accumulation factors for different elements in Lemnaceae, calculated on a dry weight basis (data which were described on a fresh weight basis have been multiplied by 20)

Species	d	Lemna minor	g	Lemna perpusilla
a Spirodela polyrrhiza	d	Lemna minor	g	Lemna perpusilla
b Spirodela punctata	e	Lemna trisulca	h	duckweed
c Lemna gibba	f	Lemna aequinoctialis		
References.				
1 SILVEY 1967	13 ALLENBY 1967			
2 MANGI et al. 1978	14 SAROSIEK and WOZAKOWSKA-NATKANIEC 1980			
3 HONDA et al. 1971	15 SKLAR 1980			
4 LEINERTE 1969	16 PIISPANEN and LAEHDENMAEKI 1983			
5 HUTCHINSON and CZYRSKA 1975	17 SZABADOS et al. 1983			
6 RODGERS et al. 1978	18 TRAPEZNIKOV and TRAPEZNIKOVA 1979			
7 GUTHRIE and CHERRY 1979a	19 NASU et al. 1985			
8 CLARK et al. 1981	20 OZINEK 1983			
9 VAN DER WERFF 1981	21 MARCIULIONIENE 1980			
10 KOVACS et al. 1984	22 VERMAAK et al. 1976			
11 TRIDECH et al 1981	23 WAYMAN et al. 1977			
12 ALLENBY 1981	24 WENTSEL and BERRY 1975			

[illegible][illegible]

ing or submerged water plants. The same observation was made by TRIDECH et al. (1981) in Louisiana. Pistia was next to Lemna but accumulated only 1/3. The same authors stated that Lemnaceae grid the following percentage of metals out of a waste water system: 17.8% B, 70.5% Hg, 11% Se (beside 55% N and 17.9% P). GLANDON and McNABB (1978) measured up to 1500 ppm B in the dry weight of L. minor. According to FERNANDEZ et al. (1983), L. aequinoctialis is more efficient in absorbing Zn and Hg than Pistia but less efficient than Eichhornia. FERRARA et al. (1985) investigated the absorption of Cd and Zn by L. minor. Though Lemna is less effective than Elodea and Eichhornia in absorbing the two metals it is thought to be useful in water purification due to its rapid multiplication. GELLINI and PICCARDI (1981) noted that L. minor is able to reduce the Cu content of water from 5 mg/l to 1/4 within 48 hours. According to NASU (1983), L. aequinoctialis removes the following percentage of Cd within a week: 2% from a solution with 1 ppm Cd, 50% from a solution with 0.1 ppm, and 70% from a solution with 0.01 ppm. STAVES (1980) propose the use of a mixture of Lemnaceae (S. polyrhiza, S. punctata, and L. gibba) in order to remove Cr from industrial waste water. However, the Cr concentrations should not exceed 10 ppm. STAVES and KNAUS (1985) discuss various types of biological treatment systems with duckweeds for the removal of Cr from waste water. At 0.1 ppm Cr S. polyrhiza, S. punctata and L. gibba exhibit the greatest percentage of Cr removal (concentration factor up to 5700 for S. punctata). In waters with a thick Lemnaceae cover, Cr^{6+} is reduced to Cr^{3+} which then readily adsorbs to organic and inorganic surfaces. FARAGO and PARSONS (1985) mention the possibility of L. minor (as well as of Eichhornia) to recover Pt from polluted waters (without giving any details).

To remove the heavy metals from the water, the Lemnaceae cover must be taken away regularly. If this is not done, the amount of heavy metals (Fe, Zn, Cr, Cu, Ni, Pb) is finally immobilized in the sediments after decaying of the Lemnaceae fronds. Only Mn is released into the water during plant decomposition and returns to the cycling system (OZIMEX 1983).

3.5.3.2. Organic toxicants

Lemnaceae are especially efficient in removing polychlorinated biphenyls (PCB), which is a rather stable toxicant, from the water. They accumu-

late the substance 6 times more than Eichhornia and 70 times more than Scirpus. In a waste water system, they were able to take out 100% PCB (ENGLAND and KAIGATE 1981, TRIDECH et al. 1981). Phenol is accumulated by duckweeds 5000 times* which is less than in most other water plants (TRIDECH et al. 1981).

MUIR et al. (1985) report a 2000- to 10000-fold accumulation of the insecticide deltamethrin by Lemnaceae from contaminated water. L. aequinoctialis bioaccumulates insecticides (e.g. DDT, endrin) from extremely low concentrations in the water. The absorbed insecticides remain in the tissue of the Lemnaceae (VROCHINSKI et al. 1970, 1971, DE LA CRUZ and YARBROUGH 1982). L. minor accumulates DDT up to c. 800 times and HCCH up to 1200 times (VROCHINSKI et al. 1970).

LOCKHART et al. (1983) measured the bioconcentration factors in L. minor for 10 herbicides. They varied between c. 80 (krenite) and 88000 (hexachlorobiphenyl). TCDD a highly toxic contaminant of the herbicide 2,4,5-T is accumulated in L. minor 20000 to 100000 times (concentration in the water 7.0-0.05 ppt) (ISENSEE and JONES 1975a). A rapid uptake of two phosphate ester flame retardants (triphenyl phosphate and 2-ethylhexyl-diphenyl phosphate) was observed by MUIR et al. (1982). The concentration factors amounted to c. 60000 for EHDPP and 42000 for TPP (10 hours after application) and 5700 and 3000, resp. (10 days after application).

* The accumulation factor is always given on a dry weight basis in this chapter. Original values on a wet weight basis have been multiplied by 20.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.